

COMPOSITIO MATHEMATICA

L. N. VASERSTEIN

**On normal subgroups of GL_2 over rings
with many units**

Compositio Mathematica, tome 74, n° 2 (1990), p. 157-164

http://www.numdam.org/item?id=CM_1990__74_2_157_0

© Foundation Compositio Mathematica, 1990, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

On normal subgroups of GL_2 over rings with many units

L.N. VASERSTEIN

Department of Mathematics, The Pennsylvania State University, University Park, PA 16802, USA

Received 5 April 1989; accepted 16 August 1989

Abstract. For several classes of rings A , we prove the “sandwich theorem” for all normal subgroups of $GL_2 A$: for every normal subgroup H of $GL_2 A$ there is a unique ideal B of A such that $[E_2 A, E_2 B] \subset H \subset G_2(A, B)$.

Introduction

For any associative ring A with 1, let $GL_1 A$ denote its multiplicative group, i.e., the group of its units, and $GL_2 A$ the group of all invertible 2 by 2 matrices over A . For any ideal B of A , let $E_2 B$ denote the subgroup of $GL_2 A$ generated by all elementary matrices $b^{1,2}$ and $b^{2,1}$ where $b \in B$. We denote by $E_2(A, B)$ the normal subgroup of $E_2 A$ generated by $E_2 B$. Let $GE_2 A$ be the subgroup of $GL_2 A$ generated by all its diagonal and elementary matrices.

Let $GL_2 B$ denote the principal congruence subgroup of $GL_2 A$, i.e., the kernel of the homomorphism $GL_2 A \rightarrow GL_2(A/B)$. Let $G_2(A, B)$ be the inverse image of the center of $GL_2(A/B)$ under this homomorphism. Clearly, the center of $GL_2(A/B)$ consists of scalar matrices over the center of the ring A/B .

We will assume that

(1) $\sum A(GL_1 A - 1) = \sum (GL_1 A - 1)A = A$, i.e., no proper one-sided ideal of A contains all $u - 1$, where $u \in GL_1 A$.

The formulas $[\text{diag}(u, 1), b^{1,2}] = (ub - b)^{1,2}$ and $[\text{diag}(u, 1), b^{2,1}] = (bu^{-1} - b)^{2,1}$ give the following result.

PROPOSITION 2. Under condition (1), $E_2(A, B) \subset [E_2 B, GE_2 A]$ for every ideal B of A .

To obtain the “sandwich theorem”, we will use one of the following conditions on A :

(3) $\text{sr}(A) \leq 1$, i.e., A satisfies the first Bass stable range condition;

(4) $A/\text{rad}(A)$ is von Neumann regular;

(5) for every element $a \in A$ there is a finite sequence x_1, \dots, x_N in A such that

$$x_1 + \dots + x_N = 1 \quad \text{and} \quad 1 - ax_i \in GL_1 A \quad \text{for all } i.$$

See [3], [23], [24], [11] about condition (3). Under this condition, $GE_2 A = GL_2 A$.

In condition (4), $\text{rad}(A)$ means the Jacobson radical of A . Recall that a ring A is called *von Neumann regular* if for every $a \in A$ there is $x \in A$ such that $axa = a$. Replacing, if necessary, x here by xax , we can have both $axa = a$ and $xax = x$. If $A/\text{rad}(A)$ is commutative, condition (4) implies (3).

Taking $x_i = 1/N$ with a large N (depending on a) in condition (5), we satisfy this condition for any Banach algebra A , as well as for the ring A of all bounded smooth functions on any smooth manifold. More generally, condition (5) holds for any connected topological ring A with 1 with $GL_1 A$ open in A .

If A is a semilocal ring, i.e., $A/\text{rad}(A)$ is the product of matrix rings over division rings (e.g., A is an Artinian ring), then both (3) and (4) hold. Condition (1) for a semilocal ring A is equivalent to condition (5) as well as to any of the following six conditions:

- A has no factor rings of two elements;
- every element of A is the sum of two units;
- condition (5) holds with $N = 2$;

$$E_2 A \subset [E_2 A, GL_2 A];$$

$$E_2 A = [E_2 A, GL_2 A];$$

(6) every element of A is a sum of elements of the form $u - v$, where $u, v \in GL_1 A$. Moreover, every semilocal ring A satisfies the following condition:

- (7) every element of A is a sum of units.

Note that condition (6) implies both (7) and (1) for an arbitrary A . Obviously, condition (5) implies (1) (because $1 - ax_i \in GL_1 A$ implies $1 - x_i a \in GL_1 A$, see [19]).

Here is the main result of this paper.

THEOREM 8. *Assume conditions (1) and (7) and one of the conditions (3), (4), or (5). Then for every subgroup H of $GL_2 A$ which is normalized by $GE_2 A$ there is a unique ideal B of A such that $[E_2 A, E_2 B] \subset H \subset G_2(A, B)$.*

EXAMPLE. Let A be a semilocal ring. Then the conditions of Theorem 8 hold if and only if A has no factor rings of two elements. Now we give an example of a commutative local ring A with $A/\text{rad}(A) = \mathbb{Z}/2\mathbb{Z}$ and a normal subgroup H of $GL_2 A = GE_2 A$ such that the conclusion of Theorem 8 is false. Let A be the commutative algebra with 1 over the field $\mathbb{Z}/2\mathbb{Z}$ with a single generator z subject to the relation $z^2 = 0$. Clearly, $\text{rad}(A) = \{0, z\}$ and $A/\text{rad}(A) = \mathbb{Z}/2\mathbb{Z}$. Let H be the subgroup of $GL_2 A$ consisting of the following 4 matrices:

$$1_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \text{diag}(1+z, 1+z) = \begin{bmatrix} 1+z & 0 \\ 0 & 1+z \end{bmatrix},$$

$$h = \begin{bmatrix} 1+z & z \\ z & 1 \end{bmatrix}, h' = \begin{bmatrix} 1 & z \\ z & 1+z \end{bmatrix}.$$

Then $g^2 = 1_2$ for all $g \in H$ and $hh' = \text{diag}(1 + z, 1 + z)$. Clearly, $[E_2A, E_2B] \subset H$ for an ideal B of A only when $B = 0$. (Indeed, otherwise $B = \text{rad}(A)$ or A , and the commutator subgroup $[E_2A, E_2B]$ contains a matrix, namely $[1^{1,2}, z^{2,1}] = \text{diag}(1 + z, 1 + z)z^{1,2}$, outside of H .) On the other hand, H is normal in $GE_2A = GL_2A$ and H is not contained in the center $G_2(A, 0) = \{1_2, \text{diag}(1 + z, 1 + z)\}$ of GL_2A . Thus, condition (1) in Theorem 8 cannot be dropped.

The question arises naturally whether the converse to the conclusion of Theorem 8 is true. To be more cautious, one may ask the following question:

(9) suppose that $E_2(A, B) \subset H \subset G_2(A, B)$ for a subgroup H of GL_2A and an ideal B of A ; is then H normalized by GE_2A ?

The positive answer for local rings A was “obtained” in [7]. However, Bass [3] pointed out a gap in [7], and [26] gives a counter-example (with a local ring A , i.e., $A/\text{rad}(A)$ is a division ring) to the results of [7]. Namely, let F be a field, and let A be the associative F -algebra with two generators x, y subject to the relations: all the 8 monomials of degree 3 in x, y are 0. Then $\dim_F A = 7$ and $\text{rad}(A)$ is the six-dimensional F -vector space with a basis x, y, yx, xy, x^2, y^2 . Let B be the four-dimensional subspace with the basis yx, xy, x^2, y^2 . Then B is an ideal of A , $F + B$ is the center of A , and the factor ring A/B is commutative. Then $[G_2(A, B), E_2A]$ is not contained in $E_2(A, B)$ which shows that the results of [7] are wrong.

Using this example, we will show that the answer to (7) is negative, in general, even for local rings A . Namely, if we define H to be the subgroup of $G_2(A, B)$ generated by GL_2B and the diagonal matrix $\text{diag}(1 + x, 1)$, then $E_2(A, B) \subset H \subset G_2(A, B)$, but H is not normalized by $\text{diag}(1 + y, 1) \in GE_2A$.

However, H is normalized by $[E_2A, E_2A]$ under the conditions of Theorem 8, by [3], [20], [22]:

PROPOSITION 10. *Suppose that one of the conditions (3), (4), or (5) holds. Then*

$$[G_2(A, B), [E_2A, E_2A]] \subset E_2(A, B)$$

for any ideal B of A . Therefore, if H is a subgroup of GL_2A such that $E_2(A, B) \subset H \subset G_2(A, B)$ for an ideal B of A , then $[H, [E_2A, E_2A]] \subset E_2(A, B)$, hence H is normalized by $[E_2A, E_2A]$.

Indeed, by [3], [20], and [22], $[GL_2B, E_2A] \subset E_2(A, B)$. Take now an arbitrary $h \in G_2(A, B)$ and consider the map $\varphi: E_2A \rightarrow (GL_2B \cap E_2B)/E_2(A, B)$ given by $\varphi(g) = [g, h]E_2(A, B)$ for any $g \in E_2A$. Then φ is a group homomorphism into an abelian group, so $\varphi([E_2A, E_2A]) = 1_2$. That is, $[[E_2A, E_2A], h] \subset E_2(A, B)$. So $[G_2(A, B), [E_2A, E_2A]] \subset E_2(A, B)$.

The question arises naturally whether

(11) $[E_2A, E_2B] \subset H \subset G_2(A, B)$ for every subgroup H of GL_2A which is normalized by E_2A ?

This question is partially answered in [15]. By [21], a necessary condition for

(11) to be true is that every quasi-ideal of A is an ideal. Recall [21, Section 4] that a quasi-ideal Y of a ring A is defined as an additive subgroup of A such that $aya, yay \in Y$ for all $a \in A$ and $y \in Y$.

If every element of A has the form $2c + d$, where c belongs to the center of A and d belongs to the ideal of A generated by the additive commutators $ab - ba$, then every quasi-ideal of A is an ideal, see [21, Section 4]. In particular, this is the case when A is semilocal and either A is commutative with $2A = A$ or A has no factor fields. Another example is any Banach algebra A with 1 which is either commutative or simple. Yet another example is any von Neumann regular ring A , see [15].

See [15] for results about subgroups H of $GL_2 A$ which are normalized by $E_2 A$.

Theorem 8 will be proved in the next section. The uniqueness of the ideal B in Theorem 8 is easy to see. It follows, for example, from the following fact which is true for an arbitrary associative ring A with 1: if H is a subgroup of $GL_2 A$ such that $[E_2 A, E_2 B] \subset H \subset G_2(A, B)$ for an ideal B of A , then B is the ideal of A consisting of all non-diagonal entries of matrices in H . Also B coincides with the set $\{x \in A: x^{1,2} x^{2,1} \in H\}$.

Some partial positive results about normal subgroups of $GL_2 A$ were known for: von Neumann regular rings A (see [20]), Banach algebras A (see [21]), rings A with $\text{sr}(A) = 1$ (see [3]), including commutative rings A with $\text{sr}(A) = 1$ (see [4], [13], [26]), commutative local A (see [1], [6], [9], [10], [12], [14], [17]), commutative semilocal rings (see [2] and [5]). If $2A \neq A$, then quasi-ideals appear even for commutative local rings [1]. In [8], Theorem 8 is proved for any semilocal ring satisfying the following two conditions: there is a unit ε in the center of A such that $1 - \varepsilon$ is also a unit; A has neither factor rings which are division algebras with centers of ≤ 5 elements nor factor rings which are isomorphic to $M_2 \mathbb{Z}/2\mathbb{Z}$.

Acknowledgements

This research was supported in part by NSF. Pere Menal helped a lot with this paper; only his modesty prevented him from being listed as a coauthor. I thank him also for hospitality during my visit in Spain, when the paper was written. I thank a referee for pointing out a mistake in a previous version of the paper, and E. Wheland for correcting many misprints.

Proof of Theorem 8

LEMMA 12. *Let G be a subgroup of $GL_1 A$. Let H be a subgroup of $GL_2 A$ which is normalized by $E_2 A$ and all diagonal matrices $\text{diag}(u, v)$ with $u, v \in G$. Set $K =$*

$\{g \in GL_2 A : [g, E_2 A] \subset H\}$. Let K contain a matrix

$$h = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

with $a \in GL_1 A$. Then $((1 - v)ca^{-1}(1 - u)(1 - w))^{2,1} \in K$ for all $u, v, w \in G$.

Proof. Note that K is a subgroup of $GL_2 A$ containing H and normalized by $E_2 A$ and all diagonal matrices $\text{diag}(u, v)$ with $u, v \in G$. We write $h = (c')^{2,1} \text{diag}(a, d')(b)^{1,2}$, where $c' = ca^{-1} \in A$, $b' = a^{-1}b$, and $d' = d - c'b$. We have

$$\begin{aligned} h_1 &= [h, \text{diag}(u^{-1}, 1)] = (c')^{2,1} \text{diag}(a, d')b^{1,2}(-u^{-1}b')^{1,2} \\ &\quad \times \text{diag}(u^{-1}a^{-1}u, (d')^{-1})(-c'u)^{2,1} \\ &= (c'')^{2,1} \text{diag}([a, u^{-1}], 1)(*)^{1,2}(-c'u)^{2,1} \in K, \end{aligned}$$

hence $h_2 = (-c'u)^{2,1} h_1 (c'u)^{2,1} = (c' - c'u')^{2,1} \text{diag}([a, u^{-1}], 1)(*) \in K$.

So $h_3 = [h_2, \text{diag}(1, v)] = (c' - c'u')^{2,1} \text{diag}([a, u^{-1}], 1)(*)^{1,2} (v(c' - c'u'))^{2,1} \in K$, hence

$$h_4 = (v(c' - c'u'))^{2,1} h_3 (-v(c' - c'u'))^{2,1} = ((1 - v)(c' - c'u'))^{2,1} (*)^{1,2} \in K.$$

Since $h_4 = ((1 - v)(c' - c'u'))^{2,1} (*)^{1,2} \in K$, $((1 - v)(c' - c'u'))^{2,1} \in K$. So

$$[((1 - v)(c' - c'u'))^{2,1}, \text{diag}(w, 1)] = ((1 - v)(c' - c'u')(1 - w))^{2,1} \in K.$$

COROLLARY 13. *Under the conditions of Lemma 12, assume that $\Sigma A(G - 1) = \Sigma(G - 1)A = A$ and that every element of A is a sum of units from G . Then $(AcA + AbA)^{2,1} \subset K$.*

Proof. Set $B = \{x \in A : x^{2,1} \in K\}$. We have to prove that $AcA + AbA \subset B$. Since K is normalized by all diagonal matrices $\text{diag}(u, v)$ with $u, v \in G$, the condition $\Sigma G = A$ gives that B is an ideal of A . By Lemma 12, $(1 - v)c'(1 - u)(1 - w) \in B$ for arbitrary units u, v and w of A , where $c' = ca^{-1} \in B$. Using the condition $\Sigma A(G - 1) = \Sigma(G - 1)A = A$, we obtain that $Ac'A = AcA \subset B$.

Now we apply this to the element tht^{-1} of K instead of h , where $t = 1^{1,2}(-1)^{2,1} 1^{1,2} \in E_2 A$. This gives that $AbA \subset B$.

LEMMA 14. *Let*

$$h = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in GL_2 A$$

commute with $E_2 A$ modulo the center $G_2(A, 0)$ of $GL_2 A$. Then $a \in GL_1 A$.

Proof. Case 1. A is commutative. Then the second entry in the first column of the matrix $[1^{1,2}, h] \in G_2(A, 0)$ is $c^2/\det(h) = 0$, hence $c^2 = 0$. So $aA = A$.

General case. Since $1^{1,2}h = h 1^{1,2} \text{diag}(\mu, \mu)$ with μ in the center of A , we conclude that $a + b = a\mu$, hence $ab = ba$. Similarly, every off-diagonal entry of h commutes with every diagonal entry of h .

Since $tht^{-1} \in G_2(A, 0)$, where $t = -t^{-1} = 1^{1,2}(-1)^{2,1} 1^{1,2} \in E_2A$, we conclude that $ad = da$ and $bc = cb$. So the subring A_0 of A generated by $1, a, b, c, d$ is commutative. Consider now the dominion of A_0 in A [16]. This is a commutative subring of A , and it contains all the entries of h^{-1} [16, Propositions 1.1 and 1.3]. So we are reduced to Case 1.

LEMMA 15. Let G, H , and K be as in Lemma 12. Suppose that $\Sigma A(G - 1) = \Sigma(G - 1)A = A$, i.e., no proper one-sided ideal of A contains all $1 - u$ with $u \in G$. Assume one of the conditions (3), (4), (5). Assume also that K is not contained in the center of GL_2A . Then there is a non-zero $y \in A$ such that $y^{1,2} \in K$.

Proof. Pick a matrix h in K outside the center $G_2(A, 0)$ of $G_2(A, A) = GL_2A$. Write

$$h = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in H. \text{ Let } h^{-1} = \begin{bmatrix} a' & b' \\ c' & d' \end{bmatrix}.$$

Case 1. $a \in GL_1A$. By Lemma 12 and the condition on G , we can assume that $b = c = 0$. Since h is outside the center, there is $x \in A$ such that $ax \neq xc$. Then $[h, x^{1,2}] = y^{1,2} \in K$ with $y = axd^{-1} - x \neq 0$. Note that we did not use any of the conditions (3), (4), (5).

General case. Under the condition (3), we argue as follows. By (3), there is $e \in A$ such that $a + xc$ is a unit of A . Replacing h by $e^{1,2}h(-e)^{1,2}$, we can assume that a is a unit. So we are reduced to Case 1.

Under condition (4), we argue as follows. If $b, c \in \text{rad}(A)$, then $a \in GL_1A$, so we are done by Case 1. Otherwise, we can assume that $c \notin \text{rad}(A)$ (the case $b \notin \text{rad}(A)$ is similar).

We pick $z \in A$ such that $czc \equiv c \pmod{\text{rad}(A)}$ and set $e = zc \in A$. Note that $e^2 \equiv e$ and $ce \equiv c \pmod{\text{rad}(A)}$. Replacing h by $(-az)^{1,2}h(az)^{1,2} \in K$, we change a by $a - azc = a(1 - e)$, and keep the same c . So we can assume that $a = a(1 - e)$.

Then $d'c \equiv d'ce \equiv (d'c + c'a)e = 0 \pmod{\text{rad}(A)}$ and the first row of the matrix $[h, e^{1,2}]$ (modulo $\text{rad}(A)$) is $(1, -e)$. Since $ce \equiv c \neq 0 \pmod{\text{rad}(A)}$, $e \neq 0 \pmod{\text{rad}(A)}$. So $[h, e^{1,2}] \in H$ is outside the center of GL_2A , and we are reduced to Case 1.

Under condition (5), having Case 1 done, we can assume, by Lemma 14, that there is an elementary matrix g which does not commute with h modulo the center. Say, $g = z^{1,2}$. By condition (5), we find $x_i \in A$ such that $1 + c'azzx_i \in GL_1A$ and

$GL_1 A$ and $\sum x_i = 1$. Since $g = z^{1,2} = \prod(zx_i)^{1,2}$, we can pick some i such that $(zx_i)^{1,2}$ does not commute with h modulo the center. The first entry of the first row of the matrix $h' = [h, (zx_i)^{1,2}] \in H$ is $1 + azx_i c' \in GL_1 A$. So we are reduced to Case 1, applied to h' instead of h .

Now we can complete our proof of Theorem 8. We define

$$K = \{g \in GL_2 A : [g, E_2 A] \subset H\}$$

as in Lemma 12 above. We want to prove that $E_2(A, B) \subset K \subset G_2(A, B)$ for an ideal B of A . Set $B = \{x \in A : x^{2,1} \in K\}$. Since K is normal in $GL_2 A$, condition (7) gives that B is an ideal of A . Since $t = -t^{-1} = 1^{1,2}(-1)^{2,1}1^{1,2} \in E_2 A$ and $tB^{2,1}t^{-1} = B^{2,1}$, we obtain that $E_2(A, B) \subset H$. We have to prove that $K \subset G_2(A, B)$, i.e. $f(H)$ is central in $GL_2(A/B)$, where $f: GL_2(A/B)$ is the canonical homomorphism.

Set $H' = f(H)$ and $K' = \{g \in GL_2(A/B) : [g, E_2(A/B)] \subset H'\}$. Note that both H' and K' are normalized by $f(GL_2 A)$. Clearly, $f(K) \subset K'$. So it suffices to show that $K' \subset G_2(A/B, 0)$, the center of $GL_2(A/B)$.

Note that the conditions of Theorem 8 on A imply the corresponding conditions on A/B . We assume now that K' is not central, and will obtain a contradiction.

Applying Lemma 15 (with G being the image of $GL_1 A$ in $GL_1(A/B)$) to H' and K' instead of H and K , we obtain that K' contains a non-trivial elementary matrix. Say, $c^{1,2} \in K'$ with $0 \neq c = a + B \in A/B$, where $a \in A \setminus B$. Then $[c^{1,2}, t] = c^{1,2}c^{2,1} \in H' = f(H)$. That is, H contains a matrix of the form hg , where $h \in GL_2 B$ and $g = a^{1,2}a^{2,1}$. Now we use that $[GL_2 B, E_2 A] \subset E_2(A, B)$ (see [3], [20], and [22]). In particular, $[1^{2,1}, h] \in E_2(A, B) \subset H$. So $[1^{2,1}, g] = [1^{2,1}, a^{1,2}] \in H$. Therefore $(-1)^{1,2}[1^{2,1}, a^{1,2}]1^{1,2} = (-a)^{2,1}(-a)^{1,2} \in H \subset K$, hence $(-a)^{2,1} \in K$ by the definition of K and $a \in B$ by the definition of B . A contradiction!

References

1. E. Abe, Chevalley groups over local rings, *Tôhoku Math. J.* 21 (1969), 474–494.
2. J.-B. An and X.-P. Tang, The structure of symplectic groups over semilocal rings, *Acta Math. Sinica (N.S.)* 1:1 (1985), 1–15.
3. H. Bass, K -theory and stable algebra, *Publ. Math. IHES* 22 (1964), 5–60.
4. D.L. Costa and G.E. Keller, On the normal subgroups of $SL(2, A)$, *J. Pure Appl. Alg.* 53:3 (1988), 201–227.
5. C.N. Chang, The structure of the symplectic group over semi-local domains, *J. Algebra* 35 (1975), 457–476.
6. W. Klingenberg, Lineare Gruppen über lokalen Ringen, *Amer. J. Math.* 83 (1961), 137–153.
7. W. Klingenberg, Die Struktur der linearen Gruppe über einen nichtkommutativen lokalen Ring, *Arch. Math.* 13 (1962), 73–81.

8. L. Yu. Kolotilina and N.A. Vavilov, Normal structure of the full linear group over semilocal rings, *J. Sov. Math.* 19:1 (1982), 998–999.
9. N.H.J. Lacroix, Two-dimensional linear groups over local rings, *Can. J. Math.* 21 (1969), 106–135.
10. N.H.J. Lacroix and C. Levesque, Sur les sous-groupes normaux de SL_2 sur un anneau local, *Can. Math. Bull.* 26:2 (1983), 209–219.
11. B.A. Magurn, W. van der Kallen, and L.N. Vaserstein, Absolute stable rank and Witt cancellation for noncommutative rings, *Inv. Math.* 91:3 (1988), 525–542.
12. A.W. Mason, On GL_2 of a local ring in which 2 is not a unit, *Can. Math. Bull.* 30:2 (1987), 165–176.
13. B.R. McDonald, GL_2 of rings with many units, *Commun. Algebra* 8:9 (1980), 869–888.
14. B.R. McDonald, Geometric algebra over local rings, New York-Basel, Marcel Dekker, 1976.
15. P. Menal and L.N. Vaserstein, On subgroups of GL_2 over Banach algebras and von Neumann regular rings which are normalized by elementary matrices, preprint, 20 p.
16. H.H. Storrer, Epimorphic extensions of non-commutative rings, *Comm. Math. Helv.* 18 (1973), 72–86.
17. S. Tazhetdinov, Subnormal structure of two-dimensional linear groups over local rings, *Alg. Logic* 22:6 (1983), 707–713.
18. S. Tazhetdinov, Subnormal structure of two-dimensional linear groups over rings that are close to fields *Alg. Logic* 24:4 (1985), 414–425.
19. L.N. Vaserstein, K_1 -theory and the congruence subgroup problem, *Mat. Zametki* 5 (1969), 233–244 = *Math. Notes* 5 (1969), 141–148.
20. L.N. Vaserstein, Normal subgroups of the general linear groups over von Neumann regular rings, *Proc. Amer. Math. Soc.* 96:2 (1986), 209–214.
21. L.N. Vaserstein, Normal subgroups of the general linear groups over Banach algebras, *J.P.A. Algebra* 41 (1986), 99–112.
22. L.N. Vaserstein, Subnormal structure of the general linear groups over Banach algebras, *J. Pure Appl. Alg.*, 52 (1988), 187–195.
23. L.N. Vaserstein, Bass's first stable range condition, *J. Pure Appl. Algebra* 34:2–3 (1984), 319–330.
24. L.N. Vaserstein, An answer to the question of M. Newman on matrix completion, *Proc. Amer. Math. Soc.* 97:2 (1986), 189–196.
25. L.N. Vaserstein, On stabilization for general linear groups over a ring, *Mat. Sbornik* 79:3 (1969), 405–424 = *Math. USSR Sbornik* 8, 383–400.
26. L.Q. Wang and Y.Z. Zhang, GL_2 over full rings, *Chin. An. Math. ser. B*, 8:4 (1987), 434–439.