

GROUPE DE TRAVAIL D'ANALYSE ULTRAMÉTRIQUE

QUIDO VAN STEEN

Decomposition of non-archimedean analytic tori

Groupe de travail d'analyse ultramétrique, tome 9, n° 3 (1981-1982), exp. n° J18, p. J1-J5

<http://www.numdam.org/item?id=GAU_1981-1982__9_3_A19_0>

© Groupe de travail d'analyse ultramétrique
(Secrétariat mathématique, Paris), 1981-1982, tous droits réservés.

L'accès aux archives de la collection « Groupe de travail d'analyse ultramétrique » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale.
Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>*

DECOMPOSITION OF NON-ARCHIMEDEAN ANALYTIC TORI

by Guido VAN STEEN (*)

[Universiteit Antwerpen]

Introduction. — In [3], SHIODA and MITANI prove that a complex abelian surface X is isogenous to the product $C \times C$ of an elliptic curve C with itself if, and only if, X is singular. This means that the rank of the Néron-Severi group of X takes its maximal value. The theorem can be generalised for higher dimensions.

In this paper, we prove an analogous result for abelian varieties which are defined over a complete non-archimedean valued field and which are (analytically) isomorphic to an analytic torus.

1. Definitions and notations.

Let K be a complete non-archimedean valued field. We assume that K is algebraically closed. The multiplicative group $(K^*)^n$ is identified with the affine group $\text{Spec}(K[z_1, z_1^{-1}, \dots, z_n^{-1}])$ where z_1, \dots, z_n are variables. This group has a canonical analytic structure (see e. g. [1], [2]).

A lattice Γ in $(K^*)^n$ is a discrete subgroup of $(K^*)^n$ without torsion. This means that

$$\Gamma \cap \{(x_1, \dots, x_n) \in (K^*)^n ; |\pi_1| \leq |x_i| \leq |\pi_2| \text{ for all } i = 1, \dots, n\}$$

is a finite set for all $\pi_1, \pi_2 \in K^*$ with $|\pi_1| \leq |\pi_2|$.

The factor group $T = (K^*)^n / \Gamma$ is given an analytic structure by requiring that the canonical map

$$\pi : (K^*)^n \rightarrow (K^*)^n / \Gamma$$

is locally biholomorphic (see e. g. [2]). The analytic space T is called a holomorphic torus of dimension n .

Let A^* be the group of nonvanishing analytic functions on $(K^*)^n$. Each $f \in A^*$, can be written as

$$f = c z_1^{r_1} \times \dots \times z_n^{r_n}$$

(*) Guido VAN STEEN, Mathematics, Rijksuniversitair centrum, 171 Groenenborgerlaan, B-2020 ANTWERPEN (Belgique).

with $c \in K^*$ and $r_1, \dots, r_n \in \mathbb{Z}$.

Let H be the character group of $(K^*)^n$; i. e.

$$H = \{z_1^{r_1} \times \dots \times z_n^{r_n}; r_1, \dots, r_n \in \mathbb{Z}\}.$$

The action of Γ on A^* , defined by $f'(x) = f(\gamma x)$ for each $f \in A^*$, $\gamma \in \Gamma$ and $x \in (K^*)^n$, makes A^* into a Γ -module.

Providing K^* and H with the trivial Γ -action, we obtain the following exact sequence of Γ -modules :

$$1 \rightarrow K^* \xrightarrow{\alpha} A^* \xrightarrow{\beta} H \rightarrow 1$$

with

$$\alpha(\lambda) = \lambda, \quad \beta(f) = (f(1))^{-1} \cdot f \quad \text{for each } \lambda \in K^* \text{ and } f \in A^*.$$

Associated is the exact sequence of cohomology groups :

$$\dots \rightarrow H^1(\Gamma, K^*) \xrightarrow{\alpha^*} H^1(\Gamma, A^*) \xrightarrow{\beta^*} H^1(\Gamma, H) \xrightarrow{\delta} H^2(\Gamma, K^*) \rightarrow \dots$$

Since K^* and H are trivial Γ -modules, we have the following isomorphisms :

$$H^1(\Gamma, K^*) \cong \text{Hom}(\Gamma, K^*), \quad H^2(\Gamma, K^*) \cong \text{Hom}(\overset{2}{\underset{\mathbb{Z}}{\text{A}}}, \Gamma, K^*)$$

and

$$H^1(\Gamma, H) \cong \text{Hom}(\Gamma, H).$$

Fixing a basis $\gamma_1, \dots, \gamma_g$ for the free abelian group Γ , we can identify

$$\text{Hom}(\overset{2}{\underset{\mathbb{Z}}{\text{A}}}, \Gamma, K^*) \text{ with } (K^*)^{\binom{n}{2}}.$$

The map δ is then defined by

$$\delta(\sigma) = [\frac{q(\gamma_i, \sigma(\gamma_j))}{q(\gamma_j, \sigma(\gamma_i))}]_{i < j, j=2, \dots, n}$$

where $q : \Gamma \times H \rightarrow K^*$ is the bilinear form defined by

$$q(\gamma, z) = z(\gamma).$$

The group $\text{Ker } \delta$ is denoted by $N(T)$. It is proved, in [1], that

$$N(T) = \{\sigma \in \text{Hom}(\Gamma, H); q(\gamma, \sigma(\gamma')) = q(\gamma', \sigma(\gamma)), \text{ for all } \gamma, \gamma' \in \Gamma\}.$$

Furthermore, T is analytically isomorphic to an n -dimensional abelian variety if, and only if, there exists some monomorphism $\sigma \in N(T)$ such that

$$|q(\gamma, \sigma(\gamma))| < 1 \quad \text{for all } \gamma \in \Gamma - \{1\}.$$

In this case, $N(T)$ is isomorphic to the Néron-Severi group of this abelian variety.

2. Decomposition of analytic tori.

We keep the notations of the previous paragraph.

LEMMA 1. - Rank($\text{Hom}(\Gamma, H)/N(T)$) $\geq \frac{n(n-1)}{2}$.

Proof. - Assume that $\gamma_i = (\gamma_{i1}, \dots, \gamma_{in})$, for all $i = 1, \dots, n$, and let $a_{ij} = -\log|\gamma_{ij}|$ for all $i, j = 1, \dots, n$.

It is easy to see that the matrix A , with entries a_{ij} , has rank n .

(a) Since A is regular, we may assume that, after renumbering the columns of A ,

$$a_{11} \neq 0.$$

For $k = 2, \dots, n$, we define $\sigma_{1k} \in \text{Hom}(\Gamma, H)$ by

$$\sigma_{1k}(z_k) = z_1^{\delta_{k,k}} \quad \text{where } \delta_{k,k} = 0 \text{ if } k \neq k \text{ and } \delta_{k,k} = 1.$$

For each $\sigma = \sigma_{1,2}^{m_2} \times \dots \times \sigma_{1,n}^{m_n} \in \text{Hom}(\Gamma, H)$, we have

$$q(\gamma_1, \sigma(\gamma_k)) = q(\gamma_1, z_1^{m_k}) = \gamma_{11}^{m_k}$$

and

$$q(\gamma_k, \sigma(\gamma_1)) = q(\gamma_k, 1) = 1$$

for all $k = 2, \dots, n$. It follows that $\sigma \notin N(T)$ unless $m_2 = \dots = m_n = 0$, and consequently $\sigma_{12}, \dots, \sigma_{1n}$ are \mathbb{Z} -independent modulo $N(T)$.

(b) Since A is regular, we may assume that, after renumbering all but the first column of A , $(\begin{smallmatrix} a_{11} \\ a_{21} \end{smallmatrix})$ and $(\begin{smallmatrix} a_{12} \\ a_{22} \end{smallmatrix})$ are linear independent over \mathbb{Z} .

For $k = 3, \dots, n$, we define $\sigma_{2k} \in \text{Hom}(\Gamma, H)$ by

$$\sigma_{2k}(z_k) = z_2^{\delta_{k,k}}.$$

One proves in an analogous way as in (a) that $\sigma_{12}, \dots, \sigma_{1n}, \sigma_{23}, \dots, \sigma_{2n}$ are linear independent modulo $N(T)$.

Repeating this construction for each $i = 3, \dots, n$, one defines for each $j > i$ a morphism $\sigma_{ij} \in \text{Hom}(\Gamma, H)$ such that $\sigma_{1,2}, \dots, \sigma_{n-1,n}$ are linear independent modulo $N(T)$. It follows that

$$\text{rank}(\text{Hom}(\Gamma, H)/N(T)) \geq (1 + 2 + \dots + (n-1)) = \frac{n(n-1)}{2}.$$

COROLLARY 2. - Rank($N(T)$) $\leq \frac{n(n-1)}{2}$.

THEOREM 3. - If $\text{rank}(N(T)) = \frac{n(n-1)}{2}$ then T is analytically isomorphic to an n -dimensional abelian variety. This variety is isogenous to the n -th self product

of an elliptic curve C of the form $C = \mathbb{K}^*/q\mathbb{Z}$ with $q \in \mathbb{K}^*$ and $|q| < 1$.

Proof. - By means of the basis $\gamma_1, \dots, \gamma_n$ of Γ , we may identify $H^2(\Gamma, \mathbb{K}^*)$ with $(\mathbb{K}^*)^{(n)}/(\mathbb{K}^*)^{(2)}$ and the map

$$H^1(\Gamma, H) = \text{Hom}(\Gamma, H) \rightarrow H^2(\Gamma, \mathbb{K}^*)$$

is defined by

$$\delta(\sigma) = [\frac{q(\gamma_i, \sigma(\gamma_j))}{q(\gamma_j, \sigma(\gamma_i))}]_{i < j, j=2, \dots, n}.$$

For each $i < j$ and $j = 1, \dots, n-1$, we define $\sigma_{ij} \in \text{Hom}(\Gamma, H)$ such as in the proof of lemma 1. Let S be the group generated by

$$\{\delta(\sigma_{ij}) ; i < j, i = 1, \dots, n-1\}.$$

For each $k = 1, \dots, n$, we define $\sigma_k \in \text{Hom}(\Gamma, H)$ by

$$\sigma_k(z_1) = z_k^{\delta_{k1}}.$$

Since $N(T)$ has rank $\frac{n(n+1)}{2}$, $\text{Im } \delta$ has rank $\frac{n(n-1)}{2}$, and hence the group S is of finite index in $\text{Im } \delta$.

It follows that $\delta(\sigma_k)^{s_k} \in S$ for some $s_k \in \mathbb{N}$.

An explicit calculation of $\delta(\sigma_{ij})$ and $\delta(\sigma_k)$ shows that

$$\gamma_{ij}^{u_{ij}} \in \gamma_{11}^Z \text{ for some } u_{ij} \in \mathbb{N}; i, j = 1, \dots, n.$$

We may assume that $|\gamma_{11}| < 1$. (If necessary replace γ_1 by γ_1^{-1} .) Consequently, there exists some $q \in \mathbb{K}^*$ with $|q| < 1$, and there exist roots of unity $\xi_{ij} \in \mathbb{K}^*$ such that

$$\gamma_{ij} = \xi_{ij} q^{m_{ij}} \text{ for some } m_{ij} \in \mathbb{Z}; i, j = 1, \dots, n.$$

Since $\gamma_1, \dots, \gamma_n$ are linear independent the matrix M with entries m_{ij} is regular. Hence there exists a matrix $B \in GL_n(\mathbb{Z})$ with entries b_{ij} such that

$$M \cdot B = \begin{bmatrix} m & & & \\ & m & & \\ & & \ddots & \\ & & & 0 \end{bmatrix} \text{ for some } m \in \mathbb{Z}.$$

Let $N \in \mathbb{N}$ such that $\xi_{ij}^N = 1$ for all $i, j = 1, \dots, n$, and let $\sigma \in \text{Hom}(\Gamma, H)$ be defined by

$$\sigma(\gamma_i) = z_1^{Nb_{1i}} \times \dots \times z_n^{Nb_{ni}} \text{ for all } i = 1, \dots, n.$$

For all $i, j = 1, \dots, n$, we have

$$q(\gamma_i, \sigma(\gamma_j)) = q^{Nb_{ij}}.$$

It follows that $\sigma \in N(T)$ and that $|q(\gamma, \sigma(\gamma))| < 1$ for all $\gamma \in \Gamma - \{1\}$. Hence T is analytically isomorphic to an abelian variety.

The endomorphism β of $(K^*)^n$, defined by

$$\beta(x_1, \dots, x_n) = (\prod_{j=1}^n x_j^{b_{j1}}, \dots, \prod_{j=1}^n x_j^{b_{jn}})$$

maps Γ onto a subgroup Γ' of $(K^*)^n$ which is generated by the elements

$$\gamma_i' = (\tau_{i1}, \dots, \tau_{ii} q^m, \tau_{ii+1}, \dots, \tau_{in})$$

where $\tau_{ij} \in K^*$ are roots of unity.

Let $M \in \mathbb{N}$ such that $\tau_{ij}^M = 1$ for all $i, j = 1, \dots, n$.

The endomorphism τ of $(K^*)^n$ defined by

$$\tau(x_1, \dots, x_n) = (x_1^M, \dots, x_n^M)$$

maps Γ' onto a subgroup Γ'' of $(K^*)^n$, defined by the elements

$$\gamma_i'' = (1, \dots, 1, q^{Mm}, 1, \dots, 1)$$

where q^{Mm} stands on the i -th entry.

Furthermore $\tau \circ \beta$ induces a morphism

$$\rho : (K^*)^n / \Gamma \rightarrow (K^*)^n / \Gamma'' .$$

It is clear that

$$(K^*)^n / \Gamma'' \simeq (K^* / (q^{Mm})^\mathbb{Z})^n ,$$

and that ρ is finite; i.e. ρ is an isogeny.

BIBLIOGRAPHY

- [1] GERRITZEN (L.). - On non-archimedean representations of abelian varieties, Math. Annalen, t. 196, 1972, p. 323-346.
- [2] GERRITZEN (L.) and VAN DER PUT (M.). - Schottky groups and Mumford curves. - Berlin, Heidelberg, New York, Springer-Verlag, 1980 (Lecture Notes in Mathematics, 817).
- [3] SHIODA (T.) and MITANI (N.). - Singular abelian surfaces and binary quadratic forms, "Classification of algebraic varieties and compact complex manifolds", p. 259-287. - Berlin, Heidelberg, New York, Springer-Verlag, 1974 (Lecture Notes in Mathematics, 412).