

GÉRARD G. EMCH

Algebraic K-Flows

Publications des séminaires de mathématiques et informatique de Rennes, 1975, fascicule S4

« International Conference on Dynamical Systems in Mathematical Physics », , p. 1-2

http://www.numdam.org/item?id=PSMIR_1975__S4_A5_0

© Département de mathématiques et informatique, université de Rennes, 1975, tous droits réservés.

L'accès aux archives de la série « Publications mathématiques et informatiques de Rennes » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

ALGEBRAIC K-FLOWS

Gérard G. Emch

Zentrum für interdisziplinäre Forschung
der Universität Bielefeld

The algebraic approach to the study of Statistical Mechanics suggests an expansion of classical ergodic theory to a noncommutative ergodic theory. We indicate here how one can proceed in this spirit to generalise the classical concept of a Kolmogorov-Sinai flow.

Firstly we recall that a classical K-flow is constituted by: a probability space (Ω, Σ, μ) , a measurable group $\{T(t) \mid t \in \mathbb{R}\}$ of measure preserving transformations of (Ω, Σ, μ) , and a partition $\xi \subset \Sigma$ such that: (i) $\xi \subseteq T(t)[\xi] \quad \forall t \geq 0$; (ii) $\bigcap_{t \in \mathbb{R}} T(t)[\xi] = \hat{0}$; (iii) $\bigvee_{t \in \mathbb{R}} T(t)[\xi] = \hat{1}$.

Secondly we construct from these elements: a separable Hilbert space $\mathfrak{H} = L^2(\Omega, \mu)$, a maximal abelian von Neumann algebra $\mathcal{N} = L^\infty(\Omega, \Sigma)$ acting on \mathfrak{H} , a cyclic and separating vector $\Phi(\omega) = 1 \quad \forall \omega \in \Omega$ for \mathcal{N} in \mathfrak{H} , a faithful normal state on $\mathcal{N} \quad \phi : f \in \mathcal{N} \mapsto \langle \Phi, f \Phi \rangle = (f \Phi, \Phi) = \int f(\omega) d\mu(\omega)$, a continuous group of automorphisms of $\mathcal{N} \quad \alpha(t) : f \in \mathcal{N} \mapsto \alpha(t)[f] = f \circ T(t)$, and a von Neumann subalgebra \mathcal{A} of $\mathcal{N} \quad \mathcal{A} = \{\chi_\Delta \mid \Delta \in \xi\}''$ with the properties: (i) $\mathcal{A} \subseteq \alpha(t)[\mathcal{A}] \quad \forall t \geq 0$; (ii) $\bigcap_{t \in \mathbb{R}} \alpha(t)[\mathcal{A}] = \text{CI}$; (iii) $\bigvee_{t \in \mathbb{R}} \alpha(t)[\mathcal{A}] = \mathcal{N}$.

Thirdly we notice that a standard representation theorem allows to get back to the original definition from $(\mathcal{N}, \phi, \alpha, \mathcal{A})$ where: \mathcal{N} is an abelian von Neumann algebra acting on a separable Hilbert space \mathfrak{H} , ϕ is a faithful normal state on \mathcal{N} , $\alpha : \mathbb{R} \rightarrow \text{Aut}(\mathcal{N}, \phi)$, and \mathcal{A} is a completely selfrefining, generating von Neumann subalgebra of \mathcal{N} .

Fourthly the generalisation to the noncommutative domain now consists exactly in taking the above as an alternative definition of a classical K-flow, and in dropping from this definition the condition that \mathcal{N} be abelian. For reasons which would be too long to make explicit here we also impose that \mathcal{A} be stable under the modular automorphism group $\{\sigma_\phi(t) \mid t \in \mathbb{R}\}$ canonically associated to ϕ , and that every maximal abelian subalgebra \mathcal{L} of the centralizer \mathcal{N}_ϕ of \mathcal{N} be already maximal abelian in \mathcal{N} . (Notice that both of the last two conditions are redundant when \mathcal{N} is abelian since $\sigma_\phi(t) = \text{id} \quad \forall t \in \mathbb{R}$ in this case.)

We now can emphasize [1] that algebraic proofs can be given to several theorems which are well-known in the classical case, and which

thus do generalize to the new situation just defined. For instance the system $(\mathcal{K}, \phi, \alpha, \mathcal{A})$ is ergodic (i.e. $N \in \mathcal{K}$ and $\alpha(t)[N] = N \quad \forall t \in \mathbb{R} \Rightarrow N = \lambda I$ with $\lambda \in \mathbb{C}$); it is mixing (i.e. $\langle \phi; N \alpha(t)[M] \rangle \rightarrow \langle \phi; N \rangle \langle \phi; M \rangle$ as $t \rightarrow \pm\infty$ for all $N, M \in \mathcal{K}$); it has Lebesgue spectrum (i.e. $\alpha(t)$ is spatial and the generator H of the corresponding unitary group on \mathcal{S} has the property $\text{Sp}(H) = \text{Sp}_d(H) \cup \text{Sp}_{ac}(H)$ with $\text{Sp}_d(H) = \{0\}$ simple and $\text{Sp}_{ac}(H) = \mathbb{R}$ has countable multiplicity). Furthermore a noncommutative entropy can be defined [2] which is strictly positive for all such systems.

We next remark that the generalisation is genuine in the sense that, in addition to the classical case where \mathcal{K} is abelian, there exist [1,3,4] K-flows where \mathcal{K} is of type II_1 , III_2 ($0 < \lambda < 1$), or III_1 .

Finally a link has been established [3] between certain quantum transport phenomena, governed by an evolution equation of the diffusion type, and the generalised K-flows presented here.

References

- [1] Nonabelian Special K-flows
Journ. Funct. Analysis 19 (1975) 1-12.
- [2] Positivity of the K-entropy on Non-Abelian K-Flows
Z. Wahrscheinlichkeitstheorie verw. Gebiete 29 (1974) 241-252.
- [3] The Minimal K-Flow associated to a Quantum Diffusion Process
in Physical Reality and Mathematical Description, Enz & Mehra, Eds.,
D. Reidel Publishing Company, Dordrecht-Holland, 1974, 477-493.
- [4] in preparation

Permanent address: Departments of Mathematics and of Physics,
The University of Rochester, Rochester N.Y. (USA)