

GARY J. KØEHLER

ANDREW B. WHINSTON

GORDON P. WRIGHT

**Fixed cost minimization over a Leontief
substitution system**

Revue française d'automatique, informatique, recherche opérationnelle. Recherche opérationnelle, tome 8, n° V2 (1974), p. 119-124

http://www.numdam.org/item?id=RO_1974__8_2_119_0

© AFCET, 1974, tous droits réservés.

L'accès aux archives de la revue « Revue française d'automatique, informatique, recherche opérationnelle. Recherche opérationnelle » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

FIXED COST MINIMIZATION OVER A LEONTIEF SUBSTITUTION SYSTEM ⁽¹⁾

par Gary J. KÖHLER, Andrew B. WHINSTON and Gordon P. WRIGHT ⁽²⁾

Résumé. — *Using Bender's decomposition method and properties of a Leontief substitution system, a formulation for solving fixed cost minimizations over Leontief substitution systems is presented.*

It has been pointed out in the literature [2] that Bender's decomposition method appears useful as a solution strategy. Using Bender's method and the properties of a Leontief substitution system, a solution procedure is specified for solving fixed cost minimization problems over a Leontief substitution system. Preliminary results indicate that for some large problems, this procedure is a viable alternative to procedures specified elsewhere [4].

A Leontief substitution system is characterized by a constraint matrix having exactly one positive element per column and non-trivial rows [3]. Substitute activities are identified by columns whose positive elements fall in the same row. A salient feature of Leontief substitution systems is that one and only one activity is optimal for each set of substitute activities.

Consider the problem

$$\begin{aligned} & \text{Min } f(x) \\ & \text{subject to} \\ & \quad Ax = b \\ & \quad x \geq 0 \\ & \quad b \geq 0 \end{aligned}$$

where A is m by n and Leontief, x is n by 1, b is m by 1, and $f(\cdot)$ is a quasi-concave function of the form :

$$f(x) = \sum_{i=1}^n f_i(x_i)$$

(1) This research was supported by the Office of Army Research under contract # DA-31-124-ARO-D-477.

(2) Krannert Graduate School of Industrial Administration Purdue University, West Lafayette.

where

$$f_i(x_i) = \begin{cases} c_i x_i + d_i & x_i > 0 \\ 0 & x_i = 0 \end{cases} \quad d_i \geq 0$$

The above problem is termed a fixed cost minimization over a Leontief substitution system.

Let the Set J_i contain the column identifications for all columns whose positive coefficients fall in row i . Associate with each continuous variable x_i a binary variable y_i and reformulate the fixed cost problem as :

$$(1) \quad \text{Min } c'x + d'y$$

subject to

$$\begin{aligned} Ax &= b \\ x - ly &\leq 0 \\ x &\geq 0 \\ b &\geq 0 \\ y &\in Y \end{aligned}$$

where

$$Y = \{ y : y \text{ is binary and } \sum_{j \in J_i} y_j = 1, i = 1, 2, \dots, m \}$$

and

$$l > x_i$$

for all possible x_i . The requirement that

$$\sum_{j \in J_i} y_j = 1 \quad i = 1, 2, \dots, m$$

is a result of the substitution property of Leontief substitution systems.

The problem given in equation (1) may be reformulated as :

$$\text{Min } d'y + \left[\begin{array}{l} \text{Min } c'x \\ x \\ \text{subject to} \\ Ax = b \\ x \leq ly \\ x \geq 0 \\ b \geq 0 \end{array} \right]$$

Using Bender's decomposition method [1], an approximation to the outer problem is now given. Let N denote the number of times the inner problem has been solved. Also, let π_1^i be the dual multipliers to the constraint set

$$Ax = b$$

at the solution of the inner problem at the i^{th} iteration. Likewise, let π_2^i be the dual multiplier to the constraint set

$$x \leq ly$$

at the i^{th} iteration. The approximation to the outer problem is

$$(2) \quad \text{Min } z + d'y$$

subject to

$$z \geq b'\pi_1^i + ly'\pi_2^i \quad i = 1, 2, \dots, N$$

$$y \in Y$$

Given a $y \in Y$, the inner problem is easily solved. Let y^i be the fixed y vector supplied to the inner problem at the beginning of iteration i . Then the solution to the inner problem is :

$$(3) \quad \pi_1^i = (A'_K)^{-1}c_K$$

where

$$K = \{j : y_j^i = 1\}$$

and A'_K is the transpose of the submatrix of A consisting of the columns in the set K . Also

$$(4) \quad \pi_2^i = [c - A'\pi_1^i]^+$$

where $[a]^+$ is a vector of $[a_i]^+$ $i = 1, 2, \dots, n$, and $[a_i]^+ = \min [a_i, 0]$.

For computational purposes, it is beneficial to note that if A_K is recursive [4], as is usually the case, then $[A'_K]^{-1}$ is easily computed.

As an example, consider an arborescence-structured production and inventory system. Let x_i^K , y_i^K , and r_i^K be, respectively, the production, end-of-period inventory, and known requirements at facility K in period i . For illustrative purposes let the number of facilities be 2 and the number of periods be 3. The arborescence model is

$$\text{Min } f(x, y)$$

subject to

$$y_{i-1}^1 + x_i^1 - y_i^1 - x_i^2 = r_i^1 \quad i = 1, 2, 3$$

$$y_{i-1}^2 + x_i^2 - y_i^2 = r_i^2 \quad i = 1, 2, 3$$

$$y_0^1 = y_0^2 = y_3^1 = y_3^2 = 0$$

$$x_i^K, y_i^K \geq 0$$

where

$$f(x, y) = \sum_{K=1}^2 \sum_{i=1}^3 \{f_i^K(x_i^K) + f_i^K(y_i^K)\}$$

and

$$f_i^K(x_i^K) = \begin{cases} c_i^K x_i^K + d_i^K & x_i^K > 0 \\ 0 & x_i^K = 0 \end{cases} \quad d_i^K \geq 0$$

and $f_i^K(y_i^K)$ is similarly defined. Let

$$r^1 = \begin{pmatrix} 3 \\ 4 \\ 2 \end{pmatrix} \quad C_x^2 = \begin{pmatrix} 5 \\ 4 \\ 7 \end{pmatrix} \quad C_y^2 = \begin{pmatrix} 5 \\ 3 \\ 0 \end{pmatrix} \quad d_x^1 = \begin{pmatrix} 1 \\ 3 \\ 1 \end{pmatrix} \quad d_y^1 = \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix}$$

$$C_x^1 = \begin{pmatrix} 3 \\ 2 \\ 4 \end{pmatrix} \quad C_y^1 = \begin{pmatrix} 1 \\ 3 \\ 0 \end{pmatrix} \quad r^2 = \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix} \quad d_x^2 = \begin{pmatrix} 1 \\ 6 \\ 4 \end{pmatrix} \quad d_y^2 = \begin{pmatrix} 2 \\ 7 \\ 0 \end{pmatrix}$$

By the structure of the problem, the set of all feasible basis is

$$K = \bigtimes_{i=1}^6 J_i$$

where

$$J_1 = \{x_1^1\} \quad J_4 = \{x_1^2\}$$

$$J_2 = \{y_1^1, x_2^1\} \quad J_5 = \{y_1^2, x_2^2\}$$

$$J_3 = \{y_2^1, x_3^1\} \quad J_6 = \{y_2^2, x_3^2\}$$

To start the solution procedure, an arbitrary initial feasible solution is specified. Let the optimal solution to the continuous portion of the problem be the initial starting basis. Then

$$K_1 = \{x_1^1, x_2^1, x_3^1, x_1^2, x_2^2, y_2^2\}$$

and the outer problem can be written as follows. Let the continuous variable identifications, x_i^K and y_i^K represent their binary counterparts.

Then equation (2) becomes :

$$\text{Min } z + d'_x x + d'_y y$$

subject to

$$z \geq 54$$

Then

$$K_2 = \{ x_1^1, y_1^1, y_2^1, x_2^2, y_1^2, x_3^2 \}$$

or

$$K_2 = \{ x_1^1, y_1^1, x_3^1, x_1^2, y_1^2, x_3^2 \}$$

and the outer problem becomes (using the first K_2) :

$$\text{Min } z + d'_x x + d'_y y$$

subject to

$$z \geq 54$$

$$z \geq 87 - 200x_2^1 - 300x_3^1 - 500x_2^2$$

where $l = 100$.

Continuing in like manner and using all alternative optimal points to form constraints, the following was found.

Iteration

3	$x_1^1, y_1^1, x_3^1, x_2^2, y_1^2, x_3^2$	$z \geq 78 - 200x_2^1 - 500x_2^2$
4	$x_1^1, x_2^1, x_3^1, x_2^2, y_1^2, x_3^2$	$z \geq 70 - 700x_2^2$
	$x_1^1, x_2^1, y_2^1, x_2^2, y_1^2, x_3^2$	$z \geq 73 - 100x_3^1 - 700x_2^2$
5	$x_1^1, y_1^1, y_2^1, x_2^2, x_2^2, x_3^2$	$z \geq 77 - 200x_2^1 - 300x_3^1 - 300y_2^2$
	$x_1^1, y_1^1, x_3^1, x_2^2, x_2^2, x_3^2$	$z \geq 68 - 200x_2^1$
6	$x_1^1, x_2^1, y_2^1, x_2^2, x_2^2, x_3^2$	$z \geq 59 - 100x_3^1 - 300y_2^2$
	$x_1^1, x_2^1, x_3^1, x_2^2, x_2^2, x_3^2$	$z \geq 56 - 200y_2^2$
7	$x_1^1, x_2^1, x_3^1, x_2^2, x_2^2, x_3^2$	

STOP

BIBLIOGRAPHY

- [1] BENDERS J. F., « Partitioning Procedures for Solving Mixed Variables Programming Problems », *Numerische Mathematik*, 4, 1962, pp. 238-252.
- [2] LASDON L. S., *Optimization Theory for Large Systems*, MacMillan Co., New York, 1970.
- [3] VEINOTT Jr., A. F. « Extreme Points of Leontief Substitution Systems », *Linear Algebra and its Applications*, 1, 1968, pp. 181-194.
- [4] VEINOTT Jr. A. F., « Minimum Concave-Cost Solution of Leontief Substitution Models of Multi Facility Inventory Systems », *Operations Research*, 17, No. 2, 1969, pp. 262-291.