

RENDICONTI *del* SEMINARIO MATEMATICO *della* UNIVERSITÀ DI PADOVA

S. K. CHATTERJEA

On a generalization of Laguerre polynomials

Rendiconti del Seminario Matematico della Università di Padova,
tome 34 (1964), p. 180-190

<http://www.numdam.org/item?id=RSMUP_1964__34__180_0>

© Rendiconti del Seminario Matematico della Università di Padova, 1964, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>*

ON A GENERALIZATION OF LAGUERRE POLYNOMIALS

*Nota *) di S. K. CHATTERJEA (a Calcutta)*

1. – In a recent paper [1], the writer has defined the polynomials $T_{kn}^{(\alpha)}(x)$ by the Rodrigues' formula

$$(1.1) \quad T_{kn}^{(\alpha)}(x) = \frac{1}{n!} x^{-x} e^{xk} D^n(x^{x+n} e^{-xk}) ,$$

where k is a natural number. The polynomials $T_{kn}^{(\alpha)}(x)$ are of exactly degree kn ($n = 0, 1, 2, \dots$). They satisfy the operational formula

$$(1.2) \quad \prod_{j=1}^n (xD - kx^k + \alpha + j) = n! \sum_{r=0}^n \frac{x^r}{r!} T_{k(n-r)}^{(\alpha+r)}(x) D^r .$$

The following are the consequences of the operational formula (1.2):

$$(1.3) \quad nT_{kn}^{(\alpha)}(x) = (xD - kx^k + \alpha + n) T_{k(n-1)}^{(\alpha)}(x)$$

$$(1.4) \quad \binom{m+n}{m} T_{k(m+n)}^{(\alpha)}(x) = \sum_{r=0}^{\min(m,n)} \frac{x^r}{r!} T_{k(m-p)}^{(\alpha+n+r)}(x) D^r T_{kn}^{(\alpha)}(x) .$$

The polynomials $T_{kn}^{(\alpha)}(x)$ are generated by the function

$$(1.5) \quad (1-t)^{-\alpha-1} \exp [x^k u(t)] = \sum_{n=0}^{\infty} T_{kn}^{(\alpha)}(x) t^n$$

*) Pervenuta in redazione il 17 giugno 1963.

Indirizzo dell'A.: Department of mathematics, Bangabasi College,
Calcutta (India).

where

$$u(t) = 1 - (1-t)^{-k}.$$

In the same paper the writer has also proved the following properties

$$(1.6) \quad \sum_{r=0}^{k+1} (-1)^r \binom{k+1}{r} (n+1-r) T_{kn+r}^{(\alpha)}(x) \\ = (\alpha+1) \sum_{r=0}^k (-1)^r \binom{k}{r} T_{kn+r}^{(\alpha)}(x) - kx^k T_{kn}^{(\alpha)}(x)$$

$$(1.7) \quad \sum_{r=0}^k (-1)^r \binom{k}{r} DT_{kn+r}^{(\alpha)}(x) = kx^{k-1} \sum_{r=1}^k (-1)^r \binom{k}{r} T_{kn+r}^{(\alpha)}(x)$$

$$(1.8) \quad T_{kn}^{(\alpha)}(x) = \sum_{r=0}^n \frac{(\alpha-\beta)_r}{r!} T_{kn+r}^{(\beta)}(x)$$

This work of the writer generalizes some properties of the Laguerre polynomials $L_n^{(\alpha)}(x)$. Indeed, when $k=1$, $T_{kn}^{(\alpha)}(x) \equiv L_n^{(\alpha)}(x)$. A similar generalization viz., $T_{kn}^{(0)}(x)$, has been previously studied by Palas [2]. The purpose of this paper is to discuss a more general class of Laguerre polynomials.

2. DEFINITION: We first make the definition

$$(2.1) \quad T_{kn}^{(\alpha)}(x, p) = \frac{1}{n!} x^{-\alpha} e^{px^k} D^n(x^{\alpha+n} e^{-px^k})$$

where k is a natural number.

We now show that the polynomial $T_{kn}^{(\alpha)}(x, p)$ is of exactly degree kn ($n = 0, 1, 2, \dots$). In this connection we know the result [3], for which I must thank Prof. H. W. Gould:

$$(2.2) \quad D_z^k f(z) = \sum_{k=0}^s \frac{(-1)^k}{k!} D_z^k f(z) \sum_{j=0}^k (-1)^j \binom{k}{j} z^{k-j} D_z^j f(z)$$

Thus we obtain from (2.1)

$$T_{kn}^{(\alpha)}(x, p) = \frac{1}{n!} x^{-\alpha} e^{px^k} \sum_{s=0}^n \binom{n}{s} (D^{n-s} x^{\alpha+n}) (D^s e^{-px^k}) =$$

$$\begin{aligned}
&= \frac{1}{n!} e^{px^k} \sum_{s=0}^n \binom{n}{s} \binom{\alpha + n}{n-s} (n-s)! x^s (D^s e^{-px^k}) \\
&= \sum_{s=0}^n \binom{\alpha + n}{n-s} \sum_{i=0}^s \frac{p^i}{i!} x^{ki} \sum_{j=0}^i (-1)^j \binom{i}{j} \binom{kj}{s} \\
&= \sum_{i=0}^n \frac{p^i}{i!} x^{ki} \sum_{j=0}^i (-1)^j \binom{i}{j} \sum_{s=i}^n \binom{\alpha + n}{n-s} \binom{kj}{s}
\end{aligned}$$

Now we know that

$$\sum_{j=0}^i (-1)^j \binom{i}{j} \sum_{s=0}^{i-1} \binom{\alpha + n}{n-s} \binom{kj}{s} = 0.$$

Thus we finally obtain

$$(2.3) \quad T_{kn}^{(\alpha)}(x, p) = \sum_{i=0}^n \frac{p^i}{i!} x^{ki} \sum_{j=0}^i (-1)^j \binom{i}{j} \binom{\alpha + n + kj}{n},$$

which is the explicit formula for $T_{kn}^{(\alpha)}(x, p)$.

In particular, when $k = 1$, and $p = 1$, we derive

$$\begin{aligned}
(2.4) \quad T_n^{(\alpha)}(x, 1) &= \sum_{i=0}^n \frac{x^i}{i!} \sum_{j=0}^i (-1)^j \binom{i}{j} \binom{\alpha + n + j}{n} = \\
&= \sum_{i=0}^n \frac{x^i}{i!} \cdot (-1)^i \binom{\alpha + n}{n-i}
\end{aligned}$$

which is the explicit formula for the general Laguerre polynomials $L_n^{(\alpha)}(x)$. Thus $T_n^{(\alpha)}(x, 1) \equiv L_n^{(\alpha)}(x)$.

3. - OPERATIONAL FORMULAE: Recently we [4] have derived the general operational formula

$$(3.1) \quad x^{-\alpha} D^n (x^{kn+\alpha} Y) = \prod_{j=1}^n \{x^{k-1}(z + \alpha + kj)\} Y, \quad (k = 1, 2, 3, \dots),$$

where $z \equiv x D$ and Y is any sufficiently differentiable function of x . The operators on the right of (3.1) commute only when $k = 1$.

Thus we derive

$$(3.2) \quad x^{-\alpha} e^{px^k} D^n (x^{\alpha+n} e^{-px^k} Y) = \prod_{j=1}^n (xD - pkx^k + \alpha + j) Y$$

Again we observe

$$\begin{aligned} D^n (x^{\alpha+n} e^{-px^k} Y) &= \sum_{r=0}^n \binom{n}{r} D^{n-r} (x^{\alpha+n} e^{-px^k}) D^r Y = \\ &= n! x^{\alpha} e^{-px^k} \sum_{r=0}^n \frac{x^r}{r!} T_{k(n-r)}^{(\alpha+r)}(x, p) D^r Y, \end{aligned}$$

whence we obtain

$$(3.3) \quad \frac{1}{n!} x^{-\alpha} e^{px^k} D^n (x^{\alpha+n} e^{-px^k} Y) = \sum_{r=0}^n \frac{x^r}{r!} T_{k(n-r)}^{(\alpha+r)}(x, p) D^r Y$$

It therefore follows from (3.2) and (3.3) that

$$(3.4) \quad \prod_{j=1}^n (xD - pkx^k + \alpha + j) Y = n! \sum_{r=0}^n \frac{x^r}{r!} T_{k(n-r)}^{(\alpha+r)}(x, p) D^r Y$$

If we set $Y = 1$, we derive from (3.4)

$$(3.5) \quad n! T_{kn}^{(\alpha)}(x, p) = \prod_{j=1}^n (xD - pkx^k + \alpha + j) \cdot 1$$

Further if $k = 1$, and $p = 1$, we obtain from (3.4)

$$(3.6) \quad \prod_{j=1}^n (xD - x + \alpha + j) Y = n! \sum_{r=0}^n \frac{x^r}{r!} T_{n-r}^{(\alpha+r)}(x, 1) D^r Y;$$

which may be compared with the operational formula for the general Laguerre polynomials, derived by Carlitz [5].

In a recent paper [6], Gould and Hopper have generalized the Hermite polynomials by the definition

$$(3.7) \quad H_n^*(x, \alpha, p) = (-1)^n x^{-\alpha} e^{px^k} D^n (x^\alpha e^{-px^k})$$

We remark that $x^n H_n^k(x, \alpha, p)$ yeilds a generalized class of polynomials of exactly degree kn ($n = 0, 1, 2, \dots$), provided k is a natural number. Consequently if we write

$$x^n H_n^k(x, \alpha, p) = H_{kn}^{(\alpha)}(x, p)$$

then

$$(3.8) \quad H_{kn}^{(\alpha)}(x, p) = (-1)^n x^{n-\alpha} e^{px^k} D^n (x^{\alpha} e^{-px^k})$$

Thus the polynomials $H_{kn}^{(\alpha)}(x, p)$ are related to our polynomials by

$$(3.9) \quad H_{kn}^{(\alpha)}(x, p) = (-1)^n n! T_{kn}^{(\alpha-n)}(x, p).$$

Now returning to the operational formula (3.5) we obtain

$$(3.10) \quad (-1)^n H_{kn}^{(\alpha)}(x, p) = \prod_{j=1}^n (xD - pkx^k + \alpha - n + j) \cdot 1$$

More generally we have from (3.4)

$$(3.11) \quad \begin{aligned} \prod_{j=1}^n (xD - pkx^k + \alpha - n + j) Y &= \\ &= \sum_{r=0}^n (-1)^{n-r} \binom{n}{r} x^r H_{k(n-r)}^{(\alpha)}(x, p) D^r Y \end{aligned}$$

This operational formula viz., (3.11) seems to be of particular interest. Indeed, using $k = 2$, $p = 1$, and $\alpha = 0$, we have

$$\begin{aligned} \prod_{j=1}^n (xD - 2x^2 - n + j) Y &= \\ &= \sum_{r=0}^n (-1)^{n-r} \binom{n}{r} x^r H_{2(n-r)}^{(0)}(x, 1) D^r Y \end{aligned}$$

Now noticing that

$$H_{2(n-r)}^{(0)}(x, 1) = x^{n-r} H_{n-r}(x),$$

where $H_n(x)$ denotes the ordinary Hermite polynomials defined by

$$H_n(x) = (-1)^n e^{x^2} D^n e^{-x^2},$$

we obtain

$$\prod_{j=1}^n (xD - 2x^2 - n + j) Y = x^n \sum_{r=0}^n (-1)^{n-r} \binom{n}{r} H_{n-r}(x) D^r Y$$

Now we note that

$$(3.12) \quad x^{-n} \prod_{j=1}^n (xD - 2x^2 - n + j) \equiv (D - 2x)^n.$$

For, (3.12) is evidently true for $n = 1$. Next assume that (3.12) is true for $n = m$. Then we have

$$\begin{aligned} x^{-(m+1)} \prod_{j=1}^{m+1} (xD - 2x^2 - m - 1 + j) &= \\ &= x^{-(m+1)} (xD - 2x^2 - m) \prod_{j=1}^m (xD - 2x^2 - m + j) = \\ &= x^{-(m+1)} (xD - 2x^2 - m) x^m (D - 2x)^m = \\ &= x^{-(m+1)} \cdot x^m (xD - 2x^2) (D - 2x)^m = \\ &= (D - 2x)^{m+1}. \end{aligned}$$

Hence by induction (3.12) is true for all positive integers n . Thus we finally derive

$$(3.13) \quad (D - 2x)^n = \sum_{r=0}^n (-1)^{n-r} \binom{n}{r} H_{n-r}(x) D^r,$$

a formula which Burchnall [7] derived some years ago.

4. - SOME APPLICATIONS OF THE OPERATIONAL FORMULA:
From (3.5) we note that

$$(4.1) \quad n T_n^{(\alpha)}(x, p) = (xD - pkx^k + \alpha + n) T_{k(n-1)}^{(\alpha)}(x, p).$$

In particular, when $k = 1$, and $p = 1$, we derive

$$(4.2) \quad n T_n^{(\alpha)}(x, 1) = (xD - x + \alpha + n) T_{n-1}^{(\alpha)}(x, 1)$$

which is well-known for the Laguerre polynomials $L_n^{(\alpha)}(x)$.

Again in terms of the polynomials of Gould and Hopper, (4.1) stands thus

$$(4.3) \quad H_{kn}^{(\alpha)}(x, p) + (xD - pkx^k + \alpha)H_{k(n-1)}^{(\alpha-1)}(x, p) = 0.$$

Next we consider

$$\begin{aligned} (m+n)! T_{k(m+n)}^{(\alpha)}(x, p) &= \\ &= \prod_{j=1}^m (xD - pkx^k + \alpha + n + j) \prod_{i=1}^n (xD - pkx^k + \alpha + i) \cdot 1 \\ &= n! \prod_{i=1}^n (xD - pkx^k + \alpha + n + i) \cdot T_{kn}^{(\alpha)}(x, p) \\ &= m! n! \sum_{r=0}^m \frac{x^r}{r!} T_{k(m-r)}^{(\alpha+m+r)}(x, p) D^r T_{kn}^{(\alpha)}(x, p); \end{aligned}$$

which implies that

$$\begin{aligned} (4.4) \quad \binom{m+n}{m} T_{k(m+n)}^{(\alpha)}(x, p) &= \\ &= \sum_{r=0}^{\min(m,n)} \frac{x^r}{r!} T_{k(m-r)}^{(\alpha+m+r)}(x, p) D^r T_{kn}^{(\alpha)}(x, p). \end{aligned}$$

The formula (4.4) readily yields the corresponding formula for the polynomials of Gould and Hopper:

$$\begin{aligned} (4.5) \quad H_{k(m+n)}^{(\alpha+m+n)}(x, p) &= \\ &= \sum_{r=0}^{\min(m,n)} (-1)^r \binom{m}{r} x^r H_{k(m-r)}^{(\alpha+m+n)}(x, p) D^r H_{kn}^{(\alpha+n)}(x, p). \end{aligned}$$

5. - GENERATING FUNCTION: We shall now show that the polynomials $T_{kn}^{(\alpha)}(x, p)$ are generated by

$$(5.1) \quad g(x, t) = (1-t)^{-\alpha-1} \exp [px^k u(t)] = \sum_{n=0}^{\infty} T_{kn}^{(\alpha)}(x, p) t^n,$$

where

$$u(t) = 1 - (1-t)^{-k}.$$

From the definition (2.1) we observe

$$(5.2) \quad T_{kn}^{(\alpha)}(x, p) = e^{px^k} \sum_{r=0}^{\infty} \frac{(-p)^r}{r!} \binom{kr + \alpha + n}{n} x^{kr}.$$

It may be noted that (2.3) is a consequence of (5.2).

Now we notice that

$$(5.3) \quad T_{kn}^{(\alpha)}(x, p) = \frac{1}{n!} \left[\frac{d^n}{dt^n} g(x, 0) \right]$$

Also

$$\begin{aligned} (5.4) \quad & \left[\frac{d^n}{dt^n} \{(1-t)^{-\alpha-1} \exp(px^k u(t))\} \right]_{t=0} \\ &= e^{px^k} \left[\frac{d^n}{dt^n} \left\{ (1-t)^{-\alpha-1} \exp \left(-p \left(\frac{x}{1-t} \right)^k \right) \right\} \right]_{t=0} \\ &= n! e^{px^k} \sum_{r=0}^{\infty} \frac{(-p)^r}{r!} \binom{kr + \alpha + n}{n} x^{kr} \end{aligned}$$

Thus a comparison of (5.3) and (5.4) with (5.2) confirms (5.1).

Now from the generating function (5.1) we easily derive the following multiplication formula:

$$(5.5) \quad T_{kn}^{(\alpha)}(xm^{1/k}, p) = T_{kn}^{(\alpha)}(x, mp),$$

which, in terms of the polynomials of Gould and Hopper, shapes into

$$(5.6) \quad H_{kn}^{(\alpha)}(xm^{1/k}, p) = H_{kn}^{(\alpha)}(x, mp),$$

which may well be compared with (3.9) of [6, p. 54].

It is also interesting to note from (5.5) that

$$(5.7) \quad T_n^{(\alpha)}(x, m) = L_n^{(\alpha)}(m.x).$$

Again we observe

$$\begin{aligned} & (1-t)^{-x-1} \exp [px^k \{1 - (1-t)^{-k}\}] \\ &= (1-t)^{-(x-\beta)} (1-t)^{-\beta-1} \exp [px^k \{1 - (1-t)^{-k}\}] \end{aligned}$$

whence we obtain

$$\sum_{n=0}^{\infty} T_{kn}^{(\alpha)}(x, p)t^n = (1-t)^{-(\alpha-\beta)} \sum_{n=0}^{\infty} T_{kn}^{(\beta)}(x, p)t^n.$$

Now comparing the coefficients of t^n on both sides we get

$$(5.8) \quad T_{kn}^{(\alpha)}(x, p) = \sum_{r=0}^n \frac{(\alpha-\beta)_r}{r!} T_{k(n-r)}^{(\beta)}(x, p),$$

where α and β are arbitrary real numbers.

Next we notice that

$$\begin{aligned} & \sum_{n=0}^{\infty} T_{kn}^{(\alpha+\beta+1)}(x, p+q)t^n \\ &= (1-t)^{-\alpha-1} e^{px^k\{1-(1-t)^{-k}\}} \cdot (1-t)^{-\beta-1} e^{qx^k\{1-(1-t)^{-k}\}} \\ &= \sum_{m=0}^{\infty} T_{km}^{(\alpha)}(x, p)t^m \cdot \sum_{n=0}^{\infty} T_{kn}^{(\beta)}(x, q)t^n \\ &= \sum_{n=0}^{\infty} \sum_{m=0}^n T_{km}^{(\alpha)}(x, p) T_{k(n-m)}^{(\beta)}(x, q)t^n. \end{aligned}$$

Thus we obtain the following ‘doubly-additive’ addition formula

$$(5.9) \quad T_{kn}^{(\alpha+\beta+1)}(x, p+q) = \sum_{m=0}^n T_{km}^{(\alpha)}(x, p) T_{k(n-m)}^{(\beta)}(x, q).$$

In particular, when $p = q = 1$, and $k = 1$, we derive

$$(5.10) \quad T_n^{(\alpha+\beta+1)}(x, 2) = \sum_{m=0}^n L_m^{(\alpha)}(x) L_{n-m}^{(\beta)}(x).$$

It follows therefore from (5.7) and (5.10) that

$$(5.11) \quad L_n^{(\alpha+\beta+1)}(2x) = \sum_{m=0}^n L_m^{(\alpha)}(x) L_{n-m}^{(\beta)}(x),$$

which is implied by the well-known formula of the Laguerre polynomials

$$(5.12) \quad L_n^{(\alpha+\beta+1)}(x+y) = \sum_{m=0}^n L_m^{(\alpha)}(x) L_{n-m}^{(\beta)}(y).$$

Again returning to (5.1) we obtain

$$(1-t)^{k+1} \frac{\partial g(x,t)}{\partial t} = [(\alpha+1)(1-t)^k - pkx^k]g(x,t)$$

whence we notice

$$(1-t)^{k+1} \sum_{n=1}^{\infty} nt^{n-1} T_{kn}^{(\alpha)}(x,p) = [(\alpha+1)(1-t)^k - pkx^k] \cdot \sum_{n=0}^{\infty} T_{kn}^{(\alpha)}(x,p)t^n.$$

Performing the indicated multiplication on both sides and comparing coefficients of t^n on both sides, we derive

$$(5.13) \quad \begin{aligned} & \sum_{r=0}^{k+1} (-1)^r \binom{k+1}{r} (n+1-r) T_{k(n+1-r)}^{(\alpha)}(x,p) \\ &= (\alpha+1) \sum_{r=0}^k (-1)^r \binom{k}{r} T_{k(n-r)}^{(\alpha)}(x,p) - pkx^k T_{kn}^{(\alpha)}(x,p). \end{aligned}$$

Lastly we observe

$$(1-t)^k \frac{\partial g(x,t)}{\partial x} = pkx^{k-1}\{(1-t)^k - 1\}g(x,t),$$

whence we obtain in like manner

$$(5.14) \quad \begin{aligned} & \sum_{r=0}^k (-1)^r \binom{k}{r} DT_{k(n-r)}^{(\alpha)}(x,p) \\ &= pkx^{k-1} \sum_{r=1}^k (-1)^r \binom{k}{r} T_{(n-r)}^{(\alpha)}(x,p). \end{aligned}$$

REFERENCES

- [1] S. K. CHATTERJEA: *A generalization of Laguerre polynomials*, Communicated to the Canad. Jour. Math.
- [2] FRANCK J. PALAS: *A Rodrigues' formula*, Amer. Math. Monthly, Vol. 66 (1959), pp. 402-404.
- [3] I. J. SCHWATT: *An introduction to the operations with series*, Philadelphia, 1924.
- [4] S. K. CHATTERJEA: *Operational formulae for certain classical polynomials*, I, to appear in the Quart. Jour. Math. (Oxford).
- [5] L. CARLITZ: *A note on Laguerre polynomials*, Michigan Math. Jour. Vol. 7 (1960), pp. 219-223.
- [6] H. W. GOULD, and A. T. HOPPER: *Operational formulas connected with two generalizations of Hermite polynomials*, Duke Math. Jour. Vol. 29 (1962), pp. 51-64.
- [7] J. L. BURCHNALL: *A note on the polynomials of Hermite*, Quart. Jour. Math. (Oxford), Vol. 12 (1941), pp. 9-11.