

RENDICONTI *del* SEMINARIO MATEMATICO *della* UNIVERSITÀ DI PADOVA

W. W. A. AL-SALAM

Remarks on some operational formulas

Rendiconti del Seminario Matematico della Università di Padova,
tome 35, n° 1 (1965), p. 128-131

<http://www.numdam.org/item?id=RSMUP_1965__35_1_128_0>

© Rendiconti del Seminario Matematico della Università di Padova, 1965, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>*

REMARKS ON SOME OPERATIONAL FORMULAS

*Nota *) di W. W. A. AL-SALAM (a Lubboch)***

Let $y_n(x, a, b)$ be the Bessel polynomial of Krall and Frink defined by means of

$$y_n(x, a, b) = b^{-n} x^{2-a} e^{b/x} D^n (x^{2n+a-2} e^{-b/x}).$$

where $D = d/dx$. Chatterjea proved in [1] the operational formula

$$(1) \quad \prod_{j=1}^n (x^2 D + (2j + a)x + b) = \sum_{r=0}^n \binom{n}{r} b^{n-r} x^{2r} y_{n-r}(x, a + 2r + 2, b) \cdot D^r$$

where the terms of the operator on the left hand side of (1) do not commute among themselves and the operator must be interpreted to mean

$$\left\{ \prod_{j=1}^{n-1} (x^2 D + (2j + a)x + b) \right\} \cdot (x^2 D + (2n + a)x + b).$$

*) Pervenuta in redazione il 21 settembre 1964.

Indirizzo dell'A.: Dept. of Maths. and Astr. Texas Technological College - Lubboch, Texas, USA.

**) Supported by National Science Foundation Grant GP-1593, while the author was at Duke University.

He also gave in [2] the formula

$$(2) \quad x^{2n} \left[D + \frac{2(nx+1)}{x^2} \right]^n = \sum_{r=0}^n \binom{n}{r} 2^{n-r} x^{2r} y_{n-r}(x, 2+2r, 2) D^r.$$

We remark that (2) can be generalized to involve the generalized Bessel polynomial $y_n(x, a, b)$. In fact we prove first the following theorem.

THEOREM 1.

$$(3) \quad x^{2n} \left\{ D + \frac{(a+2n)x+b}{x^2} \right\}^n = \prod_{j=1}^n (x^2 D + (2j+a)x + b).$$

To prove this theorem we first note that

$$(4) \quad \left\{ x^{\lambda} D + g(x) \right\}^n x^v Y = x^v \left\{ x^{\lambda} D + g(x) + \frac{v}{x} \right\}^n Y$$

which can be proved by induction.

It is now easy to prove (3) by induction making use of (4).

Combining (1) and (3) we get

$$(5) \quad x^{2n} \left\{ D + \frac{(a+2n)x+b}{x^2} \right\}^n = \sum_{r=0}^n \binom{n}{r} b^{n-r} x^{2r} y_{n-r}(x, a+2r+2, b) D^r.$$

We next show that (1) and (5) (and hence (2)) are special cases of some results of Gould and Hopper [3].

Let

$$(6) \quad \mathcal{D} = D - prx^{r-1} + \frac{a}{x}$$

and

$$(7) \quad H_n^r(x, a, p) = (-1)^n x^{-a} e^{px^r} D^n (x^a e^{-px^r}).$$

Gould and Hopper proved, among other results,

$$(8) \quad \mathfrak{D}^n = \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} H_{n-k}^r(x, a, p) D^k$$

and

$$(9) \quad x^n \mathfrak{D}^n = \prod_{j=0}^{n-1} (xD - p + x^q - a - j).$$

It is easy to see that (7) and

$$(10) \quad y_n(x, a, b) = b^{-n} x^{2-a} e^{b/x} D^n (x^{2n+a-2} e^{-b/x})$$

imply the relation

$$y_n(x, a, b) = (-1)^n b^{-n} x^{2n} H_n^{-1}(x, an + a - 2, b)$$

so that

$$(11) \quad y_{n-r}(x, a + 2r + 2, b) = (-1)^{n-r} b^{r-n} x^{2n-2r} H_{n-r}^{-1}(x, 2n + a, b).$$

Consequently putting $r = -1$ and $a = 2n + a$ in formula (8) we get

$$\left(D + \frac{(2n + \alpha)x + b}{x^2} \right)^n = \sum_{r=0}^n \binom{n}{r} b^{r-n} x^{2r-2n} y_{n-r}(x, \alpha + 2r + 2, b) D^r$$

which is formula (5). Putting $\alpha = 0$, $b = 2$ we get (2).

Another operational formula similar to (3) is contained in the following theorem.

THEOREM 2.

$$(12) \quad (x^{v+1} D + x^v g(x))^n = x^{nv} \prod_{j=1}^n (xD + g(x) + (n - j)v).$$

Proof. (By induction). Formula (12) is obviously true for $n = 1$. Assume it is true for $n = m$ and consider

$$\begin{aligned}
(x^{v+1}D + x^v g(x))^{m+1} Y &= \\
&= x^v (xD + g(x)) x^{mv} \prod_{j=1}^m (xD + g(x) + (m-j)v) Y = \\
&= x^{(m+1)v} (xD + g(x) + mv) \prod (xD + g(x) + (m-j)v) Y = \\
&= x^{(m+1)v} \prod_{j=0}^n (xD + g(x) + (m-j)v) Y = \\
&= x^{(m+1)v} \prod_{j=0}^{m+1} (xD + g(x) + (m+1-j)v) Y .
\end{aligned}$$

Thus (12) is also true for $n = m + 1$. This completes the proof of (12).

One interesting special case of (12), in terms of the Gould-Hopper operator \mathcal{D} , is obtained by taking $g(x) = a - px^q$ we get

$$(13) \quad (x^{v+1}\mathcal{D})^n = x^{nv} \prod_{j=1}^n (xD - px^r + a + (n-j)v) .$$

which may be compared with (9).

REFERENCE

- [1] CHATTERCEA S. K.: *Operational formulae for certain classical polynomials*. Quarterly Journal of Mathematics, Oxford, (2), vol. 14 (1963), pp. 241-246.
- [2] CHATTERCEA S. K.: *Operational formulae for certain classical polynomials - II*. Rendiconti del Seminario Matematico della Università di Padova, vol. 33 (1963), pp. 163-169.
- [3] GOULD H. W. and HOPPER A. T.: *Operational Formulas connected with two generalizations of Hermite polynomials*. Duke Mathematical Journal, vol. 29 (1962), pp. 51-64.