

RENDICONTI *del* SEMINARIO MATEMATICO *della* UNIVERSITÀ DI PADOVA

JYOTI CHAUDHURI

**On the operational representation of some
hypergeometric polynomials**

Rendiconti del Seminario Matematico della Università di Padova,
tome 38 (1967), p. 27-32

<http://www.numdam.org/item?id=RSMUP_1967__38__27_0>

© Rendiconti del Seminario Matematico della Università di Padova, 1967, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>*

ON THE OPERATIONAL REPRESENTATION OF SOME HYPERGEOMETRIC POLYNOMIALS

JYOTI CHAUDHURI *)

1. The object of the present paper is to obtain operational representation of some hypergeometric polynomials. We incidentally find connections between different types of classical orthogonal polynomials. We have used here the method of p -multiplied Laplace transform. The notations and formulae used are given below.

A function $f(p)$ is said to be the operational representation of a given function $h(x)$ if

$$(1.1) \quad f(p) = p \int_0^\infty e^{-px} h(x) dx, \quad p > 0,$$

provided the integral converges. The relation between the *original* $h(x)$ and its *image* $f(p)$ is denoted by

$$(1.2) \quad h(x) \xrightarrow{\cdot} f(p) \text{ or } f(p) \xleftarrow{\cdot} h(x)$$

$$(1.3) \quad x^n \xrightarrow{\cdot} \frac{\Gamma(n+1)}{p^n}.$$

2. Let $M_n(x)$ denote the polynomial

$$(2.1) \quad {}_{p+1}F_q \left[\begin{matrix} -n, a_1, \dots, a_p; \\ b_1, \dots, b_q; \end{matrix} x \right]$$

*) Indirizzo dell'A : 206/1/D, Bidhan Sarani, Calcutta 6 - India.

and $G_n(x)$ denote the Polynomial

$$(2.2) \quad {}_{p+2}F_q \left[\begin{matrix} -n, a+n, a_1, \dots, a_p; & \frac{1-x}{2} \\ b_1, \dots, b_q; & \end{matrix} \right]$$

Now,

$$x^{n+a-1} M_n(x) = \sum_{r=0}^n \frac{(-n)_r (a_1)_r \dots (a_p)_r}{r! (b_1)_r \dots (b_q)_r} x^{n+a+r-1}$$

Finding the operational representation of the right side term by term,

Left side

$$\begin{aligned} &\stackrel{\cdot}{\rightarrow} \sum_{r=0}^n \frac{(-n)_r (a_1)_r \dots (a_p)_r}{r! (b_1)_r \dots (b_q)_r} \frac{\Gamma(a+n+r)}{p^{n+a+r-1}} \\ &= \frac{\Gamma(n+a)}{p^{n+a-1}} \sum_{r=0}^n \frac{(-n)_r (a_1)_r \dots (a_p)_r}{r! (b_1)_r \dots (b_q)_r} \frac{(a+n)_r}{p^r} \\ &= \frac{\Gamma(n+a)}{p^{n+a-1}} G_n \left(1 - \frac{2}{p} \right). \end{aligned}$$

Therefore,

$$(2.3) \quad x^{n+a-1} M_n(x) \stackrel{\cdot}{\rightarrow} \frac{\Gamma(n+a)}{p^{n+a-1}} G_n \left(1 - \frac{2}{p} \right)$$

Particular cases :

- (i) Putting $p = 0, q = 1, b_1 = 1 + \alpha, a = 1 + \alpha + \beta$ in (2.3) we get

$$(2.4) \quad x^{n+\alpha+\beta} L_n^{(\alpha)}(x) \stackrel{\cdot}{\rightarrow} \frac{\Gamma(n+\alpha+\beta+1)}{p^{n+\alpha+\beta}} P_n^{(\alpha,\beta)} \left(1 - \frac{2}{p} \right)$$

where $L_n^{(\alpha)}(x)$ is the generalised Laguerre polynomial and $P_n^{(\alpha,\beta)}(x)$ is the Jacobi polynomial [[5] p. 62, p. 102.]

- (ii) Putting $\alpha = \beta = \lambda - \frac{1}{2}$ in (2.4) and using the definition of $P_n^\lambda(x)$ [[5], p. 81].

$$(2.5) \quad x^{n+2\lambda-1} L_n^{\left(\lambda-\frac{1}{2}\right)}(x) \stackrel{\cdot}{\rightarrow} \frac{\Gamma(2\lambda) \Gamma\left(n+\lambda+\frac{1}{2}\right)}{\Gamma\left(\lambda+\frac{1}{2}\right) p^{n+2\lambda-1}} P_n^\lambda \left(1 - \frac{2}{p} \right)$$

(iii) Putting $\alpha = \beta = 0$ in (2.4) we get

$$(2.6) \quad x^n L_n(x) \dot{\rightarrow} \frac{n!}{p^n} P_n\left(1 - \frac{2}{p}\right)$$

where $P_n(x)$ is the Legendre polynomial. This is a known operational representation [[2], p. 56].

(iv) Putting $\alpha = \beta = \frac{1}{2}$ in (2.4) we get

$$(2.7) \quad x^{n+\frac{1}{2}} H_{2n+1}(\sqrt{x}) \dot{\rightarrow} (-1)^n \frac{2(2n+1)!}{p^{n+1}} U_n\left(1 - \frac{2}{p}\right)$$

and putting $\alpha = \beta = -\frac{1}{2}$, we get

$$(2.8) \quad x^{n-1} H_{2n}(\sqrt{x}) \dot{\rightarrow} (-1)^n \frac{(2n)!}{np^{n-1}} T_n\left(1 - \frac{2}{p}\right)$$

where $H_n(x)$ is the Hermite polynomial [[5], p. 105] and $T_n(x)$ and $U_n(x)$ are the Tchebichef polynomials of the first and second kind [[5], p. 60].

3. We have

$$(3.1) \quad P_n^{(\alpha, \beta)}(x) = \binom{n+\alpha}{n} \left(\frac{x+1}{2}\right)^n {}_2F_1\left(-n, -n-\beta; \alpha+1; \frac{x-1}{x+1}\right)$$

Putting $\frac{x+1}{x-1} = t$ and multiplying both sides by t^β

$$t^\beta (t-1)^n P_n^{(\alpha, \beta)}\left(\frac{t+1}{t-1}\right) = \binom{n+\alpha}{n} \sum_{r=0}^n \frac{(-n)_r (-n-\beta)_r}{r! (\alpha+1)_r} t^{n+\beta-r}$$

Since

$$(-n-\beta)_r = \frac{(-1)^r \Gamma(n+\beta+1)}{\Gamma(n+\beta-r+1)},$$

the right side

$$\dot{\rightarrow} \frac{\Gamma(n+\beta+1)}{p^{n+\beta}} \binom{n+\alpha}{n} {}_1F_1(-n; \alpha+1; -p),$$

$$(3.2) \quad t^\beta (t-1)^n P_n^{(\alpha, \beta)}\left(\frac{t+1}{t-1}\right) \dot{\rightarrow} \frac{\Gamma(n+\beta+1)}{p^{n+\beta}} L_n^{(\alpha)}(-p)$$

By giving different values to α, β we get Operational representations of different kinds of polynomials.

When $\alpha = \beta = 0$, we get the known operational representation [[3], p. 52].

$$(3.3) \quad (t - 1)^n P_n \left(\frac{t+1}{t-1} \right) \dot{\rightarrow} \frac{n!}{p^n} L_n (-p)$$

4. We have

$$(4.1) \quad P_n^\lambda(x) = \sum_{r=0}^{\left[\frac{1}{2}n\right]} \frac{(-1)^r \Gamma(n-r+\lambda)}{r! \Gamma(\lambda) \Gamma(n-2r+1)} (2x)^{n-2r} \dots \left[\lambda > -\frac{1}{2} \right].$$

Putting $x = \sqrt[p]{p}$ and multiplying by $\frac{1}{p^{\lambda+\frac{1}{2}n-1}}$, we get

$$\frac{\Gamma(\lambda) P_n^\lambda\left(\frac{1}{\sqrt[p]{p}}\right)}{p^{\lambda+\frac{1}{2}n-1}} = \sum_{r=0}^{\left[\frac{1}{2}n\right]} \frac{(-1)^r}{r!} \frac{2^{n-2r}}{(n-2r)!} \frac{\Gamma(n-r+\lambda)}{p^{n-r+\lambda-1}}$$

the right side

$$\begin{aligned} &\stackrel{\leftarrow}{\dot{\rightarrow}} \sum_{r=0}^{\left[\frac{1}{2}n\right]} \frac{(-1)^r}{r!} \frac{2^{n-2r}}{(n-2r)!} x^{n-r+\lambda-1} \\ &= x^{\frac{1}{2}n+\lambda-1} \sum_{r=0}^{\left[\frac{1}{2}n\right]} \frac{(-1)^r}{r!} \frac{(2\sqrt{x})^{n-2r}}{(n-2r)!} \\ &= x^{\frac{1}{2}n+\lambda-1} \frac{H_n(\sqrt{x})}{n!} \end{aligned}$$

Therefore,

$$(4.2) \quad \frac{x^{\frac{1}{2}n+\lambda-1}}{n!} H_n(\sqrt{x}) \dot{\rightarrow} \frac{\Gamma(\lambda) P_n^\lambda\left(\frac{1}{\sqrt[p]{p}}\right)}{p^{\lambda+\frac{1}{2}n-1}}$$

By giving different values to λ we get different operational representations.

5.

$$(5.1) \quad P_n(x) = {}_2F_1\left(-n, n+1; 1; \frac{1-x}{2}\right)$$

Therefore,

$$P_n\left(1 - \frac{2\lambda^2}{p^2}\right) = \sum_{r=0}^n \frac{(-n)_r (n+1)_r}{r! (1)_r} \frac{\lambda^{2r}}{p^{2r}}$$

Since

$$(2r)! = 2^{2r} (1)_r \left(\frac{1}{2}\right)_r,$$

the right side

$$\leftarrow {}_2F_3\left(-n, n+1; 1, 1, \frac{1}{2}; \frac{1}{4} \lambda^2 x^2\right).$$

We know [[1], p. 186]

$$(5.2) \quad {}_1F_1(a; \varrho; z) {}_1F_1(a; \varrho; -z) =$$

$$= {}_2F_3\left[a, \varrho - a; \varrho, \frac{1}{2}\varrho, \frac{1}{2}(\varrho + 1); \frac{1}{4}z^2\right]$$

Putting $a = -n, \varrho = 1, z = \lambda x$, we get

$$(5.3) \quad L_n(\lambda x) L_n(-\lambda x) =$$

$$= {}_2F_3\left[-n, n+1; 1, 1, \frac{1}{2}; \frac{1}{4} \lambda^2 x^2\right] \rightarrow P_n\left(1 - \frac{2\lambda^2}{p^2}\right)$$

This was obtained by Shabde [4] by a different method.

6. From (2.4) and (4.2) we get

$$(6.1) \quad \int_0^\infty e^{-xy} x^{n+\alpha+\beta} L_n^{(\alpha)}(x) dx =$$

$$= \frac{\Gamma(n+\alpha+\beta+1)}{y^{n+\alpha+\beta+1}} P_n^{(\alpha, \beta)}\left(1 - \frac{2}{y}\right), y > 0$$

$$(6.2) \quad \int_0^{\infty} e^{-ax} x^{\frac{1}{2}n+\lambda-1} H_n(\sqrt{a}) dx = \frac{\Gamma(\lambda) \Gamma(n+1)}{a^{\lambda + \frac{1}{2}n+1}} P_n^{\lambda}\left(\frac{1}{\sqrt{a}}\right), a > 0$$

In conclusion I would like to express my gratitude to Prof. B. N. Mukherjee for his kind help and guidance in the preparation of this paper.

REFERENCES

1. A. ERDELYI. - Higher Transcendental Functions, vol. 1.
2. N. W. McLACHLAN and P. HUMBERT. - Formulaire pour le Calcul Symbolique.
3. N. W. McLACHLAN, P. HUMBERT and L. POLI - Supplément au Formulaire pour le Calcul Symbolique.
4. N. G. SHABDE - Bulletin of the Calcutta Mathematical Society, 34 (1942).
5. G. SZEGO - Orthogonal Polynomials, (1959).

Manoscritto pervenuto in redazione il 15 gennaio 1966.