

RENDICONTI
del
SEMINARIO MATEMATICO
della
UNIVERSITÀ DI PADOVA

HERBERT DIETZSCH

**On the computation of the order of Janko's
first simple group \mathfrak{J}_1**

Rendiconti del Seminario Matematico della Università di Padova,
tome 46 (1971), p. 371-374

<http://www.numdam.org/item?id=RSMUP_1971__46__371_0>

© Rendiconti del Seminario Matematico della Università di Padova, 1971, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

ON THE COMPUTATION OF THE ORDER
OF JANKO'S FIRST SIMPLE GROUP \mathfrak{F}_1

HERBERT DIETZSCH *)

Janko's computation of the order of \mathfrak{F}_1 contains a mistake as the generalized character φ'_4 of [1; p. 156] does not vanish on non special classes of $N(U)$ in case (i) of [1; Lemma 3.1]. The purpose of this note is to correct this mistake.

We use terminology and notation of [1].

Case (i) of Lemma 3.1.

Here $N(U) \subset C(t)$. Thus, $T = \langle t, t_1 \rangle$ is a S_2 -subgroup of $N(U)$, where t_1 is an involution of F which acts invertingly on U .

We have $U\langle t_1 \rangle \times \langle t \rangle = N(U)$.

The group $N(U)/\langle t \rangle$ has two linear characters: ξ'_0 (the principal character), β' , and two irreducible characters ξ'_1, ξ'_2 of degree 2.

Let ξ' be the nontrivial linear character of $N(U)/(N(U) \cap F)$. Put $U = \langle \mu \rangle$. We get the following character-table of $N(U)$.

Let the set of « special classes » of $N(U)$ in the sense of Suzuki [2] consist of the conjugate classes of $\mu, \mu^2, t\mu, t\mu^2$ in $N(U)$.

The module (over the ring of integers) of the generalized characters of $N(U)$ vanishing on non special classes of $N(U)$ has the following basis:

$$\begin{aligned}\varphi'_1 &= \xi'_2 - \xi'_1 \\ \varphi'_2 &= \xi' \cdot \xi'_2 - \xi' \cdot \xi'_1 \\ \varphi'_3 &= \xi'_0 + \beta' - \xi'_1 \\ \varphi'_4 &= \xi' + \xi' \cdot \beta' - \xi' \cdot \xi'_1.\end{aligned}$$

*) Indirizzo dell'A.: Mathematisches Institut der Universität, 65 Mainz, Germania Occ.

	ξ'_0	β'	ξ'_1	ξ'_2	ξ'	$\xi' \cdot \beta'$	$\xi' \cdot \xi'_1$	$\xi' \cdot \xi'_2$
1	1	1	2	2	1	1	2	2
μ	1	1	$\frac{1+\sqrt{5}}{2}$	$\frac{1-\sqrt{5}}{2}$	1	1	$\frac{1+\sqrt{5}}{2}$	$\frac{1-\sqrt{5}}{2}$
μ^2	1	1	$\frac{1-\sqrt{5}}{2}$	$\frac{1+\sqrt{5}}{2}$	1	1	$\frac{1-\sqrt{5}}{2}$	$\frac{1+\sqrt{5}}{2}$
t_1	1	-1	0	0	1	-1	0	0
t	1	1	2	2	-1	-1	-2	-2
$t\mu$	1	1	$\frac{1+\sqrt{5}}{2}$	$\frac{1-\sqrt{5}}{2}$	-1	-1	$\frac{1+\sqrt{5}}{2}$	$\frac{1-\sqrt{5}}{2}$
$t\mu^2$	1	1	$\frac{1-\sqrt{5}}{2}$	$\frac{1+\sqrt{5}}{2}$	-1	-1	$\frac{1-\sqrt{5}}{2}$	$\frac{1+\sqrt{5}}{2}$
tt_1	1	-1	0	0	-1	1	0	0

We have the following decomposition of the induced characters φ_i^* :

$$\varphi_1^* = \overline{\varepsilon_1} \overline{x_1} + \overline{\varepsilon_2} \overline{x_2}$$

$$\varphi_2^* = \overline{\eta_1} \overline{y_1} + \overline{\eta_2} \overline{y_2}$$

$$\varphi_3^* = 1_G + \overline{\varepsilon_1} \overline{x_1} + \overline{\zeta_2} \overline{z_2}$$

$$\varphi_4^* = \overline{\eta_1} \overline{y_1} + \overline{\theta_2} \overline{v_2} + \overline{\theta_3} \overline{v_3}$$

where \bar{x}_i , \bar{y}_i , \bar{z}_2 , \bar{v}_i are nontrivial, different irreducible characters of G and $\bar{\varepsilon}_i$, $\bar{\eta}_i$, $\bar{\zeta}_2$, $\bar{\theta}_i$ are all ± 1 .

If χ is an irreducible character of G which appears in φ_i^* with the multiplicity n_i ($1 \leq i \leq 4$), then

$$\chi(\sigma) = n_1 \bar{\xi}'_2(\sigma) + (n_2 - n_4) \bar{\xi}'(\sigma) \cdot \bar{\xi}'_2(\sigma) + n_3 - n_4 \bar{\xi}'(\sigma) \cdot \bar{\xi}'_1(\sigma)$$

where σ is an arbitrary element of the set of « special classes » of $N(U)$.

Hence we get:

$$\bar{x}_1(t\mu) = \frac{1}{2} \bar{\varepsilon}_1(1 + \sqrt{5}) \text{ and } \bar{z}_2(t\mu) = \bar{\zeta}_2$$

$$\bar{x}_1(t\mu^2) = \frac{1}{2} \bar{\varepsilon}_1(1 - \sqrt{5}) \text{ and } \bar{z}_2(t\mu^2) = \bar{\zeta}_2.$$

In Case (1) of [1; p. 155] we get $\bar{x}_1 \in \{y_1, y_2, y_3, y_4\}$ and $\bar{z}_2 \in \{x_3, v_1, v_2\}$. Since $0 = \varphi_3^*(t)$, we get $\bar{x}_1 \in \{y_1, y_2, y_3, y_4\}$ and $\bar{z}_2 \in \{v_1, v_2\}$.

In Case (2) of [1; p. 156] we get $\bar{x}_1 \in \{x_1, x_2, y_3, y_4\}$ and $\bar{z}_2 \in \{z_4, v_1, v_2\}$. Since $0 = \varphi_3^*(t)$, we get $\bar{x}_1 \in \{y_3, y_4\}$ and $\bar{z}_2 \in \{v_1, v_2\}$.

Therefore, in both cases we get: $\bar{x}_1(t) = \pm 3$ and $\bar{z}_2(t) = \pm 4$. We shall apply the Suzuki order formula of [2] for the generalized characters $\varphi'_3 = \bar{\xi}'_0 + \beta' - \bar{\xi}'_1$, $\varphi_3^* = 1_G + \bar{\varepsilon}_1 \bar{x}_1 + \bar{\zeta}_2 \bar{z}_2$ and the subgroup $N(U)$ of G .

Put $f = \bar{\varepsilon}_1 \bar{x}_1(1)$, thus $f+1 = -\bar{\zeta}_2 \bar{z}_2(1)$ as $0 = \varphi_3^*(1)$.

Denoting $|G|$ by g we obtain

$$\frac{1}{g} \left\{ \frac{g^2}{120^2} \left(1 + \frac{3^2}{f} - \frac{4^2}{f+1} \right) \right\} = \frac{1}{20} \left\{ \frac{11^2}{1} + \frac{9^2}{1} - \frac{2^2}{2} \right\} = 10.$$

Hence, $g \cdot (f-3)^2 = 2^7 \cdot 3^2 \cdot 5^3 \cdot f \cdot (f+1)$.

We know that both f and $f+1$ divide g and that f and $f+1$ are coprime. This implies that $f-3$ is a divisor of $2^3 \cdot 3 \cdot 5$.

On the other hand a S_2 -subgroup of G has order 8 and U is a S_5 -subgroup of G . This implies $2^2 \cdot 5$ divides $f-3$.

Therefore we have the following possibilities:

$$f-3 = \pm 20, \pm 40, \pm 60, \pm 120.$$

It is easy to see that all eight possibilities lead to a contradiction.
For example

$$f-3=20 \text{ implies } g \cdot 20^2 = 23 \cdot 24 \cdot 2^7 \cdot 3^2 \cdot 5^3$$

which is impossible as 7 divides g .

Similarly $f-3=-60$ implies $g=2^6 \cdot 3 \cdot 5 \cdot 7 \cdot 19$ which is against the order of a S_2 -subgroup of G .

Thus, we have ruled out Case (i) of Lemma 3.1.

REFERENCES

- [1] JANKO, Z.: *A new finite simple group with abelian Sylow-2-subgroups and its characterization*, Journal of Algebra 3 (2) (1966), 147-186.
- [2] SUZUKI, M.: *Applications of group characters*, Proc. Symp. Pure Math. 1 (1959), 88-89.

I am very grateful to Dr. DIETER HELD for suggesting this problem to me.

Manoscritto pervenuto in redazione il 10 Luglio 1971.