

RENDICONTI
del
SEMINARIO MATEMATICO
della
UNIVERSITÀ DI PADOVA

LARRY D. SHATOFF

Binary multiples of combinatorial geometries - II

Rendiconti del Seminario Matematico della Università di Padova,
tome 49 (1973), p. 237-240

http://www.numdam.org/item?id=RSMUP_1973__49__237_0

© Rendiconti del Seminario Matematico della Università di Padova, 1973, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques*
<http://www.numdam.org/>

Binary Multiples of Combinatorial Geometries - II (*).

LARRY D. SHATOFF (**)

1. Introduction.

In [2] we considered the sequence of multiples $G, 2G, \dots, mG$ of a projective geometry, where m is the smallest integer such that mG is a Boolean algebra. We showed that the only multiples that were binary were possibly G itself, mG , and $(m-1)G$ if $(n^{a-1} + n^{a-2} + \dots + n)/q$ is an integer (where G has order n and rank q). In this paper we prove similar results about affine geometries. All definitions not given here can be found in [2] and we will assume the reader is familiar with that paper.

2. Affine geometries.

Given an affine geometry on the set of points S , the lattice of flats (points, lines, etc.) is a geometric lattice, and thus defines a combinatorial geometry. We will refer to these combinatorial geometries as *affine geometries*. Recall that if f is a submodular function on a set S , then it defines a pregeometry on S as follows: $A \subseteq S$ is independent if and only if $f(A') \geq |A'|$ for all non-empty $A' \subseteq A$. [1, Prop. 7.3]. (Here $|A'|$ denotes the cardinality of A' .) If $\mathcal{G}(S)$ is a combinatorial geometry with rank function r , and k is a positive integer, then kr is submodular and thus defines a new combinatorial geometry, de-

(*) This work was supported in part by the Colgate University Research Council.

(**) Indirizzo dell'A.: Colgate University - Hamilton, N. Y., U.S.A.

noted kG . If G is an affine geometry, when are the multiples of G binary geometries? We show that kG is binary only when $k = 1$ and G is binary, or when kG is a Boolean truncation of rank one less than the cardinality of S , or when kG is a Boolean algebra. We prove this result first for affine planes and then use induction for the general case. We will make use of the following well-known results on affine geometries.

THEOREM 1. *Suppose $G(S)$ is an affine geometry. There exists an integer $n \geq 2$ such that:*

- (i) *Every flat of rank q contains exactly n^{q-1} points;*
- (ii) *Every coline of G is covered by exactly $n + 1$ copoints;*
- (iii) *There is a set of n copoints of G which partition S .*

The number n is the order of G . Notice that the rank of a flat is one more than its dimension in the affine geometry. (Thus an affine plane has rank 3.)

THEOREM 2. *If G is an affine geometry of order n and rank $q \geq 4$, and C is a copoint of G , then the subgeometry of G on the set C is an affine geometry of order n and rank $q - 1$.*

We now prove the theorem for affine planes.

THEOREM 3. *Let $G(S)$ be an affine plane of order n . kG is binary if and only if $k \geq (n^2 - 1)/3$.*

PROOF. We first consider the case $n = 2$. By Theorem 1 and [2; Theorem 1], this is the only case in which G is binary. kG , $k \geq 2$, is the Boolean algebra on S , and so kG is binary if and only if $k \geq (n^2 - 1)/3 = 1$. Now let $n \geq 3$. Since any line of G is an n -set (i.e., a set of cardinality n), kG is a Boolean truncation if $k \geq n/2$. By [2; Lemma 1] kG is binary if and only if every set of $|S| - 1 = n^2 - 1$ elements is k -independent. This is true if and only if $k \geq (n^2 - 1)/3$. Suppose $1 < k < n/2$. If $A \subseteq S$ is a $3k$ -set no $(2k + 1)$ -subset of which is contained in a line of G , then A is a basis for kG . We construct a $(3k + 2)$ -set, every $3k$ -subset of which is a basis for kG . By [2; Lemma 2] and [2; Theorem 1], this will show kG is not binary. Let C_1, C_2 be two non-intersecting lines of G . Let A consist of any $2k$ elements of C_1 and any $k + 2$ elements of C_2 . This is possible as $k + 2 \leq 2k < n$. To show any $3k$ -subset of A is a basis for kG , let $B \subseteq A$ be a $(2k + 1)$ -set. We must show that B is not contained in any line of G . If B were contained in a line of G , then $|B \cap C_1| \leq 1$.

For if $|B \cap C_1| > 1$, we would have two lines intersecting in more than one point. Similarly, $|B \cap C_2| \leq 1$. This means $|B| \leq 2$. But $2k + 1 \geq 3$, and so it is impossible for B to be contained in a line. This completes the proof.

LEMMA 1. *If n and q are positive integers such that $n \geq 2$ and $q \geq 4$, then*

$$\frac{n^{q-2} - 1}{q - 1} \geq \frac{n^{q-3}}{q - 2}$$

and

$$\frac{n^{q-2}}{q - 1} > \frac{n^{q-3}}{q - 2}.$$

THEOREM 4. *Let $G(S)$ be an affine geometry of order n and rank q . kG is binary if and only if $k \geq (n^{q-1} - 1)/q$ or both $k = 1$ and $n = 2$.*

PROOF. G is binary if and only if $n = 2$, thus we may assume $k > 1$. Because of the arithmetic details, we first note that the theorem holds where $n = 2, q = 4$. It is easy to see that $kG, k \geq 2$, is a Boolean algebra; and so kG is binary if and only if $k \geq (n^{q-1} - 1)/q = 1 \frac{3}{4}$ or $k = 1$. We now prove the theorem by induction on q . If $n \neq 2, q = 3$ is the first step of the induction. The result is then that of Theorem 3. If $n = 2$, then the case of $q = 3$ is given by Theorem 3, and we let $q = 4$ be the first step of the induction. Assume the result holds for affine geometries of order n and rank $q - 1, q \geq 4$ ($q \geq 5$ if $n = 2$). We show it holds for G . Let G' be the affine geometry of order n , rank $q - 1$ on a copoint of G (Theorem 2). kG' is a subgeometry of kG by [2; Lemma 5] and so since kG is binary we conclude that kG' is binary. By the induction hypothesis, $k \geq (n^{q-2} - 1)/(q - 1)$. Now if $k \geq n^{q-2}/(q - 1)$, then every set of n^{q-2} elements is k -independent. For if $|A| = n^{q-2}$, then $r(A) \geq q - 1$, and so $kr(A) \geq k(q - 1) \geq |A|$. If $B \subseteq A, kr(B) \geq |B|$, for

$$\frac{|B|}{r(B)} \leq \frac{|\bar{B}|}{r(\bar{B})} \leq \frac{n^m}{m + 1} \leq \frac{n^{q-2}}{q - 1} \leq k$$

(for $m = r(\bar{B}) - 1$) by Lemma 1. This means copoints of G are k -independent (i.e., independent in kG), so kG is a Boolean truncation. Thus, by [2; Lemma 1], kG is binary if and only if every $(n^{q-1} - 1)$ -set is k -independent, that is, if and only if $k \geq (n^{q-1} - 1)/q$. The only case left to consider is $k = (n^{q-2} - 1)/(q - 1)$. In this case, $k > n^{q-3}/(q - 2)$ by Lemma 1, and so by an argument like that above,

any (n^{q-3}) -set is k -independent. In fact, any $[(q-1)k]$ -set is k -independent; for if A is a subset of such a set, and if $n^{q-3} < |A| \leq k(q-1)$, then $r(A) \geq q-1$ and so $kr(A) \geq k(q-1) = n^{q-2} - 1 \geq |A|$. A copoint (which is a $[(q-1)k+1]$ -set) however is not k -independent. We conclude that if A is a qk -set no $[(q-1)k+1]$ -subset of which is contained in a copoint of G , then A is a basis for kG . We construct a $(qk+2)$ -set every qk -subset of which is a basis for kG . This, by [2; Theorem 1 and Lemma 2], will show that kG is not binary. Let C_1, C_2 be any two non-intersecting copoints of G (Theorem 1). Let A consist of any $(q-1)k$ points of C_1 and any $k+2$ points of C_2 . This is possible, as $k+2 \leq (q-1)k < n^{q-2}$. $|A| = qk+2$. To show every qk -subset of A is a basis of kG , let $B \subseteq A$ be a $[(q-1)k+1]$ -set. We show B is contained in no copoint of G . If B is contained in a copoint, then $|B \cap C_1| \leq n^{q-3}$. Otherwise we would have two copoints which intersect in more than n^{q-3} points. This is impossible by Theorem 1. Similarly $|B \cap C_2| \leq n^{q-3}$. Thus $|B| \leq 2n^{q-3} < n^{q-2}$ if $n \neq 2$. Thus it is impossible to find B unless $n = 2$. If $n = 2$, we must pick B so that $|B \cap C_1| = |B \cap C_2| = n^{q-3}$. But $|A \cap C_2| = k+2$, and it is easy to show $n^{q-3} > k+2$ for $n = 2, q \geq 5$. Since $B \cap C_2 \subseteq A \cap C_2$, this shows it is impossible to find a B in this case also. Thus the desired set A does exist, and the proof is complete.

Let G be an affine geometry with sequence of multiples $G, 2G, \dots, mG$, m the smallest integer such that mG is a Boolean algebra. mG is always binary. G is binary; its order is 2. Just as for projective geometries, we have shown that the only other multiple of an affine geometry that may be binary is $(m-1)G$. $(m-1)G$ is actually binary if and only if $k = (n^{q-1} - 1)/q$ is an integer. In this case $m-1 = k$. If n and q are relatively prime and q is a prime, it is well-known that $n^{q-1} \equiv 1 \pmod{q}$. Thus in this case $(m-1)G$ is binary.

REFERENCES

- [1] H. H. CRAPO - G.-C. ROTA, *On the Foundations of Combinatorial Theory*, preliminary edition, M.I.T. Press, 1970.
- [2] L. D. SHATOFF, *Binary multiples of combinatorial geometries*, Rendiconti del Seminario Matematico dell'Università di Padova, Vol. 48.

Manoscritto pervenuto in redazione il 1° settembre 1972.