

RENDICONTI *del* SEMINARIO MATEMATICO *della* UNIVERSITÀ DI PADOVA

GIOVANNI PORRU

θ -mappings and quasiconformal mappings in normed spaces

Rendiconti del Seminario Matematico della Università di Padova,
tome 57 (1977), p. 173-182

http://www.numdam.org/item?id=RSMUP_1977__57__173_0

© Rendiconti del Seminario Matematico della Università di Padova, 1977, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

θ -Mappings and quasiconformal mappings in normed spaces

GIOVANNI PORRU (*)

Introduction.

The θ -mappings introduced by Gehring in [2], and some definitions of quasiconformal mappings in R^n , are meaningful in every normed space. But such mappings have been investigated, till now, almost exclusively in R^n . Indeed, most of the procedures used in R^n (as Lebesgue integral) are meaningless in an infinite dimensional space H .

In this paper we consider θ -mappings in a real normed space, and we prove for them a compactness theorem. Further, we show some properties of sequences of θ -mappings. Next we consider the metric definition of quasiconformal mappings in a real normed space, and we give a relation between the θ -mappings and the quasiconformal mappings.

1. θ -MAPPINGS. Let H be a real normed space and Ω, Ω' domains of H . Following Gehring ([2], pag. 14) we give the

DEFINITION 1. *A homeomorphism T_X of Ω onto Ω' is said to be a θ -mapping if there exists a function $\theta(t)$, which is continuous and increasing in $0 \leq t < 1$ with $\theta(0) = 0$, such that the following are true.*

(*) Indirizzo dell'A.: Istituto Matematico Università di Cagliari - 09100 Cagliari, Italy.

Lavoro eseguito nell'ambito del GNAFA del CNR.

(i) If $X_0 \in \Omega$ and $\|X - X_0\| < \inf_{Y \in \partial\Omega} \|X_0 - Y\| \equiv \varrho(X_0, \partial\Omega)$, then

$$\frac{\|TX - TX_0\|}{\varrho(TX_0, \partial\Omega')} \leq \theta \left(\frac{\|X_0 - X\|}{\varrho(X_0, \partial\Omega)} \right)$$

(ii) The restriction of TX to any subdomain $\Delta \subseteq \Omega$, satisfies (i).

For the θ -mappings we now prove some properties.

THEOREM 1. Suppose that $\{T_n X\}$ is a sequence of θ -mappings (with the same distortion function $\theta(t)$) of Ω , that

$$\lim_{n \rightarrow \infty} T_n X = TX$$

uniformly on each bounded domain $\Delta \subset \Omega$ such that $\varrho(\Delta, \partial\Omega) > 0$, and that TX is a homeomorphism. Then TX is a θ -mapping.

PROOF. Let $\Delta \subset \Omega$ be a bounded domain such that $\varrho(\Delta, \partial\Omega) > 0$, and set $\Delta'_n = T_n(\Delta)$, $\Delta' = T(\Delta)$. Since $T_n X$ are θ -mappings we have, for $X_0 \in \Delta$:

$$(1) \quad \frac{\|T_n X - T_n X_0\|}{\varrho(T_n X_0, \partial\Delta'_n)} \leq \theta \left(\frac{\|X - X_0\|}{\varrho(X_0, \partial\Delta)} \right), \quad n = 1, 2, \dots$$

for $\|X - X_0\| < \varrho(X_0, \partial\Delta)$. We have to show that

$$(2) \quad \frac{\|TX - TX_0\|}{\varrho(TX_0, \partial\Delta')} \leq \theta \left(\frac{\|X - X_0\|}{\varrho(X_0, \partial\Delta)} \right)$$

for $\|X - X_0\| < \varrho(X_0, \partial\Delta)$. First we prove that

$$(3) \quad \liminf_{n \rightarrow \infty} \varrho(T_n X_0, \partial\Delta'_n) \leq \varrho(TX_0, \partial\Delta').$$

We may suppose $\varrho(TX_0, \partial\Delta') < \infty$. If (3) is not true there exists a subsequence $\{T_{n_k} X\}$ and a positive real number σ such that

$$(4) \quad \lim_{k \rightarrow \infty} \varrho(T_{n_k} X_0, \partial\Delta'_{n_k}) \geq \varrho(TX_0, \partial\Delta') + \sigma.$$

From (4) it follows that there exists a finite ν such that

$$(5) \quad \varrho(TX_0, \partial\Delta'_{n_k}) > \varrho(TX_0, \partial\Delta') + \sigma/2$$

for $k > \nu$. By (5) we may find at least one point $X' \in \partial\Delta'$ and a ball $B(X', r')$ centered in X' , radius r' , and contained in Δ'_{n_k} , for $k > \nu$. Further we may take a second ball $B(Z', r) \subset B(X', r')$ with $B(Z', r) \cap \Delta' = \emptyset$. Let Z_k be the points of Δ such that $T_{n_k}Z_k = Z'$. Since $\{T_{n_k}X\}$ converges uniformly to TX on Δ , we have

$$\|TZ_k - Z'\| = \|TZ_k - T_{n_k}Z_k\| < r$$

for $k > \nu_1$, where $\nu_1 \geq \nu$ is a suitable finite number. The latter inequality shows that TZ_k belongs to $B(Z', r)$. Hence, we obtain a contradiction and inequality (3) does hold.

Since

$$\lim_{n \rightarrow \infty} \|T_n X - T_n X_0\| = \|TX - TX_0\|,$$

from (1) and (3), we derive (2).

Let now Δ be a subdomain of Ω . If $X_0 \in \Delta$ and $\varrho(X_0, \partial\Delta) = \infty$, let $\Delta_m \equiv B(X_0, m)$. For the previous result we have, for $\|X - X_0\| < m$:

$$\frac{\|TX - TX_0\|}{\varrho(TX_0, \partial\Delta'_m)} \leq \theta\left(\frac{\|X - X_0\|}{\varrho(X_0, \partial\Delta_m)}\right) = \theta\left(\frac{\|X - X_0\|}{m}\right),$$

where $\Delta'_m = T(\Delta_m)$. Since $\lim_{m \rightarrow \infty} \theta(\|X - X_0\|/m) = 0$ it must be

$$\lim_{m \rightarrow \infty} \varrho(TX_0, \partial\Delta'_m) = \infty. \text{ As } \varrho(TX_0, \partial\Delta') \geq \varrho(TX_0, \partial\Delta'_m)$$

it must be again $\varrho(TX_0, \partial\Delta') = \infty$, and inequality (2) holds for such Δ .

Finally suppose $\Delta \subseteq \Omega$, $X_0 \in \Delta$ and such that $\varrho(X_0, \partial\Delta) = c < \infty$. Let $\Delta_m = B(X_0, c - 1/m)$ with $m > 1/c$. Inequality (2) holds for Δ_m . Further we have $\Delta' = T(\Delta) \supset T(\Delta_m) = \Delta'_m$, hence

$$\varrho(TX_0, \partial\Delta') \geq \varrho(TX_0, \partial\Delta'_m), \quad \varrho(X_0, \partial\Delta_m) = \varrho(X_0, \partial\Delta) - 1/m$$

and

$$\frac{\|TX - TX_0\|}{\varrho(TX_0, \partial\Delta')} \leq \frac{\|TX - TX_0\|}{\varrho(TX_0, \partial\Delta'_m)} \leq \theta \left(\frac{\|X - X_0\|}{\varrho(X_0, \partial\Delta_m)} \right) = \theta \left(\frac{\|X - X_0\|}{\varrho(X_0, \partial\Delta)^{-\frac{1}{m}}} \right).$$

For $m \rightarrow \infty$, because $\theta(t)$ is continuous, we obtain inequality (2) for such Δ .

THEOREM 2. *Suppose that $\{T_n X\}$ is a sequence of θ -mappings of Ω and that*

$$\lim_{n \rightarrow \infty} T_n X = TX, \quad \|TX\| < \infty$$

in Ω . Then TX is either a constant or a continuous and injective mapping.

In order to prove the Theorem we give some lemmas whose proofs are similar to those done by Gehring for $H = R^n$ ([2]).

LEMMA 1. *Suppose that $\{T_n X\}$ is a sequence of θ -mappings of Ω which are locally uniformly bounded on Ω . Then $T_n X$ are locally equicontinuous in Ω .*

PROOF. Let $Z \in \Omega$ and let $B(Z, r)$ be a ball centered in Z , radius $r < \varrho(Z, \partial\Omega)$ and such that $T_n X$ are uniformly bounded in $B(Z, r)$. Choose $X_0 \in \Omega - \bar{B}(Z, r)$ and let $\Delta = \Omega - \{X_0\}$. By hypothesis there exists a finite constant A (depending on $B(Z, r)$ and X_0) such that

$$\|T_n X - T_n X_0\| \leq \|T_n X\| + \|T_n X_0\| \leq A$$

for $X \in B(Z, r)$ and all n . Hence

$$(6) \quad \varrho(T_n X, \partial\Delta'_n) \leq \|T_n X - T_n X_0\| \leq A$$

for $X \in B(Z, r)$ and all n , where $\Delta'_n = T_n(\Delta)$. Since $r < \varrho(Z, \partial\Omega)$ and $X_0 \notin \bar{B}(Z, r)$,

$$(7) \quad \varrho(X, \partial\Delta) \geq a > 0$$

for $X \in B(Z, r)$. Now fix $X \in B(Z, r)$ and choose Y so that $\|Y -$

$\|X\| < a$. Then

$$\frac{\|T_n Y - T_n X\|}{\varrho(T_n X, \partial \Delta'_n)} \leq \theta \left(\frac{\|Y - X\|}{\varrho(X, \partial \Delta)} \right).$$

and, for (6) – (7),

$$\|T_n Y - T_n X\| \leq A \theta \left(\frac{\|Y - X\|}{a} \right).$$

Since $\lim_{t \rightarrow 0^+} \theta(t) = 0$, this implies the desired equicontinuity of $T_n X$ in $B(Z, r)$.

LEMMA 2. Suppose that $\{T_n X\}$ is a sequence of θ -mappings of Ω onto Ω'_n , that

$$\sup_n \|T_n X_0\| < \infty$$

for some $X_0 \in \Omega$, and that

$$\sup_n \varrho(0, \partial \Omega'_n) < \infty$$

where 0 denotes the null element of H . Then $T_n X$ are locally equicontinuous in Ω .

PROOF. Fix a so that $0 < a < 1$. Then, if we choose $Y \in \Omega$ and X so that $\|X - Y\| < a\varrho(Y, \partial \Omega)$ we have

$$\|T_n X - T_n Y\| \leq \theta(a) \varrho(T_n Y, \partial \Omega'_n)$$

for all n . Since

$$\varrho(T_n Y, \partial \Omega'_n) \leq \|T_n Y\| + \varrho(0, \partial \Omega'_n)$$

we thus obtain

$$(8) \quad \|T_n X\| \leq M \|T_n Y\| + N$$

where

$$M = 1 + \theta(a), \quad N = \theta(a) \sup_n \varrho(0, \partial \Omega'_n)$$

In particular we conclude that each point $Y \in \Omega$ has a neighborhood $U = U(Y) \subset \Omega$ such that (8) holds for all $X \in U$. Next, if we choose $X \in \Omega$ and Y so that $\|Y - X\| < \frac{a}{2} \varrho(X, \partial\Omega)$ we have

$$\|Y - X\| < \frac{a}{2} (\varrho(Y, \partial\Omega) + \|Y - X\|)$$

and thus

$$\|Y - X\| < \frac{a}{2 - a} \varrho(Y, \partial\Omega) < a\varrho(Y, \partial\Omega).$$

Hence we see that each point $X \in \Omega$ has a neighborhood $V = V(X) \subset \Omega$ such that (8) holds for all $Y \in V$. Now let G denote the set of points $X \in \Omega$ for which

$$\sup_n \|T_n X\| < \infty.$$

If $Y \in G$ and if U is the neighborhood described above, then (8) implies that

$$\sup_n \|T_n X\| \leq M \sup_n \|T_n Y\| + N$$

for all $X \in U$. Hence $U \subset G$ and G is open. Similarly, if $X \in \Omega - G$ and if V is the neighborhood described above, the same argument shows that $V \subset \Omega - G$, and hence that $\Omega - G$ is open. Since Ω is connected and $X_0 \in G$, we conclude that $G = \Omega$. We have shown that for $Y \in G$, $\sup_n \|T_n Y\| < \infty$. Further, by (8), follows that $T_n X$ are uniformly bounded in a suitable neighborhood $U(Y)$. The locally equicontinuity is now a consequence of Lemma 1.

LEMMA 3. *Suppose that $\{T_n X\}$ is a sequence of θ -mappings of Ω , and that*

$$\sup_n \|T_n X_0\| < \infty, \quad \sup_n \|T_n X_1\| < \infty$$

for a pair of distinct fixed points $X_0, X_1 \in \Omega$. Then $T_n X$ are locally equicontinuous in Ω .

PROOF. Let $\Delta = \Omega - \{X_1\}$ and $\Delta'_n = T_n(\Delta)$. Then $T_n X_1 \in \partial \Delta'_n$ and hence

$$\sup_n \varrho(0, \partial \Delta'_n) \leq \sup_n \|T_n X_1\| < \infty.$$

Lemma 2 implies now the locally equicontinuity of $T_n X$ in Δ . Interchanging the roles of X_0 and X_1 we obtain the desired conclusion.

LEMMA 4. *Suppose that $\{T_n X\}$ is a sequence of θ -mappings of Ω , that*

$$\lim_{n \rightarrow \infty} T_n X = TX, \quad \|TX\| < \infty$$

in Ω , that $T_n X \neq Y'_n$ in Ω , and that

$$\lim_{n \rightarrow \infty} Y'_n = Y'.$$

Then either $TX \neq Y'$ in Ω or $TX \equiv Y'$ in Ω .

PROOF. Let G be the set of points $X \in \Omega$ for which $TX = Y'$. Lemma 3 implies that $T_n X$ are locally equicontinuous in Ω . Hence TX is continuous and G is closed in Ω . Now suppose $X_0 \in G$ and let U be the set of points X for which $\|X - X_0\| < a\varrho(X_0, \partial\Omega)$, where a is some fixed constant, $0 < a < 1$. Then we have

$$\|T_n X - T_n X_0\| \leq \theta(a)\varrho(T_n X_0, \partial\Omega'_n)$$

for all $X \in U$, where $\Omega'_n = T_n(\Omega)$. Since $Y'_n \notin \Omega'_n$, we see that

$$\varrho(T_n X_0, \partial\Omega'_n) \leq \|T_n X_0 - Y'_n\|$$

and hence

$$\|TX - TX_0\| = \lim_{n \rightarrow \infty} \|T_n X_0 - T_n X\| \leq \theta(a) \lim_{n \rightarrow \infty} \|T_n X_0 - Y'_n\| = 0$$

for all $X \in U$. Hence $U \subset G$ and G is open. Since Ω is connected, we conclude, as desired, that either $G = \Omega$ or $TX \equiv Y'$ in Ω .

PROOF OF THEOREM 2. Lemma 3 implies that $T_n X$ are locally equicontinuous in Ω , hence TX is continuous. If TX is not one to

one, we can find a pair of distinct points $X_0, X_1 \in \Omega$, such that $TX_0 = TX_1 = Y'$. Let $\Delta = \Omega - \{X_1\}$ and $T_n X_1 = Y'_n$. Then $T_n X \neq Y'_n$ in Δ and

$$\lim_{n \rightarrow +\infty} Y'_n = Y'.$$

Since $X_0 \in \Delta$ and $TX_0 = Y'$, Lemma 4 implies that $TX = Y'$ in Δ and hence that TX is constant in Ω .

2. QUASICONFORMAL MAPPINGS. Let H be a real normed space, Ω, Ω' domains of H and TX a homeomorphism of Ω onto Ω' . For $X_0 \in \Omega$ and $0 < r < \rho(X_0, \partial\Omega)$ we set

$$L(X_0, r) = \sup_{\|X - X_0\| = r} \|TX - TX_0\|$$

$$l(X_0, r) = \inf_{\|X - X_0\| = r} \|TX - TX_0\|$$

$$\delta(X_0) = \limsup_{r \rightarrow 0^+} \frac{L(X_0, r)}{l(X_0, r)}$$

According to Gehring metric definition of quasiconformality for $H = R^n$, we give the following

DEFINITION 2. A homeomorphism TX of a domain Ω onto a domain Ω' is said to be K -quasiconformal if $\delta(X_0) \leq K$ for every point X_0 of Ω . TX is quasiconformal if it is K -quasiconformal for some K (finite).

If $H = R^n$, a homeomorphism is quasiconformal if and only if it is a θ -mapping ([2]). Now we show the following

THEOREM 3. A homeomorphism TX of Ω onto Ω' is quasiconformal if $T^{-1}X$ is a θ -mapping.

PROOF. Let $X_0 \in \Omega$ and c , $0 < c < \rho(X_0, \partial\Omega)$, so that

$$(9) \quad \|TX - TX_0\| < \rho(TX_0, \partial\Omega')$$

for $\|X - X_0\| < c$. Next, for each r , $0 < r < c$ and $\varepsilon > 0$, choose X_1 and X_2 so that $\|X_1 - X_0\| = \|X_2 - X_0\| = r$ and so that

$$(10) \quad \begin{cases} L(X_0, r) = \sup_{\|X - X_0\|=r} \|TX - TX_0\| < \|TX_1 - TX_0\| + \varepsilon \\ 1(X_0, r) = \inf_{\|X - X_0\|=r} \|TX - TX_0\| > \|TX_2 - TX_0\| - \varepsilon \end{cases}$$

If for some r and for each ε is

$$(11) \quad 1(X_0, r) + \varepsilon > L(X_0, r) - \varepsilon$$

then we have $1(X_0, r) = L(X_0, r)$ for such r . If r is such that (11) is not true we have, for $0 < \varepsilon < \varepsilon_0$, ε_0 suitable,

$$(12) \quad 1(X_0, r) + \varepsilon < L(X_0, r) - \varepsilon.$$

Let $\Delta = \Omega - \{X_1\}$ and $\Delta' = T(\Delta)$. Then $\Delta' = \Omega' - \{TX_1\}$. From (9), (10) and (12) we obtain

$$\|TX_2 - TX_0\| < 1(X_0, r) + \varepsilon < L(X_0, r) - \varepsilon < \|TX_1 - TX_0\| = \varrho(TX_0, \partial\Delta').$$

Since $T^{-1}X$ is a θ -mapping we have

$$1 = \frac{\|T^{-1}(TX_2) - T^{-1}(TX_0)\|}{\varrho(T^{-1}(TX_0), \partial\Delta)} \leq \theta \left(\frac{\|TX_2 - TX_0\|}{\varrho(Tx_0, \partial\Delta')} \right) < \theta \left(\frac{1(X_0, r) + \varepsilon}{L(X_0, r) - \varepsilon} \right)$$

and, as $\theta(t)$ is increasing,

$$\frac{L(X_0, r) - \varepsilon}{1(X_0, r) + \varepsilon} < \frac{1}{\theta^{-1}(1)}$$

For $\varepsilon \rightarrow 0^+$ we obtain

$$\frac{L(X_0, r)}{1(X_0, r)} \leq \frac{1}{\theta^{-1}(1)}$$

and also

$$\limsup_{r \rightarrow 0^+} \frac{L(X_0, r)}{1(X_0, r)} \leq \frac{1}{\theta^{-1}(1)} = K$$

As the latter inequality holds for each point X_0 of Ω , the theorem is proved.

Theorem 3 can be improved if TX is a diffeomorphism, that is a homeomorphism of Ω onto Ω' , such that TX and $T^{-1}X$ are differentiable. For this we have:

LEMMA 5. *Suppose that TX is a quasiconformal diffeomorphism of Ω onto Ω' . Then the diffeomorphism $T^{-1}X$ of Ω' onto Ω is quasiconformal.*

PROOF. Let $X_0 \in \Omega$. If A denotes the Fréchet derivative of TX in X_0 , we have ([3]), for some constant K ,

$$(13) \quad 0 < \sup_{Z \neq 0} \frac{\|AZ\|}{\|Z\|} \leq K \inf_{Z \neq 0} \frac{\|AZ\|}{\|Z\|}.$$

Since the derivative of $T^{-1}X$ in TX_0 is A^{-1} , (13) implies

$$(14) \quad 0 < \sup_{Y \neq 0} \frac{\|A^{-1}Y\|}{\|Y\|} \leq K \inf_{Y \neq 0} \frac{\|A^{-1}Y\|}{\|Y\|}.$$

If X_0 ranges onto Ω , then TX_0 ranges onto Ω' , and (14) does hold in every point of Ω' . Thus ([3]) $T^{-1}X$ is quasiconformal.

Theorem 3 and Lemma 5 give the following.

COROLLARY. *A diffeomorphism TX of Ω onto Ω' is quasiconformal if TX is a θ -mapping.*

BIBLIOGRAPHY

- [1] P. CARAMAN, *N-dimensional quasiconformal mappings*, Ed. Ac. Bucuresti and Abacus Press Tunbridge Wells, Kent, England (1974).
- [2] W. F. GEHRING, *The Carathéodory convergence theorem for quasiconformal mappings in space*, Ann. Acad. Sci. Fenn. Ser. AI 336/11 (1963).
- [3] G. PORRU, *Quasiconformal mappings in normed spaces*, In print by the Proceedings of the 3rd Romanian-Finnish Seminar on Complex Analysis (Bucharest, 28/6-2/7 1976).

Manoscritto pervenuto in redazione il 28 marzo 1977.