

# RENDICONTI *del* SEMINARIO MATEMATICO *della* UNIVERSITÀ DI PADOVA

LUIGI SALCE

## **The $\lambda$ -inductive topology on abelian $p$ -groups**

*Rendiconti del Seminario Matematico della Università di Padova*,  
tome 59 (1978), p. 167-177

[http://www.numdam.org/item?id=RSMUP\\_1978\\_\\_59\\_\\_167\\_0](http://www.numdam.org/item?id=RSMUP_1978__59__167_0)

© Rendiconti del Seminario Matematico della Università di Padova, 1978, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques*  
<http://www.numdam.org/>

## The $\lambda$ -Inductive Topology on Abelian $p$ -Groups (\*).

LUIGI SALCE (\*)

### 1. Introduction.

Let  $G$  be an abelian  $p$ -group. The  $p$ -adic topology on  $G$  has the family of subgroups  $\{p^n G\}_{n \in \mathbb{N}}$  as a basis of neighborhoods of 0. The Hausdorff completion  $L_\omega(G) = \varprojlim G/p^n G$  of  $G$  in this topology is not a  $p$ -group if  $G$  is unbounded; the maximal torsion subgroup of  $L_\omega(G)$ , which is a  $p$ -group, is denoted by  $T_\omega(G)$  and is called the torsion completion of  $G$ ;  $G$  is said torsion complete if the canonical homomorphism  $\delta: G \rightarrow L_\omega(G)$  sends  $G$  isomorphically upon  $T_\omega(G)$ . The class of torsion complete  $p$ -groups is one of the best known in the theory of abelian  $p$ -groups (see [4, Chap. XI]).

B. Charles made in 1967 the important observation that the torsion completion  $T_\omega(G)$  of  $G$  can be realized as the completion of  $G$  in a suitable topology, which is the inductive topology; a basis of neighborhoods of 0 for this topology is the family of large subgroups, introduced by Pierce [11], which are those fully invariant subgroups  $L$  of  $G$  such that  $G = L + B$  for every basic subgroup  $B$  of  $G$ ; for the proof of this fact see [2] or [4, 70.2]. Large subgroups have a nice description: in fact Pierce proved ([11]; see also [4, 67.2]) that, if  $G$  is unbounded, every large subgroup  $L$  of  $G$  can be realized as  $L = G(\mathbf{u})$ , where  $\mathbf{u} = (r_n)_{n \in \mathbb{N}}$  is an increasing sequence of integers, and  $G(\mathbf{u}) = \{x \in G \mid h_a(p^n x) \geq r_n, \forall n \in \mathbb{N}\}$  ( $h_a$  denotes the height in  $G$ ); note that

(\*) Indirizzo dell'A.: Seminario Matematico, Università - Via Belzoni 7 - 35100 Padova.

Lavoro eseguito nell'ambito dei Gruppi di Ricerca Matematica del Consiglio Nazionale delle Ricerche.

the subgroups  $G(\mathbf{u})$  described above are a natural generalization of the subgroups  $p^n G$  ( $n \in \mathbf{N}$ ).

The generalization of the  $p$ -adic topology from the ordinal  $\omega$  to an arbitrary limit ordinal  $\lambda$  is the  $\lambda$ -adic topology, which has as a basis of neighborhoods of 0 the family of subgroups  $\{p^\sigma G\}_{\sigma < \lambda}$ ; this topology has been studied by Mines [10]. The Hausdorff completion of  $G$  in this topology and its torsion part are denoted respectively by  $L_\lambda(G)$  and  $T_\lambda(G)$ ; if  $\lambda$  is cofinal with  $\omega$  (we shall write  $\lambda = \text{cf } \omega$ ), then, if  $\delta_\lambda: G \rightarrow L_\lambda(G)$  is the canonical homomorphism,  $L_\lambda(G)/\delta_\lambda(G)$  is divisible and the topology of the completion on  $L_\lambda(G)$  coincides with the  $\lambda$ -adic topology; if  $\lambda$  is not cofinal with  $\omega$  (we shall write  $\lambda \neq \text{cf } \omega$ ), then  $L_\lambda(G) = T_\lambda(G)$  and in general  $L_\lambda(G)/\delta_\lambda(G)$  is not divisible, and the two above topologies do not coincide. The generalization of torsion complete groups is given by those  $p$ -groups  $G$  such that  $\delta_\lambda$  sends  $G$  isomorphically upon  $T_\lambda(G)$ ; these groups coincide, if  $\lambda = \text{cf } \omega$ , with the  $p^\lambda$ -high injective groups, which have been studied by Megibben [9] and Dubois [3].

It is natural to try to generalize the inductive topology from the ordinal  $\omega$  to an arbitrary limit ordinal  $\lambda$ ; recall that, given an increasing sequence of ordinals and symbols  $\infty: \mathbf{u} = (\sigma_n)_{n \in \mathbf{N}}$  ( $\sigma < \infty$  for each ordinal  $\sigma$  and  $\infty < \infty$ ), the subgroup of  $G$

$$G(\mathbf{u}) = \{x \in G \mid h_\sigma(p^n x) \geq \sigma_n, \forall n \in \mathbf{N}\}$$

is a fully invariant subgroup of  $G$ . We define the  $\lambda$ -inductive topology on  $G$  by means of a basis  $\mathcal{B}_\lambda$  of neighborhoods of 0:  $\mathcal{B}_\lambda$  consists of those subgroups  $G(\mathbf{u})$  defined above such that  $\sigma_n < \lambda$  for every  $n \in \mathbf{N}$ . In section 2 we shall study some properties of the  $\lambda$ -inductive topology and we shall prove the analogue of Charle's result, i.e. that the completion  $L_\lambda(G)$  of  $G$  in the  $\lambda$ -inductive topology is canonically isomorphic to  $T_\lambda(G)$ ; it follows that, if  $\lambda = \text{cf } \omega$ , a reduced  $p$ -group is  $p^\lambda$ -high injective if and only if it is complete in the  $\lambda$ -inductive topology.

A more natural definition of  $\lambda$ -inductive topology would get as a basis of neighborhoods of 0 the family of  $\lambda$ -large subgroups of  $G$ , i.e. those fully invariant subgroups of  $G$  such that  $G = L + B$  for every  $\lambda$ -basic subgroup  $B$  of  $G$ ; this definition makes sense only for those  $p$ -groups which have  $\lambda$ -basic subgroups; in the non trivial cases, these groups are, as Wallace proved in [12], the  $C_\lambda$ -groups with  $\lambda = \text{cf } \omega$  (for the definitions of  $\lambda$ -basic subgroup and  $C_\lambda$ -group see Megibben [8] and Wallace [12]). In section 3 we shall prove that for a  $C_\lambda$ -group,

with  $\lambda = \text{cf } \omega$ , the  $\lambda$ -inductive topology has as a basis of neighborhoods of 0 the family of  $\lambda$ -large subgroups, by generalizing the result of Pierce which characterize large subgroups.

**2. The  $\lambda$ -inductive topology.**

All groups considered are abelian groups. Notation and terminology do not depart from those in [4]. In particular, if  $G$  is a reduced  $p$ -group,  $l(G)$  denotes the length of  $G$ , i.e. the minimal ordinal  $\sigma$  such that  $p^\sigma G = 0$ ; to every element  $x \in G$  is associated an increasing sequence of ordinals and  $\infty$ 's:  $H(x) = \{h_\sigma(p^n x) | n \in \mathbb{N}\}$ , which is said the indicator of  $x$ . Let  $\lambda$  be a limit ordinal; let  $\mathcal{B}_\lambda$  be the family of subgroups  $G(\mathbf{u})$  of  $G$ , where  $\mathbf{u} = (\sigma_n)_{n \in \mathbb{N}}$ , such that  $\sigma_n < \lambda$  for every  $n \in \mathbb{N}$ ; if  $\mathbf{u}$  and  $\mathbf{v}$  are two such sequences of ordinals, the equality  $G(\mathbf{u}) \cap G(\mathbf{v}) = G(\mathbf{u} \vee \mathbf{v})$ , where  $\mathbf{u} \vee \mathbf{v}$  denotes the pointwise supremum, ensures that  $\mathcal{B}_\lambda$  is a basis of neighborhoods of 0 for a topology, which is said the  $\lambda$ -inductive topology. We shall consider now some properties of the  $\lambda$ -inductive topology.

1) Every  $G(\mathbf{u}) \in \mathcal{B}_\lambda$  is closed in the  $\lambda$ -adic topology.

It follows from the easily proved equality:  $G(\mathbf{u}) = \bigcap_{\tau < \lambda} (p^\tau G + G(\mathbf{u}))$ .

2)  $G$  is discrete in the  $\lambda$ -inductive topology if and only if  $l(G) < \lambda$  and it is Hausdorff if and only if  $l(G) \leq \lambda$ .

$G$  is discrete if and only if there exists  $G(\mathbf{u}) \in \mathcal{B}_\lambda$  such that  $G(\mathbf{u}) = 0$ ; this implies that  $p_{\sigma_0} G[p] = 0$ , where  $\sigma_0$  is the first ordinal of  $\mathbf{u}$ , hence  $p^{\sigma_0} G = 0$  and  $l(G) \leq \sigma_0 < \lambda$ . The viceversa is obvious. The second claim follows from the following property.

3)  $p^\lambda G = \bigcap \{G(\mathbf{u}) | G(\mathbf{u}) \in \mathcal{B}_\lambda\}$ .

If  $\mathbf{u} = (\sigma_n)_{n \in \mathbb{N}}$  with  $\sigma_n < \lambda$  for every  $n \in \mathbb{N}$ , then obviously  $p^\lambda G \subseteq G(\mathbf{u})$ , hence  $p^\lambda G \subseteq \bigcap \{G(\mathbf{u}) | G(\mathbf{u}) \in \mathcal{B}_\lambda\}$ ; the opposite inclusion follows from the fact that, if  $\sigma < \lambda$ , then  $p^\sigma G \in \mathcal{B}_\lambda$ : in fact  $p^\sigma G = G(\sigma, \sigma + 1, \sigma + 2, \dots)$ .

4) The  $\lambda$ -adic topology is coarser than the  $\lambda$ -inductive topology; the two topologies coincide if  $\lambda \neq \text{cf } \omega$ .

We have already seen that, if  $\sigma < \lambda$ , then  $p^\sigma G \in \mathcal{B}_\lambda$ ; if  $\lambda \neq \text{cf } \omega$ , given  $G(\mathbf{u}) \in \mathcal{B}_\lambda$  let  $\sigma = \sup \{\sigma_n | n \in \mathbb{N}\}$ ; then  $\sigma < \lambda$  and  $G(\mathbf{u}) \supseteq p^\sigma G$  imply that  $G(\mathbf{u})$  is open in the  $\lambda$ -adic topology.

The above property, with the following theorem 2.1, clarify the role of the cofinality with  $\omega$  of the ordinal  $\lambda$  in the  $\lambda$ -adic topology. Let now  $A$  be a subgroup of the reduced  $p$ -group  $G$ . We say that  $A$  satisfies the  $\lambda$ -th *Pierce's condition* if, for every  $n \in \mathbb{N}$ , there exists an ordinal  $\tau_n < \lambda$  such that

$$(1) \quad p^{\tau_n} G[p^n] \subseteq A .$$

5) *A subgroup  $A$  of  $G$  is open in the  $\lambda$ -inductive topology if and only if it satisfies the  $\lambda$ -th Pierce's condition.*

If  $A$  is open there exists  $G(\mathbf{u}) \in \mathfrak{B}_\lambda$ ,  $\mathbf{u} = (\sigma_n)_{n \in \mathbb{N}}$ , such that  $G(\mathbf{u}) \subseteq A$ ; then  $A$  satisfies (1): just take  $\tau_n = \sigma_n$ . Viceversa, let  $\sigma_n = \tau_{n+1} + n$  for each  $n \in \mathbb{N}$ . If  $\mathbf{u} = (\sigma_n)_{n \in \mathbb{N}}$  and  $x \in G(\mathbf{u}) \cap G[p]$ , then  $px = 0$  and  $h_\alpha(x) \geq \sigma_0 = \tau_1$  imply  $x \in A$ ; now we suppose by induction that, given  $k \geq 1$ ,  $G(\mathbf{u}) \cap G[p^{k-1}] \subseteq A$  and let  $p^k x = 0$ ,  $x \in G(\mathbf{u})$ ; then  $h_\alpha(p^{k-1}x) \geq \sigma_{k-1} = \tau_k + k - 1$ , therefore there exists  $y \in G$  such that  $p^{k-1}x = p^{k-1}y$  and  $h_\alpha(y) \geq \tau_k$ ; it follows that  $y \in A$  and  $h_\alpha(p^n(x-y)) \geq \inf \{ \sigma_n, \tau_k + n \} = \sigma_n$  for  $n = 0, 1, \dots, k-2$ ; hence  $x-y \in G(\mathbf{u}) \cap G[p^{k-1}] \subseteq A$  and  $x \in A$ .

6) *The  $\lambda$ -inductive topology on  $G/p^\lambda G$  coincides with the quotient topology of the  $\lambda$ -inductive topology on  $G$ .*

It follows easily from the equality  $G(\mathbf{u})/p^\lambda G = (G/p^\lambda G)(\mathbf{u})$ , which holds if  $\mathbf{u} = (\sigma_n)_{n \in \mathbb{N}}$  with  $\sigma_n < \lambda$  for every  $n \in \mathbb{N}$ .

Let now  $I_\lambda(G)$  be the Hausdorff completion of  $G$  in the  $\lambda$ -inductive topology; if  $l(G) < \lambda$ , then  $I_\lambda(G) = G$ ; if  $l(G) > \lambda$ , from property 6 it follows that  $I_\lambda(G) = I_\lambda(G/p^\lambda G)$ ; therefore we can suppose  $l(G) = \lambda$ . Recall that  $I_\lambda(G) = \varprojlim_{G(\mathbf{u}) \in \mathfrak{B}_\lambda} G/G(\mathbf{u})$  and  $L_\lambda(G) = \varprojlim_{\sigma < \lambda} G/p^\sigma G$ ; but  $p^\sigma G \in \mathfrak{B}_\lambda$  if  $\sigma < \lambda$ , therefore we can define a homomorphism

$$\varphi: I_\lambda(G) \rightarrow L_\lambda(G)$$

in the following way: if  $(g_u + G(\mathbf{u}))_u \in I_\lambda(G)$ , we consider only the cosets modulo the subgroups  $p^\sigma G$ , where  $\sigma < \lambda$ , denote them by  $g_\sigma + p^\sigma G$  and put

$$(2) \quad \varphi((g_u + G(\mathbf{u}))_u) = (g_\sigma + p^\sigma G)_\sigma;$$

with this notation we can prove the following theorem.

**THEOREM 2.1.** *The homomorphism  $\varphi$  sends  $I_\lambda(G)$  isomorphically upon  $T_\lambda(G)$ .*

**PROOF.** We prove first that, if in (2)  $g_\sigma \in p^\sigma G$  for each  $\sigma < \lambda$  and if  $G(\mathbf{u}) \in \mathcal{B}_\lambda$ , then  $g_{\mathbf{u}} \in G(\mathbf{u})$ , hence  $\varphi$  is injective. Let  $\tau$  be an arbitrary ordinal less than  $\lambda$  and  $\mathbf{v} = p^\tau G = (\tau, \tau + 1, \tau + 2, \dots)$ ; put  $\mathbf{w} = \mathbf{u} \vee \mathbf{v}$ ; then  $g_{\mathbf{w}} - g_{\mathbf{v}} \in p^\tau G$ , therefore  $g_{\mathbf{w}} \in p^\tau G$ ; it follows that  $g_{\mathbf{u}} \in p^\tau G + G(\mathbf{u})$ , for every  $\tau < \lambda$ ; by property 1)  $g_{\mathbf{u}} \in G(\mathbf{u})$ . We shall prove now that  $I_\lambda(G)$  is a  $p$ -group; let exist, by way of contradiction,  $(g_{\mathbf{u}} + G(\mathbf{u}))_{\mathbf{u}} \in I_\lambda(G)$  such that, for a suitable increasing sequence of indexes:  $\mathbf{u}_1 < \mathbf{u}_2 < \mathbf{u}_3 < \dots$  the orders  $p^{m_k}$  of  $g_{\mathbf{u}_k} + G(\mathbf{u}_k)$  give an increasing sequence of positive integers; we can suppose that  $m_{k+1} \geq m_k + k$  for every  $k \in \mathbb{N}$ . Let  $\mathbf{u}_k$  be fixed; then  $p^{m_k-1}g_{\mathbf{u}_k} + G(\mathbf{u}_k)$  has order  $p$  and, if  $G(\mathbf{u}) \subseteq G(\mathbf{u}_k)$ , then  $p^{m_k-1}g_{\mathbf{u}} + G(\mathbf{u}_k)$  has also order  $p$ . Then  $p^{m_k-1}g_{\mathbf{u}_k} \notin G(\mathbf{u}_k)$ , therefore, by property 1), there exists  $\tau_k < \lambda$  such that

$$\tau_k = \sup \{h_\sigma(p^{m_k-1}g_{\mathbf{u}}) | \mathbf{u} \geq \mathbf{u}_k, G(\mathbf{u}) \in \mathcal{B}_\lambda\}.$$

For every  $k \in \mathbb{N}$  let  $\varrho_k$  be ordinals such that  $\tau_{k+1} < \varrho_k < \lambda$  and  $\varrho_k < \varrho_{k+1}$ ; consider now  $G(\mathbf{v}) \in \mathcal{B}_\lambda$  where  $\mathbf{v} = (\varrho_n)_{n \in \mathbb{N}}$ . If  $g_{\mathbf{v}} + G(\mathbf{v})$  has order  $p^n$ , then  $p^n g_{\mathbf{u}} \in G(\mathbf{v})$  for every  $\mathbf{u} \geq \mathbf{v}$  with  $G(\mathbf{u}) \in \mathcal{B}_\lambda$ . Let  $r \in \mathbb{N}$  be such that  $m_r \geq n$ . If  $\mathbf{u} \geq \mathbf{v} \vee \mathbf{u}_{k+1}$  and  $G(\mathbf{u}) \in \mathcal{B}_\lambda$ , then  $p^{m_r}g_{\mathbf{u}} \in G(\mathbf{v})$  implies

$$h_\sigma(p^{m_{r+1}-1}g_{\mathbf{u}}) \geq \varrho_{m_{r+1}-m_r-1};$$

but  $m_{r+1} - m_r - 1 \geq r$  implies  $\varrho_{m_{r+1}-m_r-1} \geq \varrho_r > \tau_{r+1}$  and  $\mathbf{u} \geq \mathbf{u}_{r+1}$  implies

$$h_\sigma(p^{m_{r+1}-1}g_{\mathbf{u}}) \leq \tau_{r+1}$$

which is absurd. From the above proof we get the inclusion  $\varphi(I_\lambda(G)) \subseteq T_\lambda(G)$ ; let now  $(g'_\sigma + p^\sigma G)_\sigma \in T_\lambda(G)$  be an element of order  $m$ ; if  $G(\mathbf{u}) \in \mathcal{B}_\lambda$  let  $g_{\mathbf{u}} = g'_{\sigma_m}$ ; it is easy to see that  $\varphi((g_{\mathbf{u}} + G(\mathbf{u}))_{\mathbf{u}}) = (g'_\sigma + p^\sigma G)_\sigma$  and that  $(g_{\mathbf{u}} + G(\mathbf{u}))_{\mathbf{u}} \in I_\lambda(G)$ , therefore the image of  $\varphi$  is the whole of  $T_\lambda(G)$ .

An obvious consequence of theorem 2.1 is the following

**COROLLARY 2.2.** *A reduced  $p$ -group  $G$  is complete in the  $\lambda$ -inductive topology if and only if  $p^\lambda G = 0$  and  $\delta_\lambda(G) = T_\lambda(G)$ .*

By property 4) we are interested in the  $\lambda$ -inductive topology on a reduced  $p$ -group only if  $\lambda = \text{cf } \omega$ ; in this case the  $\lambda$ -adic topology

on  $I_\lambda(G) \cong T_\lambda(G)$  coincides with the topology of the completion of  $G$  endowed with the  $\lambda$ -adic topology, hence the subgroups  $p^\sigma I_\lambda(G)$  with  $\sigma < \lambda$ , are open in the topology of the completion of  $G$  endowed with the  $\lambda$ -inductive topology; it follows that, if  $G(\mathbf{u}) \in \mathcal{B}_\lambda$ , then  $I_\lambda(G)(\mathbf{u})$  is closed in the above topology, hence it contains the closure of  $\delta_\lambda(G(\mathbf{u}))$  which is open; therefore  $I_\lambda(G)(\mathbf{u})$  is open in the topology of the completion of  $G$  endowed with the  $\lambda$ -inductive topology. A subbase of neighborhoods of 0 for this topology is given by the subgroups  $I_\lambda(G) \cap \text{Ker } \pi_{\mathbf{u}}$ , where  $G(\mathbf{u}) \in \mathcal{B}_\lambda$  and

$$\pi_{\mathbf{u}}: \prod_{G(\mathbf{v}) \in \mathcal{B}_\lambda} G/G(\mathbf{v}) \rightarrow G/G(\mathbf{u}),$$

is the canonical surjection; we shall prove that  $I_\lambda(G)(\mathbf{u})$  is contained in  $I_\lambda(G) \cap \text{Ker } \pi_{\mathbf{u}}$ ; let  $(g_{\mathbf{v}} + G(\mathbf{v}))_{\mathbf{v}} \in I_\lambda(G)(\mathbf{u}) \subseteq \prod_{G(\mathbf{v}) \in \mathcal{B}_\lambda} (G/G(\mathbf{v}))(\mathbf{u})$ , let  $r$  be the order of  $g_{\mathbf{u}}$  and, if  $\mathbf{u} = (\sigma_n)_{n \in \mathbb{N}}$ , put  $\mathbf{w} = (\sigma_r, \sigma_{r+1}, \sigma_{r+2}, \dots)$ . If  $\sigma < \sigma_r$  we have;  $p^\sigma(G/G(\mathbf{w})) = p^\sigma G/G(\mathbf{w})$ , therefore, for  $i = 0, 1, \dots, \dots, r-1$ , the following inequalities hold:  $h_G(p^i g_{\mathbf{w}}) \geq \sigma_i$ . But  $\mathbf{w} \geq \mathbf{u}$  implies  $g_{\mathbf{w}} - g_{\mathbf{u}} \in G(\mathbf{u})$ , hence  $h_G(p^i g_{\mathbf{u}}) \geq \sigma_i$  for  $i = 0, 1, \dots, r-1$ , which means that  $g_{\mathbf{u}} \in G(\mathbf{u})$ . Thus we have proved the following

**THEOREM 2.3.** *Let  $G$  be a  $p$ -group and  $\lambda$  a limit ordinal cofinal with  $\omega$ . Then the topology of the completion (of  $G$  endowed with the  $\lambda$ -inductive topology) on  $I_\lambda(G)$  coincides with its own  $\lambda$ -inductive topology.*

**3.  $\lambda$ -large subgroups of  $C_\lambda$ -groups.**

A  $p$ -group  $G$  is a  $C_\lambda$ -group, where  $\lambda$  is a limit ordinal, if  $G/p^\sigma G$  is totally projective for every ordinal  $\sigma < \lambda$ . The  $C_\lambda$ -groups have been introduced by Megibben [8] with the restriction on  $\lambda$  to be countable; Wallace studied [12] the general case; in particular, he proved that a reduced  $p$ -group  $G$  of length  $\geq \lambda$  contains a proper  $\lambda$ -basic subgroup  $B$  if and only if  $G$  is a  $C_\lambda$ -group and  $\lambda = \text{cf } \omega$  (recall that  $B$  is a  $\lambda$ -basic subgroup of  $G$  if it is totally projective of length  $\leq \lambda$  and isotype and dense in the  $\lambda$ -adic topology in  $G$ ). We are interested on groups with proper  $\lambda$ -basic subgroups, therefore from now on  $G$  will denote a  $C_\lambda$ -group with  $\lambda = \text{cf } \omega$ . A subgroup  $L$  of  $G$  is said  $\lambda$ -large if it is fully invariant and  $G = L + B$  for every  $\lambda$ -basic subgroup  $B$  of  $G$ ; this definition has been introduced by Linton (see [6]). If  $l(G) < \lambda$ , then

the unique  $\lambda$ -basic subgroup of  $G$  is  $G$  itself and every fully invariant subgroup is  $\lambda$ -large; to exclude this trivial case, we suppose that  $l(G) \geq \lambda$ . If  $L$  is  $\lambda$ -large in  $G$ , then  $p^n L$  is also  $\lambda$ -large for every  $n \in \mathbb{N}$ .

LEMMA 3.1. *Let  $L$  be a  $\lambda$ -large subgroup of the  $C_\lambda$ -group  $G$ , with  $\lambda = \text{cf } \omega$ . Then  $p^\lambda G$  is contained in  $L$ .*

PROOF. We can suppose  $l(G) > \lambda$ . Let  $x \in p^\lambda G$ ; if  $G = L + B$  for a  $\lambda$ -basic subgroup  $B$  of  $G$ , then  $x = b + y$ , with  $b \in B$ ,  $y \in L$ . There exists a direct summand  $B'$  of  $B$  such that  $b \in B'$  and  $l(B') < \lambda$ . By [1, 2.3] we have:  $G = B' \oplus G'$ ; if  $\pi$  is the projection of  $G$  upon  $G'$  with respect to this decomposition, then  $\pi(x) = \pi(y) \in L$  because  $L$  is fully invariant, and  $(1 - \pi)(x) = 0$ , because  $h_a(x) = h_B(x) \geq \lambda$ ; it follows that  $x = \pi(y) \in L$ .

We will need in the following some results on  $C_\lambda$ -groups.

LEMMA 3.2. *Let  $G$  be a  $C_\lambda$ -group,  $\lambda = \text{cf } \omega$ ; let  $x \in G$  such that  $\langle x \rangle \cap p^\lambda G = 0$ ; then there exists a direct summand  $X$  of  $G$  containing  $x$  such that  $l(X) < \lambda$ .*

PROOF. The lemma is proved in [1, 2.7] if  $l(G) = \lambda$ ; if  $l(G) > \lambda$  the proof is quite similar and it is omitted.

The fact that every  $C_\lambda$ -group of length  $\leq \lambda$  is fully transitive, which is proved by Le Borgne [5] (see also Linton [6]), is generalized in the following

LEMMA 3.3. *Let  $G$  be a  $C_\lambda$ -group,  $\lambda = \text{cf } \omega$ . Let  $x, y \in G$  such that  $H(x) \leq H(y)$  and  $\langle x \rangle \cap p^\lambda G = 0 = \langle y \rangle \cap p^\lambda G$ . Then there exists an endomorphism  $f \in \text{End}(G)$  such that  $f(x) = y$ .*

PROOF. From lemma 3.2 one easily deduces a decomposition  $G = X \oplus G'$ , with  $x, y \in X$  and  $l(X) < \lambda$ ; since  $X$  is totally projective, hence fully transitive, the claim is now obvious.

LEMMA 3.4. *Let  $C$  be a  $C_\lambda$ -group,  $\lambda = \text{cf } \omega$ , and  $L$  a fully invariant subgroup of  $G$  such that  $p^\lambda G \subseteq p^\omega L$ . If  $\sigma_n = \inf \{h_a(p^n x) | x \in L\} < \lambda$  for every  $n \in \mathbb{N}$ , there exists, for each  $r \in \mathbb{N}$ , an element  $g \in L$  such that  $h_a(p^i g) = \sigma_i$  for  $i = 0, 1, \dots, r$ .*

PROOF. We induct on  $r$ . If  $r = 0$  the claim is obvious; suppose  $r \geq 1$ ; there exists an element  $g' \in L$  such that

$$H(g') = (\sigma_0, \sigma_1, \dots, \sigma_{r-1}, \tau_r, \tau_{r+1}, \dots)$$



where we can suppose  $\tau_r > \sigma_r$ , otherwise  $g'$  is the wanted element. There exists an element  $t \in L$  such that  $h_g(p^r t) = \sigma_r$ . Let  $k(t)$  be the minimal non negative integer such that  $h_g(p^i t) = \sigma_i$  if  $k(t) \leq i \leq r$ . We choose  $t \in L$  such that  $k(t)$  is minimal; let us observe that  $k(t) \leq r-1$  (otherwise consider  $t + g'$ ). Our goal is to prove that  $k(t) = 0$ . Among the elements  $t$  with  $k(t)$  minimal, choose one of minimal order  $m+1$  ( $m \geq r$ ); then  $h_g(p^m t) < \lambda$  otherwise  $p^m t \in p^\lambda G \subseteq p^\omega L$ , hence  $p^m t = p^{m+1} t'$  for a  $t' \in L$ ; in this case  $p^m(t - pt') = 0$  and  $h_g(p^i(t - pt')) = \sigma_i$  if  $k(t) \leq i \leq r$ , because  $h_g(p^i t) = \sigma_i$  and  $h_g(p^{i+1} t') \geq \sigma_{i+1} > \sigma_i$ . Thus, without loss of generality, we can assume, if  $k = k(t)$ :

$$H(t) = (\sigma'_0, \sigma'_1, \dots, \sigma'_{k-1}, \sigma_k, \sigma_{k+1}, \dots, \sigma_r, \sigma'_{r+1}, \dots, \sigma'_m, \infty, \dots)$$

where  $\sigma'_{k-1} > \sigma_{k-1}$ ,  $\sigma'_m < \lambda$  and  $k \leq r-1$ ; suppose, by way of contradiction, that  $k > 0$ . There exist elements  $b \in L$  such that  $h_g(p^{k-1} b) = \sigma_{k-1}$  and  $h_g(p^r b) = \sigma_r$  (consider for instance  $t + g'$ ); choose  $b$  among these elements such that  $k(b)$  is minimal; it is enough to prove that  $k(b) \leq k-1$ . Let, by absurd,  $k(b) \geq k$  and put  $k(b) = k'$ . Then  $h_g(p^{k'} b) = \sigma_{k'}$  and  $h_g(p^{k'-1} b) > \sigma_{k'-1}$  imply that  $\sigma_{k'} > \sigma_{k'-1} + 1$ ; therefore there exists an element  $w \in G$  such that  $p^{k'} t = p^{k'} w$  and  $h_g(p^{k'-1} w) > \sigma_{k'-1}$ ; of course we can choose  $w$  in such a way that  $H(w) \geq H(t)$ ; because of  $\langle t \rangle \cap p^\lambda G = 0 = \langle w \rangle \cap p^\lambda G$ , we can apply lemma 3.3 and deduce that  $w \in L$ . Then we have:  $p^{k'}(t - w) = 0$  and  $h_g(p^{k'-1}(t - w)) = \sigma_{k'-1}$ . Consider now the element  $b - t + w \in L$ ; then  $k(b - t + w) < k(b) = k'$ , because  $h_g(p^{k'-1}(b - t + w)) = \sigma_{k'-1}$ , which contradicts the choice of  $b$ .

We can now prove the main result of this section, which generalizes the characterization of large subgroups given by Pierce [11].

**THEOREM 3.5.** *Let  $G$  be a  $C_\lambda$ -group,  $\lambda = \text{cf } \omega$ , with  $l(G) \geq \lambda$ . Let  $L$  be a fully invariant subgroup of  $G$ . Then the following are equivalent:*

- 1)  $L = G(\mathbf{u})$ , where  $\mathbf{u} = (\sigma_n)_{n \in \mathbb{N}}$  is an increasing sequence of ordinals and  $\sigma_n < \lambda$  for every  $n \in \mathbb{N}$ .
- 2)  $L$  satisfies the  $\lambda$ -th Pierce's condition.
- 3)  $L$  is  $\lambda$ -large.

**PROOF.** 1)  $\rightarrow$  2) See property 5) of section 2.

2)  $\rightarrow$  3) Let  $B$  be a  $\lambda$ -basic subgroup of  $G$  and let  $g \in G$  be an element of order  $k \in \mathbb{N}$ ; there exists an ordinal  $\sigma < \lambda$  such that  $p^\sigma G[p^k] \subseteq L$ ;

but  $G = p^\sigma G + B$ , hence  $g = g' + b$  with  $g' \in p^\sigma G$ ,  $b \in B$ ; then  $0 = p^k g = p^k g' - p^k b$ , and  $p^k b \in p^{\sigma+k} G \cap B = p^{\sigma+k} B$ ; therefore there exists a  $b' \in p^\sigma B$  such that  $p^k b = p^k b'$ ; thus we obtain:  $g = (g' + b') + (b - b')$  where  $b - b' \in B$  and  $p^k(b - b') = 0$ ; it follows that  $g' + b' \in p^\sigma G[p^k] \subseteq L$  and  $g \in L + B$ .

3)  $\rightarrow$  1) If  $l(G) = \lambda$ , then  $G$  is fully transitive, therefore every fully invariant subgroup  $L$  is equal to  $G(\mathbf{u})$ , for a suitable increasing sequence  $\mathbf{u} = (\sigma_n)_{n \in \mathbb{N}}$  of ordinals and  $\infty$ 's. If  $\sigma_n \geq \lambda$  for some  $n$ , then  $p^n L \subseteq p^\lambda G = 0$ , which is absurd, because  $p^n L$  is  $\lambda$ -large while  $0$  is not  $\lambda$ -large. Suppose now that  $l(G) > \lambda$ . For every  $n \in \mathbb{N}$  we put:

$$\sigma_n = \inf \{h_\alpha(p^n x) \mid x \in L\};$$

obviously  $\sigma_n < \sigma_{n+1}$  and  $\sigma_n < \lambda$  for each  $n$ , otherwise  $p^n L = p^\lambda G$ , which is absurd because  $p^\lambda G$  is not  $\lambda$ -large; the inclusion  $L \subseteq G(\mathbf{u})$ , where  $\mathbf{u} = (\sigma_n)_{n \in \mathbb{N}}$  is obvious; we suppose by absurd that this inclusion is proper; let  $y$  be an element of minimal order among the elements of  $G(\mathbf{u}) \setminus L$ ; if  $H(y) = (\tau_0, \tau_1, \dots, \tau_k, \infty, \infty, \dots)$  with  $\tau_k < \infty$ , then  $\tau_k < \lambda$ : in fact, if  $\tau_k \geq \lambda$ , then  $p^k y \in p^\lambda G$ , which is contained in  $\bigcap_{n \in \mathbb{N}} p^n L = p^\omega L$ ,

by lemma 3.1; therefore there exists an element  $z \in L$  such that  $p^k(y - z) = 0$  and  $y - z \in G(\mathbf{u}) \setminus L$ , contradicting the choice of  $y$ . Let now  $x$  be an element of minimal order among the elements of  $L$  such that  $h_\alpha(p^i x) = \sigma_i$  for  $i = 0, 1, \dots, k$ ; such an element does exist, as lemma 2.4 shows; let  $H(x) = (\sigma_0, \sigma_1, \dots, \sigma_k, \sigma'_{k+1}, \dots, \sigma'_n, \infty, \dots)$ , with  $\sigma'_n < \infty$ ; then  $\sigma'_n < \lambda$ , otherwise, fixed an integer  $m \geq 1$ , there exists an element  $a \in L$  such that  $p^n x = p^{n+m} a$ , which implies that  $p^n(x - p^m a) = 0$ ; but if  $i = 0, 1, \dots, k$ , then  $h_\alpha(p^i(x - p^m a)) = \inf \{\sigma_i, h_\alpha(p^{m+i} a)\} = \sigma_i$ , contradicting the choice of  $x$ . We are now in the hypotheses of lemma 3.3, therefore there exists an endomorphism  $f$  of  $G$  such that  $f(x) = y$ ; but  $x \in L$  and  $L$  is fully invariant, hence  $y \in L$  and  $L = G(\mathbf{u})$ .

Theorem 3.5 enables us to improve theorems 4 and 5 of [6] to  $C_\lambda$ -groups of arbitrary length.

**COROLLARY 3.6.** *Let  $G$  be a  $C_\lambda$ -group,  $\lambda = \text{cf } \omega$ , and  $L = G(\mathbf{u})$  a  $\lambda$ -large subgroup, where  $\mathbf{u} = (\sigma_n)_{n \in \mathbb{N}}$  and  $\sigma_n < \lambda$  for each  $n \in \mathbb{N}$ . Then  $G/L$  is totally projective and, if  $\sigma = \sup \sigma_n$  and  $\lambda = \sigma + \tau$ , then  $L$  is a  $C_{\omega+\tau}$ -group.*

PROOF. If  $B$  is a  $\lambda$ -basic subgroup of  $G$ , the first fact follows from [6, Thm. 1] and the following isomorphism

$$G/L = G/G(\mathbf{u}) = G(\mathbf{u}) + B/G(\mathbf{u}) \cong B/G(\mathbf{u}) \cap B = B/B(\mathbf{u}).$$

It is easy to show that  $p^\omega L = p^\sigma G$ ; then the second fact follows from the easily proved equalities, which hold for every ordinal  $\varrho$

$$L/p^{\omega+\varrho} L = G(\mathbf{u})/p^{\omega+\varrho} G(\mathbf{u}) = G(\mathbf{u})/p^{\sigma+\varrho} G = (G/p^{\sigma+\varrho} G)(\mathbf{u}).$$

COROLLARY 3.7. *Let  $G$  be a  $C_\lambda$ -group,  $\lambda = \text{cf } \omega$ , and  $L$  a  $\lambda$ -large subgroup of  $G$ . Then  $L$  is totally projective if and only if  $G$  is totally projective.*

PROOF. The sufficiency follows from [6, Thm. 1]. The necessity follows from [7, Thm. 3], because  $G/L$  is totally projective by corollary 3.6.

Let us finally remark that, in the class of  $C_\lambda$ -groups,  $\lambda = \text{cf } \omega$ , two groups complete in the  $\lambda$ -inductive topology, i.e. two reduced  $p^\lambda$ -high injective groups, are isomorphic if and only if they have the same Ulm-Kaplansky invariants. This result has been proved by Megibben [8, Thm. 7] in the case of  $\lambda$  countable and by Crawley [1] in the general case; it has only to be pointed out that, if  $\lambda = \text{cf } \omega$ , the  $C_\lambda$ -groups torsion complete defined by Crawley are essentially the same as the reduced  $p^\lambda$ -high injective  $C_\lambda$ -groups.

*Added in proof.* While this paper was submitted for publication, appeared R. C. LINTON's paper:  $\lambda$ -large subgroups of  $C_\lambda$ -groups, in Pacific J. Math., **78**, no. 2 (1978), in which the equivalence of 1) and 3) of our theorem 3.5 and corollary 3.6 are proved. However his proof is quite different and is based on topological methods.

## REFERENCES

- [1] P. CRAWLEY, *On the structure of a class of abelian  $p$ -groups*, Symposia Math., **20** (1978).
- [2] D. O. CUTLER - R. W. STRINGALL, *A topology for primary abelian groups*, Studies on Abelian Groups, Paris (1968), pp. 93-100.

- [3] P. F. DUBOIS, *Generally  $p^\alpha$ -torsion complete abelian groups*, Trans. Amer. Math. Soc., **139** (1971), pp. 245-255.
- [4] L. FUCHS, *Infinite Abelian Groups*, vol. 2, New York-London, 1973.
- [5] M. M. LE BORGNE, *Groupes  $\lambda$ -separables*, C. R. Acad. Sci. Paris, **281** (1975), pp. 415-417.
- [6] R. C. LINTON, *On fully invariant subgroups of primary abelian groups*, Michigan Math. J., **22** (1975), pp. 281-284.
- [7] R. C. LINTON - C. MEGIBBEN, *Extensions of totally projective groups*, Proc. Amer. Math. Soc., **61-1** (1977), pp. 35-38.
- [8] C. MEGIBBEN, *A generalization of the classical theory of primary groups*, Tohoku Math. J., **22** (1970), pp. 347-356.
- [9] C. MEGIBBEN, *On  $p^\alpha$ -high injectives*, Math. Z., **122** (1971), pp. 104-110.
- [10] R. MINES, *A family of functor defined on generalized primary groups*, Pacific J. Math., **26** (1968), pp. 349-360.
- [11] R. S. PIERCE, *Homomorphisms of primary groups*, Topics in Abelian Groups, Chicago (1963), pp. 215-310.
- [12] K. D. WALLACE,  *$C_\lambda$ -groups and  $\lambda$ -basic subgroups*, Pacific J. Math., **43** (1972), pp. 799-809.

Manoscritto pervenuto in redazione il 7 ottobre 1978.