

RENDICONTI
del
SEMINARIO MATEMATICO
della
UNIVERSITÀ DI PADOVA

M. D. PÉREZ-RAMOS

A p -stable action of the automorphism group of a group

Rendiconti del Seminario Matematico della Università di Padova,
tome 79 (1988), p. 11-14

http://www.numdam.org/item?id=RSMUP_1988__79__11_0

© Rendiconti del Seminario Matematico della Università di Padova, 1988, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

A p -Stable Action of the Automorphism Group of a Group.

M. D. PÉREZ-RAMOS (*)

SUMMARY - The following Theorem is proved: « Let G be a group and $\text{Aut}(G)$ its automorphism group. If $\text{Aut}(G)$ is not p -stable over G , p an odd prime, then $\text{Aut}(G)$ involves $SL(2, p)$ ». This Theorem is a generalization of a classic result on stability ([4], Th. 3.8.3).

Introduction. Notation.

In this note all groups will be finite. p will denote a prime number, and $O_p(G \bmod N)$ the inverse image in G of the p -radical of G/N . If we assume that $A \leq \text{Aut}(G)$ and H is a subgroup of G which is fixed by A , then, whenever $a \in A$, we denote $[H, a] = \langle [x, a] = x^{-1}x^a/x \in H \rangle$, $[H, a, a] = [[H, a], a]$, $[H, A] = \langle [H, a]/a \in A \rangle$, and $[H, A, A] = [[H, A], A]$. The remainder of the notation is standard, and is taken mainly from [4]. In particular, $\Pi(G)$ is the set of all primes which divide the order of the group G .

The first concept concerning stability is the following:

DEFINITION A ([4], 3.8). Let G be a group with no nontrivial normal p -subgroups, p odd. A faithful representation ψ of G on a vector space V over $GF(p^n)$ will be called p -stable provided no p -element of $(G)\psi$ has a quadratic minimal polynomial on V . Moreover, G is said to be p -stable if all such faithful representations of G are p -stable. (We will say G is p -stable linear to distinguish it from later definitions of p -stability).

(*) Indirizzo dell'A.: Dept. de Algebra, Universidad de Valencia, Doctor Moliner 50, 46100 Burjassot (Spagna).

And the main result on p -stability linear:

THEOREM A ([4], 3.8.3). Let G be a group with no nontrivial normal p -subgroups, p odd. If G is not p -stable linear, then G involves $SL(2, p)$.

Later we can find other definitions of p -stable groups, ([3], [5]) π -stable groups, π a set of primes, ([1]), and in general, \mathcal{F} -stable groups, where \mathcal{F} is either a saturated Formation, ([2]), or a Fitting class, ([8]). As well, we find the close relationship between the stability of a group G , and the fact that G does not involve the special affin groups $SA(2, p)$, for adequated primes p , ([3], [5], [2], [7]).

The aim of this paper is to study the structure of the automorphisms group of a group by using these ideas.

We give the following Definition:

DEFINITION 1. Let G be a group. We say that the automorphisms group of G , $\text{Aut}(G)$, is p -estable over G if, whenever A is a p -subgroup of $\text{Aut}(G)$ and B is a p -subgroup of G , such that B is fixed by A , (that is, $A \leq N_{\text{Aut}(G)}(B)$), and $[B, A, A] = 1$, then

$$A \leq O_p(N_{\text{Aut}(G)}(B) \text{ mod } C_{\text{Aut}(G)}(B)).$$

We obtain the following result:

THEOREM 1. Let G be a group. If $\text{Aut}(G)$ is not p -stable over G , p odd, then $\text{Aut}(G)$ involves the special linear group $SL(2, p)$.

First, we give a preliminary result:

LEMMA. For a group G , the following are equivalent:

- i) $\text{Aut}(G)$ is p -stable over G .
- ii) Whenever B is a p -subgroup of G , and x is a p -automorphism of G which fixes B , and such that $[B, x, x] = 1$, then

$$x \in O_p(N_{\text{Aut}(G)}(B) \text{ mod } C_{\text{Aut}(G)}(B)).$$

- iii) Whenever B is a p -subgroup of G , and x is an automorphism of G which fixes B , and such that $[B, x, x] = 1$, then

$$x \in O_p(N_{\text{Aut}(G)}(B) \text{ mod } C_{\text{Aut}(G)}(B)).$$

PROOF. i) \rightarrow ii) Let B and x be as in (ii). The conclusion follows easily by taking $A = \langle x \rangle$, and using (i).

ii) \rightarrow iii) Let B and x be as in (iii). Let $\Pi(\langle x \rangle) = \{p_1, \dots, p_n\}$. Then $\langle x \rangle = \langle x_1 \rangle \times \dots \times \langle x_n \rangle$, where $\langle x_i \rangle$ is a p_i -group, for each $i = 1, \dots, n$. Obviously, each x_i , $i = 1, \dots, n$, fixes B and satisfies $[B, \langle x_i \rangle, \langle x_i \rangle] = 1$.

If $p_i \neq p$, then $x_i \in C_{\text{Aut}(G)}(B)$. Otherwise, we have

$$x_i \in O_p(\text{Aut}(G)(B) \bmod C_{\text{Aut}(G)}(B))$$

by using (ii). So,

$$x \in O_p(N_{\text{Aut}(G)}(B) \bmod C_{\text{Aut}(G)}(B)).$$

iii) \rightarrow i) Let A and B be as in Def. 1. For every $x \in A$, B and x satisfy the assumptions in (iii). Then, the conclusion is clear.

PROOF OF THEOREM 1. Because of the previous Lemma, if $\text{Aut}(G)$ is not p -stable over G , there exist a p -subgroup P of G , and a p -element x in $N_{\text{Aut}(G)}(P)$ such that $[P, x, x] = 1$, and

$$x \notin O_p(N_{\text{Aut}(G)}(P) \bmod C_{\text{Aut}(G)}(P)).$$

But, from ([5], Th. IX.7.8), this is equivalent to the existence of an element $g \in N_{\text{Aut}(G)}(P)$ such that $\langle x, x^g \rangle C_{\text{Aut}(G)}(P) / C_{\text{Aut}(G)}(P)$ is not a p -group.

Let $T = \langle x, x^g \rangle$. Let us refine the T -series $P \triangleright 1$ to a T -chief series $P = P_0 \triangleright P_1 \triangleright \dots \triangleright P_r = 1$ (*)

For each $i = 0, \dots, r-1$, let $\bar{P}_i = P_i / P_{i+1}$: So, it is clear that each \bar{P}_i , $i = 0, \dots, r-1$, is an abelian p -element group, that is, it is a vector space over $GF(p)$, and each $TC_{\text{Aut}(G)}(\bar{P}_i) / C_{\text{Aut}(G)}(\bar{P}_i) = \bar{T}_i$, $i = 0, \dots, r-1$, acts faithfully and irreducibly on \bar{P}_i : Therefore, since ([4], Th. 3.1.3), it follows that $O_p(\bar{T}_i) = 1$.

If \bar{T}_i was a p -group for every $i = 0, \dots, r-1$, that is, $\bar{T}_i = 1$, then $T \leq \bigcap_{i=0}^{r-1} C_{\text{Aut}(G)}(\bar{P}_i)$. Whence T would stabilize the series (*) of P , and we would get a contradiction because $TC_{\text{Aut}(G)}(P) / C_{\text{Aut}(G)}(P)$ is not a p -group.

Therefore, there exists an $i \in \{0, \dots, r-1\}$ such that $\bar{T}_i \neq 1$, that is, \bar{T}_i is not p -stable linear, (Def. A). Now, because of Th. A, \bar{T}_i involves $SL(2, p)$, and clearly $\text{Aut}(G)$ involves $SL(2, p)$.

Using an argument similar to ([4], Th. 3.8.4), and Th. 1, we get the following:

COROLLARY. Let G be a group. If one of the following conditions is verified in $\text{Aut}(G)$:

- i) $\text{Aut}(G)$ is of odd order.
- ii) A Sylow 2-subgroup of $\text{Aut}(G)$ is abelian.
- iii) A Sylow 2-subgroup of $\text{Aut}(G)$ is dihedral.
- iv) $\text{Aut}(G)$ is isomorphic to $L_2(q)$.
- v) $\text{Aut}(G)$ is soluble and either $p \geq 5$ or $p = 3$ and $SL(2, 3)$ is not involved in $\text{Aut}(G)$.

Then, $\text{Aut}(G)$ is p -stable over G .

Acknowledgments. This note is part of the author's Doctoral Thesis at the University of Valencia (Spain). The author wishes to express her gratitude to her supervisor Professor Pérez Monasor and also to Professor Iranzo for their devoted guidance and encouragement. The author was supported by a scholarship from the « Ministerio de Educación y Ciencia » of Spain.

REFERENCES

- [1] Z. ARAD, *A characteristic subgroup of π -stable groups*, *Canad. J. Math.*, **26** (1974), pp. 1509-1514.
- [2] L. M. EZQUERRO, *\mathcal{F} -stability of finite groups*, *J. Alg.*, **92**, N. 2 (1985), pp. 395-399.
- [3] G. GLAUBERMAN, *A characteristic subgroup of a p -stable group*, *Canad. J. Math.*, **20** (1968), pp. 1101-1135.
- [4] D. GORENSTEIN, *Finite Groups*, Harper & Row, New York, 1968.
- [5] B. HUPPERT - N. BLACKBURN, *Finite Groups II*, Springer-Verlag, Berlin, 1982.
- [6] M. D. PÉREZ-RAMOS, *Estabilidad con respecto de Nilpotentes, de Cuasinilpotentes y de clases de Fitting en general*, Tesis Doctoral, U. de Valencia, 1986.
- [7] M. D. PÉREZ-RAMOS, *Stability with respect to a Fitting class*, preprint.
- [8] M. D. PÉREZ-RAMOS, *A characteristic subgroup of \mathcal{N} -stable groups*, *Isr. J. Math.*, **54**, N. 1 (1986), pp. 51-59.

Manoscritto pervenuto in redazione il 10 ottobre 1986.