

RENDICONTI
del
SEMINARIO MATEMATICO
della
UNIVERSITÀ DI PADOVA

RICHARD E. PHILLIPS

On absolutely simple locally finite groups

Rendiconti del Seminario Matematico della Università di Padova,
tome 79 (1988), p. 213-220

http://www.numdam.org/item?id=RSMUP_1988__79__213_0

© Rendiconti del Seminario Matematico della Università di Padova, 1988, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

On Absolutely Simple Locally Finite Groups.

RICHARD E. PHILLIPS (*)

1. Introduction.

A well-known result of Kegel [5; pp. 172-173] (or [6; p. 115]) asserts that if G is a countably infinite locally finite simple group then

(1.1) there is an ascending chain $F_1 \subseteq \dots \subseteq F_n \subseteq \dots$ of finite subgroups of G satisfying

(a) $\bigcup F_n = G$, and

(b) for each $n > 1$ there is a maximal normal subgroup M_n of F_n such that $F_{n-1} \cap M_n = 1$.

The import of this result lies in the display of finite simple sections of unbounded orders in the finite subgroups of a countably infinite locally finite simple G . In general, the condition (1.1) does not imply simplicity [6; p. 116]. Indeed, there are countably infinite residually finite groups satisfying (1.1).

A minor adaptation of Kegel's arguments can be used to strengthen (1.1) to a condition equivalent to simplicity, and we will give such a condition in Theorems 1 and 2. In these same theorems we give a similar criteria for the absolute simplicity of G .

Recall that G is *absolutely simple* if the only composition series of G is the one consisting of 1 and G only; equivalently, G is absolutely

(*) Indirizzo dell'A.: Dept. of Mathematics, Michigan State University, East Lansing, MI. 48824, U.S.A.

simple if the only serial subgroups of G are 1 and G (see [11; I, p. 12, p. 16] or our § 2.1 for the relevant definitions). Obviously, every absolutely simple group is simple and, in general, the absolutely simple groups form a proper subclass of the class of simple groups ([1] or [11; II, 3.4]). However, it is not known whether or not every *locally finite* simple group is absolutely simple. Our Theorems 1 and 2 put the (possible) differences between these two (locally finite) classes in a «local» context.

In the sequel, we frequently encounter ascending chains $F_1 \subseteq \dots \subseteq F_n \subseteq \dots$ of finite subgroups of the countably infinite locally finite G with $\bigcup F_n = G$. Such a chain is called an *approximating sequence* of G (caution; this term is used in a different way in [6; p. 116]).

Part of Theorem 1 is stated in terms of subnormal subgroups. Recall that if $M \subseteq G$, the *standard series* of M in G is defined inductively by

$$M(0, G) = G \quad \text{and for } n \leq 1, \quad M(n, G) = M^{M(n-1), G}.$$

We also have occasion to use the subgroups

$$M(\omega, G) = \bigcap \{M(n, G) : n \geq 0\}.$$

THEOREM 1. Let $\{D_n\}$ be an approximating sequence of the countably infinite locally finite G .

- a) If G is simple, there is a subsequence $\{D_{n_k}\}$ of $\{D_n\}$ and an approximating sequence $\{F_k\}$ of G satisfying
- i) $F_1 = D_{n_1}$ and for $k > 1$, $F_k = D_{n_{k-1}}(1, D_{n_k}) = D_{n_{k-1}}^{D_{n_k}}$ and
 - ii) for $k > 1$, if $V \not\subseteq F_k$ then there is an $x \in N_{F_{k-1}}(F_k)$ such that $V^x \cap F_{k-1} = 1$.
- b) If G is absolutely simple, there is a subsequence $\{D_{n_k}\}$ of $\{D_n\}$ and an approximating sequence $\{F_k\}$ of G satisfying
- i) $F_1 = D_{n_1}$, and for $k > 1$, $F_k = D_{n_{k-1}}(\omega, D_{n_k})$, and
 - ii) for $k > 1$, if $V \not\subseteq F_k$, then $V \cap F_{k-1} = 1$.

Obviously, the condition in (ii) of part (a) implies (1.1).

A type of converse is provided in

THEOREM 2. Suppose $\{F_k\}$ is an approximating sequence of the countably infinite locally finite G .

- a) If $\{F_k\}$ satisfies the property (ii) of Theorem 1(a), then G is simple.
- b) If $\{F_k\}$ satisfies the property (ii) of Theorem 1(b), then G is absolutely simple.

An interesting interplay between Theorems 1 and 2 is that the existence of a single approximating sequence $\{F_k\}$ of G satisfying condition (ii) of Theorem 1(a) (or Theorem 1(b)) implies that a sequence with similar properties can be extracted from any approximating sequence (as in part a(i) of Theorem 1). The conditions a(ii) and b(ii) of Theorem 1 accentuate the possible differences between the countable « simple » and « absolutely simple » locally finite groups.

We also note that the condition (ii) of Theorem 1(b) is equivalent to

$$(1.2) \quad \text{for all } k \geq 1, 1 \neq x \in F_k \text{ implies } x^{F_{k+1}} = F_{k+1}.$$

While the above results are stated for countable groups, they can be extended to higher cardinalities by employing « countable » local theorems for the classes of simple and absolutely simple groups. The available theorems are recorded in

- (1.3) The infinite group G is simple (absolutely simple) if and only if G has a local system of countable simple (absolutely simple) subgroups (see [6; p. 114], [9; p. 190], [7; p. 131] for the simple case and [3; p. 529] for the absolutely simple case).

It follows immediately that Theorems 1 and 2 can be formulated in terms of the countable subgroups of the locally finite G .

Our final result gives a sufficient condition for absolute simplicity.

THEOREM 3. Let G be a countably infinite locally finite simple group and $\{D_n\}$ an approximating sequence of G . If there is a $d \geq 0$ such that every perfect subnormal subgroup of D_n has defect at most d in D_n , then G is absolutely simple.

The definition of the defect of a subnormal subgroup is given later in § 2.1; see also [11; I, p. 173].

It is not difficult to see that the perfect subnormal subgroups of a finite group L all have defect d or less if and only if the perfect subnormal subgroups of L/H have defect d or less, where H is the solvable

radical of L . Since all known countably infinite locally finite simple groups have approximating sequences $\{D_n\}$ where for $n \geq 1$, $D_n/\zeta(D_n)$ is a direct product of non-Abelian simple groups (see [10; p. 385]) we have, as a consequence of Theorem 3 and the above remarks on extensions to higher cardinals,

(1.4) all known locally finite simple groups are absolutely simple.

As a final remark, we point out, that with minor modifications, the groups $\{F_k\}$ of Theorem 1 can always be chosen to be perfect. To see this, let $\{D_n\}$ be an approximating sequence of the simple G and denote by D_n^ω the intersection of the members of the derived series of D_n . Since G cannot be locally solvable [11; p. 154] and $\bigcup \{D_n^\omega: n \geq 1\}$ is a normal subgroup of G , we must have $\bigcup \{D_n^\omega: n \geq 1\} = G$. Thus, $\{D_n^\omega\}$ is an approximating sequence of perfect subgroups of G . If the groups $\{F_k\}$ are chosen relative to $\{D_n^\omega\}$ (rather than $\{D_n\}$), the F_k 's will also be perfect.

2. Proofs.

2.1. Remarks on serial and subnormal subgroups.

The standard series of M in G has been defined in § 1. It is frequently easier to work with the commutator form

$$M(n, G) = M[G, nM]$$

where $[G, nM]$ is defined inductively by $[G, 0M] = G$ and for $n \geq 1$, $[G, nM] = [[G, (n-1)M], M]$ (c.f. [11; I, p. 173]). The subgroup M of G is subnormal in G written $M \triangleleft\triangleleft G$ if and only if $M = M(n, G)$ for some $0 \leq n < \omega$. Equivalently, $M \triangleleft\triangleleft G$ if and only if $[G, nM] \subseteq M$ for some $n \geq 0$. If $M \triangleleft\triangleleft G$, the minimal n for which $M = M(n, G)$ is called the *defect* of M in G . The symbol $M \triangleleft\triangleleft_n G$ will mean that M is subnormal in G of defect n or less.

Several useful facts are given in

- (2.1.1) (a) if $N \subseteq M \triangleleft\triangleleft_n G$, then $N(n, G) \subseteq M$.
 (b) If $N \subseteq M \triangleleft\triangleleft G$, then $N(\omega, G) \subseteq M$.
 (c) If G is finite and $M \subseteq G$, then $M^{M(\omega, G)} = M(\omega, G)$.

Our use of the term normal series coincides with the normal systems of Kurosh [8; p. 171] and is essentially equivalent to the series of Robinson [11; I, pp. 9-10]. A subgroup H of G is a serial subgroup of G (written H ser G) if there is a normal series \mathcal{C} of G with $H \in \mathcal{C}$. We will need the following «local» characterization of serial subgroups [4; Theorem 2] (or in the locally finite case [2; Lemma 2]).

(2.1.2) If $H \subseteq G$, then H ser G if and only if for every finitely generated $F \subseteq G$, $F \subseteq H^F$ implies $F \subseteq H$.

An essential lemma for our arguments is

LEMMA 1. Let $\{D_n\}$ be an approximating sequence of the countably infinite locally finite G and suppose that for each $n \geq 1$ we have a subgroup $M_n \triangleleft\triangleleft D_n$ and that $n > m$ implies $M_m \subseteq M_n$. Then $M = \bigcup \{M_n : n \geq 1\}$ ser G . Further, if there is a $d \geq 0$ such that $M_n \triangleleft\triangleleft_d D_n$ for all $n \geq 1$ then $M \triangleleft\triangleleft_d G$.

PROOF. For the first part, we use the criterion (2.1.2). Let F be a finite subgroup of G and suppose that $F \subseteq M^F$. Then there is an n such that $F \subseteq D_n$ and $F \subseteq M_n^F$. Since $M_n \triangleleft\triangleleft \langle M_n, F \rangle \subseteq D_n$ we have $F \subseteq M_n \subseteq M$ as desired.

Suppose now that $M_n \triangleleft\triangleleft_d D_n$ for all n . Then, for $n \geq 1$, $[D_n, dM_n] \subseteq M_n$ and so

$$[G, dM] = \bigcup \{[D_n, dM_n] : n \geq 1\} \subseteq M.$$

Thus, $M \triangleleft\triangleleft_d G$ and this completes the proof.

2.2. For the proof of Theorem 1 we require the following lemma. The proof follows the lines of argument given in [6; pp. 112-114].

LEMMA 2. Let $\{D_n\}$ be an approximating sequence of the countably infinite locally finite G and put $D = D_1$.

- a) If G is simple and $d \geq 0$ then
 - i) $\{D(d, D_n) : n \geq 1\}$ is an approximating sequence of G , and
 - ii) there is a positive integer j such that for $n \geq j$, $Y \triangleleft\triangleleft_d D_n$ implies $Y \cap D \in \{1, D\}$.
- b) If G is absolutely simple, then
 - i) $\{D(\omega, D_n) : n \geq 1\}$ is an approximating sequence of G , and

ii) there is a positive integer j such that for $n \geq j$, $Y \triangleleft\triangleleft D_n$ implies $Y \cap D \in \{1, D\}$.

PROOF. For the proof of (i) of part (a), note first that for $n \geq 1$, $D(d, D_n) \subseteq D(d, D_{n+1})$. From Lemma 1 we have $V = \bigcup \{D(d, D_n) : n \geq 1\} \triangleleft\triangleleft_a G$ and the simplicity of G forces $V = G$.

Part (i) of (b) follows similarly; in this case we have $V = \bigcup \{D(\omega, D_n) : n \geq 1\}$ ser G (by Lemma 1) and since G is absolutely simple, $V = G$.

Proceeding to part (ii) of (a), suppose that there is no j with the asserted property. There is then an approximating sequence $\{P_n\} \subseteq \{D_n\}$ and subgroups $Y_n \triangleleft\triangleleft_a P_n$ such that $Y_n \cap D \notin \{1, D\}$. Since D is finite there is a subgroup M of D with $M \notin \{1, D\}$ and an approximating sequence $\{E_n\} \subseteq \{P_n\}$ such that for $n \geq 1$ there are subgroups $X_n \triangleleft\triangleleft_a E_n$ with $X_n \cap D = M$. Now for $n \geq 1$, $M(d, E_n) \subseteq X_n$ (by (2.1.1)(a)) and so $M = D \cap M(d, E_n)$. From part (i) we have $G = \bigcup \{M(d, E_n) : n \geq 1\}$ and the contradiction $M = D \cap G$ now follows.

The proof of b(ii) is identical with that of a(ii); in the same manner we arrive at an approximating sequence $\{E_n\} \subseteq \{D_n\}$ and subgroups $X_n \triangleleft\triangleleft E_n$ with $M \notin \{1, D\}$. The fact that $\bigcup \{M(\omega, E_n) : n \geq 1\} = G$ (part (i) of (b)) together with $M = D \cap M(\omega, E_n)$ for $n \geq 1$ again yields the contradiction $D = M$.

2.3 PROOF OF THEOREM 1. Let $\{D_n\}$ be an approximating sequence of G and suppose G is simple. If F, S are finite subgroups of G there is, by Lemma 2(a) a positive integer $j = j(F, S)$ such that $\langle F, S \rangle \subseteq F(1, D_j) = \mu(F, S)$ and $Y \triangleleft\triangleleft_2 D_j$ implies $Y \cap F \in \{1, F\}$. If $V \triangleleft\triangleleft \mu(F, S)$ then $V \triangleleft\triangleleft_2 D_j$ and so $V \cap F \in \{1, F\}$. Further, if $F \subseteq V$ and $L = \text{Core}_{D_j}(V)$ then $L \cap F \in \{1, F\}$. If $F \subseteq L$, then $L = F^{D_j} = \mu(F, S)$ which contradicts the fact that $V \neq \mu(F, S)$. Thus, $L \cap F = 1$ and since for every $x \in D_j$ we have $V^x \cap F \in \{1, F\}$, there must be an $x \in D_j$ with $V^x \cap F = 1$.

Now for the construction of the desired subsequence $\{F_k\}$. Put $F_1 = D_1$, $F_2 = \mu(F_1, D_2)$ and $F_3 = \mu(F_2, D_{j_3})$ where $j_3 = \max \{3, j(F_1, D_2)\}$; for $k > 3$, let $F_k = \mu(F_{k-1}, D_{j_k})$ where $j_k = \max \{k, j(F_{k-1}, D_{j_{k-1}})\}$. One checks easily that $\{F_k\}$ has the properties listed in Theorem 1(a).

For the proof of (b), let $\{D_n\}$ be an approximating sequence of the absolutely simple G and F and S finite subgroups of G . Using Lemma 2(b) there is a positive integer $j = j(F, S)$ such that

$\langle F, S \rangle \subseteq F(\omega, D_j) = \mu(F, S)$ and $Y \triangleleft \triangleleft D_j$ implies $Y \cap F \in \{1, F\}$. If $V \not\subseteq \mu(F, S)$ and $F \subseteq V$ we have $F \subseteq V \triangleleft \triangleleft \mu(F, S)$ which forces $\mu(F, S) = V$ (by (2.1.1)(b)). From this we conclude that $F \cap V = 1$.

The sequence $\{F_k\}$ satisfying (i) and (ii) of Theorem 1(b) may now be constructed as follows:

$$F_1 = D_1, \dots, \quad F_n = \mu(F_{n-1}, D_n), \dots$$

2.4 PROOF OF THEOREM 2. Suppose G has an approximating sequence $\{F_k\}$ satisfying the property (ii) of Theorem 1(a) and let $1 \neq H \triangleleft G$. Then for some k_0 , $k \geq k_0$ implies $H \cap F_k \neq 1$. For any such k , $f_k \cap (F_{k+1} \cap H)^x = F_k \cap H = 1$ for any $x \in N(F_{k+1})$. Thus, $F_{k+1} \cap H = F_{k+1}$ and this forces $H = G$. We have shown that G is simple and this concludes the proof of part (a).

For part (b), suppose the approximating sequence $\{F_k\}$ satisfies the property (ii) of Theorem 1(b) and that $1 \neq H \text{ ser } G$. As above, there is a k_0 such that $k \geq k_0$ implies $H \cap F_k \neq 1$. Thus, if $k \geq k_0$, $1 \neq H \cap F_k \triangleleft \triangleleft F_k$ and $(H \cap F_k) \cap (H \cap F_{k+1})^{F_{k+1}} \neq 1$. This gives $(H \cap F_{k+1})^{F_{k+1}} = F_{k+1}$ and we conclude that $(H \cap F_{k+1}) = F_{k+1}$. It follows that $H = G$ and that G is absolutely simple.

2.5. Prior to our proof of Theorem 3, we need

(2.5.1) If H is a serial locally solvable subgroup of a locally finite G , then H^G is also locally solvable.

The proof of (2.5.1) is straightforward and will not be given here. For the proof of Theorem 3, let G be a countable simple locally finite group and $\{D_n\}$ an approximating sequence such that for each n , the perfect subnormal subgroups of D_n are of defect d or less. Now let $H \text{ ser } G$ with $1 \neq H$; from (2.5.1) and the fact that simple locally solvable groups are finite [11; I, p. 154], we see that H is not locally solvable. Thus, there is an n_0 such that for $n \geq n_0$, $H \cap D_n$ is not solvable. Consequently, if $n \geq n_0$, the subgroup $(H \cap D_n)^\omega$, the intersection of the terms of the derived series of $H \cap D_n$, is a non-trivial perfect subnormal subgroup of D_n . From our assumptions, we have $(H \cap D_n)^\omega \triangleleft \triangleleft_d D_n$. Lemma 1 now implies that $V = \bigcup \{(H \cap D_n)^\omega : n \geq 1\} \triangleleft \triangleleft_d G$ and the simplicity of G forces $V = G$. Since $V \subseteq H$, we have $H = G$ also, and G is absolutely simple.

REFERENCES

- [1] S. CHEHATA, *An algebraically simple ordered group*, Proc. London Math. Soc., (3) **2** (1952), pp. 183-197.
- [2] B. HARTLEY, *Serial subgroups of locally finite groups*, Proc. Cambridge Philos. Soc., **71** (1972), pp. 199-201.
- [3] K. HICKIN, *Countable type local theorems in algebra*, J. of Algebra, **27** (1973), pp. 523-537.
- [4] K. HICKIN - R. E. PHILLIPS, *On classes of groups defined by systems of subgroups*, Arch. der Math., **24** (1973), pp. 346-350.
- [5] O. KEGEL, *Über einfache, lokal endliche Gruppen*, Math. Z., **95** (1967), pp. 169-195.
- [6] O. KEGEL - B. A. F. WEHRFRITZ, *Locally Finite Groups*, North Holland, London (1973).
- [7] R. D. KOPPERMAN - A. R. D. MATHIAS, *Some problems in group theory*, «The Syntax and Semantics of Infinitary Languages», Lecture Notes in Math., **72**, Springer-Verlag, New York (1968).
- [8] A. G. KUROSH, *The Theory of Groups*, Vol. II, Chelsea, New York (1956).
- [9] R. E. PHILLIPS, *Countably recognizable classes of groups*, Rocky Mt. J. of Math., **1** (1971), pp. 489-497.
- [10] R. E. PHILLIPS, *Existentially closed locally finite central extensions; multipliers and local systems*, Math. Z., **487** (1984), pp. 383-392.
- [11] D. J. S. ROBINSON, *Finiteness Conditions and Generalized Soluble Groups*, Vol. I and II, Springer-Verlag, New York (1972).

Manoscritto pervenuto in redazione il 9 aprile 1987.