

RENDICONTI
del
SEMINARIO MATEMATICO
della
UNIVERSITÀ DI PADOVA

H. M. SRIVASTAVA

M. A. PATHAN-YASMEEN

**Some reduction formulas for multiple
hypergeometric series**

Rendiconti del Seminario Matematico della Università di Padova,
tome 82 (1989), p. 1-7

http://www.numdam.org/item?id=RSMUP_1989__82__1_0

© Rendiconti del Seminario Matematico della Università di Padova, 1989, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

Some Reduction Formulas for Multiple Hypergeometric Series.

H. M. SRIVASTAVA (*) - M. A. PATHAN - YASMEEN (**)

SUMMARY - The authors prove a general theorem on the reducibility of a certain multiple series and apply it to derive numerous reduction formulas for hypergeometric functions of several variables. Some applications of this theorem, involving Jacobi polynomials, are also presented.

1. Introduction and the main result.

For Appell's double hypergeometric function F_4 defined by (cf., e.g., [1, p. 14])

$$(1.1) \quad F_4[\alpha, \beta; \gamma, \delta; x, y] = \sum_{l, m=0}^{\infty} \frac{(\alpha)_{l+m}(\beta)_{l+m} x^l y^m}{(\gamma)_l(\delta)_m l! m!},$$

where, and throughout this paper,

$$(1.2) \quad (\lambda)_l = \frac{\Gamma(\lambda + l)}{\Gamma(\lambda)} = \begin{cases} 1, & \text{if } l = 0, \\ \lambda(\lambda + 1) \dots (\lambda + l - 1), & \forall l \in \{1, 2, 3, \dots\}, \end{cases}$$

(*) The work of this author was supported, in part, by the *Natural Sciences and Engineering Research Council of Canada* under Grant A-7353.

(**) Indirizzo degli AA.: H. M. SRIVASTAVA: Department of Mathematics, University of Victoria, Victoria, British Columbia V8W 2Y2, Canada; M.A. PATHAN: Department of Mathematics, Aligarh Muslim University, Aligarh 202001, Uttar Pradesh, India; YASMEEN: Department of Mathematics, Government College, Kota 324001, Rajasthan, India.

Bailey [3, p. 42, Equation (4.2)] proved the reduction formula:

$$(1.3) \quad F_4 \left[\alpha, \beta; \beta, \alpha; -\frac{x}{(1-x)(1-y)}, -\frac{y}{(1-x)(1-y)} \right] = \\ = (1-x)^\alpha (1-y)^\beta (1-xy)^{-1}.$$

Formula (1.3) is known to play a rather crucial rôle in one of many interesting proofs of Jacobi's generating function:

$$(1.4) \quad \sum_{n=0}^{\infty} P_n^{(\alpha, \beta)}(x) t^n = 2^{\alpha+\beta} R^{-1} (1-t+R)^{-\alpha} (1+t+R)^{-\beta}, \\ R = (1-2xt+t^2)^{\frac{1}{2}}$$

for the classical Jacobi polynomials $P_n^{(\alpha, \beta)}(x)$ defined, in terms of the Gaussian hypergeometric function, by

$$(1.5) \quad P_n^{(\alpha, \beta)}(x) = \binom{\alpha+n}{n} {}_2F_1 \left[-n, \quad \alpha + \beta + n + 1; \frac{1-x}{\alpha + 1} \right].$$

See, for example, Rainville [8, p. 269, Section 140] and Carlitz [4]; see also Askey [2, p. 246] and Srivastava [9, p. 201].

An interesting extension of Bailey's formula (1.3) was given recently by Cohen [6, p. 272, Theorem 1] who indeed applied his result to deduce several new or known generating functions for Jacobi polynomials including, for instance, a remarkable special case ($\lambda = 0$, $\mu = s + 1$) of the following generalization of (1.4) due to Srivastava and Singhal [12, p. 749, Equation (8)]:

$$(1.6) \quad \sum_{n=0}^{\infty} P_n^{(\alpha-\lambda n, \beta-\mu n)}(x) t^n = \\ = (1+\xi)^{\alpha+1} (1+\eta)^{\beta+1} [1+\lambda\xi + \mu\eta - (1-\lambda-\mu)\xi\eta]^{-1},$$

where ξ and η are functions of x and t defined implicitly by

$$(1.7) \quad (x+1)^{-1}\xi = (x-1)^{-1}\eta = \frac{1}{2}t(1+\xi)^{1-\lambda}(1+\eta)^{1-\mu}.$$

The object of the present paper is to prove a multidimensional extension of Cohen's theorem and to show how our general result

can be applied with a view to deriving, for example, reduction formulas for multiple hypergeometric functions. Our main result is contained in the following

THEOREM. *Let r_1, \dots, r_n and $\alpha_1, \dots, \alpha_n$ denote arbitrary complex numbers. Also, for convenience, let*

$$(1.8) \quad k_{n+1} = k_1,$$

where n is a fixed positive integer.

Then

$$(1.9) \quad \sum_{k_1, \dots, k_n=0}^{\infty} \prod_{i=1}^n \left\{ \frac{(\alpha_i + r_i k_{i+1})_{k_i} (-z_i)^{k_i}}{(1 - z_i)^{k_i + r_i k_{i+1}} k_i!} \right\} = \frac{(1 - z_1)^{\alpha_1} \dots (1 - z_n)^{\alpha_n}}{1 - (-1)^n r_1 z_1 \dots r_n z_n},$$

provided that

$$|z_1| < 1, \dots, |z_n| < 1 \quad \text{and} \quad |r_1 z_1 \dots r_n z_n| < 1.$$

2. Proof of the Theorem.

To prove the multidimensional reduction formula (1.9), we begin by considering the multiple series:

$$(2.1) \quad \Omega(x; z_1, \dots, z_n) = \sum_{m_1, \dots, m_n=0}^{\infty} \frac{z_1^{m_1}}{m_1!} \dots \frac{z_n^{m_n}}{m_n!} \cdot D_x^{m_1} \{x^{\alpha_1 + m_1 - 1} (1 - x^{r_1})^{m_2}\} \dots D_x^{m_n} \{x^{\alpha_n + m_n - 1} (1 - x^{r_n})^{m_1}\},$$

where, as usual, $D_x = d/dx$.

Applying the binomial theorem and operating upon the resulting sums term-by-term, we find from (2.1) that

$$(2.2) \quad \Omega(x; z_1, \dots, z_n) = \sum_{m_1, \dots, m_n=0}^{\infty} z_1^{m_1} \dots z_n^{m_n} \cdot \sum_{k_1=0}^{m_1} \dots \sum_{k_n=0}^{m_n} \prod_{i=1}^n \left\{ (-1)^{k_i} \frac{(\alpha_i + r_i k_{i+1})_{m_i}}{k_i! (m_i - k_i)!} x^{\alpha_i + r_i k_{i+1} - 1} \right\},$$

where we have followed the convention (1.8).

Now we set $x = 1$ in (2.1) and (2.2). Obviously, (2.1) contributes only when

$$m_1 = \dots = m_n,$$

and we have

$$(2.3) \quad \Omega(1; z_1, \dots, z_n) = \sum_{m=0}^{\infty} (-1)^{mn} (r_1 z_1 \dots r_n z_n)^m = [1 - (-1)^n r_1 z_1 \dots r_n z_n]^{-1},$$

provided that

$$|r_1 z_1 \dots r_n z_n| < 1.$$

On the other hand, by inverting the order of summation in (2.2) with $x = 1$, we readily observe that

$$(2.4) \quad \Omega(1; z_1, \dots, z_n) = \sum_{k_1, \dots, k_n=0}^{\infty} \prod_{i=1}^n \left\{ (\alpha_i + r_i k_{i+1})_{k_i} \frac{(-z_i)^{k_i}}{k_i!} \right\} \cdot \prod_{i=1}^n \left\{ \sum_{m_i=0}^{\infty} (\alpha_i + k_i + r_i k_{i+1})_{m_i} \frac{z_i^{m_i}}{m_i!} \right\}.$$

Since

$$(2.5) \quad \sum_{m=0}^{\infty} (\lambda)_m \frac{z^m}{m!} = (1-z)^{-\lambda}, \quad |z| < 1,$$

we find from (2.4) that

$$(2.6) \quad \Omega(1; z_1, \dots, z_n) = (1-z_1)^{-\alpha_1} \dots (1-z_n)^{-\alpha_n} \cdot \sum_{k_1, \dots, k_n=0}^{\infty} \prod_{i=1}^n \left\{ \frac{(\alpha_i + r_i k_{i+1})_{k_i} (-z_i)^{k_i}}{(1-z_i)^{k_i + r_i k_{i+1}} k_i!} \right\},$$

provided that

$$\max \{|z_1|, \dots, |z_n|\} < 1.$$

The reduction formula (1.9) would follow immediately upon equating the two expressions for

$$\Omega(1; z_1, \dots, z_n)$$

given by (2.3) and (2.6), thus completing the proof of the theorem.

3. Applications.

The case $n = 2$ of the reduction formula (1.9) was given earlier by Cohen [6, p. 272, Theorem 1], and Bailey's formula (1.3) follows immediately from (1.9) upon setting $n = 2$ and $r_1 = r_2 = 1$. We shall show how our general result (1.9) can be applied, when $n > 2$, in order to derive various new or known results involving multiple hypergeometric functions and Jacobi polynomials.

I. For $r_1 = \dots = r_n = 1$, (1.9) can at once be rewritten as a reduction formula for the hypergeometric function $H_B^{(n)}$ of n variables, and we are thus led to a known result due to Khichi ([7, p. 23, Equation (3.2)]; see also Srivastava and Karlsson [10, p. 308, Equation (122)]) who incidentally derived it first for $n = 3$ by appealing to the proof of Bailey's formula (1.3) by Bailey [3, pp. 41-42] and Rainville [8, pp. 266-267] *mutatis mutandis*. It should be remarked in passing that $H_B^{(2)}$ is the Appell function F_4 defined by (1.1), and $H_B^{(3)}$ is Srivastava's function H_B [10, p. 43, Equation (12)].

II. For the triple hypergeometric series F_K of Lauricella's set (see, e.g., [12, p. 42, Equation (4)]), our theorem with $n = 4$, $r_1 = r_3 = r_4 = 1$, and $r_2 = \rho$ yields

$$\begin{aligned}
 (3.1) \quad & \sum_{l=0}^{\infty} \frac{(c)_l}{l!} \left\{ -\frac{z}{(1-z)(1-y)^e} \right\}^l F_K \left[b + \rho l, d, d, a, c + l, a; a, c, d; \right. \\
 & \left. -\frac{y}{(1-y)(1-x)}, -\frac{t}{(1-t)(1-z)}, -\frac{x}{(1-x)(1-t)} \right] = \\
 & = (1-x)^a (1-y)^b (1-z)^c (1-t)^d (1-\rho x y z t)^{-1}.
 \end{aligned}$$

Some further consequences of (3.1) are worthy of mention. First of all, since F_K is expressible as an infinite series of products of two Gaussian hypergeometric ${}_2F_1$ functions (*cf.* [11, p. 243, Equation (13)]), each of which can be rewritten fairly easily as a Jacobi polynomial in view of (1.5), setting

$$a = \alpha + 1, \quad b = -\beta, \quad c = \gamma + 1, \quad d = -\delta,$$

and

$$\frac{y}{1-y} = -\frac{1}{2}(1-x)(1-u), \quad \frac{t}{1-t} = -\frac{1}{2}(1-z)(1-v),$$

and replacing z by y , we find from (3.1) that

$$(3.2) \quad \sum_{l,m=0}^{\infty} \left\{ -\frac{2x[2-(1-y)(1-v)]}{(1-x)(1+u)(1+v)} \right\}^l \left\{ -\frac{2y[2-(1-x)(1-u)]^e}{(1-y)(1+u)^e(1+v)} \right\}^m \\ \cdot P_l^{\alpha, \beta-l-em}(u) P_m^{\gamma, \delta-l-m}(v) = \\ = \frac{(1-x)^{\alpha+1}(1-y)^{\gamma+1}[2-(1-x)(1-u)]^{\beta+1}[2-(1-y)(1-v)]^{\delta+1}}{(1+u)^{\beta}(1+v)^{\delta}[4-2\{(1-x)(1-u)+(1-y)(1-v)\}+(1-x)(1-y)(1-u)(1-v)(1-\rho xy)]},$$

which, for $x = 0$ and $\rho = s + 1$, corresponds to the aforementioned special case ($\lambda = 0$, $\mu = s + 1$) of (1.6).

The triple hypergeometric function F_K is expressible also as an infinite series of Appell's function F_2 which can, in turn, be specialized to give a ${}_2F_1$ either directly (cf. [10, p. 305, Equation (108)]) or through Appell's function F_1 (cf. [10, p. 305, Equation (106); p. 304, Equation (99)]). In this manner, (3.1) with $a = d$ and $\rho = 1$ yields an interesting reduction formula for the triple Gaussian hypergeometric series $10c$ or $18b$ [10, pp. 77 and 79], and (3.1) with $a = d$ and $\rho = 0$ leads us to a special case of a result due to Chaundy ([5, p. 62, Equation (25)]; see also [11, p. 107, Equation (14)]).

III. In terms of Horn's double hypergeometric function H_2 (see, e.g., [10, p. 24, Equation (10)]), our theorem with $n = 4$, $r_1 = -1$, $r_2 = \rho$, and $r_3 = r_4 = 1$ yields

$$(3.3) \quad \sum_{l,m=0}^{\infty} \frac{(c)_{l+m}(d)_m}{l!m!(c)_m} \left\{ -\frac{z}{(1-z)(1-y)^e} \right\}^l \left\{ -\frac{t}{(1-t)(1-z)} \right\}^m \\ \cdot H_2 \left[a, d+m, 1-a, b+\rho l; d; -\frac{x}{(1-x)(1-t)}, \frac{y(1-x)}{1-y} \right] = \\ = (1-x)^a(1-y)^b(1-z)^c(1-t)^d(1+\rho xyzt)^{-1}.$$

In its special case when $\rho = 0$ and $d = a + b$, this last result (3.3) can easily be rewritten in terms of the Horn function G_2 by appealing to a known transformation [10, p. 319, Equation (170)]. If we further set $c = a + b$, we shall be led eventually to a reduction formula involving the triple Gaussian hypergeometric series $4h$ [10, p. 75].

We conclude by remarking that analogous results involving various other classes of multiple hypergeometric functions can be deduced, in a manner described above, for other appropriate choices of n and r_1, \dots, r_n .

REFERENCES

- [1] P. APPELL - J. KAMPÉ DE FÉRIET, *Fonctions Hypergéométriques et Hypersphériques; Polynômes d'Hermite*, Gauthier-Villars, Paris (1926).
- [2] R. ASKEY, *Jacobi's generating function for Jacobi polynomials*, Proc. Amer. Math. Soc., **71** (1978), pp. 243-246.
- [3] W. N. BAILEY, *Some infinite integrals involving Bessel functions*, Proc. London Math. Soc. (2), **40** (1936), pp. 37-48.
- [4] L. CARLITZ, *The generating function for the Jacobi polynomial*, Rend. Sem. Mat. Univ. Padova, **38** (1967), pp. 86-88.
- [5] T. W. CHAUNDY, *An extension of hypergeometric functions (I)*, Quart. J. Math. Oxford Ser., **14** (1943), pp. 55-78.
- [6] M. E. COHEN, *Generating functions for the Jacobi polynomial*, Proc. Amer. Math. Soc., **57** (1976), pp. 271-275.
- [7] K. S. KHICHI, *A note on Srivastava's triple hypergeometric function H_B* , Ganita, **19**, no. 1 (1968), pp. 21-24.
- [8] E. D. RAINVILLE, *Special Functions*, Macmillan, New York (1960).
- [9] H. M. SRIVASTAVA, *A note on Jacobi's generating function for the Jacobi polynomials*, Proc. Japan Acad. Ser. A Math. Sci., **61** (1985), pp. 201-202.
- [10] H. M. SRIVASTAVA - P. W. KARLSSON, *Multiple Gaussian Hypergeometric Series*, Halsted Press (Ellis Horwood Limited, Chichester), John Wiley and Sons, New York (1985).
- [11] H. M. SRIVASTAVA - H. L. MANOCHA, *A Treatise on Generating Functions*, Halsted Press (Ellis Horwood Limited, Chichester), John Wiley and Sons, New York (1984).
- [12] H. M. SRIVASTAVA - J. P. SINGHAL, *New generating functions for Jacobi and related polynomials*, J. Math. Anal. Appl., **41** (1973), pp. 748-752.

Manoscritto pervenuto in redazione il 3 settembre 1987.