

# RENDICONTI *del* SEMINARIO MATEMATICO *della* UNIVERSITÀ DI PADOVA

DĂNUȚ MARCU

## **A note on the projective representations of finite groups**

*Rendiconti del Seminario Matematico della Università di Padova*,  
tome 102 (1999), p. 23-27

[http://www.numdam.org/item?id=RSMUP\\_1999\\_\\_102\\_\\_23\\_0](http://www.numdam.org/item?id=RSMUP_1999__102__23_0)

© Rendiconti del Seminario Matematico della Università di Padova, 1999, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques*  
<http://www.numdam.org/>

## A Note on the Projective Representations of Finite Groups.

DĂNUȚ MARCU (\*)

ABSTRACT - In this paper, we describe exactly the greatest common divisor of the degrees of projective representations of finite groups.

### 1. Introduction.

All representations and characters, studied in this paper, are taken over the complex numbers, and all considered groups are finite. For basic definitions concerning projective representations, see [1]. If  $G$  is a group and  $\alpha$  is a cocycle of  $G$ , we denote by  $\text{Proj}(G, \alpha) = \{\tau_1, \tau_2, \dots, \tau_t\}$  the set of irreducible projective characters of  $G$  with cocycle  $\alpha$ , where (of course)  $t$  is the number of  $\alpha$ -regular conjugacy classes of  $G$ ,  $\tau_i(1)$  being called the *degree* of  $\tau_i$ . Also as normal,  $M(G)$  will denote the Schur multiplier of  $G$ ,  $[\alpha]$  the cohomology class of  $\alpha$ , and  $[1, G]$  the cohomology class of the trivial cocycle of  $G$ .

The main result of this paper exactly describes the greatest common divisor of the degrees of  $\text{Proj}(G, \alpha)$ .

### 2. Main result.

Our main result is the following

**THEOREM.** *Let  $p_1, p_2, \dots, p_n$  be the prime divisors of  $|G|$ , with  $P_1, P_2, \dots, P_n$  corresponding Sylow  $p_i$ -subgroups of  $G$ . Let  $M_i$  be a sub-*

(\*) Indirizzo dell'A.: Str. Pasului 3, Sect. 2, 70241-Bucharest, Romania.

group of  $P_i$  of minimal index such that  $[a_{M_i}] = [1, G]$ . Then, the greatest common divisor of the degrees of  $\text{Proj}(G, a)$  is equal to  $\prod_{i=1}^n [P_i: M_i]$ . ■

We start by defining

$$s(G, a) = \min \{ \tau_i(1): 1 \leq i \leq t \}$$

and

$$c(G, a) = \text{g.c.d.} \{ \tau_i(1): 1 \leq i \leq t \}.$$

It is obvious that if  $[a] = [1, G]$ , then  $c(G, a) = s(G, a) = 1$ . Thus, we are only really interested in non-trivial cocycles of  $G$ .

Now, we quote the following well-known result, which, to my known, belongs to the *folklore* of this topic (e.g., for the part (a), see [1, 6.2.6]).

LEMMA 1. *Let  $a$  be a cocycle of  $G$  with  $o([a]) = e$  in  $M(G)$ . Then,*

(a)  $e | c(G, a)$ ;

(b) *if  $p$  is a prime number such that  $p | c(G, a)$ , then  $p | e$ .* ■

We note here that it is not true, in general, that  $c(G, a) = e$ , or indeed that, if some integer  $m$  divides  $c(G, a)$ , then  $m | e$ . For, according to [2], there exists a cocycle  $a$  of  $G = 2^4$  with  $o([a]) = 2$ , but  $c(G, a) = 4$ . In other words, the corresponding central extension of the elementary abelian group  $G$ , an extraspecial group of order 32, has an unique ordinary irreducible nonlinear character, which is of a degree 4.

Now, we show that to analyse  $c(G, a)$  we should consider the prime divisors of  $o([a])$  and  $s(P, \alpha_P)$  for the corresponding Sylow subgroups  $P$  of  $G$ .

PROPOSITION 1. *Let  $c = c(G, a)$ . Then, the  $p$ -th part of  $c$ ,  $c_p$ , is equal to  $s(P, \alpha_P)$  for  $P$  a Sylow  $p$ -subgroup of  $G$ .*

PROOF. Let  $P \in \text{Syl}_p(G)$  and  $\text{Proj}(P, \alpha_P) = \{\gamma_1, \gamma_2, \dots, \gamma_r\}$ . Now, let  $\tau \in \text{Proj}(G, \alpha)$  such that  $(\tau(1))_p = c_p$ . Then,  $\tau_P = \sum_{j=1}^r b_j \gamma_j$ , where the  $b_j$ 's are non-negative integers so that

$$c_p = s(P, \alpha_P) \left( \sum_{j=1}^r b_j (\gamma_j(1) / s(P, \alpha_P)) \right)_p.$$

Hence,  $s(P, \alpha_P) \mid c_p$ .

On the other hand, let  $\gamma \in \text{Proj}(P, \alpha_P)$  be such that  $\gamma(1) = s(P, \alpha_P)$ . Then,  $\gamma^G = \sum_{i=1}^t a_i \tau_i$  for some non-negative integers  $a_i$ , and so, comparing the  $p$ -th parts of the degrees, we obtain

$$s(P, \alpha_P) = c_p \left( \sum_{i=1}^t a_i (\tau_i(1) / c) \right)_p.$$

Hence,  $c_p \mid s(P, \alpha_P)$ . ■

Thus, we are left with the task of describing  $s(P, \alpha_P) = c(P, \alpha_P)$  for  $P \in \text{Syl}_p(G)$ . However, we shall actually consider a more general situation than this. Recall that  $\tau \in \text{Proj}(G, \alpha)$  is called *monomial*, if it is induced from a projective character of degree 1 of a subgroup, and  $G$  is said to be a *PM-group* if all its irreducible projective characters are monomial.

PROPOSITION 2. *Let  $M$  be a subgroup of  $G$  of minimal index such that  $[a_M] = [1, G]$ . Then, we have:*

- (a)  $s(G, \alpha) \leq [G: M]$  and  $c(G, \alpha) \mid [G: M]$ ;
- (b) if  $c(G, \alpha) = [G: M]$ , then  $c(G, \alpha) = s(G, \alpha)$ ;
- (c)  $s(G, \alpha) = [G: M]$  if and only if there exists a monomial character  $\tau \in \text{Proj}(G, \alpha)$  with  $\tau(1) = s(G, \alpha)$ .

PROOF. Let  $\tau' \in \text{Proj}(G, \alpha)$  such that  $\tau'(1) = s(G, \alpha)$ , and  $\lambda \in \text{Proj}(M, \alpha_M)$  with  $\lambda(1) = 1$ . Then,  $\lambda^G = \sum_{i=1}^t a_i \tau_i$ , for some non-negative integers  $a_i$ , and so

$$(1) \quad \lambda^G(1) = [G: M] = c(G, \alpha) \left( \sum_{i=1}^t a_i (\tau_i(1) / c(G, \alpha)) \right) \geq \tau'(1),$$

proving (a). Since  $c(G, \alpha) \mid s(G, \alpha)$ , we have that (b) is immediate from (a).

Now, suppose that equality holds in (1). Then, we must have that  $\lambda^G$  is irreducible. Conversely, if  $\tau \in \text{Proj}(G, \alpha)$  is monomial and  $\tau(1) = s(G, \alpha)$ , then, by definition, there exists a subgroup  $L$  of  $G$  and  $\mu \in \text{Proj}(L, \alpha_L)$  with  $\mu(1) = 1$  such that  $\mu^G = \tau$ . Obviously, then  $[\alpha_L] = [1, G]$ , from Lemma 1(a). Also,  $[G : L] = s(G, \alpha) \leq [G : M]$ , by (a), and, hence, by hypothesis,  $[G : L] = [G : M]$ . ■

Of course, equality in Proposition 2(c) does occur when  $G$  is a *PM*-group and, in particular, when  $G$  is supersolvable (see [1, 6.5.11]). However, if  $G = A_4$ ,  $o([\alpha]) = 2$ , then  $s(G, \alpha) = c(G, \alpha) = 2$ , but  $A_4$  has no subgroup of index 2, so that equality does not always hold.

The proof of the main theorem is now yielded by the above remarks in conjunction with Propositions 1 and 2.

We mention three applications of the above results.

**COROLLARY 1.** *Let  $L$  be a cyclic subgroup of  $G$ . Then,  $s(G, \alpha) \leq [G : L]$  and  $c(G, \alpha) \mid [G : L]$  for all cocycles  $\alpha$  of  $G$ .*

**PROOF.** Since  $L$  is cyclic,  $M(L)$  is trivial, and, hence,  $[\alpha_L] = [1, G]$  for all cocycles  $\alpha$  of  $G$ . Thus, the result is immediate, from Proposition 2(a). ■

Now, we show that a slightly weaker version of Proposition 2(a) gives an alternative proof for the final assertion of [1, 4.1.9].

**COROLLARY 2.** *Let  $e$  denote the exponent of  $M(G)$ ,  $\alpha$  be a cocycle of  $G$  with  $o([\alpha]) = e$ , and  $L$  be a subgroup of  $G$  such that  $[\alpha_L] = [1, G]$ . Then,  $e \mid [G : L]$ . In particular,  $e$  divides the index of each cyclic subgroup of  $G$ .*

**PROOF.** By Lemma 1(a) and Proposition 2(a), we have  $e \mid c(G, \alpha) \mid [G : L]$ . ■

Finally, the following type of result is useful in constructing the projective representations of a given group with specified Sylow structure.

COROLLARY 3. *Let  $\alpha$  be a cocycle of  $G$  with  $2 \mid o([\alpha])$ , and suppose that  $G$  has a dihedral Sylow 2-subgroup. Then,  $(c(G, \alpha))_2 = 2$ .*

PROOF. Let  $P \in \text{Syl}_2(G)$ . The restriction mapping from  $\text{Syl}_2(M(G))$  into  $M(P)$  is a monomorphism. Hence, since  $\mathcal{P}$  has a cyclic subgroup of index 2, we have, by Proposition 1 and Corollary 1, that  $(c(G, \alpha))_2 = s(P, \alpha_P) = 2$ . ■

*Acknowledgement.* I wish to thank the referee for his kindness and interest concerning this paper.

#### REFERENCES

- [1] G. KARPILOVSKY, *Projective Representations of Finite Groups*, Monographs and Textbooks in Pure and Applied Mathematics, 94, Marcel Dekker (1985).
- [2] A. O. MORRIS, *Projective representations of abelian groups*, J. London Math. Soc., 7 (1973), pp. 235-238.

Manoscritto pervenuto in redazione il 4 giugno 1997.