

# SÉMINAIRE DE PROBABILITÉS (STRASBOURG)

P. MC GILL

## A direct proof of the Ray-Knight theorem

*Séminaire de probabilités (Strasbourg), tome 15 (1981), p. 206-209*

<[http://www.numdam.org/item?id=SPS\\_1981\\_\\_15\\_\\_206\\_0](http://www.numdam.org/item?id=SPS_1981__15__206_0)>

© Springer-Verlag, Berlin Heidelberg New York, 1981, tous droits réservés.

L'accès aux archives du séminaire de probabilités (Strasbourg) (<http://portail.mathdoc.fr/SemProba/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques  
<http://www.numdam.org/>*

A DIRECT PROOF OF THE RAY-KNIGHT THEOREM

P. Mc GILL (\*)

Of the several known proofs of the Ray-Knight theorem, the martingale stopping argument of ([3], problem 5, p. 74) and [2] is arguably the most elementary. Here we exploit what is essentially the same idea to give a proof which avoids explicit computations.

Let  $(B_t)_{t \geq 0}$  be a Brownian motion started at zero. Its local time  $L_t^a$  is given by the Doob-Meyer decomposition

$$|B_t - a| = |a| + L_t^a + \beta_t^a$$

where  $\beta_t^a$  is a Brownian motion. The occupation density formula [1] gives

$$\int_0^t g(B_s) ds = \int_R g(a) L_t^a da.$$

Here  $g$  is a bounded Borel function and we always assume that  $L_t^a$  is the jointly continuous version. In the following,  $T$  will always denote the stopping time :  $\inf\{t / B_t = 1\}$ .

Now define a process  $(z_a)_{a \geq 0}$  as the unique (positive) solution of the S.D.E

$$z_a = 2 \int_0^a \sqrt{|z_b|} d\beta_b + 2 \int_0^a 1_{\{0 \leq b \leq 1\}} db$$

Here  $\beta_a$  is a Brownian motion and by [8] this equation has a unique  $\beta_a$  adapted solution.  $(z_a)_{a \geq 0}$  is a diffusion. Let

$$m_0 = \inf\{a > 0 : z_a = 0\}.$$

As remarked in [9],  $(z_a)_{0 \leq a \leq 1}$  is equivalent in law to  $(X_t)_{0 \leq t \leq 1}$  where  $X_t$  is a  $BES^2(2)$  process. Therefore, by path continuity,  $m_0$  is greater than one almost surely.

Lemma 1 :  $m_0$  is finite almost surely.

Proof : For  $\alpha > 0$ , let  $u$  be the decreasing (strictly positive on  $R^+$ ) solution of

$$2a \frac{d^2 u}{da^2} = \alpha u$$

---

(\*)

Dept. of Mathematics, New University of Ulster, Coleraine, N. Ireland

We can take  $u(a) = \sqrt{2\alpha a} K_1(\sqrt{2\alpha a})$  where  $K_1$  is the modified Bessel function so that (see [6] 5.7.12)  $\lim_{a \rightarrow 0} u(a) = 1$ . Now for  $0 < \varepsilon < c$  if

$m_\varepsilon^1 = \inf\{b \geq 1 ; z_b = \varepsilon\}$ , we can use Itô's formula to check that

$$\mathbb{E}\left[\exp\left\{-\alpha m_\varepsilon^1\right\} ; c < z_1\right] = \mathbb{E}\left[\frac{u(z_1)}{u(\varepsilon)} ; c < z_1\right]$$

is a bounded martingale. By stopping at  $m_\varepsilon^1$ , we get

$$\mathbb{E}\left[\exp\left\{-\alpha m_\varepsilon^1\right\} ; c < z_1\right] = e^{-\alpha} \mathbb{E}\left[\frac{u(z_1)}{u(\varepsilon)} ; c < z_1\right]$$

Letting  $\varepsilon \rightarrow 0$ ,  $\alpha \rightarrow 0$  and  $c \rightarrow 0$  we find that  $P[m_0^1 < +\infty] = 1$ .

The next result is well-known.

Lemma 2 : Let  $g \geq 0$  be continuous with compact support on  $\mathbb{R}$ . The equation

$$f'' = 2fg ; f'(-\infty) = 0 ; f(0) = \delta > 0$$

has a unique (strictly positive, convex, increasing) solution.

Theorem (Ray [7], Knight [5]) : The process  $(L_T^{1-a}, a \geq 0)$  has the same law as  $(z_a, a \geq 0)$ .

Proof : We show that for every continuous function  $g \geq 0$  with compact support in  $(-\infty, 1)$ ,

$$\mathbb{E}\left[\exp\left\{-\int_{-\infty}^1 g(a) L_T^a da\right\}\right] = \mathbb{E}\left[\exp\left\{-\int_0^\infty g(1-a) z_a da\right\}\right]$$

To do this, we use martingale stopping to calculate a more explicit form for each side of this equation.

L.H.S. : By Lemma 2, find  $f$  with

$$f'' = 2gf ; f'(-\infty) = 0 ; f(0) = 1$$

By convexity, the martingale

$$f(B_t) \exp\left\{-\int_0^t g(B_s) ds\right\}$$

is uniformly integrable up to time  $T$  so by the occupation density formula we get

$$\mathbb{E}\left[\exp\left\{-\int_{-\infty}^1 g(a) L_T^a da\right\}\right] = \frac{1}{f(1)}$$

R.H.S. : By Lemma 2 choose  $v$  with

$$v'' = 2g(1-a)v ; v(1) = 1 ; v'(+\infty) = 0$$

Then  $v(a) = f(1-a)$  for  $a \geq 0$  and by Itô's formula

$$\frac{1}{v(a \wedge 1)} \exp\left\{z_a \frac{v'}{2v}(a) - \int_0^a g(1-b) z_b db\right\}$$

is a local martingale. It is uniformly integrable since  $v' \leq 0$  hence by stopping at  $m_0$  we have (since  $z_0 = 0$ ,  $m_0 > 1$ )

$$E\left[\exp\left\{-\int_0^\infty g(1-b) z_b db\right\}\right] = \frac{1}{v(0)}$$

This completes the proof.

Final Remark : The above method applies equally well to any suitable diffusion  $X_t$  with generator

$$G_f = \frac{1}{2} \sigma^2(x) \frac{d^2}{dx^2} + \tau(x) \frac{d}{dx}$$

In this case we choose  $f$  such that

$$\frac{1}{2} \sigma^2 f'' + \tau f' = fg \sigma^2 ; \quad f(0) = 1 ; \quad f'(-\infty) = 0$$

and we replace  $z_a$  by the solution of

$$W_a = 2 \int_0^a \sqrt{|W_b|} d\beta_b + 2 \int_0^a \left[ 1_{\{0 \leq b \leq 1\}} - W_b \frac{\tau(b)}{\sigma^2(b)} \right] db$$

The argument now proceeds as before. See [4], Proposition 5.

Acknowledgement : I wish to thank T. Jeulin for his invaluable help with both the proofs and the presentation.

REFERENCES :

- [1] J. AZEMA and M. YOR : "En guise d'introduction". Soc. Math. France Astérisque 52-53, 3-16, (1978).
- [2] J. AZEMA and M. YOR : "Une solution simple au problème de Skorokhod". Sem. Probab. XIII, Lecture Notes in Mathematics 721, 90-115, Springer (1979).
- [3] K. ITO and H.P. Mc KEAN : "Diffusion Processes and their Sample Paths". Springer (1965).
- [4] T. JEULIN and M. YOR : "Autour d'un théorème de Ray". Soc. Math. France Astérisque 52-53, 145-158, (1978).
- [5] F. KNIGHT : "Random Walks and a sojourn density of Brownian motion". TAMS 109, 56-86, (1963).
- [6] N. LEBEDEV : "Special functions and their applications". Prentice Hall (1965).
- [7] D. RAY : "Sojourn times of diffusion processes". Ill. Journ. Math. 7, 615-630, (1963).
- [8] T. YAMADA and S. WATANABE : "On the uniqueness of solutions of stochastic differential equations". J. Math. Kyoto Univ. 11, 155-167, (1971).
- [9] T. SHIGA and S. WATANABE : "Bessel diffusions as a one-parameter family of diffusion processes". Z. für Wahr., 27, 37-46, (1973).