

# SÉMINAIRE DE PROBABILITÉS (STRASBOURG)

YAO-ZHONG HU

## **A unified approach to several inequalities for gaussian and diffusion measures**

*Séminaire de probabilités (Strasbourg)*, tome 34 (2000), p. 329-335

[http://www.numdam.org/item?id=SPS\\_2000\\_\\_34\\_\\_329\\_0](http://www.numdam.org/item?id=SPS_2000__34__329_0)

© Springer-Verlag, Berlin Heidelberg New York, 2000, tous droits réservés.

L'accès aux archives du séminaire de probabilités (Strasbourg) (<http://portail.mathdoc.fr/SemProba/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

# A Unified Approach to Several Inequalities for Gaussian and Diffusion Measures

Yaozhong Hu

Department of Mathematics, University of Kansas  
405 Snow Hall, Lawrence, KS 66045-2142

## Abstract

This paper presents a simple unified approach to several inequalities for Gaussian and diffusion measures. They include hypercontractive inequalities, logarithmic Sobolev inequalities, FKG inequalities, and correlation inequalities.

## 1 Introduction

This paper is concerned with several inequalities associated with Gaussian measures and measures generated by diffusions. We are mainly interested in hypercontractive inequalities, Poincaré inequalities (spectral gap inequalities), logarithmic Sobolev inequalities, FKG inequalities, and correlation inequalities. A unified approach is developed to produce all these inequalities at the same time. This approach is the so-called semigroup approach [9], [10], [6]. However, one usually obtains hypercontractive inequalities from logarithmic Sobolev inequalities. An interesting point of this paper is to illustrate that the hypercontractive inequalities can be obtained in a somewhat simpler way. We prove hypercontractivity by differentiating an auxiliary function once and (while if we differentiate it twice we obtain the logarithmic Sobolev inequalities). This approach is natural since it is well-known that the logarithmic Sobolev inequality is the infinitesimal form of hypercontractivity. If we are only interested in Gaussian measures, our approach for hypercontractivity is as simple as the famous Neveu approach ([11], see also [7]).

## 2 Hypercontractivity for Diffusions

Let  $(E, \mathcal{E}, \mu)$  be a measure space and let  $L^2(\mu)$  be the set of all square integrable functions from  $E$  to  $\mathbb{R}$ . Let  $L$  be the Markov generator associated with a continuous Markov semigroup  $P_t$  in  $L^2(\mu)$ . Denote  $\mu(f) = \int_E f(x)\mu(dx)$ .

**Assumptions:**  $\mu$  is invariant with respect to  $L$  and  $(P_t)_{t \geq 0}$ . There is a nice algebra  $\mathcal{A}$  of bounded functions on  $E$  which is dense in the  $L^2$  domain of  $L$ , stable by  $L$ , by  $P_t$  and by the action of composition with  $C^\infty$  real functions which are 0 at 0. Let also  $\mathcal{A}$  be dense in  $L^p(\mu)$  for any  $1 < p < \infty$ .  $P_t$  is ergodic in the sense that for any  $f$ ,  $P_t f \rightarrow \mu(f)$   $\mu$ -almost surely.

Following P.-A. Meyer, we introduce the so-called ‘‘carré du champ’’ operator  $\Gamma$  as the symmetric bilinear form on  $\mathcal{A}$  defined by

$$\Gamma(f, g) = \frac{1}{2} \{L(fg) - fLg - gLf\}, \quad \forall f, g \in \mathcal{A}. \quad (2.1)$$

Denote  $\Gamma(f) = \Gamma(f, f)$ . It is well-known that  $\Gamma$  is positive-definite.

We shall consider a Markov semigroup whose generator is a diffusion in the following sense:

For every  $C^\infty$  function  $\psi$  on  $\mathbb{R}^n$ , and for every finite family  $F = (f_1, \dots, f_n)$ , where  $f_1, \dots, f_n \in \mathcal{A}$ ,

$$L\psi(F) = \sum_{i=1}^n \frac{\partial \psi}{\partial x_i}(f_1, \dots, f_n) L(f_i) + \sum_{i,j=1}^n \frac{\partial^2 \psi}{\partial x_i \partial x_j}(f_1, \dots, f_n) \Gamma(f_i, f_j). \quad (2.2)$$

The iterated ‘‘carré du champ’’ operator is defined as

$$\Gamma_2(f, g) := \frac{1}{2} \{L\Gamma(f, g) - \Gamma(f, Lg) - \Gamma(g, Lf)\}, \quad \forall f, g \in \mathcal{A}. \quad (2.3)$$

Let  $\text{Rg}(f)$  denote the closure of the range of  $f$ .

The main result of this work is the following statement.

**Theorem 2.1** *Let  $R, \tau$  be fixed real numbers. Let  $\phi$  and  $\psi$  be  $C^\infty$  functions on an open interval  $J \subset \mathbb{R}$ . If  $\Gamma_2(f) \geq R\Gamma(f)$ ,  $\forall f \in \mathcal{A}$ , and if  $\phi(x) \geq 0$ ,  $\psi(x) \geq 0$ ,  $\phi''(x) \leq 0$ ,  $\psi''(x) \leq 0$ , and*

$$(e^{-R\tau} \phi'(x) \psi'(y))^2 \leq \phi(x) \phi''(x) \psi(y) \psi''(y), \quad \forall x, y \in J, \quad (2.4)$$

then for all  $f, g \in \mathcal{A}$  such that  $\text{Rg}(P_t(f)) \subset J$ ,  $\text{Rg}(P_t(g)) \subset J$ ,

$$\mu \{ \psi(g) P_\tau(\phi(f)) \} \leq \phi(\mu(f)) \psi(\mu(g)). \quad (2.5)$$

*Proof* It is easy to see that for every such  $\phi$ , and  $f \in \mathcal{A}$  with  $\text{Rg}(f) \subset J$ ,  $\phi(f)$  is well-defined and (2.2) holds. Denote  $F_t = P_t f$  and  $G_t = P_t g$ . For any  $T > 0$ , introduce an auxiliary function of  $t$ ,

$$h_t := P_{T-t} \{ [P_\tau(\phi(F_t))] \psi(G_t) \}, \quad 0 \leq t \leq T.$$

Let us compute its derivative. Let  $L_t = L - \frac{\partial}{\partial t}$ . By (2.2) it is easy to check that for any  $f_t, g_t$  defined on  $\mathbb{R}_+ \times E$ ,

$$L_t \phi(f_t) = \phi''(f_t) \Gamma(f_t), \quad L_t(f_t g_t) = f_t L_t g_t + g_t L_t f_t + 2\Gamma(f_t, g_t), \quad (2.6)$$

Set  $\Phi = \phi(F_t)$ ,  $\Psi = \psi(G_t)$ ,  $\Phi' = \phi'(F_t)$ ,  $\Psi' = \psi'(G_t)$ ,  $\Phi'' = \phi''(F_t)$ , and  $\Psi'' = \psi''(G_t)$ . By (2.6),

$$\begin{aligned} \frac{dh_t}{dt} &= -P_{T-t} \{L_t [(P_\tau(\Phi))\Psi]\} \\ &= -P_{T-t} \{L_t (P_\tau(\Phi)) \Psi + P_\tau(\Phi)L_t (\Psi) + 2\Gamma(P_\tau(\Phi), \Psi)\} \\ &= -P_{T-t} \{P_\tau[\Phi''\Gamma(F_t)] \Psi + P_\tau[\Phi]\Psi''\Gamma(G_t) + 2\Gamma(P_\tau(\Phi), \Psi)\}. \end{aligned} \tag{2.7}$$

Since  $\Gamma$  is positive definite, we have that

$$|\Gamma(P_\tau(\Phi), \Psi)|^2 \leq \Gamma(P_\tau(\Phi))\Gamma(\Psi).$$

It is proved in [1] (page 149) that if  $\Gamma_2(f) \geq R\Gamma(f)$  for all  $f \in \mathcal{A}$ , then

$$\sqrt{\Gamma(P_T f)} \leq e^{-RT} P_T \sqrt{\Gamma(f)}, \quad \forall f \in \mathcal{A}. \tag{2.8}$$

Thus we have

$$\begin{aligned} |\Gamma(P_\tau(\Phi), \Psi)| &\leq \sqrt{P_\tau[\Gamma(\Phi)]}\sqrt{\Gamma(\Psi)} \leq e^{-R\tau} P_\tau \sqrt{\Gamma(\Phi)}\sqrt{\Gamma(\Psi)} \\ &= e^{-R\tau} P_\tau \left[ |\Phi'| \sqrt{\Gamma(F_t)} \right] |\Psi'| \sqrt{\Gamma(G_t)}. \end{aligned}$$

The last inequality follows from  $\Gamma(\phi(f)) = \phi'(f)^2\Gamma(f)$ , which is an easy consequence of (2.1) and (2.2). Let us then introduce the operator  $\tilde{P}_\tau$  acting on  $\mathcal{A} \times \mathcal{A}$  by  $\tilde{P}_\tau(f \times g) = P_\tau(f)(x)g(x)$  (where  $(f \times g)(x, y) = f(x)g(y)$ ,  $x, y \in E$ ). Since  $P_\tau$  is positivity preserving, the same is true for  $\tilde{P}_\tau$ . Then we have

$$\begin{aligned} \frac{dh_t}{dt} &\geq -P_{T-t} \{ \tilde{P}_\tau[\Phi''\Gamma(F_t) \times \Psi + \Phi \times \Psi''\Gamma(G_t)] \\ &\quad + 2e^{-R\tau} |\Phi'| |\Psi'| \sqrt{\Gamma(F_t)} \times \sqrt{\Gamma(G_t)} \}. \end{aligned} \tag{2.9}$$

If (2.4) holds, then  $\frac{dh_t}{dt} \geq 0$ . Thus  $h_0 \leq h_T$ . Computing  $h_T$  and  $h_0$ , we obtain the inequality

$$P_T \{ (P_\tau(\phi(f)))\psi(g) \} \leq P_\tau(\phi(P_T(f)))\psi(P_T g). \tag{2.10}$$

Letting  $T \rightarrow \infty$ , the theorem is proven. □

This theorem is a generalization of the hypercontractivity theorem. Set  $\|f\|_p^p = \mu(|f|^p)$ .

**Corollary 2.2** *Let  $p > 1$  and  $q > 1$ . If  $\Gamma_2(f) \geq R\Gamma(f)$  and if  $e^{-2R\tau} \leq \frac{p-1}{q-1}$ , then  $\|P_\tau f\|_q \leq \|f\|_p$ ,  $\forall f \in \mathcal{A}$ .*

*Proof* Let us take  $J_\varepsilon = (-\varepsilon/2, +\infty)$ ,  $\phi_\varepsilon(x) = |x + \varepsilon|^{1/p}$ , and  $\psi_\varepsilon(y) = |y + \varepsilon|^{1/q'}$  with  $1/q + 1/q' = 1$ . Therefore (2.4) is true for  $\phi_\varepsilon$  and  $\psi_\varepsilon$  if and only if  $e^{-2R\tau} \leq (p-1)/(q-1)$ . Thus (2.5) holds for  $\phi_\varepsilon$  and  $\psi_\varepsilon$ . Letting  $\varepsilon \rightarrow 0$ , we obtain that for all  $f, g \in \mathcal{A}$  and  $f, g \geq 0$ ,

$$\mu \left\{ P_\tau(f^{1/p}g^{1/q'}) \right\} \leq (\mu(f))^{1/p}(\mu(g))^{1/q'}.$$

In the above inequality replacing  $f$  by  $(f^2 + \varepsilon)^{p/2} - \varepsilon^{p/2}$  and  $g$  by  $(g^2 + \varepsilon)^{q'/2} - \varepsilon^{q'/2}$  and then letting  $\varepsilon \rightarrow 0$ , we get

$$\mu((P_\tau |f|)|g|) \leq (\mu(|f|^p))^{1/p} (\mu(|g|^{q'}))^{1/q'}, \quad \forall f, g \in \mathcal{A}.$$

The corollary is thus established. □

### 3 Logarithmic Sobolev Inequalities

It is not so easy to compute the second derivative of  $h_t$  introduced in the previous section. We shall compute it in the next section for a Gaussian measure. Here we take  $\psi(x) = x$  and  $\tau = 0$  and compute  $h'_t$ . In this case we have  $\frac{dh_t}{dt} = P_{T-t}\{\phi''\Gamma(F)\}$ . Differentiating this identity once more with respect to  $t$  and using (2.2) for  $\Psi(x, y) = \phi''(x)y$  and  $F_1 = F$  and  $F_2 = \Gamma(F)$ , we obtain

$$\begin{aligned} \frac{d^2 h_t}{dt^2} &= P_{T-t} \{L_t [\phi''\Gamma(F)]\} \\ &= P_{T-t} \left\{ \phi^{(4)}\Gamma(F)^2 + \phi''L_t\Gamma(F) + 2\phi^{(3)}\Gamma(F, \Gamma(F)) \right\} \\ &= P_{T-t} \left\{ \phi^{(4)}\Gamma(F)^2 + 2\phi''\Gamma_2(F) + 2\phi^{(3)}\Gamma(F, \Gamma(F)) \right\} \\ &= P_{T-t} \{ \text{Tr}(AB) + 2R\phi''\Gamma(F) \}, \end{aligned}$$

where  $A = \begin{pmatrix} \phi''/2 & \phi^{(3)} \\ \phi^{(3)} & \phi^{(4)} \end{pmatrix}$  and  $B = \begin{pmatrix} 4[\Gamma_2(F) - R\Gamma(F)] & \Gamma(F, \Gamma(F)) \\ \Gamma(F, \Gamma(F)) & \Gamma(F)^2 \end{pmatrix}$ .  $B$  is positive definite by [1], p.149. If (3.2) below holds,  $A$  is also positive definite. Thus we have

$$\frac{d^2 h_t}{dt^2} \geq 2RP_{T-t} [\phi''(F)\Gamma(F)] = 2R\frac{dh_t}{dt}. \tag{3.1}$$

This implies that the derivative of  $h_t - h'_t \frac{1 - e^{-2Rt}}{2R}$  is non-negative. Hence

$$h_T - h_0 \leq h'_T(1 - e^{-2RT})/(2R).$$

Thus we have

**Theorem 3.1** *Let  $\phi$  be a  $C^\infty$  function on some open interval  $J \subset \mathbb{R}$  satisfying*

$$\phi''(x) \geq 0, \quad \phi^{(4)}(x) \geq 0, \quad 2(\phi^{(3)}(x))^2 \leq \phi''(x)\phi^{(4)}(x), \quad \forall x \in J. \tag{3.2}$$

*Assume that  $\Gamma_2(f) \geq R\Gamma(f)$ ,  $\forall f \in \mathcal{A}$ . Then for every  $f \in \mathcal{A}$  such that  $\text{Rg}(P_t f) \subset J$ ,*

$$P_T(\phi(f)) \leq \phi(P_T(f)) + \frac{1 - e^{-2RT}}{2R} P_T(\phi''(f)\Gamma(f)), \quad \forall T > 0. \tag{3.3}$$

*Letting  $T \rightarrow \infty$ , we have*

$$\mu(\phi(f)) \leq \phi(\mu(f)) + \frac{1}{2R}\mu(\phi''(f)\Gamma(f)). \tag{3.4}$$

**Remark 1** In (3.4) letting  $\phi(x) = x^2$ , one gets the Poincaré inequality and letting  $\phi(x) = x \log x$  (using the argument of Corollary 2.2), one gets the logarithmic Sobolev inequality, see [6]. For the Ornstein-Uhlenbeck semigroup, (2.8) is obvious so that the preceding proof of hypercontractivity for Gaussian measures is indeed rather simple.

## 4 Gaussian Measures

For a Gaussian measure, we may also use the heat semigroup [8] instead of the Ornstein-Uhlenbeck semigroup. Let  $\mu$  be the standard Gaussian measure on  $\mathbb{R}^d$ . Let  $P_t$  be heat kernel semigroup associated with the standard Laplacian:  $\frac{\partial P_t}{\partial t} = \Delta P_t$  (We omit the factor 1/2 for simplicity). Let  $\Delta_t = \Delta - \frac{\partial}{\partial t}$ . Denote by  $\langle \cdot, \cdot \rangle$  the inner product of two vectors or the Hilbert-Schmidt product of two matrices. The same computation rule (2.2) applies to  $\Delta_t$  and moreover, for any functions  $f$  and  $g$  on  $\mathbb{R}^d$  (in this section,  $f$  and  $g$  always denote  $C^\infty$  functions with compact supports except otherwise stated), if  $F_t = P_t f$  and  $G_t = P_t g$  as before,

$$\begin{aligned} \Delta_t \langle \nabla F_t, \nabla G_t \rangle &= \langle \nabla(\Delta_t F_t), \nabla G_t \rangle + \langle \nabla F_t, \nabla(\Delta_t G_t) \rangle + \langle \nabla^2 F_t, \nabla^2 G_t \rangle \\ &= 2 \langle \nabla^2 F_t, \nabla^2 G_t \rangle. \end{aligned} \tag{4.1}$$

Fix  $T \geq 0$ . Consider  $h_t = P_{T-t} \{ \phi(F_t) \psi(G_t) \}$ ,  $0 \leq t \leq T$ . Then it is easy to see that

$$\begin{aligned} \frac{dh_t}{dt} &= -P_{T-t} \{ \phi''(F_t) \psi |\nabla F_t|^2 + \phi(F_t) \psi''(G_t) |\nabla G_t|^2 \\ &\quad + \phi'(F_t) \psi'(G_t) \langle \nabla F_t, \nabla G_t \rangle \}. \end{aligned}$$

If we take  $\phi(x) = \psi(x) = x$ , and if  $\nabla f \geq 0$  and  $\nabla g \geq 0$  component-wise, then we see  $h'_t \geq 0$ . Thus  $h_T \geq h_0$ . Thus we obtain in this way the classical FKG inequality for Gaussian measures.

**Proposition 4.1** *If  $\nabla f \geq 0$ ,  $\nabla g \geq 0$  component-wise, then*

$$\mu(f)\mu(g) \leq \mu(fg).$$

Now, compute the second derivative of  $h_t$  for general  $\phi$  and  $\psi$ . We have that

$$\begin{aligned} \frac{d^2 h_t}{dt^2} &= P_{T-t} \left\{ \Delta_t (\phi''(F_t) \psi(G_t) |\nabla F_t|^2) + \Delta_t (\phi(F_t) \psi''(G_t) |\nabla G_t|^2) \right. \\ &\quad \left. + 2 \Delta_t (\phi'(F_t) \psi'(G_t) \langle \nabla F_t, \nabla G_t \rangle) \right\}. \end{aligned} \tag{4.2}$$

Apply then (2.2) to  $\Psi = \phi''(x_1) \psi(x_2) x_3$  and use (4.1) to compute  $\Delta_t |\nabla F_t|^2$ . It follows that

$$\begin{aligned} \Delta_t (\phi''(F_t) \psi(G_t) |\nabla F_t|^2) &= 2 \phi'' \psi \langle \nabla^2 F_t, \nabla^2 F_t \rangle + \phi^{(4)} \psi |\nabla F_t|^4 \\ &\quad + \phi'' \psi'' |\nabla F_t|^2 |\nabla G_t|^2 + 2 \phi^{(3)} \psi' \langle \nabla F_t, \nabla G_t \rangle |\nabla F_t|^2 \\ &\quad + 4 \phi^{(3)} \psi \langle \nabla F_t \otimes \nabla F_t, \nabla^2 F_t \rangle \\ &\quad + 4 \phi'' \psi' \langle \nabla F_t \otimes \nabla G_t, \nabla^2 F_t \rangle. \end{aligned}$$

In a similar way we can compute the other terms in (4.2). If we introduce  $X^T = [X_1, X_2, X_3, X_4, X_5]$ , where

$$X_1 = \nabla F_t \otimes \nabla F_t, \quad X_2 = \nabla G_t \otimes \nabla G_t, \quad X_3 = \nabla F_t \otimes \nabla G_t, \\ X_4 = \nabla^2 F_t, \quad X_5 = \nabla^2 G_t,$$

then we can write

$$\frac{d^2 h_t}{dt^2} = \langle X, AX \rangle, \tag{4.3}$$

where  $A = B + C$ ,  $B$  is a  $5 \times 5$  matrix with  $B_{45} = B_{54} = 2\phi'\psi'$  and the other elements 0 and  $C$  is defined as

$$C = \begin{pmatrix} \phi^{(4)}\psi & 2\phi''\psi'' & 2\phi^{(3)}\psi' & 2\phi^{(3)}\psi & 2\phi''\psi' \\ 2\phi''\psi'' & \phi\psi^{(4)} & 2\phi'\psi^{(3)} & 2\phi'\psi'' & 2\phi\psi^{(3)} \\ 2\phi^{(3)}\psi' & 2\phi'\psi^{(3)} & 2\phi''\psi'' & 4\phi''\psi' & 4\phi'\psi'' \\ 2\phi^{(3)}\psi & 2\phi'\psi'' & 4\phi''\psi' & 2\phi''\psi & 0 \\ 2\phi''\psi' & 2\phi\psi^{(3)} & 4\phi'\psi'' & 0 & 2\phi\psi'' \end{pmatrix}. \tag{4.4}$$

If  $A$  is positive definite, then  $h_t'' \geq 0$ . Namely,  $h_0 \leq h_T - h_0'$ . Thus we proved

**Theorem 4.2** *If  $\phi$  and  $\psi$  are  $C^\infty$  functions on some open interval  $J \subset \mathbb{R}$  such that  $A$  is positive definite on  $J$ , then for all  $f, g$  such that  $\text{Rg}(P_t(f)) \subset J$ ,  $\text{Rg}(P_t(g)) \subset J$ ,*

$$\mu(\phi(f)\psi(g)) \leq \phi(\mu(f))\psi(\mu(g)) + \mu\left\{ \phi''(f)\psi(g)|\nabla f|^2 + \phi(f)\psi''(g)|\nabla g|^2 \right. \\ \left. + 2\phi'(f)\psi'(g)\langle \nabla f, \nabla g \rangle \right\}. \tag{4.5}$$

**Remark 2** *1) Take  $\phi(x) = \psi(x) = x$ . If  $f$  and  $g$  are convex functions and if  $\mu(\nabla f) = 0$  and  $\mu(\nabla g) = 0$ , then (4.5) implies the following correlation inequality [8]:*

$$\mu(fg) \geq \mu(f)\mu(g).$$

*2) Let  $\psi(x) \equiv 1$  and let  $\phi$  satisfy (3.2). Then it is easy to check that  $A$  is positive definite. Thus (4.5) implies*

$$\mu(\phi(f)) \leq \phi(\mu(f)) + \mu(\phi''(f)|\nabla f|^2). \tag{4.6}$$

*As we mentioned, this implies the Poincaré and the logarithmic Sobolev inequalities. In fact, if  $\phi(x) = x^2$ , then  $\phi$  satisfies (3.2). (4.6) becomes the Poincaré inequality*

$$\mu(|f|^2) \leq \mu(f)^2 + \mu(|\nabla f|^2).$$

*If  $\phi(x) = x \log^+ x$ , then  $\phi$  satisfies (3.2) (using the argument of Corollary 2.2). (4.6) implies the logarithmic Sobolev inequality*

$$\mu(f \log f) \leq \mu(f) \log \mu(f) + \mu\left(\frac{1}{f} |\nabla f|^2\right), \quad \forall f \geq 0.$$

## References

- [1] Bakry, D. Transformations de Riesz pour les semigroupes symétriques. Seconde partie: Etude sous la condition  $\Gamma_2 \geq 0$ , Séminaire de probabilités, XIX, 1983/84, 145-174, Lecture Notes in Math., 1123, Springer, Berlin-New York, 1985.
  - [2] Bakry, D. L'hypercontractivité et son utilisation en théorie des semigroupes. Lectures on probability theory (Saint-Flour, 1992), 1-114, Lecture Notes in Math., 1581, Springer, Berlin, 1994.
  - [3] Bakry, D.; Emery, M. Diffusions hypercontractives. Séminaire de probabilités, XIX, 1983/84, 177-206, Lecture Notes in Math., 1123, Springer, Berlin-New York, 1985.
  - [4] Bakry, D.; Ledoux, M. Lévy-Gromov's isoperimetric inequality for an infinite-dimensional diffusion generator. *Invent. math.* 123 (1996), 259-281.
  - [5] Davies, E. B.; Gross, L.; Simon, B. Hypercontractivity: a bibliographic review. *Ideas and methods in quantum and statistical physics (Oslo, 1988)*, 370-389, Cambridge Univ. Press, Cambridge, 1992.
  - [6] Driver, B. K.; and Hu, Y.Z. On heat kernel logarithmic Sobolev inequalities. *Stochastic analysis and applications (Powys, 1995)*, 189-200, World Sci. Publishing, River Edge, NJ, 1996.
  - [7] Dellacherie, C.; Meyer, P.-A.; Maisonneuve, B. *Probabilités et Potentiel. V.* Hermann, Paris, 1992.
  - [8] Hu, Y.Z. Itô-Wiener chaos expansion with exact residual and correlation, variance inequalities. *J. Theoret. Probab.* 10 (1997), no. 4, 835-848.
  - [9] Ledoux, M. L'algèbre de Lie des gradients itérés d'un générateur markovien - Développements de moyennes et entropies. *Ann. Sci. École Norm. Sup.* 28 (1995), no. 4, 435-460.
  - [10] Ledoux, M. Semigroup proofs of the isoperimetric inequality in Euclidean and Gauss space. *Bull. Sci. Math.* 118 (1994), 485-510.
  - [11] Neveu, J. Sur l'espérance conditionnelle par rapport à un mouvement brownien. *Ann. Inst. H. Poincaré Sect. B (N.S.)* 12 (1976), 105-109.
- Note: More complete references can be found in [2], [5].