

SÉMINAIRE DE PROBABILITÉS (STRASBOURG)

DAVAR KHOSHNEVISAN

ZHAN SHI

Fast sets and points for fractional brownian motion

Séminaire de probabilités (Strasbourg), tome 34 (2000), p. 393-416

http://www.numdam.org/item?id=SPS_2000__34__393_0

© Springer-Verlag, Berlin Heidelberg New York, 2000, tous droits réservés.

L'accès aux archives du séminaire de probabilités (Strasbourg) (<http://portail.mathdoc.fr/SemProba/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

FAST SETS AND POINTS FOR FRACTIONAL BROWNIAN MOTION

BY

DAVAR KHOSHNEVISAN* & ZHAN SHI
University of Utah & Université Paris VI

Summary. In their classic paper, S. Orey and S.J. Taylor compute the Hausdorff dimension of the set of points at which the law of the iterated logarithm fails for Brownian motion. By introducing “fast sets”, we describe a converse to this problem for fractional Brownian motion. Our result is in the form of a limit theorem. From this, we can deduce refinements to the aforementioned dimension result of Orey and Taylor as well as the work of R. Kaufman. This is achieved via establishing relations between stochastic co-dimension of a set and its Hausdorff dimension along the lines suggested by a theorem of S.J. Taylor.

Keywords and Phrases. Fast point, fast set, fractional Brownian motion, Hausdorff dimension.

A.M.S. 1991 Subject Classification. 60G15, 60G17, 60J65.

§1. INTRODUCTION

Suppose $W \triangleq (W(t); t \geq 0)$ is standard one-dimensional Brownian motion starting at 0. Continuity properties of the process W form a large part of classical probability theory. In particular, we mention A. Khintchine’s law of the iterated logarithm (see, for example, [21, Theorem II.1.9]): for each $t \geq 0$, there exists a null set $\mathcal{N}_1(t)$ such that for all $\omega \notin \mathcal{N}_1(t)$,

$$\limsup_{h \rightarrow 0^+} \frac{|W(t+h) - W(t)|}{\sqrt{h \ln \ln(1/h)}} = \sqrt{2}. \quad (1.1)$$

Later on, P. Lévy showed that $\cup_{t \geq 0} \mathcal{N}_1(t)$ is not a null set. Indeed, he showed the existence of a null set \mathcal{N}_2 outside which

$$\limsup_{h \rightarrow 0^+} \sup_{0 \leq r \leq 1} \frac{|W(r+h) - W(r)|}{\sqrt{h \ln(1/h)}} = \sqrt{2}. \quad (1.2)$$

* Research supported by a grant from the National Security Agency

See [13, p. 168] or [21, Theorem I.2.7], for example. It was observed in [18] that the limsup is actually a limit. Further results in this direction can be found in [2, p. 18]. The apparent discrepancy between (1.1) and (1.2) led S. Orey and S.J. Taylor to further study the so-called fast (or rapid) points of W . To describe this work, for all $\lambda > 0$, define $F_1(\lambda)$ to be the collection of all times $t \geq 0$ at which

$$\limsup_{h \rightarrow 0^+} \frac{|W(t+h) - W(t)|}{\sqrt{h \ln(1/h)}} \geq \lambda \sqrt{2}.$$

The main result of [18] is that with probability one,

$$\dim(F_1(\lambda)) = 1 - \lambda^2. \tag{1.3}$$

One can think of this as the multi-fractal analysis of white noise. Above and throughout, “ $\dim(A)$ ” refers to the Hausdorff dimension of A . Furthermore, whenever $\dim(A)$ is (strictly) negative, we really mean $A = \emptyset$. Orey and Taylor’s discovery of Eq. (1.3) relied on special properties of Brownian motion. In particular, they used the strong Markov property in an essential way. This approach has been refined in [3, 4, 11], in order to extend (1.3) in several different directions.

Our goal is to provide an alternative proof of Eq. (1.3) which is robust enough to apply to non-Markovian situations. We will do so by (i) viewing $F_1(\lambda)$ as a random set and considering its hitting probabilities; and (ii) establishing (within these proofs) links between Eqs. (1.2) and (1.3).

To keep from generalities, we restrict our attention to fractional Brownian motion. With this in mind, let us fix some $\alpha \in]0, 2[$ and define $X \triangleq (X(t); t \geq 0)$ to be a one-dimensional Gaussian process with stationary increments, mean zero and incremental standard deviation given by,

$$\|X(t) - X(s)\|_2 = |t - s|^{\alpha/2}.$$

See (1.8) for our notation on $L^p(\mathbb{P})$ norms.

The process X is called **fractional Brownian motion** with index α — hereforth written as $\text{fBM}(\alpha)$. We point out that when $\alpha = 1$, X is Brownian motion.

Let $\overline{\dim}_M(E)$ denote the upper Minkowski dimension of a Borel set $E \subset \mathbb{R}^1$; see references [17, 24]. Our first result, which is a fractal analogue of Eq. (1.2), is the following limit theorem:

Theorem 1.1. *Suppose X is $\text{fBM}(\alpha)$ and $E \subset [0, 1]$ is closed. With probability one,*

$$\limsup_{h \rightarrow 0^+} \sup_{t \in E} \frac{|X(t+h) - X(t)|}{h^{\alpha/2} \sqrt{\ln(1/h)}} \leq \sqrt{2 \overline{\dim}_M(E)}. \tag{1.4}$$

On the other hand, with probability one,

$$\sup_{t \in E} \limsup_{h \rightarrow 0^+} \frac{|X(t+h) - X(t)|}{h^{\alpha/2} \sqrt{\ln(1/h)}} \geq \sqrt{2 \dim(E)}. \tag{1.5}$$

Loosely speaking, when $\alpha = 1$, Theorem 1.1 is a converse to (1.3). For all $\lambda \geq 0$, define $\mathcal{F}_\alpha(\lambda)$ to be the collection of all closed sets $E \subset [0, 1]$ such that

$$\limsup_{h \rightarrow 0^+} \sup_{t \in E} \frac{|X(t+h) - X(t)|}{h^{\alpha/2} \sqrt{\ln(1/h)}} \geq \lambda \sqrt{2}.$$

One can think of the elements of $\mathcal{F}_\alpha(\lambda)$ as λ -fast sets. Theorem 1.1 can be recast in the following way.

Corollary 1.2. *Suppose X is fBM(α) and $E \subset [0, 1]$ is closed. If $\overline{\dim}_M(E) < \lambda^2$, then $E \notin \mathcal{F}_\alpha(\lambda)$ almost surely. On the other hand, if $\dim(E) \geq \lambda^2$, then $E \in \mathcal{F}_\alpha(\lambda)$.*

Remark 1.2.1. An immediate consequence of Theorem 1.1 is the following extension of (1.2):

$$\limsup_{h \rightarrow 0^+} \sup_{0 \leq t \leq 1} \frac{|X(t+h) - X(t)|}{h^{\alpha/2} \sqrt{\ln(1/h)}} = \sqrt{2}, \quad \text{a.s.}$$

When $\alpha \in]0, 1[$, this is a consequence of [15, Theorem 7]. When $\alpha \in]1, 2[$, the existence of such a modulus of continuity is mentioned in [16, Section 5].

A natural question is: can one replace E by a random set? The first random set that comes to our mind is the zero set. When $\alpha = 1$, the process is Brownian motion. Its Markovian structure will be used to demonstrate the following.

Theorem 1.3. *Suppose W is Brownian motion. Let $\mathcal{Z} \triangleq \{s \in [0, 1] : W(s) = 0\}$. Then, with probability one,*

$$\limsup_{h \rightarrow 0^+} \sup_{t \in \mathcal{Z}} \frac{|W(t+h)|}{\sqrt{h \ln(1/h)}} = \sup_{t \in \mathcal{Z}} \limsup_{h \rightarrow 0^+} \frac{|W(t+h)|}{\sqrt{h \ln(1/h)}} = 1.$$

Thus, the escape of Brownian motion from zero is slower than Lévy’s modulus (1.2).

Next, we come to dimension theorems; see (1.3) for an example. Define the λ -fast points for fBM(α) as follows:

$$F_\alpha(\lambda) \triangleq \left\{ t \in [0, 1] : \limsup_{h \rightarrow 0^+} \frac{|X(t+h) - X(t)|}{h^{\alpha/2} \sqrt{\ln(1/h)}} \geq \lambda \sqrt{2} \right\}. \quad (1.6)$$

In particular, $F_1(\lambda)$ denotes the λ -fast points of Brownian motion. In [9], R. Kaufman has shown that for any closed $E \subset [0, 1]$ and every $\lambda > 0$, with probability one,

$$\dim(E) - \lambda^2 \leq \dim(E \cap F_1(\lambda)). \quad (1.7)$$

Moreover, there exists a certain fixed closed set $E \subset [0, 1]$ for which the above is an equality. Our next aim is to show that the inequality in (1.7) is an equality for many sets $E \subset [0, 1]$, and that this holds for all fBM(α)'s. More precisely, we offer the following:

Theorem 1.4. *Suppose X is fBM(α), $E \subset [0, 1]$ is closed and $\lambda > 0$. Then, with probability one,*

$$\dim(E) - \lambda^2 \leq \dim(E \cap F_\alpha(\lambda)) \leq \dim_{\mathbb{P}}(E) - \lambda^2,$$

where $\dim_{\mathbb{P}}$ denotes packing dimension.

See [17] and [24] for definitions and properties of $\dim_{\mathbb{P}}$.

In particular, (1.3) holds for the fast points of any fractional Brownian motion. Moreover, when $E = \mathcal{Z}$ and $\alpha = 1$, we have the following dimension analogue of Theorem 1.3. Note that $F_1(\lambda)$ is the set of fast points of W as defined earlier. In other words, it is defined by (1.6) with X replaced by W .

Theorem 1.5. *Suppose W is Brownian motion and $\mathcal{Z} \triangleq \{s \in [0, 1] : W(s) = 0\}$. Then, for all $\lambda > 0$, with probability one,*

$$\dim(\mathcal{Z} \cap F_1(\lambda)) = \frac{1}{2} - \lambda^2.$$

A natural question which we have not been able to answer is the following:

Problem 1.5.1. *Does Theorem 1.5 have a general analogue for all fBM(α)'s?*

The proofs of Theorems 1.1, 1.3, 1.4 and 1.5 rely on parabolic capacity techniques and entropy arguments. The entropy methods follow the arguments of [18] closely. On the other hand, the parabolic capacity arguments rely on relationships between the Hausdorff dimension of random sets and stochastic co-dimension (see §2). The latter is a formalization of a particular application of [23, Theorem 4], which can be found in various forms within the proofs of [1, 7, 14, 19]. We suspect our formulation has other applications. In §3, we demonstrate (1.4) while (1.5) and the first inequality (i.e., the lower bound) of Theorem 1.4 are derived in §4. The proof of the upper bound of Theorem 1.3 appears in §5; the upper bounds of Theorems 1.4 and 1.5 can be found in §6 and §7, respectively; and the lower bounds for Theorems 1.3 and 1.5 are proved simultaneously in §8.

We conclude the Introduction by mentioning some notation which will be utilized throughout this article. Define the function ψ as

$$\psi(h) \triangleq \sqrt{2 \ln(1/h)}, \quad h \in]0, 1[.$$

By $\bar{\Phi}$ we mean the tail of a standard normal distribution, i.e.

$$\bar{\Phi}(x) \triangleq \frac{1}{\sqrt{2\pi}} \int_x^\infty e^{-u^2/2} du, \quad x \in]-\infty, \infty[.$$

Furthermore, $(\Omega, \mathcal{F}, \mathbb{P})$ denotes our underlying probability space. For any real variable Z on $(\Omega, \mathcal{F}, \mathbb{P})$ and for every $p > 0$ we write the $L^p(\mathbb{P})$ -norm of Z by,

$$\|Z\|_p \triangleq \left(\int_\Omega |Z(\omega)|^p \mathbb{P}(d\omega) \right)^{1/p}. \quad (1.8)$$

Throughout, X denotes fBM(α) for any $\alpha \in]0, 2[$. However, when we wish to discuss Brownian motion specifically (i.e., when $\alpha = 1$), we write W instead. In accordance with the notation of Theorem 1.3, \mathcal{Z} will always denote the zero set of W restricted to $[0, 1]$. Finally, the collection of all atomless probability measures on a set E is denoted by $\mathcal{P}_+(E)$.

Remark. Since the first circulation of this paper, many of the ‘gaps’ in the inequalities of this paper have been bridged. For instance, in Theorem 1.1, both constants of (1.4) and (1.5) can be computed. This and related material can be found in [10].

Acknowledgements. Much of this work was done while the first author was visiting Université Paris VI. We thank L.S.T.A. and Laboratoire de Probabilités for their generous hospitality. Our warmest thanks are extended to Steve Evans, Mike Marcus, Yuval Peres and Yimin Xiao. They have provided us with countless suggestions, references, corrections and their invaluable counsel. In particular, it was Yuval Peres who showed us the rôle played by packing dimensions as well as allowing us to use his argument (cf. [10]) in the proof of Theorem 2.5.

§2. PRELIMINARIES ON DIMENSION

In this section, we discuss a useful approach to estimating Hausdorff dimensions of random sets via intersection probabilities.

Let \mathbf{S}_0^1 denote the collection of all Borel measurable subsets of $[0, 1]$. We say that $E : \Omega \mapsto \mathbf{S}_0^1$ is a **random set**, if $\mathbf{1}_E(\omega) : \Omega \times \mathbf{S}_0^1 \ni (\omega, E) \mapsto \{0, 1\}$ is a random variable in the product measure space. An important class of random sets are the closed stochastic images $E \triangleq \overline{S[0, 1]} \triangleq \{S(t); t \in [0, 1]\}$, where S is a stochastic process with càdlàg sample paths.

Let us begin with some preliminaries on Hausdorff dimension; see [8, 16, 24] for definitions and further details. Given $s \geq 0$ and a Borel set $E \subset [0, 1]$, let $\Lambda^s(E)$ denote the s -dimensional Hausdorff measure of E . Recall that the **Hausdorff dimension** — $\dim(E)$ — of E is defined by: $\dim(E) \triangleq \inf \{s > 0 : \Lambda^s(E) < \infty\}$. When it is finite, $\Lambda^s(E)$ extends nicely to a Carathéodory outer

measure on analytic subsets of $[0, 1]$. By a slight abuse of notation, we continue to denote this outer measure by Λ^s .

Suppose E is a random set. Since we can economically cover E with intervals with rational endpoints, $\dim(E)$ is a random variable. We will use this without further mention.

Typically, computing upper bounds for $\dim(E)$ is not very difficult: find an economical cover (I_j) of E , whose diameter is h or less and compute $\sum_j |I_j|^s$. Obtaining good lower bounds for $\dim(E)$ is the harder of the two bounds. The standard method for doing this is to utilize the connections between Hausdorff dimension and potential theory. For any $\mu \in \mathcal{P}_+(E)$ and all $\beta > 0$, define,

$$A_\beta(\mu) \triangleq \sup_{0 < h \leq 1/2} \sup_{t \in [h, 1-h]} \frac{\mu[t-h, t+h]}{h^\beta}. \tag{2.1}$$

(It is possible that $A_\beta(\mu) = \infty$.) We need the following connection between Hausdorff dimension and potential theory; while it is only half of Frostman's lemma of classical potential theory, we refer to it as 'Frostman's lemma' for brevity.

Lemma 2.1. (Frostman's Lemma; [8, p. 130]) *Suppose $E \in \mathbf{S}_0^1$ satisfies $\beta < \dim(E)$. Then there exists $\mu \in \mathcal{P}_+(E)$ for which $A_\beta(\mu) < \infty$.*

Thus, a method for obtaining lower bounds on $\dim(E)$ is this: find a probability measure μ which lives on E and show that $A_\beta(\mu) < \infty$. If this can be done for some $\beta > 0$, then $\dim(E) \geq \beta$. In general, this is all which can be said. However, if the set E in question is a random set in the sense of the first paragraph of this section, there is an abstract version of [23, Theorem 4] which can be used to bound $\dim(E)$ from below; see also [1]. We shall develop this next. Define the **upper stochastic co-dimension** ($\overline{\text{co-dim}}$) of a random set E by

$$\overline{\text{co-dim}}(E) \triangleq \inf \{ \beta \in [0, 1] : \forall G \in \mathbf{S}_0^1 \text{ with } \dim(G) > \beta, \mathbb{P}(E \cap G \neq \emptyset) = 1 \}. \tag{2.2}$$

In order to make our definition sensible and complete, we need to define $\inf \emptyset \triangleq 1$.

Remark 2.1.1. In applications, we often need the following fact: if $G \in \mathbf{S}_0^1$ satisfies $\dim(G) > \overline{\text{co-dim}}(E)$, then $\mathbb{P}(E \cap G \neq \emptyset) = 1$.

In this section we present two results about stochastic co-dimension, the first of which is the following.

Theorem 2.2. *Suppose E is a random set. Then, for all $G \in \mathbf{S}_0^1$,*

$$\dim(E \cap G) \geq \dim(G) - \overline{\text{co-dim}}(E), \quad \text{a.s.}$$

As mentioned earlier, Theorem 2.2 is an abstract form of a part of [23, Theorem 4]. This kind of result has been implicitly used in several works. For

example, see [7, 14, 19, 20]. To prove it, let us introduce an independent symmetric stable Lévy process $S_\gamma \triangleq (S_\gamma(t); t \geq 0)$ of index $\gamma \in]0, 1[$. The following two facts are due to J. Hawkes; cf. [6].

Lemma 2.3. *Suppose $G \in \mathbf{S}_0^1$ satisfies $\dim(G) < 1 - \gamma$. Then, with probability one, $\overline{S_\gamma[0, 1]} \cap G = \emptyset$.*

Lemma 2.4. *Suppose $G \in \mathbf{S}_0^1$ satisfies $\dim(G) > 1 - \gamma$. Then, $\mathbb{P}(\overline{S_\gamma[0, 1]} \cap G \neq \emptyset) > 0$. Furthermore, on $(\overline{S_\gamma[0, 1]} \cap G \neq \emptyset)$,*

$$\dim(\overline{S_\gamma[0, 1]} \cap G) = \dim(G) + \gamma - 1, \quad \text{a.s.} \tag{2.3}$$

Historically, the above results are stated with $\overline{S_\gamma[0, 1]}$ replaced by $S_\gamma[0, 1]$. By symmetry, semi-polar sets are polar for S_γ . Therefore, the same facts hold for $\overline{S_\gamma[0, 1]}$.

We can now proceed with Theorem 2.2.

Proof of Theorem 2.2. Without loss of generality, we can assume that the compact set G satisfies $\dim(G) > \overline{\text{co-dim}}(E)$. With this reduction in mind, let us choose a number $\gamma \in]0, 1[$ satisfying,

$$\gamma > 1 - \dim(G) + \overline{\text{co-dim}}(E). \tag{2.4}$$

Choose the process S_γ as in the earlier part of this section. Since $\gamma > 1 - \dim(G)$, it follows from Lemma 2.4 that $\kappa \triangleq \mathbb{P}(\overline{S_\gamma[0, 1]} \cap G \neq \emptyset) > 0$. By (2.3),

$$\begin{aligned} \kappa &= \mathbb{P}(\overline{S_\gamma[0, 1]} \cap G \neq \emptyset, \dim(\overline{S_\gamma[0, 1]} \cap G) = \dim(G) + \gamma - 1) \\ &\leq \mathbb{P}(\overline{S_\gamma[0, 1]} \cap G \neq \emptyset, \dim(\overline{S_\gamma[0, 1]} \cap G) > \overline{\text{co-dim}}(E)), \end{aligned}$$

where we have used (2.4) in the last inequality. In view of Remark 2.1.1, κ is bounded above by $\mathbb{P}(\overline{S_\gamma[0, 1]} \cap G \cap E \neq \emptyset)$. Applying Lemma 2.3 gives

$$\begin{aligned} \kappa &\leq \mathbb{P}(\overline{S_\gamma[0, 1]} \cap G \cap E \neq \emptyset, \dim(G \cap E) \geq 1 - \gamma) \\ &\leq \mathbb{P}(\overline{S_\gamma[0, 1]} \cap G \neq \emptyset, \dim(G \cap E) \geq 1 - \gamma) \\ &= \kappa \mathbb{P}(\dim(G \cap E) \geq 1 - \gamma). \end{aligned}$$

The last line utilizes the independence of S_γ and E . Since $\kappa > 0$, it follows that for all γ satisfying (2.4), $\dim(G \cap E) \geq 1 - \gamma$, almost surely. Let $\gamma \downarrow 1 - \dim(G) + \overline{\text{co-dim}}(E)$ along a rational sequence to obtain the result. \diamond

Next, we present the second result of this Section. It is an immediate consequence of the estimates of [10, Section 3] and Theorem 2.2 above.

Theorem 2.5. *Suppose $(E_n; n \geq 1)$ is a countable collection of open random sets. If $\sup_{n \geq 1} \overline{\text{co-dim}}(E_n) < 1$, then*

$$\overline{\text{co-dim}}\left(\bigcap_{n=1}^{\infty} E_n\right) = \sup_{n \geq 1} \overline{\text{co-dim}}(E_n).$$

In particular, $\mathbb{P}(\bigcap_{n \geq 1} E_n \neq \emptyset) = 1$.

Informally speaking, this is a dual to the fact that for all $F_n \in \mathbf{S}_0^1$ ($n = 1, 2, \dots$), $\dim(\bigcup_{n=1}^{\infty} F_n) = \sup_{n \geq 0} \dim(F_n)$.

§3. THEOREM 1.1: UPPER BOUND

Define the set of “near-fast points” as follows: for all $\lambda, h > 0$,

$$F_\alpha(\lambda, h) \triangleq \{t \in [0, 1] : \sup_{t \leq s \leq t+h} |X(s) - X(t)| \geq \lambda h^{\alpha/2} \psi(h)\}. \tag{3.1}$$

Next, for any $R, \eta > 1$, all integers $j \geq 1$ and every integer $0 \leq m < R^{\eta j} + 1$, define

$$I_{m,j}^\eta \triangleq [mR^{-\eta j}, (m+1)R^{-\eta j}]. \tag{3.2}$$

Finally, define for the above parameters,

$$P_{m,j}^{\lambda,\alpha} \triangleq \mathbb{P}(I_{m,j}^\eta \cap F_\alpha(\lambda, R^{-j}) \neq \emptyset). \tag{3.3}$$

The main technical estimate which we shall need in this section is the following:

Lemma 3.1. *Let X be fBM(α), where $\alpha \in]0, 2[$. For all $\lambda > 0, \varepsilon \in]0, 1[, \eta > 1$ and all $R > 1$, there exists $J_1 = J_1(\varepsilon, \alpha, \eta, \lambda, R) \in [2, \infty[$ such that for all $j \geq J_1$ and all $m \geq 0$,*

$$P_{m,j}^{\lambda,\alpha} \leq R^{-\lambda^2(1-\varepsilon)j}.$$

Remark 3.1.1. Part of the assertion is that J_1 does not depend on the choice of m .

Proof. By stationarity and scaling,

$$P_{m,j}^{\lambda,\alpha} = \mathbb{P}\left(\sup_{0 \leq t \leq R^{-(1-\eta)j}} \sup_{0 \leq s \leq 1} |X(s+t) - X(s)| \geq \lambda \psi(R^{-j})\right).$$

We obtain the lemma by applying standard estimates and [12, Lemma 3.1] to the Gaussian process $(X(s+t) - X(t); s, t \geq 0)$. \diamond

The proof of the upper bound is close to its counterpart [18]; cf. the first part of Theorem 2 therein.

Proof of Theorem 1.1: upper bound. Recall (3.1) and (3.2). Consider a fixed closed set $E \subset [0, 1]$. Fix $R, \eta > 1$ and $\lambda > 0$. Define for all integers $m \geq 0$,

$$\mathcal{J}_k \triangleq \sum_{j \geq k} \sum_{m \geq 0} \mathbf{1}_{\{I_{m,j}^n \cap F_\alpha(\lambda, R^{-j}) \neq \emptyset\}} \mathbf{1}_{\{I_{m,j}^n \cap E \neq \emptyset\}}. \tag{3.4}$$

By (3.3) and Lemma 3.1, for all $\varepsilon \in]0, 1[$, there exists $J_1 = J_1(\varepsilon, \alpha, \eta, \lambda, R) \in [2, \infty[$ such that for all $k \geq J_1$,

$$\begin{aligned} \|\mathcal{J}_k\|_1 &= \sum_{j \geq k} \sum_{m \geq 0} P_{m,j}^{\lambda, \alpha} \mathbf{1}_{\{I_{m,j}^n \cap E \neq \emptyset\}} \leq \sum_{j \geq k} \sum_{m \geq 0} R^{-\lambda^2(1-\varepsilon)j} \mathbf{1}_{\{I_{m,j}^n \cap E \neq \emptyset\}} \\ &\leq \sum_{j \geq k} R^{-\lambda^2(1-\varepsilon)j} M(R^{-\eta j}; E), \end{aligned}$$

where $M(\varepsilon; E)$ denotes the ε -capacity of E . That is, it is the maximal number of points in E which are at least ε apart; see [5]. On the other hand, by definition, for all $\delta \in]0, 1[$, there exists $J_2 = J_2(\delta, R, \eta) \in [2, \infty[$, such that for all $j \geq J_2$, $M(R^{-\eta j}; E) \leq R^{\eta j(1+\delta)} \overline{\dim}_M(E)$. Hence, for all $k \geq J_1 \vee J_2$,

$$\|\mathcal{J}_k\|_1 \leq \sum_{j \geq k} R^{-j(\lambda^2(1-\varepsilon) - \eta(1+\delta) \overline{\dim}_M(E))}.$$

It may help to recall that $J_1 \vee J_2$ depends only on the parameters $(\lambda, \alpha, R, \eta, \delta, \varepsilon)$. Let us pick these parameters so that $\lambda^2(1 - \varepsilon) > \eta(1 + \delta) \overline{\dim}_M(E)$. It is easy to see that for this choice of parameters, $\sum_k \|\mathcal{J}_k\|_1 < \infty$. By the Borel–Cantelli lemma, with probability one, there exists a finite random variable k_0 such that for all $k \geq k_0$, $\mathcal{J}_k = 0$. In other words, with probability one, for all $j \geq k_0$, $F_\alpha(\lambda, R^{-j}) \cap E = \emptyset$. Rewriting the above, we see that with probability one, for all $j \geq k_0$, $\sup_{t \in E} \sup_{t \leq s \leq t+R^{-j}} |X(s) - X(t)| \leq \lambda R^{-j\alpha/2} \psi(R^{-j})$. Take any $h \leq R^{-k_0}$. There exists a $j \geq k_0$, so that $R^{-j-1} \leq h \leq R^{-j}$. It follows that for all $h \leq R^{-k_0}$,

$$\begin{aligned} \sup_{t \in E} |X(t+h) - X(t)| &\leq \sup_{t \in E} \sup_{t \leq s \leq t+R^{-j}} |X(s) - X(t)| \\ &\leq \lambda R^{-j\alpha/2} \psi(R^{-j}) \leq \lambda R^{\alpha/2} h^{\alpha/2} \psi(h), \end{aligned}$$

since ψ is monotone decreasing. This implies that whenever $\lambda^2(1 - \varepsilon) > \eta(1 + \delta) \overline{\dim}_M(E)$,

$$\limsup_{h \rightarrow 0^+} \sup_{t \in E} \frac{|X(t+h) - X(t)|}{h^{\alpha/2} \psi(h)} \leq \lambda R^{\alpha/2}, \quad \text{a.s.}$$

Along rational sequences, let $\varepsilon, \delta \rightarrow 0^+$, $\eta, R \rightarrow 1^+$ and $\lambda^2 \downarrow \overline{\dim}_M(E)$ — in this order — to see that with probability one,

$$\limsup_{h \rightarrow 0^+} \sup_{t \in E} \frac{|X(t+h) - X(t)|}{h^{\alpha/2} \psi(h)} \leq \sqrt{\overline{\dim}_M(E)}.$$



This proves the desired upper bound. ◇

§4. THEOREMS 1.1 AND 1.4: LOWER BOUNDS

For each closed set $E \subset [0, 1]$ and for every $\mu \in \mathcal{P}_+(E)$ and $h, \lambda > 0$, define,

$$I_\mu(h; \lambda) \triangleq \int_0^1 \mu(dt) \mathbf{1}_{\{X(t+h) - X(t) > \lambda h^{\alpha/2} \psi(h)\}}. \tag{4.1}$$

The key technical estimate of this section is the following:

Lemma 4.1. *Suppose $E \subset [0, 1]$ is compact, $\mu \in \mathcal{P}_+(E)$. For any $\varepsilon \in]0, 1[$ and $\lambda > 0$, there exists a small $h_0 \in]0, 1[$ such that for all $h \in]0, h_0[$,*

$$\frac{\|I_\mu(h; \lambda)\|_2^2}{\|I_\mu(h; \lambda)\|_1^2} \leq 1 + \varepsilon + 4 \sup_{h^{1-\varepsilon} \leq t \leq 1-h^{1-\varepsilon}} \frac{\mu([t - h^{1-\varepsilon}, t + h^{1+\varepsilon}])}{\overline{\Phi}(\lambda\psi(h))}.$$

Proof. Since X has stationary increments,

$$\|I_\mu(h; \lambda)\|_1 = \overline{\Phi}(\lambda\psi(h)). \tag{4.2}$$

We proceed with the estimate for the second moment. Define,

$$\begin{aligned} a &\triangleq \lambda\psi(h), \\ U &\triangleq \frac{X(s+h) - X(s)}{h^{\alpha/2}}, \\ V &\triangleq \frac{X(t+h) - X(t)}{h^{\alpha/2}}, \\ \rho &\triangleq \|UV\|_1. \end{aligned}$$

Then, ignoring the dependence on (h, s, t) ,

$$\|I_\mu(h; \lambda)\|_2^2 = Q_1 + Q_2 + Q_3, \tag{4.3}$$

where,

$$\begin{aligned} Q_1 &\triangleq \int_0^1 \mu(dt) \int_0^1 \mu(ds) \mathbf{1}_{\{\rho \leq (\ln(1/h))^{-2}\}} \mathbb{P}(U \geq a, V \geq a), \\ Q_2 &\triangleq 2 \int_0^1 \mu(dt) \int_0^t \mu(ds) \mathbf{1}_{\{\rho > (\ln(1/h))^{-2}\}} \mathbf{1}_{\{t-s > 2h\}} \mathbb{P}(U \geq a, V \geq a), \\ Q_3 &\triangleq 2 \int_0^1 \mu(dt) \int_0^t \mu(ds) \mathbf{1}_{\{\rho > (\ln(1/h))^{-2}\}} \mathbf{1}_{\{t-s \leq 2h\}} \mathbb{P}(U \geq a, V \geq a). \end{aligned}$$

We estimate each term separately. The critical term is Q_1 . Write $x^+ = \max(x, 0)$ for any x . If $\rho < 1/4$,

$$\begin{aligned} \mathbb{P}(U \geq a, V \geq a) &= \frac{1}{2\pi\sqrt{1-\rho^2}} \int_a^\infty \int_a^\infty \exp\left(-\frac{x^2+y^2-2\rho xy}{2(1-\rho^2)}\right) dx dy \\ &\leq \frac{1}{2\pi\sqrt{1-\rho^2}} \int_a^\infty \int_a^\infty \exp\left(-\frac{(1-4\rho^+)(x^2+y^2)}{2(1-\rho^2)}\right) dx dy \\ &= \frac{\sqrt{1-\rho^2}}{1-4\rho^+} \left(\bar{\Phi}\left(a\sqrt{\frac{1-4\rho^+}{1-\rho^2}}\right)\right)^2. \end{aligned}$$

According to Mill's ratio for Gaussian tails ([22, p. 850]), for any $x > 1$,

$$\frac{1-x^{-2}}{\sqrt{2\pi}x} \exp\left(-\frac{x^2}{2}\right) \leq \bar{\Phi}(x) \leq \frac{1}{\sqrt{2\pi}x} \exp\left(-\frac{x^2}{2}\right). \tag{4.4}$$

Therefore, using the fact that μ is an atomless probability measure, we have,

$$\begin{aligned} Q_1 &\leq \sup_{r: -1 < r \leq (\ln(1/h))^{-2}} \frac{\sqrt{1-r^2}}{1-4r^+} \left(\bar{\Phi}\left(\lambda\psi(h)\sqrt{\frac{1-4r^+}{1-r^2}}\right)\right)^2 \\ &\leq [\bar{\Phi}(\lambda\psi(h))]^2 \sup_{r: -1 < r \leq (\ln(1/h))^{-2}} \left(1 - \frac{1}{(\lambda\psi(h))^2}\right)^{-2} \frac{(1-r^2)^{3/2}}{(1-4r^+)^2} \times \\ &\quad \times \exp\left(\frac{(\lambda\psi(h))^2(4r^+ - r^2)}{1-r^2}\right). \end{aligned}$$

Since $(\psi(h))^2(\ln(1/h))^{-2} = o(1)$ (as h goes to 0), this leads to:

$$Q_1 \leq B(h; \lambda) [\bar{\Phi}(\lambda\psi(h))]^2, \tag{4.5}$$

where $B(h; \lambda)$ is such that, for any $\lambda > 0$,

$$\lim_{h \rightarrow 0^+} B(h; \lambda) = 1. \tag{4.6}$$

To estimate Q_3 , use the trivial inequality $\mathbb{P}(U \geq a, V \geq a) \leq \mathbb{P}(U \geq a)$, to see that

$$\begin{aligned} Q_3 &\leq 2 \sup_{2h \leq t \leq 1} \mu([t-2h, t]) \bar{\Phi}(\lambda\psi(h)) \\ &\leq 2 \sup_{2h \leq t \leq 1-2h} \mu([t-2h, t+2h]) \bar{\Phi}(\lambda\psi(h)). \end{aligned} \tag{4.7}$$

Finally, we need to approximate Q_2 . Directly computing, note that when $s < t - 2h$,

$$\begin{aligned} \rho &= \frac{|t-s-h|^\alpha + |t-s+h|^\alpha - 2|t-s|^\alpha}{2h^\alpha} \\ &= \frac{1}{2} \left(\frac{t-s}{h}\right)^\alpha \left[\left(1 - \frac{h}{t-s}\right)^\alpha + \left(1 + \frac{h}{t-s}\right)^\alpha - 2\right]. \end{aligned}$$

By Taylor expansion of $(1 \pm x)^\alpha$, we see that for all $|x| \leq \frac{1}{2}$,

$$(1 - x)^\alpha + (1 + x)^\alpha - 2 = \frac{\alpha(\alpha - 1)}{2} \left[(1 - \xi_1)^{\alpha-2} + (1 - \xi_2)^{\alpha-2} \right] x^2,$$

where $|\xi_i| \leq \frac{1}{2}$ for $i = 1, 2$. In particular, for all $|x| \leq \frac{1}{2}$,

$$\left| (1 - x)^\alpha + (1 + x)^\alpha - 2 \right| \leq 2^{3-\alpha} x^2.$$

In other words, when $s < t - 2h$, $\rho \leq \{2h/(t - s)\}^{2-\alpha}$. On the other hand, if we also know that $\rho > (\ln(1/h))^{-2}$, it follows that for all h small, $|t - s| \leq 2h(\ln(1/h))^{2/(2-\alpha)} \leq h^{1-\varepsilon}$. Since $\mathbb{P}(U \geq a, V \geq a) \leq \overline{\Phi}(\lambda\psi(h))$, we obtain the following: for all $\varepsilon > 0$, there exists $h_1 \in]0, 1[$ such that for all $h \in]0, h_1[$,

$$Q_2 \leq 2 \sup_{h^{1-\varepsilon} \leq t \leq 1-h^{1-\varepsilon}} \mu([t - h^{1-\varepsilon}, t + h^{1-\varepsilon}]) \overline{\Phi}(\lambda\psi(h)).$$

Together with (4.7), we obtain: for all $\varepsilon > 0$, there exists $h_2 \in]0, 1[$ such that for all $h \in]0, h_2[$,

$$Q_2 + Q_3 \leq 4 \sup_{h^{1-\varepsilon} \leq t \leq 1-h^{1-\varepsilon}} \mu([t - h^{1-\varepsilon}, t + h^{1-\varepsilon}]) \overline{\Phi}(\lambda\psi(h)).$$

Combining this with (4.2)–(4.3) and (4.5)–(4.7), we obtain the result. ◇

Now we can prove the lower bounds in Theorems 1.1 and 1.4.

Proof of Theorems 1.1 and 1.4: lower bounds. By Frostman’s lemma (Lemma 2.1), for any $\beta < \dim(E)$, there exists a $\mu \in \mathcal{P}_+(E)$, such that for all $h \in]0, 1[$ small,

$$\sup_{h \leq t \leq 1-h} \mu([t - h, t + h]) \leq h^\beta.$$

For such a μ , use Lemma 4.1 to see that for all $\varepsilon \in]0, 1[$ and $\lambda > 0$, there exists $h_3 \in]0, 1[$ such that whenever $h \in]0, h_3[$,

$$\frac{\|I_\mu(h; \lambda)\|_2^2}{\|I_\mu(h; \lambda)\|_1^2} \leq 1 + \varepsilon + \frac{4h^{(1-\varepsilon)\beta}}{\overline{\Phi}(\lambda\psi(h))}.$$

According to Mill’s ratio for Gaussian tails (see (4.4)), for any $\lambda > 0$, there exists a small $h_4 > 0$, such that for all $h \in]0, h_4[$, $\overline{\Phi}(\lambda\psi(h)) \geq h^{\lambda^2}/4\lambda\sqrt{\ln(1/h)}$. Hence, for all $h \in]0, h_3 \wedge h_4[$,

$$\frac{\|I_\mu(h; \lambda)\|_2^2}{\|I_\mu(h; \lambda)\|_1^2} \leq 1 + \varepsilon + 16\lambda h^{(1-\varepsilon)\beta - \lambda^2} \sqrt{\ln(1/h)}.$$

By choosing λ such that $(1 - \varepsilon)\beta > \lambda^2$ we can deduce that for all $h \in]0, h_3 \wedge h_4[$, $\|I_\mu(h; \lambda)\|_2^2 \|I_\mu(h; \lambda)\|_1^{-2} \leq 1 + 2\varepsilon$. Applying the Paley–Zygmund inequality ([8, p. 8]), we see that

$$\mathbb{P}(I_\mu(h; \lambda) > 0) \geq \frac{1}{1 + 2\varepsilon}. \tag{4.8}$$

We are ready to complete the proof. Define,

$$\mathcal{A}_\alpha(\lambda, h) \triangleq \left\{ t \in [0, 1] : \sup_{0 \leq r \leq h} \frac{|X(t+r) - X(t)|}{r^{\alpha/2}\psi(r)} > \lambda \right\}. \tag{4.9}$$

This is the collection of all **h -approximate λ -fast points**. Observe that for each $h > 0$, $\mathcal{A}_\alpha(\lambda, h)$ is an open subset of $[0, 1]$. If $I_\mu(h; \lambda) > 0$ ($\mu \in \mathcal{P}_+(E)$), then $E \cap \mathcal{A}_\alpha(\lambda, h) \neq \emptyset$. By (4.8), we see that as long as $(1 - \varepsilon)\beta > \lambda^2$, then for all $h \in]0, h_3 \wedge h_4[$, $\mathbb{P}(E \cap \mathcal{A}_\alpha(\lambda, h) \neq \emptyset) \geq (1 + 2\varepsilon)^{-1}$. Note that if $h \leq h'$, then $\mathcal{A}_\alpha(\lambda, h) \subset \mathcal{A}_\alpha(\lambda, h')$. Hence, for all $\lambda^2 \in [0, \dim(E)[$, $\mathbb{P}(\mathcal{A}_\alpha(\lambda, h) \cap E \neq \emptyset, \forall h > 0) = 1$. By Theorem 2.5, for all $\lambda^2 \in [0, \dim(E)[$, $\mathbb{P}(\bigcap_{h>0} \mathcal{A}_\alpha(\lambda, h) \cap E \neq \emptyset) = 1$. Since $\bigcap_{h>0} \mathcal{A}_\alpha(\lambda, h) = F_\alpha(\lambda)$, we have shown that

$$\overline{\text{co-dim}}(F_\alpha(\lambda)) \leq \lambda^2.$$

Unravelling the notation, this implies Eq. (1.5) (i.e., the lower bound in Theorem 1.1). It also implies the lower bound in Theorem 1.4. \diamond

§5. THEOREM 1.3: UPPER BOUND

For $\eta, R > 1$, and $j \geq 1$, define,

$$\mathcal{G}(j) \triangleq \{0 \leq m \leq R^{nj} : |W(mR^{-nj})| \leq 2\sqrt{\eta j R^{-nj} \ln R}\}. \tag{5.1}$$

The notation is motivated by the following description: we think of $I_{m,j}^\eta$ (see (3.2)) as “good” if $m \in \mathcal{G}(j)$. Otherwise, $I_{m,j}^\eta$ is deemed “bad”. We also recall Eq. (3.1) with $\alpha = 1$ (thus replacing X by W in (3.1)). In analogy with the definition of \mathcal{J}_k (see (3.4)), we define,

$$\mathcal{J}'_k \triangleq \sum_{j \geq k} \sum_{m \in \mathcal{G}(j)} \mathbf{1}_{\{I_{m,j}^\eta \cap F_1(\lambda, R^{-j}) \neq \emptyset\}}. \tag{5.2}$$

By the independence of the increments of W , $\{I_{m,j}^\eta \cap F_1(\lambda, R^{-j}) \neq \emptyset\}$ is independent of $\{m \in \mathcal{G}(j)\}$. Recalling (3.3), we see that

$$\|\mathcal{J}'_k\|_1 = \sum_{j \geq k} \sum_{m \geq 0} \mathbb{P}(m \in \mathcal{G}(j)) P_{m,j}^{\lambda,1}.$$

Since $|W(1)|$ has a probability density which is uniformly bounded above by 1, for all $0 \leq m \leq R^{\eta j}$,

$$\mathbb{P}(m \in \mathcal{G}(j)) \leq 4 \left(\sqrt{\frac{\eta j \ln R}{m}} \wedge 1 \right). \tag{5.3}$$

Next, fix $\varepsilon \in]0, 1[$. By Lemma 3.1, there exists $J_3 \triangleq J_1(\varepsilon, 1, \eta, \lambda, R) \in [2, \infty[$, such that for all $j \geq J_3$ and all $m \geq 0$, $P_{m,j}^{\lambda,1} \leq R^{-\lambda^2(1-\varepsilon)j}$. Using (5.3), we see that for all $k \geq J_3$,

$$\|\mathcal{J}'_k\|_1 \leq 4 \sum_{j \geq k} \sum_{m=0}^{\lfloor R^{\eta j} \rfloor} \left(\sqrt{\frac{\eta j \ln R}{m}} \wedge 1 \right) R^{-\lambda^2(1-\varepsilon)j}.$$

If we choose $\lambda^2(1 - \varepsilon) > \eta/2$, then a few lines of calculations reveal that $\sum_k \|\mathcal{J}'_k\|_1 < \infty$. By the Borel–Cantelli lemma, there exists a finite random variable k_1 , such that with probability one, for all $k \geq k_1$, $\mathcal{J}'_k = 0$. In particular, with probability one, for all $j \geq k_1$,

$$\mathbf{1}_{\{m \in \mathcal{G}(j)\}} \mathbf{1}_{\{I_{m,j}^{\eta} \cap F_1(\lambda, R^{-j}) \neq \emptyset\}} = 0. \tag{5.4}$$

By Lévy’s modulus of continuity for W (cf. (1.2)), there exists a finite random variable $k_2(\eta, R)$, such that with probability one, for all $j \geq k_2(\eta, R)$,

$$\mathbf{1}_{\{m \in \mathcal{G}(j)\}} \geq \mathbf{1}_{\{I_{m,j}^{\eta} \cap \mathcal{Z} \neq \emptyset\}}. \tag{5.5}$$

Eq. (5.4) shows that with probability one, for all $j \geq k_3 \triangleq k_1 \vee k_2(\eta, R)$, $F_1(\lambda, R^{-j}) \cap \mathcal{Z} = \emptyset$. That is, almost surely, for all $j \geq k_3$,

$$\sup_{t \in \mathcal{Z}} \sup_{t \leq s \leq t+R^{-j}} |W(t+s)| \leq \lambda R^{-j/2} \psi(R^{-j}).$$

If $h \leq R^{-k_3}$, there exists $j \geq k_3$, such that $R^{-j-1} \leq h \leq R^{-j}$. By monotonicity,

$$\sup_{t \in \mathcal{Z}} \sup_{t \leq s \leq t+h} |W(t+s)| \leq \lambda R^{-j/2} \psi(R^{-j}) \leq \lambda R^{1/2} h^{1/2} \psi(h).$$

In particular, we have shown that as long as $\lambda^2(1 - \varepsilon) > \eta/2$, then almost surely,

$$\limsup_{h \rightarrow 0^+} \sup_{t \in \mathcal{Z}} \frac{|W(t+h)|}{h^{1/2} \psi(h)} \leq \lambda R^{1/2}.$$

Along rational sequences (and in this order), let $\eta, R \rightarrow 1^+$, $\varepsilon \rightarrow 0^+$ and $\lambda^2 \downarrow \frac{1}{2}$ to see that with probability one,

$$\limsup_{h \rightarrow 0^+} \sup_{t \in \mathcal{Z}} \frac{|W(t+h)|}{h^{1/2} \psi(h)} \leq \frac{1}{\sqrt{2}}.$$

This proves the upper bound in Theorem 1.3. ◇

§6. THEOREM 1.4: UPPER BOUND

Recall Eqs. (1.6) and (3.1). We begin with a “regularization scheme” which is used later in our good covering for dimension calculations.

Lemma 6.1. *For all integers $k \geq 1$ and all reals $\theta \in]0, 1[$ and $R > 1$,*

$$F_\alpha(\lambda) \subset \bigcup_{j \geq k} F_\alpha(\theta \lambda R^{-\alpha/2}, R^{-j+1}).$$

Proof. From first principles, it follows that for all $\beta < \lambda$, $F_\alpha(\lambda) \subset \bigcup_{h>0} \bigcup_{0<\delta<h} F_\alpha(\beta, \delta)$. Let $\delta < h < 1$, say $R^{-j} \leq \delta \leq R^{-j+1}$. For all $t \in F_\alpha(\beta, \delta)$,

$$\sup_{t \leq s \leq t+R^{-j+1}} |X(s) - X(t)| \geq \beta R^{-j\alpha/2} \psi(R^{-j+1}) = \beta R^{-\alpha/2} R^{-(j-1)\alpha/2} \psi(R^{-j+1}).$$

We have used the monotonicity properties of ψ . This implies that $t \in F_\alpha(\beta R^{-\alpha/2}, R^{-(j-1)})$. The result follows. ◇

We are prepared to demonstrate the upper bound in Theorem 1.4.

Proof of Theorem 1.4: upper bound. Fix $R, \eta > 1$, $\theta \in]0, 1[$ and recall $I_{m,j}^\eta$ from (3.2). From Lemma 6.1, it is apparent that for any integer $k \geq 1$, we have the following covering of $F_\alpha(\lambda)$:

$$F_\alpha(\lambda) \subset \bigcup_{j=k}^\infty \bigcup_{m=1}^\infty I_{m,j}^\eta \cap F_\alpha(\theta \lambda R^{-\alpha/2}, R^{-j+1}). \tag{6.1}$$

By (3.3) and Lemma 3.1, for all $\varepsilon > 0$, there exists $J_4 = J_4(\varepsilon, \alpha, \eta, \lambda, R, \theta) \in [2, \infty[$, such that for all $j \geq J_4$,

$$\mathbb{P}(I_{m,j}^\eta \cap F_\alpha(\theta \lambda R^{-\alpha/2}, R^{-j+1}) \neq \emptyset) \leq R^{-(1-\varepsilon)\theta^2 \lambda^2 R^{-2\alpha j}}. \tag{6.2}$$

For any $s \geq 0$ and every integer $k \geq 1$, define,

$$\mathcal{J}_k(s) \triangleq \sum_{j \geq k} \sum_{m \leq R^{\eta j+1}} |I_{m,j}^\eta|^s \mathbf{1}_{\{I_{m,j}^\eta \cap E \neq \emptyset\}} \mathbf{1}_{\{I_{m,j}^\eta \cap F_\alpha(\theta \lambda R^{-\alpha/2}, R^{-j+1}) \neq \emptyset\}}.$$

By (6.2), for all $k \geq J_4$,

$$\|\mathcal{J}_k(s)\|_1 \leq \sum_{j \geq k} M((1 + R^{\eta j})^{-1}; E) R^{-\eta s j} R^{-(1-\varepsilon)\theta^2 \lambda^2 R^{-2\alpha j}}.$$

(Recall from §3 that $M(\varepsilon; E)$ is the ε -capacity of E .) Note that if we enlarge J_4 further, then for all $\beta > \overline{\dim}_M(E)$ and all $j \geq J_4$, we can also ensure that $M((1 + R^{\eta j})^{-1}; E) \leq R^{\eta \beta j}$. Suppose $\beta > \overline{\dim}_M(E)$ and

$$\eta s > \eta \beta - (1 - \varepsilon)\theta^2 \lambda^2 R^{-2\alpha}. \tag{6.3}$$

It follows that $\sum_k \|\mathcal{J}_k(s)\|_1 < \infty$. By the Borel–Cantelli lemma, $\lim_{k \rightarrow \infty} \mathcal{J}_k(s) = 0$, a.s. From (6.1) and the definition of Hausdorff dimension, it follows that for any s satisfying (6.3),

$$\Lambda^s(E \cap F_\alpha(\lambda)) \leq \lim_{k \rightarrow \infty} \mathcal{J}_k(s) = 0, \quad \text{a.s.}$$

Therefore, almost surely,

$$\dim(E \cap F_\alpha(\lambda)) \leq s.$$

Letting $\beta \downarrow \overline{\dim}_M(E)$, $\varepsilon \downarrow 0$, $\eta, R \downarrow 1$, $\theta \uparrow 1$ in this order and along rational sequences, we obtain

$$\dim(E \cap F_\alpha(\lambda)) \leq \overline{\dim}_M(E) - \lambda^2, \quad \text{a.s.}$$

By [17, pp. 57 and 81], for every $G \in \mathbb{S}_0^1$,

$$\begin{aligned} \dim(G) &= \inf_{G = \bigcup_{i=1}^\infty G_i} \sup_i \dim(G_i), \\ \dim_P(G) &= \inf_{G = \bigcup_{i=1}^\infty G_i} \sup_i \overline{\dim}_M(G_i), \end{aligned}$$

where the G_i 's are assumed to be bounded. Thus,

$$\dim(E \cap F_\alpha(\lambda)) \leq \dim_P(E) - \lambda^2, \quad \text{a.s.},$$

which is the desired upper bound. ◇

§7. PROOF OF THEOREM 1.5: UPPER BOUND

Recall the notation of §3 and §6, and $\mathcal{G}(j)$ from (5.1). By Lemma 6.1 and (5.5), for any $\eta, R > 1$ and $\theta \in]0, 1[$, there exists a finite random variable K such that almost surely for all $k \geq K$,

$$Z \cap F_1(\lambda) \subset \bigcup_{j \geq k} \bigcup_{m \in \mathcal{G}(j)} I_{m,j}^\eta \cap F_1(\theta \lambda R^{-1/2}, R^{-j+1}). \tag{7.1}$$

Next, we show that the above is a fairly economical covering. Since W has independent increments, for any $s > 0$,

$$\begin{aligned} &\sum_{j \geq k} \sum_{0 \leq m < R^{\eta j} + 1} |I_{m,j}^\eta|^s \mathbb{P}(I_{m,j}^\eta \cap F_1(\theta \lambda R^{-1/2}, R^{-j+1}) \neq \emptyset, m \in \mathcal{G}(j)) \\ &= \sum_{j \geq k} \sum_{0 \leq m < R^{\eta j} + 1} |I_{m,j}^\eta|^s \mathbb{P}(I_{m,j}^\eta \cap F_1(\theta \lambda R^{-1/2}, R^{-j+1}) \neq \emptyset) \mathbb{P}(m \in \mathcal{G}(j)). \end{aligned}$$

By (5.3),

$$\begin{aligned} & \sum_{j \geq k} \sum_{0 \leq m < R^{j+1}} |I_{m,j}^\eta|^s \mathbb{P}(I_{m,j}^\eta \cap F_1(\theta \lambda R^{-1/2}, R^{-j+1}) \neq \emptyset, m \in \mathcal{G}(j)) \\ & \leq 4 \sum_{j \geq k} \sum_{m=0}^{\lfloor R^{j+1} \rfloor} |I_{m,j}^\eta|^s \mathbb{P}(I_{m,j}^\eta \cap F_1(\theta \lambda R^{-1/2}, R^{-j+1}) \neq \emptyset) \left(\sqrt{\frac{\eta j \ln R}{m}} \wedge 1 \right). \end{aligned}$$

Using Lemma 3.1, we see that for all $\varepsilon \in]0, 1[$, there exists $J_5 = J_5(\varepsilon, \theta, \eta, \lambda, R) \in [2, \infty[$, such that for all $k \geq J_5$,

$$\begin{aligned} & \sum_{j \geq k} \sum_{0 \leq m < R^{j+1}} |I_{m,j}^\eta|^s \mathbb{P}(I_{m,j}^\eta \cap F_1(\theta \lambda R^{-1/2}, R^{-j+1}) \neq \emptyset, m \in \mathcal{G}(j)) \\ & \leq 4 \sum_{j \geq k} R^{-\eta s j} R^{-(1-\varepsilon)\theta^2 \lambda^2 R^{-1} j} \sum_{m=0}^{\lfloor R^{j+1} \rfloor} \left(\sqrt{\frac{\eta j \ln R}{m}} \wedge 1 \right). \end{aligned}$$

In particular, if

$$\eta s > \frac{\eta}{2} - (1 - \varepsilon)\theta^2 \lambda^2 R^{-1}, \tag{7.2}$$

then,

$$\lim_{k \rightarrow \infty} \sum_{j \geq k} \sum_{m \in \mathcal{G}(j)} |I_{m,j}^\eta|^s \mathbf{1}_{\{I_{m,j}^\eta \cap F_1(\theta \lambda R^{-1/2}, R^{-j+1}) \neq \emptyset\}} = 0, \quad \text{a.s.}$$

Thanks to (7.1), we can deduce that for any s satisfying (7.2), $\Lambda^s(\mathcal{Z} \cap F_1(\lambda)) = 0$, almost surely. In particular, almost surely, $\dim(\mathcal{Z} \cap F_1(\lambda)) \leq s$. Let $\varepsilon \downarrow 0$, $\theta \uparrow 1$ and $R, \eta \downarrow 1$ in this order to see that with probability one, $\dim(\mathcal{Z} \cap F_1(\lambda)) \leq \frac{1}{2} - \lambda^2$. This is the desired upper bound. \diamond

§8. PROOFS OF THEOREMS 1.3 AND 1.5: LOWER BOUNDS

The main result of this section is the following which may be of independent interest.

Theorem 8.1. *Fix a compact set $E \subset [0, 1]$. Then,*

$$\dim(E) > \lambda^2 + \frac{1}{2} \implies \mathbb{P}(\mathcal{Z} \cap F_1(\lambda) \cap E \neq \emptyset) = 1.$$

As the lower bounds in Theorems 1.3 and 1.5 follow immediately from the above, the rest of this section is devoted to proving Theorem 8.1.

Suppose $E \subset [0, 1]$ is compact, $\mu \in \mathcal{P}_+(E)$ and $h, \lambda > 0$. Define,

$$J_\mu(h; \lambda) \triangleq \int_0^1 \mu(ds) \mathbf{1}_{\{|W(s)| < h\}} \mathbf{1}_{\{W(s+h) - W(s) > \lambda h^{1/2} \psi(h)\}}.$$

For all $\mu \in \mathcal{P}_+(E)$ and any $h, \beta > 0$, define the following:

$$S_h(\mu) \triangleq \sup_{0 \leq s \leq h} \int_s^h \frac{\mu(dt)}{\sqrt{t-s}},$$

$$\tilde{S}_h(\mu) \triangleq \sup_{0 \leq s \leq 1} \int_s^{(s+h) \wedge 1} \frac{\mu(dt)}{\sqrt{t-s}}.$$

The proof of Theorem 8.1 is divided into several steps. In analytical terms, our first result is an estimate, uniform in $s \in [0, h]$, for the $1/2$ -potential of a measure μ restricted to an interval $[s, h]$. Indeed, recalling $A_\beta(\mu)$ from (2.1), we have the following:

Lemma 8.2. *Suppose $\mu \in \mathcal{P}_+(E)$. Then for all $\beta > 1/2$ and all $h > 0$,*

$$S_h(\mu) \leq \frac{2e^\beta}{2\beta - 1} A_\beta(\mu) h^{\beta - \frac{1}{2}}, \tag{8.1}$$

$$\tilde{S}_h(\mu) \leq \frac{2e^\beta}{2\beta - 1} A_\beta(\mu) h^{\beta - \frac{1}{2}}. \tag{8.2}$$

Proof. Without loss of generality, we can assume that $A_\beta(\mu) < \infty$ for some $\beta > 1/2$ (otherwise, there is nothing to prove). We proceed with an approximate integration by parts: for all $0 \leq s \leq h$,

$$\begin{aligned} \int_s^h \mu(dt) (t-s)^{-1/2} &= \sum_{j=0}^{\infty} \int_{s+(h-s)e^{-j-1}}^{s+(h-s)e^{-j}} \mu(dt) (t-s)^{-1/2} \\ &\leq (h-s)^{-1/2} \sum_{j=0}^{\infty} e^{(j+1)/2} \mu[s + (h-s)e^{-j-1}, s + (h-s)e^{-j}] \\ &\leq e^{1/2} (h-s)^{\beta-1/2} A_\beta(\mu) \sum_{j=0}^{\infty} \exp\left(-j\left(\beta - \frac{1}{2}\right)\right), \end{aligned}$$

which yields (8.1). The estimate (8.2) can be checked in the same way. ◇

Our next two lemmas are moment estimates for J_μ .

Lemma 8.3. *Suppose $\mu \in \mathcal{P}_+(E)$ is fixed. For every $\varepsilon > 0$, there exists an $h_\varepsilon > 0$, such that for all $h \in]0, h_\varepsilon[$, and all $\lambda > 0$,*

$$\|J_\mu(h; \lambda)\|_1 \geq \sqrt{\frac{2}{\pi e}} (1 - \varepsilon) h \bar{\Phi}(\lambda \psi(h)).$$

Proof. By the independence of the increments of W ,

$$\|J_\mu(h; \lambda)\|_1 = \bar{\Phi}(\lambda\psi(h)) \int_0^1 \mu(ds) \mathbb{P}(|W(s)| \leq h).$$

A direct calculation reveals that if $r \in]0, 1[$, $\mathbb{P}(|W(1)| \leq r) \geq \sqrt{2/(\pi e)} r$. Therefore, by Brownian scaling, for every $h \in]0, 1[$,

$$\begin{aligned} \|J_\mu(h; \lambda)\|_1 &\geq \sqrt{\frac{2}{\pi e}} h \bar{\Phi}(\lambda\psi(h)) \int_{h^2}^1 \frac{\mu(ds)}{\sqrt{s}} \\ &\geq \sqrt{\frac{2}{\pi e}} \mu[h^2, 1] h \bar{\Phi}(\lambda\psi(h)). \end{aligned}$$

The lemma follows upon taking $h > 0$ small enough. ◇

Lemma 8.4. Fix $\mu \in \mathcal{P}_+(E)$. Suppose $A_\beta(\mu) < \infty$ for some $\beta > 1/2$. Then, for all $h, \lambda > 0$,

$$\|J_\mu(h; \lambda)\|_2^2 \leq \frac{8e^{2\beta}}{(2\beta - 1)^2} A_\beta^2(\mu) \left\{ 2h^{\beta+3/2} \bar{\Phi}(\lambda\psi(h)) + h^2 \bar{\Phi}^2(\lambda\psi(h)) \right\}.$$

Proof. To save space, for all $h, t \geq 0$, define,

$$\Delta_h W(t) \triangleq W(t+h) - W(t).$$

By the independence of the increments of W ,

$$\begin{aligned} \|J_\mu(h; \lambda)\|_2^2 &= 2 \left\| \int_0^1 \mu(dt) \mathbf{1}_{\{|W(t)| \leq h\}} \mathbf{1}_{\{\Delta_h W(t) \geq \lambda h^{1/2} \psi(h)\}} \right\| \times \\ &\quad \times \left\| \int_0^t \mu(ds) \mathbf{1}_{\{|W(s)| \leq h\}} \mathbf{1}_{\{\Delta_h W(s) \geq \lambda h^{1/2} \psi(h)\}} \right\|_1 \\ &= 2 \bar{\Phi}(\lambda\psi(h)) \left\| \int_0^1 \mu(dt) \mathbf{1}_{\{|W(t)| \leq h\}} \int_0^t \mu(ds) \mathbf{1}_{\{|W(s)| \leq h\}} \mathbf{1}_{\{\Delta_h W(s) \geq \lambda h^{1/2} \psi(h)\}} \right\|_1 \\ &\leq 2 \bar{\Phi}(\lambda\psi(h)) [T_1 + T_2], \end{aligned} \tag{8.3}$$

where,

$$\begin{aligned} T_1 &\triangleq \left\| \int_h^1 \mu(dt) \mathbf{1}_{\{|W(t)| \leq h\}} \int_0^{(t-h)^+} \mu(ds) \mathbf{1}_{\{|W(s)| \leq h\}} \mathbf{1}_{\{\Delta_h W(s) \geq \lambda h^{1/2} \psi(h)\}} \right\|_1, \\ T_2 &\triangleq \left\| \int_0^1 \mu(dt) \mathbf{1}_{\{|W(t)| \leq h\}} \int_{(t-h)^+}^t \mu(ds) \mathbf{1}_{\{|W(s)| \leq h\}} \right\|_1. \end{aligned}$$

We will estimate T_1 and T_2 in turn.

We will need the following consequence of Gaussian laws: for all $s, h > 0$,

$$\mathbb{P}(|W(s)| \leq h) \leq s^{-1/2}h. \quad (8.4)$$

First, we estimate T_1 . Note that,

$$T_1 = \int_h^1 \mu(dt) \int_0^{t-h} \mu(ds) \mathbb{P}(|W(t)| \leq h, |W(s)| \leq h, \Delta_h W(s) \geq \lambda h^{1/2} \psi(h)). \quad (8.5)$$

Suppose $t \in [h, 1]$ and $s \in [0, t-h]$. Then $s \leq s+h \leq t$, and we have,

$$\begin{aligned} \mathbb{P}(|W(t)| \leq h \mid W(r); r \leq s+h) \\ &= \mathbb{P}(|W(t) - W(s+h) + W(s+h)| \leq h \mid W(r); r \leq s+h) \\ &\leq \sup_{\zeta \in \mathbb{R}} \mathbb{P}(|W(t-s-h) + \zeta| \leq h). \end{aligned}$$

On the other hand, $W(t-s-h)$ has a Gaussian distribution. By unimodality of the latter,

$$\mathbb{P}(|W(t)| \leq h \mid W(r); r \leq s+h) \leq \mathbb{P}(|W(t-s-h)| \leq h).$$

(This actually is a particular case of T.W. Anderson's inequality for general Gaussian shifted balls). Using (8.5) and the principle of conditioning,

$$\begin{aligned} T_1 &\leq \int_h^1 \mu(dt) \int_0^{t-h} \mu(ds) \mathbb{P}(|W(t-s-h)| \leq h) \times \\ &\quad \times \mathbb{P}(|W(s)| \leq h, \Delta_h W(s) \geq \lambda h^{1/2} \psi(h)) \\ &\leq \bar{\Phi}(\lambda \psi(h)) \int_h^1 \mu(dt) \int_0^{t-h} \mu(ds) \mathbb{P}(|W(t-s-h)| \leq h) \mathbb{P}(|W(s)| \leq h). \end{aligned}$$

By (8.4),

$$T_1 \leq h^2 \bar{\Phi}(\lambda \psi(h)) \int_h^1 \mu(dt) \int_0^{t-h} \mu(ds) \frac{1}{\sqrt{s(t-s-h)}}.$$

Changing the order of integration, we arrive at the following estimate:

$$\begin{aligned} T_1 &\leq h^2 \bar{\Phi}(\lambda \psi(h)) \int_0^{1-h} \frac{\mu(ds)}{\sqrt{s}} \int_{s+h}^1 \frac{\mu(dt)}{\sqrt{t-s-h}} \\ &\leq h^2 \bar{\Phi}(\lambda \psi(h)) S_1^2(\mu) \\ &\leq \frac{4e^{2\beta}}{(2\beta-1)^2} A_\beta^2(\mu) h^2 \bar{\Phi}(\lambda \psi(h)), \end{aligned} \quad (8.6)$$

by Lemma 8.2. Next, we estimate T_2 . By another unimodality argument, for all $t \geq s$ and all $h > 0$,

$$\mathbb{P}(|W(s)| \leq h, |W(t)| \leq h) \leq \mathbb{P}(|W(s)| \leq h) \mathbb{P}(|W(t-s)| \leq h).$$

Applying (8.4), $\mathbb{P}(|W(s)| \leq h, |W(t)| \leq h) \leq h^2 / \sqrt{s(t-s)}$. Therefore,

$$\begin{aligned} T_2 &\leq h^2 \int_0^1 \mu(dt) \int_{(t-h)^+}^t \mu(ds) \frac{1}{\sqrt{s(t-s)}} \\ &= h^2 (T_{2,1} + T_{2,2}), \end{aligned} \tag{8.7}$$

where,

$$\begin{aligned} T_{2,1} &\triangleq \int_0^h \mu(dt) \int_0^t \mu(ds) \frac{1}{\sqrt{s(t-s)}} \\ T_{2,2} &\triangleq \int_h^1 \mu(dt) \int_{t-h}^t \mu(ds) \frac{1}{\sqrt{s(t-s)}}. \end{aligned}$$

To estimate $T_{2,1}$, reverse the order of integration:

$$\begin{aligned} T_{2,1} &= \int_0^h \frac{\mu(ds)}{\sqrt{s}} \int_s^h \frac{\mu(dt)}{\sqrt{t-s}} \leq S_h^2(\mu) \\ &\leq \frac{4e^{2\beta}}{(2\beta-1)^2} A_\beta^2(\mu) h^{2\beta-1}, \end{aligned} \tag{8.8}$$

by Lemma 8.2. Similarly,

$$\begin{aligned} T_{2,2} &= \int_0^1 \frac{\mu(ds)}{\sqrt{s}} \int_s^{(s+h) \wedge 1} \frac{\mu(dt)}{\sqrt{t-s}} \\ &\leq S_1(\mu) \tilde{S}_h(\mu) \\ &\leq \frac{4e^{2\beta}}{(2\beta-1)^2} A_\beta^2(\mu) h^{\beta-1/2}. \end{aligned}$$

Since $h < 1$ and $\beta > 1/2$, using the above, (8.8) and (8.7), we arrive at the following:

$$T_2 \leq \frac{8e^{2\beta}}{(2\beta-1)^2} A_\beta^2(\mu) h^{\beta+3/2}.$$

Use this, together with (8.6) and (8.3) in this order to get the result. \diamond

We are ready for the main result of this section:

Proof of Theorem 8.1. For $\lambda, h > 0$, recall (4.9) and define,

$$\mathcal{Z}(h) \triangleq \{s \in [0, 1] : |W(s)| < h\}.$$

Path continuity of W alone implies that $\mathcal{Z}(h) \cap \mathcal{A}_1(\lambda, h)$ is an open random set. We estimate the probability that it intersects E . By Frostman's lemma, for all $\beta < \dim(E)$, there exists $\mu \in \mathcal{P}_+(E)$ such that $A_\beta(\mu) < \infty$. Let us fix a μ corresponding to an arbitrary but fixed choice of β satisfying:

$$\lambda^2 + \frac{1}{2} < \beta < \dim(E). \tag{8.9}$$

Applying Lemmas 8.3 and 8.4 to this choice of μ , we see that for all $\varepsilon > 0$, there exists $h_\varepsilon > 0$, such that for all $h \in]0, h_\varepsilon[$,

$$\frac{\|J_\mu(h; \lambda)\|_1^2}{\|J_\mu(h; \lambda)\|_2^2} \geq \gamma_{\varepsilon, \beta} \left[\frac{2h^{\beta-1/2}}{\overline{\Phi}(\lambda\psi(h))} + 1 \right]^{-1},$$

where,

$$\gamma_{\varepsilon, \beta} \triangleq \frac{(2\beta - 1)^2(1 - \varepsilon)^2}{4\pi e^{1+2\beta} A_\beta^2(\mu)}.$$

According to Mill's ratio for Gaussian tails (see (4.4)), we can pick h_λ so small that for each and every $h \in]0, h_\lambda[$, $\overline{\Phi}(\lambda\psi(h)) \geq h^{\lambda^2}/4\lambda\sqrt{\ln(1/h)}$. Therefore, for all $h \in]0, h_\varepsilon \wedge h_\lambda[$,

$$\frac{\|J_\mu(h; \lambda)\|_1^2}{\|J_\mu(h; \lambda)\|_2^2} \geq \gamma_{\varepsilon, \beta} \left[8\lambda h^{\beta-\lambda^2-1/2} \sqrt{\ln(1/h)} + 1 \right]^{-1}.$$

By (8.9), $\liminf_{h \rightarrow 0^+} \|J_\mu(h; \lambda)\|_1^2 \|J_\mu(h; \lambda)\|_2^{-2} \geq \gamma_{\varepsilon, \beta} > 0$. By the Paley-Zygmund inequality ([8, p. 8]),

$$\liminf_{h \rightarrow 0^+} \mathbb{P}(J_\mu(h; \lambda) > 0) \geq \gamma_{\varepsilon, \beta} > 0.$$

Note that the event $(J_\mu(h; \lambda) > 0)$ implies that $\mathcal{Z}(h) \cap \mathcal{A}_1(\lambda, h)$ intersects E . Hence,

$$\liminf_{h \rightarrow 0^+} \mathbb{P}(\mathcal{Z}(h) \cap \mathcal{A}_1(\lambda, h) \cap E \neq \emptyset) \geq \gamma_{\varepsilon, \beta} > 0.$$

However, as $h \downarrow 0$, the (random) open set $\mathcal{Z}(h) \cap \mathcal{A}_1(\lambda, h)$ decreases to $\mathcal{Z} \cap F_1(\lambda)$. Adapting Theorem 2.5 to the positive probability case, we can conclude that $\mathbb{P}(\mathcal{Z} \cap F_1(\lambda) \cap E \neq \emptyset) \geq \gamma_{\varepsilon, \beta} > 0$. Note that the only requirement on E was that $\dim(E) > \lambda^2 + 1/2$. Since Hausdorff dimension is scale invariant, we see that for any real number $s \in]0, 1[$,

$$\mathbb{P}(\mathcal{Z} \cap F_1(\lambda) \cap s^{-1}E \neq \emptyset) \geq \gamma_{\varepsilon, \beta},$$

where $s^{-1}E \triangleq \{r/s : r \in E\}$. We finish the proof by showing that this probability is actually 1. Fix $s \in]0, 1[$ and observe from Brownian scaling that $\mathcal{Z} \cap F_1(\lambda)$ has the same distribution as $\mathcal{Z} \cap F_1(\lambda) \cap [0, s]$ in the sense that for all $G \in \mathcal{S}_0^1$,

$$\mathbb{P}(\mathcal{Z} \cap F_1(\lambda) \cap G \neq \emptyset) = \mathbb{P}(\mathcal{Z} \cap F_1(\lambda) \cap [0, s] \cap sG \neq \emptyset).$$

In particular, for all $s \in]0, 1[$, $\mathbb{P}(\mathcal{Z} \cap F_1(\lambda) \cap [0, s] \cap E \neq \emptyset) \geq \gamma_{\varepsilon, \beta} > 0$. Note that $\mathcal{Z} \cap F_1(\lambda) \cap E \cap [0, s]$ is increasing in s . Thus,

$$\mathbb{P}\left(\bigcap_{s \in]0, 1[} \left\{ \mathcal{Z} \cap F_1(\lambda) \cap E \cap [0, s] \neq \emptyset \right\}\right) \geq \gamma_{\varepsilon, \beta}.$$

Let $C \triangleq \bigcap_{s \in]0, 1[} (\mathcal{Z} \cap F_1(\lambda) \cap E \cap [0, s] \neq \emptyset)$. Observe that C is measurable with respect to the germ field of W at 0 and we have just argued that $\mathbb{P}(C) \geq \gamma_{\varepsilon, \beta} > 0$. By Blumenthal's 0-1 law, $\mathbb{P}(C) = 1$. Since $(\mathcal{Z} \cap F_1(\lambda) \cap E \neq \emptyset) \supset C$, the result follows for F_1 . \diamond

REFERENCES

- [1] M.T. BARLOW AND E. PERKINS (1984). Levels at which every Brownian excursion is exceptional. *Sém. Prob. XVIII, Lecture Notes in Math.* **1059**, 1–28, Springer–Verlag, New York.
- [2] M. CSÖRGŐ AND P. RÉVÉSZ (1981). *Strong Approximations in Probability and Statistics*, Academic Press, New York.
- [3] P. DEHEUVELS AND M.A. LIFSHITS (1997). On the Hausdorff dimension of the set generated by exceptional oscillations of a Wiener process. *Studia Sci. Math. Hung.*, **33**, 75–110.
- [4] P. DEHEUVELS AND D.M. MASON (1997). Random fractal functional laws of the iterated logarithm. (preprint)
- [5] R.M. DUDLEY (1984). A Course on Empirical Processes. *École d'Été de St. Flour 1982. Lecture Notes in Mathematics* **1097**. Springer, Berlin.
- [6] J. HAWKES (1971). On the Hausdorff dimension of the range of a stable process with a Borel set. *Z. Wahr. verw. Geb.*, **19**, 90–102.
- [7] J. HAWKES (1981). Trees generated by a simple branching process. *J. London Math. Soc.*, **24**, 373–384.
- [8] J.-P. KAHANE (1985). *Some Random Series of Functions*, second edition. Cambridge University Press, Cambridge.
- [9] R. KAUFMAN (1974). Large increments of Brownian Motion. *Nagoya Math. J.*, **56**, 139–145.
- [10] D. KHOSHNEVISAN, Y. PERES AND Y. XIAO (1998). Limsup random fractals. In preparation.
- [11] N. KÔNO (1977). The exact Hausdorff measure of irregularity points for a Brownian path. *Z. Wahr. verw. Geb.*, **40**, 257–282.
- [12] M. LEDOUX AND M. TALAGRAND (1991). *Probability in Banach Space, Isoperimetry and Processes*, Springer–Verlag, Heidelberg–New York.
- [13] P. LÉVY (1937). *Théorie de l'Addition des Variables Aléatoires*. Gauthier–Villars, Paris.
- [14] R. LYONS (1980). Random walks and percolation on trees. *Ann. Prob.*, **18**, 931–958.
- [15] M.B. MARCUS (1968). Hölder conditions for Gaussian processes with stationary increments. *Trans. Amer. Math. Soc.*, **134**, 29–52.
- [16] M.B. MARCUS AND J. ROSEN (1992). Moduli of continuity of local times of strongly symmetric Markov processes via Gaussian processes. *J. Theoretical Prob.*, **5**, 791–825.

- [17] P. MATILLA (1995). *Geometry of Sets and Measures in Euclidean Spaces, Fractals and Rectifiability*, Cambridge University Press, Cambridge.
- [18] S. OREY AND S.J. TAYLOR (1974). How often on a Brownian path does the law of the iterated logarithm fail? *Proc. London Math. Soc.*, **28**, 174–192.
- [19] Y. PERES (1996). Remarks on intersection–equivalence and capacity–equivalence. *Ann. Inst. Henri Poincaré: Physique Théorique*, **64**, 339–347.
- [20] E. PERKINS AND S.J. TAYLOR (1988), Measuring close approaches on a Brownian path, *Ann. Prob.*, **16**, 1458–1480.
- [21] D. REVUZ AND M. YOR (1994). *Continuous Martingales and Brownian Motion*, second edition. Springer, Berlin.
- [22] G.R. SHORACK AND J.A. WELLNER (1986). *Empirical Processes with Applications to Statistics*. Wiley, New York.
- [23] S.J. TAYLOR (1966). Multiple points for the sample paths of the symmetric stable process, *Z. Wahr. ver. Geb.*, **5**, 247–64.
- [24] S.J. TAYLOR (1986). The measure theory of random fractals. *Math. Proc. Camb. Phil. Soc.*, **100**, 383–406.

Davar Khoshnevisan
Department of Mathematics
University of Utah
Salt Lake City, UT. 84112
U.S.A.
E-mail: davar@math.utah.edu

Zhan Shi
Laboratoire de Probabilités
Université Paris VI
4, Place Jussieu
75252 Paris Cedex 05, France
E-mail: shi@ccr.jussieu.fr