SÉMINAIRE DE PROBABILITÉS (STRASBOURG)

NORIHIKO KAZAMAKI

Examples on local martingales

Séminaire de probabilités (Strasbourg), tome 6 (1972), p. 98-100

<http://www.numdam.org/item?id=SPS_1972__6__98_0>

© Springer-Verlag, Berlin Heidelberg New York, 1972, tous droits réservés.

L'accès aux archives du séminaire de probabilités (Strasbourg) (http://portail.mathdoc.fr/SemProba/) implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/conditions). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
http://www.numdam.org/

EXAMPLES ON LOCAL MARTINGALES

by N.KAZAMAKI

In this note we shall give two remarks relative to changes of time for local martingales. Example 2 shows that the local martingale property is not invariant through changes of time.

We assume that the reader knows the usual definitions; for example, local martingales, stopping times, etc. By a change of time $A=(\underline{\underline{F}}_t, a_t)$ we means a family of stopping times of the $\underline{\underline{F}}_t$ family, finite valued, such that for $\omega \in \Omega$ the sample function $a_\cdot(\omega)$ is right continuous and increasing. All martingales below are assumed to be right continuous.

EXAMPLE 1.- Let $\Omega = R_+$, $\underline{\underline{F}}^o$ the class of all linear Borel sets in Ω and we designate by S the identity function of Ω into R_+. Let $\underline{\underline{F}}^o_t$ be the Borel field generated by $S \wedge t$. We define the probability measure P on Ω by $P(S>t)=e^{-t}$. Let $\underline{\underline{F}}_t$ be the P-completed Borel field of $\underline{\underline{F}}^o_t$. Note that the family $(\underline{\underline{F}}_t)$ is right continuous and quasi-left continuous.

PROPOSITION 1.- Let $A=(\underline{\underline{F}}_t, a_t)$ be a change of time such that $P(S > a_t) > 0$ for each t. Then for any martingale $M=(M_t, \underline{\underline{F}}_t)$, the process $AM=(M_{a_t}, \underline{\underline{F}}_{a_t})$ is also a martingale.

PROOF.- According to THEOREM 1 of [1], it follows that for each t there exists some $s_t \in \bar{R}$ such that

$$(1) \qquad \begin{cases} \text{(i)} \quad a_t \geqq S & \text{if } S \leqq s_t \\ \text{(ii)} \quad a_t = s_t & \text{if } S > s_t \end{cases}.$$

Obviously s_t is right continuous. As $P(S > a_t) > 0$ for each t from the assumption, each s_t is finite. On the other hand, there exists a constant process $(c_t, \underline{\underline{F}}_t)$ such that

$$(2) \qquad M_t = M_S I_{[S \leqq t]} + c_t I_{[S>t]} .$$

It follows from (1) that we have

$$M_{a_t} = M_S I_{[S \leqq a_t]} + c_{a_t} I_{[S>a_t]} = M_S I_{[S \leqq s_t]} + c_{s_t} I_{[S>s_t]} = M_{s_t} .$$

Furthermore it is easy to show that for each t we have $\underline{\underline{F}}_{a_t} = \underline{\underline{F}}_{a_t \wedge S}$ and $\underline{\underline{F}}_{s_t} = \underline{\underline{F}}_{s_t \wedge S}$. This implies that $\underline{\underline{F}}_{s_t} = \underline{\underline{F}}_{a_t}$ for each t. Consequently TM is also a martingale. This completes the proof.

PROPOSITION 2.- If $M=(M_t, \underline{\underline{F}}_t)$ is a continuous martingale, then $M_t \equiv C$, where C is a constant.

PROOF.- From formule(2) the continuity of c_t can be deduced by noting that M is continuous. Clearly we then have $M_S(\omega) = c_\omega$ a.s. On the other hand, it follows from formule(2) that the martingale equality implies the following:

$$(3) \qquad \int_s^t M_S \, dP = c_s e^{-s} - c_t e^{-t} \quad (s<t).$$

Then an easy computation shows

$$(4) \qquad \frac{d}{dt} c_t = 0 \qquad \text{i.e} \quad c_t = C.$$

Consequently we have $M_t \equiv C$.

The above proposition implies that any non-constant martingale on this probability space is quasi-left continuous but not continuous.

EXAMPLE 2.- Let $(\Omega, \underline{\underline{F}}, P)$ be a complete probability space, given an increasing right continuous family $(\underline{\underline{F}}_t)$ of Borel subfields of $\underline{\underline{F}}$ as usal. Note that if M is a weak martingale, for any change of time $A=(\underline{\underline{F}}_t, a_t)$ the process AM is also a weak martingale.(see[2]).

We suppose now that there exists a continuous martingale $M = (M_t, \underline{\underline{F}}_t)$, $M_0 = 0$, with the property $P(\limsup_{t \to \infty} M_t = \infty) = 1$; for example one dimensinal Brownian motion. Then the random variable a_t defined by

$$(5) \qquad a_t = \inf(u: M_u > t)$$

is a finite stopping time of the family $(\underline{\underline{F}}_t)$. Clearly $a_0 = 0$ and $a_\infty = \infty$ a.s. It is easy to see that the change of time A satisfies $M_{a_t} = t$ from the continuity of M. The process $AM = (t, \underline{\underline{F}}_{a_t})$ is not a local martingale. Thus the local martingale property is not invariant through changes of time. This fact should be noted.

REFERENCES

[1] C.DELLACHERIE ; Un exemple de la théorie général des processus. Séminaire de Probabilités IV, Université de Strasbourg , Lecture Notes in Mathematics vol.124, Springer , Heidelberg 1970.

[2] N.KAZAMAKI ; Changes of time, Stochastic integrals and weak martingales, Zeitschrift für W-theorie (to appear).

MATHEMATICAL INSTITUTE
TÔHOKU UNIVERSITY
SENDAI, JAPAN