SÉMINAIRE DE PROBABILITÉS (STRASBOURG)

SAUL D. JACKA

A local time inequality for martingales

Séminaire de probabilités (Strasbourg), tome 17 (1983), p. 106-116

<http://www.numdam.org/item?id=SPS_1983__17__106_0>

© Springer-Verlag, Berlin Heidelberg New York, 1983, tous droits réservés.

L'accès aux archives du séminaire de probabilités (Strasbourg) (http://portail.mathdoc.fr/SemProba/) implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/conditions). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
http://www.numdam.org/

A Local Time Inequality For Martingales

by S.D. Jacka*

M.T. Barlow and M. Yor [1] have established the existence of universal constants $c_p, C_p > 0$ such that, for all continuous martingales M, with $M_0 = 0$:

$$c_p \| \langle M \rangle_\infty^{\frac{1}{2}} \|_p \leq \| \sup_a L_\infty^a(M) \|_p \leq C_p \| \langle M \rangle_\infty^{\frac{1}{2}} \|_p . \qquad (A)$$

One is naturally led to consider possible extensions of these inequalities involving the term $\sup_a \sup_t |L_t^a(M) - L_t^a(N)|$ and in this paper we establish the existence of a universal constant c_p such that

$$\| (\langle M-N \rangle_\infty - \langle M-N \rangle_0)^{\frac{1}{2}} \|_p \leq C_p \| \sup_a \sup_t |L_t^a(M) - L_t^a(N)| \|_p$$

for all continuous martingales M and N (Theorem 1).

Conversely, Barlow and Yor [2], have recently established the inequality:

$$\| \sup_a \sup_t |L_t^a(M) - L_t^a(N)| \|_p \leq C_p \| (M-N)_\infty^* \|_p^{\frac{1}{2}} \| M_\infty^* + N_\infty^* \|_p^{\frac{1}{2}} \left\{ 1 \vee \ln \left(\frac{\| M_\infty^* + N_\infty^* \|_p}{\| (M-N)_\infty^* \|_p} \right) \right\}^{\frac{1}{2}} \qquad (B)$$

We also establish (Theorem 2) the ess sup equality:

$$\text{ess sup} \sup_t |L_t^a(M) - L_t^a(N)| = \text{ess sup} |L_\infty^a(M) - L_\infty^a(N)|$$

for each $a \in \mathbb{R}$.

---

* This research was supported by the SERC

Let $(\Omega,F,(F_t;t\geq 0))$ be a filtered probability space satisfying the usual conditions. For any random variable f and any $p\in(0,\infty)$ we set $\|f\|_p = (E[|f|^p])^{1/p}$ and we set $\|f\|_\infty = \text{ess sup}|f|$. For any continuous local F_t-martingale X and any $p\in(0,\infty)$ we set $\|X\|_{H^p} = \| \langle X\rangle_\infty^{\frac{1}{2}} \|_p$ where $\langle X\rangle_t$ is the unique, increasing adapted process such that $\langle X\rangle_0 = X_0^2$ and $X_t^2 - \langle X\rangle_t$ is a local F_t-martingale, and define $H^p = \{X : \|X\|_{H^p}<\infty\}$.

We recall the Burkholder-Davis-Gundy inequalities which state that for each $p\in(0,\infty)$ there exist universal constants $c_p, C_p>0$ such that, for all $X\in H^p$

$$c_p \|X_\infty^*\|_p \leq \|\langle X\rangle_\infty^{\frac{1}{2}}\|_p \leq C_p \|X_\infty^*\|_p$$

where $X_t^* = \sup_{s\leq t}|X_s|$.

Following [5] we define the local time of X by Tanaka's formula:

$$|X_t-a| = |X_0-a| + \int_{0+}^t \text{sgn}(X_s-a)dX_s + L_t^a(X),$$

we recall that, for each a, $L_t^a(X)$ is increasing in t, [6], and the support of the measure dL_t^a is contained in $\{t : X_t=a\}$. Furthermore, since we are working with continuous local martingales we may take a version of $(L_t^a(X);\ a\in\mathbb{R},\ t\geq 0)$ which is jointly continuous in a and t, [3].

For any $X\in H^p$ set $\hat{X} = X-X_0$. Finally we recall two definitions : if $F : \mathbb{R}^+\to\mathbb{R}^+$ is an increasing function with $F(0)=0$, $F(x)\neq 0$ for $x\neq 0$ we say that F is <u>moderate</u> if there exists an $\alpha>1$ such that

$$\sup_{x>0}\frac{F(\alpha x)}{F(x)} < \infty$$

and that F is slowly increasing if there exists an $\alpha>1$ such that

$$\sup_{x>0} \frac{F(\alpha x)}{F(x)} < \alpha.$$

Theorem 1 For each $p>0$ there exists a universal constant c_p such that

$$c_p \left\| \sup_a \sup_t |L_t^a(M) - L_t^a(N)| \right\|_p \geq \left\| (\langle M-N\rangle_\infty - \langle M-N\rangle_0)^{\frac{1}{2}} \right\|_p \quad (1)$$

for all M and N in H^p.

The proof is obtained via several lemmas.

For $M,N\in H^p$ define, for each $c>0$, the stopping time

$$\tau_c = \inf\{t\geq 0 : |M_t-N_t| \geq |M_0-N_0|+c\}$$

where the infimum of the empty set is taken as $+\infty$.

Lemma 2 For M and N in H^1

$$8E\left[\sup_a \sup_t |L_t^a(M)-L_t^a(N)| I_{(\tau_c<\infty)}\right] \geq cP(\tau_{2c}<\infty) \quad (2)$$

Proof Define

$$\sigma_c = \inf\{t\geq\tau_c : |M_t-M_{\tau_c}| \vee |N_t-N_{\tau_c}| \geq \tfrac{1}{2}c\}$$

Now, by the continuity of M and N, $|M_{\tau_c}-N_{\tau_c}| = |M_0-N_0|+c$ on $(\tau_c<\infty)$, and so N_t does not hit M_{τ_c} and M_t does not hit N_{τ_c} on the interval $[\tau_c,\sigma_c]$; therefore

$$\left.\begin{aligned} L_{\sigma_c}^{M_{\tau_c}}(N) &= L_{\tau_c}^{M_{\tau_c}}(N) \\ L_{\sigma_c}^{N_{\tau_c}}(M) &= L_{\tau_c}^{N_{\tau_c}}(M) \end{aligned}\right\} \qquad (3)$$

setting

$$U(a,t) = L_t^a(M) - L_t^a(N)$$

$$D_t = \sup_a \sup_{s\le t} U(a,s)$$

we see that

$$4D_{\sigma_c} I_{(\tau_c<\infty)} \ge [U(M_{\tau_c},\sigma_c) - U(M_{\tau_c},\tau_c)] - [U(N_{\tau_c},\sigma_c) - U(N_{\tau_c},\tau_c)]$$

Using (3) we obtain

$$4D_\infty I_{(\tau_c<\infty)} \ge (L_{\sigma_c}^{M_{\tau_c}}(M) - L_{\tau_c}^{M_{\tau_c}}(M)) + (L_{\sigma_c}^{N_{\tau_c}}(N) - L_{\tau_c}^{N_{\tau_c}}(N)) \qquad (4)$$

Applying Tanaka's formula we see that the right-hand side of (4) is

$$|M_{\sigma_c} - M_{\tau_c}| + |N_{\sigma_c} - N_{\tau_c}| - \int_{\tau_c}^{\sigma_c} \operatorname{sgn}(M_s - M_{\tau_c})\, dM_s - \int_{\tau_c}^{\sigma_c} \operatorname{sgn}(N_s - N_{\tau_c})\, dN_s \qquad (5)$$

The two stochastic integrals in (5) are martingales in H^1, as M and N are in H^1, and so, applying the optional sampling theorem we obtain

$$4E(D_\infty I_{(\tau_c<\infty)}) \ge E[|M_{\sigma_c} - M_{\tau_c}| + |N_{\sigma_c} - N_{\tau_c}|] \qquad (6)$$

Finally, $\sigma_c<\tau_{2c}$, and on $(\sigma_c<\infty)$, $|M_{\sigma_c}-M_{\tau_c}|+|N_{\sigma_c}-N_{\tau_c}|\geq c/2$ so, substituting in (6) we obtain (2) □.

The following is a slight adaptation of lemma 1.4 of [4].

Lemma 3 [4, lemma 1.4] If X is a positive, right-continuous adapted process and B is an increasing, previsible process with $X_0=B_0=0$, such that for all finite stopping times T; $E[X_T] \leq E[B_T]$, then for each slowly increasing function F there exists a constant C_F such that

$$C_F\, E[F(X^*_\infty)] \leq E[F(B_\infty)] \quad .$$

Lemma 4 There exists a universal constant K such that for all $M,N\in H^1$

$$KE\left[\sup_a \sup_t |L^a_t(M)-L^a_t(N)|\right] \geq E[(M-N)^*_\infty - |M_0-N_0|] \qquad (7)$$

Proof Integrating the inequality (2) with respect to c we obtain

$$8E[D_\infty[(M-N)^*_\infty - |M_0-N_0|]] = 8\int_0^\infty E[D_\infty I_{(\tau_c<\infty)}]dc$$

$$\geq \int_0^\infty c\, P(\tau_{2c}<\infty)dc = \tfrac{1}{2}E[((M-N)^*_\infty - |M_0-N_0|)^2]$$

which gives, using Hölder's inequality

$$KED^2_\infty \geq E[((M-N)^*_\infty - |M_0-N_0|)^2] \qquad (8)$$

$(M-N)^*_t - |M_0-N_0|$ is a positive right-continuous adapted process whilst D is continuous (and so previsible) as a consequence of the joint continuity in (a,s) of $(L^a_s(M)$ and $(L^a_s(N).)$ Applying (8) to the martingales M^T and N^T we see that $[(M-N)^*_t - |M_0-N_0|]^2$ and KD^2_t satisfy the conditions of lemma 3 so setting $F(x) = x^{\frac{1}{2}}$ we obtain (7) □.

Lemma 5 There exists a universal constant c such that

$$c\,E[\sup_a \sup_t |L_t^a(M)-L_t^a(N)|] \geq E[\langle \widehat{M-N} \rangle_\infty^{\frac{1}{2}}] \qquad (9)$$

Proof Set

$$\nu = \inf\{t\geq 0 : |M_t-M_0| \vee |N_t-N_0| \geq \tfrac{1}{2}|M_0-N_0|\}.$$

As the ranges of $(M_t ; t\leq\nu)$ and $(N_t; t\leq\nu)$ are disjoint $L_t^a(M)\wedge L_t^a(N)=0$ for each a , for $t\leq\nu$. Thus

$$D_\nu = (\sup_a L_\nu^a(M)) \vee (\sup_a L_\nu^a(N)) \geq \tfrac{1}{2}(\sup_a L_\nu^a(M)) + \tfrac{1}{2}(\sup_a L_\nu^a(N))$$

and so by theorem 3.1 of [1]

$$c\,ED_\infty \geq E(\hat{M}_\nu^* + \hat{N}_\nu^*)$$

which leads to

$$4c\,ED_\infty \geq 4E((\hat{M}_\nu^*+\hat{N}_\nu^*)I_{(\nu<\infty)}) + 4E((\hat{M}_\infty^*+\hat{N}_\infty^*)I_{(\nu=\infty)})$$

$$\geq 2E(|M_0-N_0|I_{(\nu<\infty)}) + E((\hat{M}-\hat{N})_\infty^* I_{(\nu=\infty)}) \qquad (10)$$

Adding (7) and (10) we obtain

$$CED_\infty \geq E((M-N)_\infty^* - |M_0-N_0|) + 2E(|M_0-N_0|I_{(\nu<\infty)}) + E((\hat{M}-\hat{N})_\infty^* I_{(\nu=\infty)})$$

$$= E[((M-N)_\infty^* - |M_0-N_0|)I_{(\nu=\infty)} + ((M-N)_\infty^* + |M_0-N_0|)I_{(\nu<\infty)}]$$

$$+ E((\hat{M}-\hat{N})_\infty^* I_{(\nu=\infty)})$$

$$\geq E[(\hat{M}-\hat{N})_\infty^*]$$

We obtain (9) by observing that $(\hat{M}-\hat{N}) = (\widehat{M-N})$ and by applying the Burkholder-Davis-Gundy inequality with $p=1$ □.

Lemma 6 [4, lemma 1.1] If A and B are increasing, previsible processes and there exist $a,q>0$ such that for all pairs of finite stopping times $S\leq T$

$$E[(A_T I_{(T>0)} - A_S I_{(S>0)})^q] \leq aE[B_T^q I_{(T>S)}]$$

then for every moderate function F there exists a $c=c(a,q,F)$ such that

$$E[F(A_\infty)] \leq c\, E[F(B_\infty)]$$

Proof of theorem 1 For $M,N\in H^1$ set

$$m_t = M^T_{(S+t)}$$

$$n_t = N^T_{(S+t)}$$

We see that

$$L^a_t(m) = L^a_{S+t}(M^T) - L^a_S(M^T)$$

$$L^a_t(n) = L^a_{S+t}(N^T) - L^a_S(N^T)$$

and, applying lemma 5 to these $(\mathcal{F}_{S+t})$-martingales we obtain, with some simple manipulation

$$2ED_T I_{(S<T)} \geq E[\sup_{S\leq s\leq T}\ \sup_a\ |(L^a_s(M) - L^a_S(M)) - (L^a_s(N) - L^a_S(N))|]$$

$$\geq cE[\langle \hat{m}-\hat{n}\rangle^{\frac{1}{2}}_{T-S}]$$

$$\geq cE[\langle \hat{M}-\hat{N}\rangle^{\frac{1}{2}}_T - \langle \hat{M}-\hat{N}\rangle^{\frac{1}{2}}_S]$$

So we obtain (1) by lemma 6 with $F(x) = x^p$. To complete the proof in the case $p<1$, we apply the above inequality to M^{S_n} and N^{S_n} , where $S_n = \inf\{t : |M_t| \vee |N_t| \geq n\}$. and then use monotone convergence to obtain the result. □

Corollary 7 If F is a moderate function there exists a universal constant C_F such that

$$C_F\, E(F(\sup_a \sup_t |L_t^a(M) - L_t^a(N)|) \geq E(F((<M-N>_\infty - <M-N>_0)^{\frac{1}{2}}))$$

for all continuous local martingales M and N .

The proof follows immediately from the above.

Remark Inequality (B) [Barlow and Yor] leads one to ask whether there exists a universal c such that

$$c\, E[\sup_a \sup_t |L_t^a(M) - L_t^a(N)|] \geq \|(M-N)^*_\infty\|_1^{1-\varepsilon} \|M^*_\infty + N^*_\infty\|_1^{\varepsilon}$$

for some $\varepsilon>0$. The answer is no. For, take a brownian motion B with $B_0=0$, let $T = \inf\{t\geq 0 : |B_t|=1\}$ and take $\delta>0$; setting $M = B^{T+\delta}$ $N = B^T$ we find that $D_\infty = \sup_a \sup_{T\leq t\leq T+\delta} |L_t^a(B) - L_T^a(B)| = \sup_a (L^a_{T+\delta}(B) - L^a_T(B))$ and so, by [1] , $ED_\infty \leq c\delta^{\frac{1}{2}}$ whilst $E(M^*_\infty + N^*_\infty) \geq 2$ and $E[(M-N)^*_\infty] \geq C\delta^{\frac{1}{2}}$ so that

$$\frac{\|(M-N)^*_\infty\|_1^{1-\varepsilon} \|M^*_\infty + N^*_\infty\|_1^{\varepsilon}}{ED_\infty} \geq K\delta^{-\frac{1}{2}\varepsilon} \longrightarrow \infty \quad \text{as} \quad \delta \downarrow 0 .$$

We now present our second result.

Theorem 8 If M and N are in H^1 then, for each $a \in \mathbb{R}$,

$$\|L_\infty^a(M) - L_\infty^a(N)\|_\infty = \|\sup_t |L_t^a(M) - L_t^a(N)|\|_\infty$$

Proof Let $\eta = \text{ess sup}|L_\infty^a(M) - L_\infty^a(N)|$, and define

$$\sigma = \inf\{t\geq 0 : L_t^a(M) - L_t^a(N) \geq \eta + 2\varepsilon\}$$

$$\tau = \inf\{t\geq \sigma : L_t^a(M) - L_t^a(N) \leq \eta + \varepsilon\}$$

Since $L_\infty^a(M) - L_\infty^a(N) \leq \eta$ we see that $(\sigma<\infty)=(\tau<\infty)$. Consider

$$|N_\tau - a| - |N_\sigma - a| = (|N_\tau - a| - |N_\sigma - a|)I_{(\sigma<\infty)} =$$

$$L_\tau^a(N) - L_\sigma^a(N) - \int_\sigma^\tau \text{sgn}(N_s - a)\,dN_s \qquad (11)$$

$N\in H^1$ so the stochastic integral in (11) is uniformly integrable so, by the optional sampling theorem

$$E[(|N_\tau - a| - |N_\sigma - a|)I_{(\sigma<\infty)}] = E[L_\tau^a(N) - L_\sigma^a(N)]$$

But on $(\tau<\infty)=(\sigma<\infty)$, $N_\tau = a$ so

$$0 \geq E[(|N_\tau - a| - |N_\sigma - a|)I_{(\sigma<\infty)}] = E[L_\tau^a(N) - L_\sigma^a(N)] \geq 0 \qquad (12)$$

Now

$$[L_\tau^a(N) - L_\sigma^a(N)]I_{(\sigma<\infty)} = [(L_\tau^a(M) - (\eta+\varepsilon)) - (L_\sigma^a(M) - (\eta+2\varepsilon))]I_{(\sigma<\infty)}$$

$$\geq \varepsilon I_{(\sigma<\infty)},$$

so we conclude from (12) that $0 \geq \varepsilon P(\sigma<\infty)$. As ε is arbitrary

$$\sup_t (L_t^a(M) - L_t^a(N)) \overset{a.s.}{\leq} \eta$$

and we may deduce the same inequality with M and N reversed. □

Corollary 9 If M and N are in H^1 with $M \neq M_0$ then $M=N$ if and only if for each a $L^a_\infty(M)=L^a_\infty(N)$.

Proof The reverse implication is clear. Now suppose $M_0=N_0$ then, since $D_\infty=0$, theorem 1 implies that $E(M-N)^*_\infty=0$ so that $M=N$. Suppose now $M_0 \neq N_0$, set $\nu=\inf\{t\geq 0 : |M_t-M_0| \vee |N_t-N_0| = \frac{1}{2}|M_0-N_0|\}$ then, since the ranges of $(M_t;t\leq\nu)$ and $(N_t;t\leq\nu)$ are distinct we may conclude that $L^a_\nu(M) \wedge L^a_\nu(N)=0$ but $L^a_\nu(M)=L^a_\nu(N)$ so $L^a_\nu(M)=L^a_\nu(N)=0$ $a\in\mathbb{R}$ and so we conclude that $E((M-M_0)^*_\infty)=0$ and so $M_0=M_t$ for $t\leq\nu$ and thus $(\nu=\infty)$ and $M=M_0$ which contradicts the initial assumption. □

Remark In fact, to conclude that M=N, it is sufficient that $L^a_\infty(M)=L^a_\infty(N)$ holds for all $a\in$range (M); the proof is left to the reader.

Acknowledgements

The author would like to thank M.T. Barlow for suggesting this problem and would like to thank M.T. Barlow, M. Yor and D.P. Kennedy for helpful criticism on its presentation.

References

[1] Barlow, M.T. and Yor, M.: (Semi-) Martingale inequalities and local times. Z. Wahrscheinlichkeitstheorie verw. Gebiete 55, 237-254 (1981).

[2] Barlow, M.T. and Yor, M.: Semimartingale inequalities via the Garsia-Rodemich-Rumsey lemma, and applications to local times. (To appear in Journal of Funct. Anal.)

[3] Yor, M.: Sur la continuité des temps locaux associés à certaines semi-martingales. Temps Locaux. Astérisque. 52-53, 23-36 (1978).

[4] Lenglart, E., Lépingle, D., Pratelli, M.: Présentation unifiée de certaines inégalités de la théorie des

martingales. Sém. Probab. XIV, Lect. Notes in Maths. 784, Berlin-Heidelberg-New York: Springer (1980).

[5] Azéma, J. and Yor, M.: En guise d'introduction. Temps Locaux. Astérisque 52-53, 3-16 (1978)

[6] Yor, M.: Rappels et préliminaires généraux. Temps Locaux. Astérisque 52-53, 17-22 (1978)

[7] Ray, D.: Sojourn times of diffusion processes. Ill.J.Math. 7, 615-630 (1963).

[8] Knight, F.: Random walks and the sojourn density process of brownian motion. Trans. Amer. Math. Soc. 109, 58-86 (1963).

[9] Meyer, P.A.: Un cours sur les intégrales stochastiques. Sém. Probab. Strasbourg X Lect. Notes in Maths. 511, Berlin-Heidelberg-New York: Springer (1976)

[10] Jacod, J.: Calcul Stochastique et Problèmes de Martingales Lect. Notes in Maths 714, Berlin-Heidelberg-New York: Springer (1979).

[11] Dellacherie, C. and Meyer, P.A.: Probability and Potential Amsterdam-New York-Oxford: North Holland (1978).

Statistical Laboratory,
16 Mill Lane,
University of Cambridge,
CAMBRIDGE CB2 1SB
England.