

Local solutions of the Landau equation with rough, slowly decaying initial data [☆]

Christopher Henderson ^{a,*}, Stanley Snelson ^b, Andrei Tarfulea ^c

^a Department of Mathematics, University of Arizona, Tucson, AZ 85721, United States of America

^b Department of Mathematical Sciences, Florida Institute of Technology, Melbourne, FL 32901, United States of America

^c Department of Mathematics, Louisiana State University, Baton Rouge, LA 70803, United States of America

Received 11 December 2019; received in revised form 17 April 2020; accepted 30 April 2020

Available online 12 May 2020

Abstract

We consider the Cauchy problem for the spatially inhomogeneous Landau equation with soft potentials in the case of large (i.e. non-perturbative) initial data. We construct a solution for any bounded, measurable initial data with uniform polynomial decay in the velocity variable, and that satisfies a technical lower bound assumption (but can have vacuum regions). For uniqueness in this weak class, we have to make the additional assumption that the initial data is Hölder continuous. Our hypotheses are much weaker, in terms of regularity and decay, than previous large-data well-posedness results in the literature. We also derive a continuation criterion for our solutions that is, for the case of very soft potentials, an improvement over the previous state of the art.

© 2020 L'Association Publications de l'Institut Henri Poincaré. Published by Elsevier B.V. All rights reserved.

Résumé

Nous considérons le problème de Cauchy pour l'équation de Landau non-homogène en espace avec potentiels mous dans le cas de grandes (i.e. non perturbatrices) données initiales. Nous construisons une solution pour toute donnée initiale bornée et mesurable à décroissance polynomiale uniforme dans la variable de vitesse, et qui satisfait une hypothèse technique de borne inférieure (il est toujours permis d'avoir des régions de vide). Pour être unique dans cette famille générale, nous devons supposer que les données initiales sont Hölder continues. Nos hypothèses sont beaucoup plus faibles, en termes de régularité et de décroissance, que les résultats de la littérature sur le caractère bien-posé dans le cas de grandes données. Nous dérivons également un critère de continuation pour nos solutions qui est, pour les potentiels très mous, une amélioration par rapport à l'état de l'art.

© 2020 L'Association Publications de l'Institut Henri Poincaré. Published by Elsevier B.V. All rights reserved.

Keywords: Landau equation; Large data well-posedness; Vacuum regions; Classical solutions; Kinetic equations

[☆] AT was partially supported by NSF grant DMS-1816643. SS was partially supported by a Ralph E. Powe Award from ORAU. CH was partially supported by NSF grant DMS-2003110.

* Corresponding author.

E-mail addresses: ckhenderson@math.arizona.edu (C. Henderson), ssnelson@fit.edu (S. Snelson), tarfulea@lsu.edu (A. Tarfulea).

1. Introduction

We consider the spatially inhomogeneous Landau equation in the whole space: for $(t, x, v) \in \mathbb{R}_+ \times \mathbb{R}^3 \times \mathbb{R}^3$, the solution $f(t, x, v) \geq 0$ satisfies

$$(\partial_t + v \cdot \nabla_x) f = Q_L(f, f) = \nabla_v \cdot (\bar{a}^f \nabla_v f) + \bar{b}^f \cdot \nabla_v f + \bar{c}^f f, \quad (1.1)$$

where the nonlocal coefficients are defined by

$$\begin{aligned} \bar{a}^f(t, x, v) &:= a_\gamma \int_{\mathbb{R}^3} \left(I - \frac{w}{|w|} \otimes \frac{w}{|w|} \right) |w|^{\gamma+2} f(t, x, v-w) dw, \\ \bar{b}^f(t, x, v) &:= b_\gamma \int_{\mathbb{R}^3} |w|^\gamma w f(t, x, v-w) dw, \\ \bar{c}^f(t, x, v) &:= c_\gamma \int_{\mathbb{R}^3} |w|^\gamma f(t, x, v-w) dw, \end{aligned} \quad (1.2)$$

and $a_\gamma, b_\gamma, c_\gamma$ are constants with $a_\gamma, c_\gamma > 0$. We are concerned with the case of *soft potentials*, i.e. $\gamma \in [-3, 0)$. When $\gamma = -3$, the formula for \bar{c}^f must be replaced by $\bar{c}^f = cf$. In fact, the constants are such that $\bar{b}_i^f = -\sum \partial_{v_j} \bar{a}_{ij}^f$, so that if $f \in C_v^2$, (1.1) may equivalently be written in non-divergence form:

$$(\partial_t + v \cdot \nabla_x) f = \text{tr}(\bar{a}^f D_v^2 f) + \bar{c}^f f. \quad (1.3)$$

We use both forms of the equation.

The $\gamma = -3$ case of (1.1), called the Landau-Coulomb equation, is used in plasma physics to model the density $f(t, x, v)$ of charged particles in phase space as it evolves in time. For $\gamma > -3$, the equation arises as a formal limit of the Boltzmann equation as grazing collisions (such that the angle between pre- and post-collisional velocities goes to zero) predominate. In addition to its own interest, part of the motivation for studying the Landau equation (particularly the non-Coulomb cases) is to shed light on the Boltzmann equation without angular cutoff (see, e.g., [30,10] or [5] for the physical background).

Until relatively recently, the theory of classical solutions of the inhomogeneous Landau equation focused mainly on global-in-time solutions that are perturbations of a (global) Maxwellian equilibrium state (i.e. $\mu(t, x, v) = c_1 e^{-c_2 |v|^2}$ with $c_1, c_2 > 0$). This study began with the work of Guo [21] in 2002 and was subsequently extended by many authors; see Section 1.3 below for a partial bibliography.

Regarding the non-perturbative case, it is an outstanding open problem to construct global-in-time solutions for large (i.e. non-perturbative) initial data. This problem is extremely difficult, perhaps out of reach for the foreseeable future. As such, our focus is on two more modest but still nontrivial goals: local well-posedness among a “reasonable” class of large initial data, and physically meaningful conditional assumptions on f under which global well-posedness holds.

There are two previous large-data existence results for the inhomogeneous Landau equation we are aware of: He-Yang [22] constructed solutions for the case $\gamma = -3$ with spatially periodic initial data satisfying $|v|^P f_{\text{in}} \in H^7(\mathbb{T}^3 \times \mathbb{R}^3)$, for P an exponent on the order of 100, and in our previous work [24], we constructed solutions for initial data satisfying $e^{\rho|v|^2} f_{\text{in}}$ in an H^4 space¹ on \mathbb{R}^6 for some $\rho > 0$.² The strength of these hypotheses, in terms of regularity and decay, is comparable to the state of the art for the non-cutoff Boltzmann equation [2–4] but is especially unsatisfactory when compared to (i) what is needed to make sense of the equation and (ii) the conditions required for good *a priori* control of solutions, which are zeroth-order and impose only mild polynomial decay in v (see Section 1.2). The main results of this paper go a long way toward bridging this gap, by proving existence of solutions in a merely polynomially-weighted L^∞ space, which requires no control on derivatives or exponential moments of

¹ More precisely, [24] took f_{in} in a “uniformly local” space H_{ul}^4 , which imposes no decay as $|x| \rightarrow \infty$.

² Shortly after the present work was submitted, Chaturvedi [11] proved local existence for the Landau equation with hard and Maxwellian potentials ($\gamma \in [0, 1]$), with similarly strong assumptions, namely that f_{in} has exponential decay in v and lies in a tenth-order Sobolev space.

f_{in} , as well as uniqueness, under the additional assumption that the initial data is Hölder continuous. We also slightly improve the conditions under which solutions can be extended past a given time.

1.1. Main results

In order to state our results, we define weighted Lebesgue spaces as follows: let $\langle v \rangle = \sqrt{1 + |v|^2}$, $1 \leq p \leq \infty$, $k \in \mathbb{R}$, and Ω be any set of the form \mathbb{R}^3 or $\omega \times \mathbb{R}^3$ with the \mathbb{R}^3 variable denoted v , then define

$$\|g\|_{L^{p,k}(\Omega)} = \|\langle v \rangle^k g\|_{L^p(\Omega)}.$$

We also require the kinetic Hölder spaces, denoted C_{kin}^α and $C_{\text{kin}}^{2,\alpha}$ for $\alpha \in (0, 1)$ and defined in Section 1.5, that are analogous to the standard parabolic Hölder spaces. We use the standard Hölder spaces, C^α and $C^{2,\alpha}$, as well.

One quantity that plays a large role in our analysis is

$$\Psi(t) := \begin{cases} \|f\|_{L_{t,x}^\infty L_v^{1,2}([0,t] \times \mathbb{R}^6)}, & \gamma \in (-2, 0), \\ \|f\|_{L_{t,x}^\infty L_v^1([0,t] \times \mathbb{R}^6)} + \|f\|_{L_{t,x}^\infty L_v^p(\mathbb{R}^6)}, & \gamma \in [-3, -2], \end{cases} \quad (1.4)$$

where $p > 3/(3 + \gamma)$ and $p = \infty$ for $\gamma = -3$. In the case $\gamma \in [0, 2]$, Ψ is related to the hydrodynamic quantities of mass density and energy density (see Section 1.2 below). Notice that Ψ is bounded by the $L^{\infty,k}$ norm of f if $k > 5$.

We require the following condition on our initial data, recalled from [24]. It ensures that the self-generating lower bounds on our solution are uniform in x (after a small time has passed):

Definition 1.1. A function $g : \mathbb{R}^3 \times \mathbb{R}^3 \rightarrow [0, \infty)$ is *well-distributed with parameters* $R, \delta, r > 0$ if, for every $x \in \mathbb{R}^3$, there exists $x_m \in B_R(x)$ and $v_m \in B_R(0)$ such that $g \geq \delta \mathbb{1}_{B_r(x_m, v_m)}$.

Our first result establishes the existence of solutions in $L^{\infty,k}([0, T] \times \mathbb{R}^6)$:

Theorem 1.2. Let $k > \max\{5, 15/(5 + \gamma)\}$. Assume $\|f_{\text{in}}\|_{L^{\infty,k}(\mathbb{R}^6)} \leq K$, $f_{\text{in}}(x, v) \geq 0$ in \mathbb{R}^6 , and f_{in} is well-distributed with parameters δ, r , and R . Then:

- (i) **(Existence)** There exists $T > 0$ and $f \geq 0$, such that, for any compact $\Omega \subset (0, T] \times \mathbb{R}^6$, $f \in C_{\text{kin}}^{2,\alpha}(\Omega)$ for some $\alpha \in (0, 1)$, and f satisfies (1.1) classically. Each of T and $\|f\|_{L^{\infty,k}([0,T] \times \mathbb{R}^6)}$ depend only on γ, K, δ, r , and R , and α depends on the same quantities plus Ω .
- (ii) **(Matching with initial data)** For any compact $K_x \subset \mathbb{R}^3$ and any $\phi(t, x, v)$ satisfying, for some $\eta > 0$,

$$\begin{aligned} \phi &\in C_t L_{x,v}^{2,-7/2+\eta}([0, T] \times K_x \times \mathbb{R}^3), \quad (\partial_t + v \cdot \nabla_x)\phi \in L^{2,-7/2+\eta}([0, T] \times K_x \times \mathbb{R}^3) \\ \nabla_v \phi &\in L^{2,1+\gamma/2}([0, T] \times K_x \times \mathbb{R}^3) \quad \text{and} \quad \text{supp}(\phi) \subset [0, T] \times K_x \times \mathbb{R}^3, \end{aligned}$$

the solution f constructed in (i) satisfies

$$\begin{aligned} &\int_{\mathbb{R}^6} f_{\text{in}}(x, v) \phi(0, x, v) \, dx \, dv \\ &= \int_{[0,T] \times \mathbb{R}^6} \left(f(\partial_t + v \cdot \nabla_x)\phi - \nabla_v \phi \cdot (\bar{a}^f \nabla_v f) - f \bar{b}^f \cdot \nabla_v \phi \right) \, dx \, dv \, dt. \end{aligned} \quad (1.5)$$

- (iii) **(Higher regularity)** Let T_E be the maximal time of existence of the solution constructed above, and let $\bar{T} \in (0, T_E]$. For any partial derivative $\partial_t^j \partial_x^\beta \partial_v^\eta$, there exists $k_M > 0$ and ℓ_M depending only on $\Psi(\bar{T}), \bar{T}, \gamma, \delta, r, R$, and $M = 2j + 3|\beta| + |\eta|$, such that if $f_{\text{in}} \in L^{\infty,k_M}(\mathbb{R}^6)$, then

$$\partial_t^j \partial_x^\beta \partial_v^\eta f \in L^{\infty,k_M-\ell_M}([0, \bar{T}] \times \mathbb{R}^6)$$

and is continuous for all positive times.

A few comments are in order. Firstly, if the initial data is continuous, we can show that f does match the initial data in the pointwise sense (see Proposition 3.1). Secondly, by “classical” solutions, we mean those elements of $C_{\text{kin},\text{loc}}^{2,\alpha}$ that satisfy (1.1) pointwise; this does not necessitate that $\partial_t f$ and $\nabla_x f$ exist pointwise but rather that $(\partial_t + v \cdot \nabla_x)f$ does (see Section 1.5 for a discussion of the kinetic Hölder spaces). Lastly, we point out that Theorem 1.2.(iii) implies that if $f_{\text{in}} \in L^{\infty,k}$ for all k , then f is smooth and has infinitely many moments, that is, $\partial_t^j \partial_x^\beta \partial_v^\eta \in L^{\infty,k}$ for any j, β, η , and k , for as long as Ψ remains bounded.

Next, we extend our continuation criterion from [24] to the solutions of Theorem 1.2, and sharpen it in the case of very soft potentials:

Theorem 1.3. *The solution f constructed in Theorem 1.2 can be extended for as long as the quantity Ψ remains finite, where $p > 3/(3 + \gamma)$ and $p = \infty$ for $\gamma = -3$. More formally, $T_E = \sup\{t \geq 0 : \Psi(t) < \infty\}$, where T_E is defined in Theorem 1.2.*

Because the solution constructed in Theorem 1.2 lies in a weak space relative to the order of the equation, uniqueness is a challenging issue, and in fact our proof requires stronger assumptions on f_{in} : Hölder continuity and a lower bound assumption that rules out vacuum regions in x (see Section 1.4 below for an explanation of these extra hypotheses). The uniqueness or non-uniqueness of the solutions of Theorem 1.2, without any extra assumptions on f_{in} , remains an open question.

In nonlinear equations where the initial data has low regularity, it is often more difficult to prove uniqueness than existence of solutions even when they are smooth immediately. For another example of such a situation, where uniqueness has not been established without extra hypotheses, even though the system regularizes instantaneously, see the work of Kiselev-Nazarov-Shterenberg [29] on the fractal Burgers equation (i.e., Burgers equation with fractional dissipation).

Theorem 1.4. *Assume that f_{in} satisfies:*

- $f_{\text{in}} \in C^\alpha$ for some $\alpha \in (0, 1)$, and $f_{\text{in}} \in L^{\infty,k}$ for some k ;
- there exist $r, \delta, R > 0$ such that for each $x \in \mathbb{R}^3$, there exists $|v_x| < R$ such that $f_{\text{in}} \geq \delta \mathbb{1}_{B_r(x, v_x)}$.

Let f be the solution constructed in Theorem 1.2, and let $T > 0$ be any time such that $\Psi(T) < \infty$. Then there exists k_α and $T_H \in (0, T]$ such that if $k \geq k_\alpha$, then $f \in C_{\text{kin}}^\alpha([0, T_H] \times \mathbb{R}^6)$ and, for any uniformly continuous $g \in L^{\infty, 5+\gamma+\eta}([0, T_H] \times \mathbb{R}^6)$ with $\eta > 0$ such that g solves (1.1) weakly and $g(t, x, v) \rightarrow f_{\text{in}}$ as $t \rightarrow 0+$, we have $f = g$. Moreover, if $f_{\text{in}} \in L^{\infty,k}$ for all $k \geq 0$, then the above holds with $T_H = T$. We have that k_α depends only on α and γ and T_H depends only on $r, \delta, R, k, \alpha, \gamma, \|f_{\text{in}}\|_{L^{\infty,k}}$, and $\|f_{\text{in}}\|_{C^\alpha}$.

Let us make some brief comments on Theorem 1.4. First, our notion of weak solution is made precise in Section 5 (see the comment after Proposition 5.2). Second, our assumptions on f_{in} imply, via interpolation between C^α and $L^{\infty,k}$, that

$$\langle v \rangle^m f \in C^\beta(\mathbb{R}^6) \quad \text{if } m \leq k \left(1 - \frac{\beta}{\alpha}\right),$$

see Lemma B.2. For the sake of convenience, we prove that the conclusion of Theorem 1.4 holds under the assumption that $f_{\text{in}} \in L^{\infty,k}(\mathbb{R}^6)$ and $\langle v \rangle^m f_{\text{in}} \in C^\alpha(\mathbb{R}^6)$ for m and k sufficiently large, depending on α and γ (see Proposition 4.4), which, by the above, is enough to establish Theorem 1.4. Finally, we note that the positivity condition, that is, the existence of δ, r , and R , is met in many standard cases such as for initial data that is continuous, periodic, and positive.

1.2. Conditional regularity and continuation

This paper fits within the program of seeking weaker conditions under which solutions of the Landau equation remain smooth and can be extended past a given time. To describe this recent thread of research in more detail, let us recall the following hydrodynamic quantities associated to the solution f :

$$M_f(t, x) = \int_{\mathbb{R}^3} f(t, x, v) \, dv, \quad (\text{mass density})$$

$$E_f(t, x) = \int_{\mathbb{R}^3} |v|^2 f(t, x, v) \, dv, \quad (\text{energy density})$$

$$H_f(t, x) = \int_{\mathbb{R}^3} f(t, x, v) \log f(t, x, v) \, dv. \quad (\text{entropy density})$$

We note that, in the homogeneous (i.e., x -independent) setting, $M_f(t)$ and $E_f(t)$ are conserved and $H_f(t)$ is non-increasing, and this behavior is crucial to the well-posedness theory.

In [18], Golse-Imbert-Mouhot-Vasseur established local Hölder regularity for solutions of (1.1) with M_f bounded above and away from zero, and E_f , H_f , and $\|f(t, x, \cdot)\|_{L_v^\infty}$ bounded above, uniformly in t and x . Under these conditions, the coefficients in (1.2) are all bounded above, and \tilde{a}^f is uniformly elliptic. The equation can then be treated as a linear kinetic Fokker-Planck equation with bounded measurable coefficients, and the main contribution of [18] was in adapting De Giorgi's method to such linear equations (see also [36,44].) In [7], Cameron-Silvestre-Snelson derived a conditional global L^∞ estimate in terms of the upper bounds for M_f , E_f , and H_f , and the lower bound for M_f , in the case $\gamma \in (-2, 0]$ (see [39] for a similar result for $\gamma \in (0, 1]$). Next, Henderson-Snelson [23] established C^∞ regularity in the same conditional regime, with the additional assumption that infinitely many L^1 moments of f in v are finite. This study also relied on the linear theory via Schauder estimates for kinetic equations, but made essential use of the coupling between f and the coefficients in the bootstrapping procedure. (See also the earlier result [12], which proved C^∞ regularity under stronger conditional assumptions.)

Most recently, in [24] the present authors combined local well-posedness with these conditional results to derive a continuation criterion and improved this criterion by establishing self-generating lower bounds for f that suffice to remove the assumptions that M_f is bounded below and H_f is bounded above. For very soft potentials ($\gamma \in [-3, -2]$), the methods of [23] and [24] all go through, but under stricter conditions on f that, unlike the hydrodynamic quantities above, are not physically meaningful. In the above notation, the continuation criterion established in [24] was

$$\tilde{\Psi}(t) := \begin{cases} \|f\|_{L_{t,x}^\infty L_v^{1,2}([0,t] \times \mathbb{R}^6)}, & \gamma \in (-2, 0), \\ \|f\|_{L_{t,x}^\infty L_v^{1,\ell}([0,t] \times \mathbb{R}^6)} + \|f\|_{L_{t,x,v}^\infty([0,t] \times \mathbb{R}^6)}, & \gamma \in [-3, -2], \end{cases} \quad (1.6)$$

with $\ell > 3|\gamma|/(5 + \gamma)$. For $\gamma \in [-3, -2]$, Theorem 1.3 above improves this criterion in two ways. The improvement from $\|f\|_{L_{t,x,v}^\infty}$ to $\|f\|_{L_{t,x}^\infty L_v^p}$ with p as in Theorem 1.3 was attainable with the methods of [24], but improving $\|f\|_{L_{t,x}^\infty L_v^{1,\ell}}$ to $\|f\|_{L_{t,x}^\infty L_v^1}$ requires the more general decay estimates of the current paper. Theorem 1.3 is also an improvement over [24] because it applies to solutions with more general initial data.

The significance of lower bounds for f , besides implying that vacuum regions in the initial data are instantly filled, is in getting a lower ellipticity constant for \tilde{a}^f , which is needed to apply local estimates (either De Giorgi or Schauder type). Taking f_{in} to be well-distributed (see Definition 1.1) guarantees this ellipticity constant is uniform in x , after a short time has passed. The extra lower bound assumption of Theorem 1.4 ensures this ellipticity constant does not degenerate as $t \rightarrow 0$. It would be interesting to remove these structural lower bounds for f_{in} from the results in the current paper, both for a more robust local well-posedness theory, and because doing so may lead to better understanding of continuation for states that are not controlled uniformly in x . Doing so would, however, require a completely new approach that does not depend on the regularity estimates near $t = 0$.

There is a parallel program of conditional regularity for the non-cutoff Boltzmann equation: see [37,28,26,25,27]. To pair these conditional estimates with a compatible local existence result, one would need to work with polynomially-decaying initial data. This has only been accomplished for certain ranges of the physical parameters (see [33]). In a forthcoming article, we plan to extend this result to a less restrictive setting. See also the review [34] for more on the conditional regularity of both equations.

1.3. Related work

There are many existence and regularity results for the spatially homogeneous (x -independent) Landau equation, see [6,43,15,1,45,19,20,14,38,17] and the references therein. In this setting, large-data global solutions are known to exist in the cases $\gamma \in [-2, 1]$, but for $\gamma \in [-3, -2)$, the problem remains open.

In the inhomogeneous setting considered here, a suitable notion of weak solution has been defined. Global solutions in this class have been established by Villani [42] for general initial data (see also Lions [31] and Alexandre-Villani [5]). The uniqueness and regularity of these solutions are not understood.

As mentioned above, there is a large literature on close-to-Maxwellian solutions of the Landau equation that exist globally and converge to equilibrium as $t \rightarrow \infty$: see for example [21,35,40,41,9,8,16]. The majority of these papers work with initial data that is close to a Maxwellian in an exponentially-weighted (in v) norm, which implies f_{in} decays exponentially at worst, but the work of Carrapatoso-Mischler [8] improves this to a polynomially-weighted $H_x^2 L_v^2$ norm (see also [9], which works with polynomially-weighted norms in the case $\gamma > 0$.) In terms of regularity, early results took f_{in} in a high-order Sobolev space, and subsequent works gradually enlarged the allowable space.

The recent interesting work of Duan-Liu-Sakamoto-Strain [16] constructs mild solutions with initial data close to a Maxwellian in an exponentially-weighted $L_k^1 L_v^2$ space (here, L_k^1 refers to the Wiener algebra in x which contains all C^1 functions but not all Hölder continuous functions) on $\mathbb{T}^3 \times \mathbb{R}^3$. When working on a bounded spatial domain instead of \mathbb{T}^3 , they require an additional derivative in this space. Uniqueness is also shown, as is convergence to the Maxwellian, which is their main interest.

A somewhat different setting was considered by Luk in [32]. For $\gamma \in (-2, 0)$, he has shown the global existence of a solution with initial data close to the vacuum state $f \equiv 0$.

To our knowledge, polynomially decaying solutions have not been established previously for the Landau equation except in $\mathbb{T}_x^3 \times \mathbb{R}_v^3$ (as in [22,8,16]) or when f_{in} is “localized” in x (e.g., $f_{\text{in}} \in L^2(\mathbb{R}^6)$ as in [11]). This distinction is important because, in these settings, no moment loss can occur due to the transport operator $(\partial_t + v \cdot \nabla_x)$; however, on $\mathbb{R}_x^3 \times \mathbb{R}_v^3$ without any assumption of decay for large x , the transport operator causes moment loss. Indeed, it is easy to construct solutions to $(\partial_t + v \cdot \nabla_x)g = 0$ on \mathbb{R}^6 such that $g(0, \cdot, \cdot) \in L^2$ but $g(t, \cdot, \cdot) \notin L^2$ for some $t > 0$.

1.4. Proof ideas

1.4.1. Existence

First, let us point out some disadvantages of the usual method of L^2 -based energy estimates like the one pursued in [24]. Because of their physical relevance, we wish for our class of initial data to include (global) Maxwellians, which do not have finite $L_{x,v}^2(\mathbb{R}^6)$ norm. Thus, it makes sense to estimate $\|\varphi f(t)\|_{L_{x,v}^2}$ for cut-off functions φ in x . To this end, after multiplying (1.1) by $\varphi^2 f$ and integrating over \mathbb{R}^6 , we find, after some formal computations,

$$\frac{1}{2} \frac{d}{dt} \|\varphi f\|_{L_{x,v}^2}^2 = \int_{\mathbb{R}^6} \left(-\frac{1}{2} f^2 v \cdot \nabla_x (\varphi^2) - \varphi^2 \nabla_v f \cdot (\tilde{a}^f \nabla_v f) + \frac{1}{2} \tilde{c}^f \varphi^2 f^2 \right) dx dv. \quad (1.7)$$

The first difficulty is that the $f^2 v \cdot \nabla_x (\varphi^2)$ term may be unbounded for large $|v|$. Thus, this term cannot be controlled by the unweighted $L_{x,v}^2$ norm of f . In [24], we avoid this issue by dividing the solution f by a time-dependent Gaussian, i.e. studying the equation for $g = e^{(\rho - \kappa t)(v)^2} f$, which has an extra term of $\kappa \langle v \rangle^2 g$ with the right sign to absorb the other terms with growth in v in the energy estimates (this approach was applied earlier to the Boltzmann equation in [2–4]). However, this method requires the initial data f_{in} to have Gaussian decay in v , and we want to get around this requirement.

If the spatial domain were the torus \mathbb{T}^3 instead of \mathbb{R}^3 , no cut-off would be necessary so this term would disappear from the energy estimates, which is one reason the spatially periodic case is simpler. It is important to note that this is not a purely technical issue; as discussed in Section 1.3, transport can cause moment loss when working on the whole space. See Morimoto-Yang [33] for a different approach to this obstacle, in the context of the Boltzmann equation.

The second difficulty is that the coefficients are not adequately controlled by the L^2 -norm of f . For example, \tilde{c}^f clearly must be bounded in L^∞ in order to make the last term in (1.7) finite for φf merely in L^2 . Unfortunately, $\|\tilde{c}^f\|_{L^\infty}$ is not controlled by $\|\varphi f\|_{L^2}$, and one instead needs a bound on $\|\varphi f\|_{L_{x,v}^p}$ for $p \geq 2$ depending on γ . This necessitates bounds on higher derivatives of f in order to use an embedding theorem.

When searching for bounds on derivatives of f , we encounter another issue. To obtain such bounds, one might be tempted to differentiate the equation and then follow a similar strategy as above to obtain estimates. Unfortunately, differentiating the equation brings up new technical difficulties—for example, when derivatives fall on \bar{a}^f , positive-definiteness is lost, so the corresponding term does not have a good sign anymore. Thus, it is significantly harder to obtain higher order energy estimates.

The key observation that allows us to overcome the above issues is to work in $L^{\infty,k}([0, T] \times \mathbb{R}^6)$. The associated norm is not affected by the moment loss due to the transport operator and the coefficients \bar{a}^f and \bar{c}^f can be bounded in terms of it when $k > 5 + \gamma$. The basis of our construction is an estimate (Lemma 2.3) on $\|f\|_{L^{\infty,k}}$ that follows from a comparison principle argument with barriers of the form $e^{\beta t} \langle v \rangle^{-k}$. It is interesting to note that Lemma 2.3 does not rely on lower bounds for the matrix \bar{a}^f or on the anisotropy of \bar{a}^f in v . The $L^{\infty,k}$ estimate gives good upper bounds on the coefficients, which then allows us to adapt the mass-spreading theorem of [24] to get lower bounds of f that imply coercivity of \bar{a}^f .

The $L^{\infty,k}$ -estimates established above are not sufficient to ensure that the constructed solution f is classical; however we can obtain higher regularity via (properly rescaled versions of) the De Giorgi and Schauder estimates previously established in [18,23]. To pass from these *a priori* estimates—which require good smoothness and decay—to an existence theorem, we must approximate $f_{\text{in}} \in L^{\infty,k}(\mathbb{R}^6)$ by smooth, compactly supported functions $f_{\text{in}}^\varepsilon$ and apply our previous existence theorem from [24] for H^4 , rapidly-decaying data. Using the $L^{\infty,k}$ estimate of Lemma 2.3 and—crucially—our continuation criterion from [24], we can extend these approximate solutions up to a time independent of ε . This step is where the restriction $k > \max\{5, 15/(5 + \gamma)\}$ comes from, because for such k , $\|f\|_{L^{\infty,k}}$ controls the quantities in the continuation criterion of [24] (see (1.6)). Finally, we can apply the local regularity estimates of [18,23] to obtain a solution f in $C_{\text{kin,loc}}^{2,\alpha}$ by compactness.

1.4.2. Uniqueness

To demonstrate some of the difficulties in proving uniqueness in a weak space, let us consider two solutions f and g with the same initial data. Then $w := f - g$ satisfies

$$\partial_t w + v \cdot \nabla_x w = \text{tr}(\bar{a}^g D_v^2 w) + \text{tr}(\bar{a}^w D_v^2 f) + \bar{c}^g w + \bar{c}^w f.$$

Ignoring the growth of terms on the right for large v (which we can deal with by multiplying w by a polynomial weight), the most difficult term in this equation to bound in terms of w is $\text{tr}(\bar{a}^w D_v^2 f)$. With initial data only in $L^{\infty,k}$, we certainly cannot expect a uniform-in-time bound on $D_v^2 f$, but for a Grönwall-style argument, an upper bound that is integrable in t is good enough. Schauder estimates for kinetic equations, along with a standard interpolation between $C^{2,\alpha}$ and C^α , provide a bound like

$$\|D_v^2 f\|_{L^\infty([t/2, t] \times \mathbb{R}^6)} \lesssim t^{-1+r(\alpha)} \|f\|_{C_{\text{kin}}^\alpha([t/2, t] \times \mathbb{R}^6)},$$

where $r(\alpha) = \alpha^2/(6 - \alpha)$. (Again, we are ignoring velocity weights. See Lemma 4.1 for the precise statement.) At this point, to bound the C_{kin}^α norm of f , one could try to apply the De Giorgi-type estimate of [18] in a kinetic cylinder of radius $\sim t^{-1/2}$ centered at each (t, x, v) , but the constant in this estimate degenerates like $t^{-\alpha/2}$, giving a total decay like $D_v^2 f(t, x, v) \lesssim t^{-1-\alpha/2+r(\alpha)}$, which is not integrable.

We are therefore led to take initial data f_{in} that is Hölder continuous, and try to propagate the Hölder modulus forward in time. This is the subject of Section 4. Our method of proof adapts an idea used in [13] for the forced critical SQG equation: for $(t, x, v, \chi, \nu) \in \mathbb{R}_+ \times \mathbb{R}^6 \times B_1(0)^2$ and $\ell > 0$, define

$$g(t, x, v, \chi, \nu) = \frac{|f(t, x + \chi, v + \nu) - f(t, x, v)|^2}{(|\chi|^2 + |\nu|^2)^\alpha} \langle v \rangle^\ell.$$

The function g is chosen so that the size of g in $L_{x,v,\chi,\nu}^\infty$ controls the weighted $C_{x,v}^\alpha$ norm³ of f . The factor $\langle v \rangle^\ell$ is there to account for polynomial decay in the Hölder modulus for large $|v|$. Calculating the equation satisfied by g , it can be shown that all the terms either respect a maximum principle, or can be bounded by a constant times g , so the

³ in the Euclidean Hölder metric, i.e. without the kinetic scaling of $\rho(z, z')$. This choice is imposed on us by the proof—the reason is explained in Section 4.

L^∞ norm of g is bounded for some positive amount of time. Since this argument only gives Hölder continuity in x and v for f , we also have to show, via the equation, that this implies Hölder continuity in all three variables. This is established by a general technique that may be of independent interest (see Appendix A). This argument provides a method to establish Hölder continuity of f —in the case that f_{in} happens to be Hölder continuous—without appealing to the De Giorgi estimate of [18].

1.5. Notation

To aid the reader, whenever possible we denote supersolutions with overlines and subsolutions with underlines, e.g. \overline{f} and \underline{f} .

We often use z to refer to a point $(t, x, v) \in [0, T] \times \mathbb{R}^3 \times \mathbb{R}^3$, and if z is decorated by a symbol the coordinates are as well, e.g. $\tilde{z} = (\tilde{t}, \tilde{x}, \tilde{v})$.

For z, z' , define the kinetic distance

$$\rho(z, z') = |t' - t|^{1/2} + |x' - x - (t' - t)v|^{1/3} + |v' - v|. \quad (1.8)$$

It is not a metric since the triangle inequality is not satisfied and it is not necessarily symmetric; however, it is straightforward to check that $\rho(z_1, z_2) \lesssim \rho(z_2, z_1)$ for any z_1 and z_2 .

The kinetic distance ρ gives rise to kinetic Hölder norms, which give rise to the kinetic Hölder spaces in the obvious way. For $Q \subset [0, T] \times \mathbb{R}^6$, we define the Hölder seminorm

$$[u]_{C_{\text{kin}}^\alpha(Q)} = \sup_{z \neq z' \in Q} \frac{|u(z) - u(z')|}{\rho(z, z')^\alpha}, \quad (1.9)$$

and the norm $\|u\|_{C_{\text{kin}}^\alpha(Q)} := [u]_{C_{\text{kin}}^\alpha(Q)} + \sup_Q |u|$. In addition, we define the second order norm

$$\|u\|_{C_{\text{kin}}^{2,\alpha}(Q)} := \sup_Q |u| + \sup_Q |\nabla u| + \|D_v^2 u\|_{C_{\text{kin}}^\alpha(Q)} + \|(\partial_t + v \cdot \nabla_x)u\|_{C_{\text{kin}}^\alpha(Q)} < \infty.$$

The differential operator $\partial_t + v \cdot \nabla_x$ has been extended to the space $C_{\text{kin}}^{2,\alpha}(Q)$ by density, and even though $(\partial_t + v \cdot \nabla_x)u$ is continuous, $\partial_t u$ and $v \cdot \nabla_x u$ need not exist in a classical sense.

We denote by $Q_r(z_0)$ a ball under ρ , i.e. a kinetic cylinder:

$$Q_r(z_0) = \{z \in (-\infty, t_0] \times \mathbb{R}^6 : \rho(z_0, z) < r\}.$$

Notice that this includes only times *before* t_0 . If the center point z_0 is omitted in the notation, it is assumed that $z_0 = 0$. We also use $B_r(p)$ to denote a ball in the standard metric.

We use the notation $A \lesssim B$ if there is a constant C such that $A \leq CB$. In each section we clarify the dependencies of C , but in general, C may depend on γ, δ, r, R , and k . We use $A \approx B$ if $A \lesssim B$ and $B \lesssim A$.

1.6. Outline

The rest of the paper is organized as follows. In Section 2, we establish the existence of solutions (Theorem 1.2) and prove our continuation criterion (Theorem 1.3). In Section 3, we prove that f is continuous up to $t = 0$, as long as f_{in} is continuous. In Section 4, we show propagation of Hölder regularity and a time-integrable bound for $D_v^2 f$, and in Section 5, we prove uniqueness of solutions (Theorem 1.4). In Appendix A, we prove that Hölder continuity in x and v implies Hölder continuity in t , and in Appendix B, we prove some interpolation lemmas.

2. Existence

The purpose of this section is to prove Theorem 1.2 by constructing a classical solution to (1.1). As discussed in Section 1.4, the key ideas are: (i) an *a priori* estimate in the space $L^{\infty,k}([0, T] \times \mathbb{R}^6)$ (Lemma 2.3), and (ii) an argument that extends conditional regularity and continuation results to solutions in this weak space. Some steps of this construction—which we will indicate—involve adapting results and proof techniques from [7], [23], and [24], which are previous works on the Landau equation involving the present authors. We avoid giving full details of these steps, in order expedite the presentation and to emphasize the more novel aspects of the argument. However, for the reader's convenience, we outline the omitted proofs when possible.

2.1. Coefficient bounds

Our first step is to obtain bounds on the coefficients that are compatible with the $L^{\infty,k}$ norm.

Lemma 2.1. *If $k > \gamma + 5$, then, with all norms over \mathbb{R}^3 , we have*

$$\begin{aligned}\bar{a}^f(t, x, v) &\lesssim \langle v \rangle^{(\gamma+2)+} \|f(t, x, \cdot)\|_{L_v^{\infty,k}}, \\ |\bar{b}^f(t, x, v)| &\lesssim \langle v \rangle^{(\gamma+1)+} \|f(t, x, \cdot)\|_{L_v^{\infty,k}}, \\ \bar{c}^f(t, x, v) &\lesssim \begin{cases} \|f(t, x, \cdot)\|_{L_v^{\infty}}, & \gamma = -3, \\ \|f(t, \cdot, \cdot)\|_{L_v^{\infty,k}}, & \gamma \in (-3, 0), \end{cases}\end{aligned}$$

whenever the right-hand sides are finite, with implied constants depending on γ and k .

Proof. It is elementary to show that for $r > -3$ and $g : \mathbb{R}^3 \rightarrow \mathbb{R}_+$, there holds for $v \in \mathbb{R}^3$,

$$(g * |\cdot|^r)(v) \lesssim \langle v \rangle^{r+} \|g\|_{L^{\infty,k}},$$

where $k > r + 3$ and $r_+ = \max\{r, 0\}$. The statement of the lemma follows from this convolution estimate and the formulas for \bar{a}^f , \bar{b}^f , and \bar{c}^f in (1.2). \square

For the proof of the continuation criterion, we also require bounds on the coefficients in terms of L_v^1 -based norms:

Lemma 2.2. *For f such that the right-hand sides are finite, we have, with all norms over \mathbb{R}^3 ,*

$$\begin{aligned}|\bar{a}^f(t, x, v)| &\lesssim C \begin{cases} \langle v \rangle^{\gamma+2} \|f(t, x, \cdot)\|_{L_v^{1,2}}, & \gamma \in (-2, 0), \\ \|f(t, x, \cdot)\|_{L_v^{3/(5+\gamma)+\eta}} + \|f(t, x, \cdot)\|_{L_v^1}, & \gamma \in [-3, -2], \end{cases} \\ |\bar{b}^f(t, x, v)| &\lesssim \begin{cases} \langle v \rangle^{\gamma+1} \|f(t, x, \cdot)\|_{L_v^{1,1}}, & \gamma \in (-1, 0), \\ \|f(t, x, \cdot)\|_{L_v^{3/(4+\gamma)+\eta}} + \|f(t, x, \cdot)\|_{L_v^1}, & \gamma \in [-3, -1], \end{cases} \\ |\bar{c}^f(t, x, v)| &\lesssim \begin{cases} \|f(t, x, \cdot)\|_{L_v^{3/(3+\gamma)+\eta}} + \|f(t, x, \cdot)\|_{L_v^1}, & \gamma \in (-3, 0), \\ \|f(t, x, \cdot)\|_{L_v^{\infty}}, & \gamma = -3, \end{cases}\end{aligned}$$

with $\eta > 0$ any small constant. If f has even more decay, then the quadratic form associated to \bar{a}^f satisfies the following anisotropic upper bounds for $e \in \mathbb{S}^2$: If $\gamma \in [-3, -2]$, then

$$\bar{a}^f(t, x, v) e_i e_j \leq C \left(\|f(t, x, \cdot)\|_{L_v^{1,\ell}} + \|f(t, x, \cdot)\|_{L_v^p} \right) \begin{cases} \langle v \rangle^{\gamma+2}, & e \perp v, \\ \langle v \rangle^{\gamma}, & e \parallel v, \end{cases} \quad (2.1)$$

where $p > 3/(5 + \gamma)$ and $\ell > 3|\gamma|/(5 + \gamma)$. If $\gamma \in (-2, 0)$, then the same estimate holds without the L_v^p bound, and with 2 replacing ℓ .

Proof. The isotropic upper bounds follow from standard integral estimates and are omitted. For the proof of (2.1), see [7, Lemma 2.1] for the case $\gamma \in (-2, 0)$ and [23, Lemma A.3] for the case $\gamma \in [-3, -2]$. The statement of [23, Lemma A.3] uses the L_v^{∞} norm rather than the L_v^p norm, but the same proof works with L_v^p . \square

2.2. A priori estimates

A key element of our construction is the following closed estimate in the space $L^{\infty,k}(\mathbb{R}^6)$ for suitable k , which is proven using the maximum principle for the linear Landau equation.

Lemma 2.3. *Let $k_0 > \gamma + 5$. If f is a smooth solution to (1.1) on $[0, T] \times \mathbb{R}^6$, $\|f_{\text{in}}\|_{L^{\infty, k_0}(\mathbb{R}^6)} + \|f\|_{L^{\infty, k_0}([0, T] \times \mathbb{R}^6)} < \infty$, and $f \in C_t L^{\infty, k_0}_{\mathbb{R}^6}([0, T] \times \mathbb{R}^6)$, then there exists $T_0 > 0$ depending on $\|f_{\text{in}}\|_{L^{\infty, k_0}(\mathbb{R}^6)}$ and γ , and $C > 0$ depending only on γ , such that*

$$\|f(t, \cdot, \cdot)\|_{L^{\infty, k_0}(\mathbb{R}^6)} \leq C \|f_{\text{in}}\|_{L^{\infty, k_0}(\mathbb{R}^6)}, \quad 0 \leq t \leq \min(T, T_0).$$

Furthermore, if $\|f_{\text{in}}\|_{L^{\infty, k}(\mathbb{R}^6)} < \infty$ for any $k \geq k_0$, and either $\|f\|_{L^{\infty, k_0}([0, T] \times \mathbb{R}^6)}$ or $\Psi(t) + \|f\|_{L^{\infty, x}_v L^p_v([0, T] \times \mathbb{R}^6)}$ is finite (recall the definition of Ψ in (1.4)), with $p > 3/(3 + \gamma)$, then the inequality

$$\|f(t, \cdot, \cdot)\|_{L^{\infty, k}(\mathbb{R}^6)} \leq \|f_{\text{in}}\|_{L^{\infty, k}(\mathbb{R}^6)} e^{CKt}, \quad 0 \leq t \leq T,$$

holds for either $K = \|f\|_{L^{\infty, k_0}([0, T] \times \mathbb{R}^6)}$ or $K = \Psi(t) + \|f\|_{L^{\infty, x}_v L^p_v([0, T] \times \mathbb{R}^6)}$, where C depends on γ and k .

Although this estimate depends on $\|f\|_{L^{\infty, x}_v L^p_v([0, T] \times \mathbb{R}^6)}$ for all $\gamma \in [-3, 0)$, this dependence is removed in the case $\gamma \in (-2, 0)$ during the proof of the continuation criterion (Theorem 1.3).

Proof. Define the linear operator L by

$$Lg = \partial_t g + v \cdot \nabla_x g - \text{tr}(\bar{a}^f D_v^2 g) - \bar{c}^f g.$$

With $\beta > 0$ and $k > 0$ to be chosen, define $\bar{\phi}(t, x, v) = e^{\beta t} \langle v \rangle^{-k}$. Then

$$|\partial_{ij} \bar{\phi}| = e^{\beta t} \left| k(k+2) \langle v \rangle^{-k-4} v_i v_j - k \langle v \rangle^{-k-2} \delta_{ij} \right| \lesssim \langle v \rangle^{-2} \bar{\phi},$$

and

$$\begin{aligned} L\bar{\phi} &= \beta \bar{\phi} - \text{tr}(\bar{a}^f D_v^2 \bar{\phi}) - \bar{c}^f \bar{\phi} \\ &\geq \beta \bar{\phi} - CK \langle v \rangle^{(\gamma+2)+} \langle v \rangle^{-2} \bar{\phi} - CK \bar{\phi} \geq (\beta - C_0 K) \bar{\phi}, \end{aligned}$$

where K is any quantity such that $|\bar{a}^f| \leq K \langle v \rangle^{(\gamma+2)+}$ and $|\bar{c}^f| \leq K$, and C_0 depends on γ and k . With $\beta = C_0 K$, we have $L\bar{\phi} \geq 0$. If k is such that $\|f_{\text{in}}\|_{L^{\infty, k}(\mathbb{R}^6)} < \infty$, we can apply the maximum principle to $\|f_{\text{in}}\|_{L^{\infty, k}} \bar{\phi} - f$ and conclude

$$\|f(t, \cdot, \cdot)\|_{L^{\infty, k}(\mathbb{R}^6)} \leq \|f_{\text{in}}\|_{L^{\infty, k}(\mathbb{R}^6)} \exp(C_0 K t), \quad t \in [0, T]. \quad (2.2)$$

Now we set $k = k_0$, and for $t \in [0, T]$, define $H(t) = \|f\|_{L^{\infty, k_0}([0, t] \times \mathbb{R}^6)}$. Since H is increasing, (2.2) applied on $[0, t]$ with $K = H(t)$ implies $H(t) \leq \|f_{\text{in}}\|_{L^{\infty, k_0}} \exp(C_0 t H(t))$ for $t \in [0, T]$. From this inequality and Lemma 2.4 below, we have $H(t) \leq C \|f_{\text{in}}\|_{L^{\infty, k_0}}$ if $C_0 \|f_{\text{in}}\|_{L^{\infty, k_0}} t \leq 1/e$, and we can choose $T_0 = (e C_0 \|f_{\text{in}}\|_{L^{\infty, k_0}})^{-1}$.

For the second conclusion of the lemma, we apply (2.2) for any $k \geq k_0$. We may choose $K = \|f\|_{L^{\infty, k_0}([0, T] \times \mathbb{R}^6)}$ by Lemma 2.1, or $K = \Psi(t) + \|f\|_{L^{\infty, x}_v L^p_v([0, T] \times \mathbb{R}^6)}$ by Lemma 2.2. \square

Lemma 2.4. *If $H : [0, T] \rightarrow \mathbb{R}_+$ is a continuous increasing function and $H(t) \leq A e^{BtH(t)}$ for all $t \in [0, T]$ and some positive constants A and B , then*

$$H(t) \leq eA \quad \text{for} \quad 0 \leq t \leq T_* := \min\left(T, \frac{1}{eAB}\right).$$

Proof. First, we may assume that $T_* = 1/(eAB)$ by simply extending $H(t)$ to be constant after $t = T$. For each $t \in (0, 1/(eAB))$, let $\phi_t(x) = A e^{Btx} - x$ and let $x_{\min}(t) = -(Bt)^{-1} \log(ABt)$. A simple computation implies that $\phi_t(x_{\min}(t)) < 0$. We claim that $H(1/(eAB)) \leq x_{\min}(1/(eAB))$, which establishes the claim because H is increasing. We argue by contradiction supposing $H(1/(eAB)) > x_{\min}(1/(eAB))$. Since $H(0) < x_{\min}(0)$, the intermediate value theorem implies that $H(t_0) = x_{\min}(t_0)$ for some $t_0 < 1/(eAB)$, which implies that $\phi(H(t_0)) = \phi(x_{\min}(t_0)) < 0$. This is a contradiction since $\phi_t(H(t)) \geq 0$ for all $t \in [0, T_*]$ by hypothesis. This finishes the proof. \square

We require the following result guaranteeing a lower ellipticity bound for \bar{a}^f that is controlled from below whenever \bar{a}^f is bounded above. Lemma 2.5 follows from the work in [24], although it is not explicitly stated as a lemma in that paper. We require the following definition: for $k \geq 0$, the uniformly local Sobolev norm H_{ul}^k is defined by

$$\|g\|_{H_{\text{ul}}^k(\mathbb{R}^6)}^2 = \sum_{|\alpha|+|\beta| \leq k} \sup_{a \in \mathbb{R}^3} \int_{\mathbb{R}^6} |\phi(x-a) \partial_x^\alpha \partial_v^\beta g(x,v)|^2 dx dv,$$

where $\phi \in C_0^\infty(\mathbb{R}^3)$ is a cutoff satisfying $0 \leq \phi \leq 1$, $\phi \equiv 1$ in B_1 , and $\phi \equiv 0$ in $\mathbb{R}^3 \setminus B_2$. A bound on f in the H_{ul}^4 norm is needed to apply the results of [24] directly, but none of the conclusions depend quantitatively on this norm.

Lemma 2.5. *Let $f \geq 0$ be a solution of (1.1) in $[0, T] \times \mathbb{R}^6$ with initial data f_{in} well-distributed with parameters $\delta, r, R > 0$, and such that $e^{\rho|v|^2} f_{\text{in}} \in H_{\text{ul}}^4(\mathbb{R}^6)$ for some $\rho > 0$. If K is a constant such that $|\bar{a}^f(t, x, v)| \leq K \langle v \rangle^{(\gamma+2)_+}$ in $[0, T] \times \mathbb{R}^6$, then f satisfies the pointwise lower bound*

$$f(t, x, v) \geq c_1(t) e^{-|v|^{2-\gamma}/c_1(t)}, \quad (t, x, v) \in E,$$

where $c_1(t) > 0$ depends only on γ, δ, r, R , and K . Furthermore, the matrix \bar{a}^f satisfies

$$\bar{a}_{ij}^f(t, x, v) e_i e_j \geq c_2(t) \begin{cases} \langle v \rangle^\gamma, & e \in \mathbb{S}^2, \\ \langle v \rangle^{\gamma+2}, & e \cdot v = 0, \end{cases} \quad (2.3)$$

for $0 \leq t \leq T$, with $c_2(t)$ depending on γ, δ, r, R , and K . The functions $c_1(t)$ and $c_2(t)$ may degenerate as $t \rightarrow 0$ but are otherwise uniformly positive.

If, in addition, f_{in} is such that for all x , there is some $|v_m| < R$ such that $f_{\text{in}}(\cdot, \cdot) \geq \delta \mathbb{1}_{B_r(x, v_m)}$, then (2.3) holds with $c_2(t)$ replaced by a uniform positive constant $c_T > 0$ depending on γ, δ, r, R, K , and T , that are positive for any $T < \infty$.

Proof. The first statement follows directly from [24, Theorem 1.3]. As stated, that theorem requires a bound on the hydrodynamic quantities (precisely $\tilde{\Psi}(t)$ defined in (1.6)), but the only role these quantities play in the proof is providing an upper bound of the form $|\bar{a}^f(t, x, v)| \leq K \langle v \rangle^{(\gamma+2)_+}$, so any such K suffices (see [24, Section 4]).

The lower ellipticity bound (2.3) for \bar{a}^f follows from the lower bound for f and [24, Lemma 4.3].

The last statement follows from an examination of the proof of [24, Proposition 4.1]. For any $x \in \mathbb{R}^3$, let v_m be such that $f_{\text{in}}(x, \cdot) \geq \delta \mathbb{1}_{B_r(v_m)}$, where $|v_m| < R$. Step 1 of the proof of [24, Proposition 4.1] establishes the existence of some $T_* \in (0, T]$ depending on γ, K , and r , such that for all $t \in [0, T_*]$,

$$f(t, \cdot, \cdot) \geq \frac{\delta}{2} \mathbb{1}_{B_{r/2}(x) \times B_{r/2}(v_m)}.$$

The proper v -dependence for $t \in [0, T_*]$ is then implied by the proof of [24, Theorem 1.3(ii)]. We can take c_1 to be the minimum of $\delta/2$ and $\inf_{t \in [T_*, T]} c_1(t)$, and similarly for $c(t)$ in (2.3). \square

Remark. It is seen from the proof of Theorem 1.2.(i) that the quantitative lower bounds of Lemma 2.5 also apply to the solution constructed in Theorem 1.2.

2.3. A convenient transformation and new coefficients

Since the ellipticity ratio of \bar{a}^f degenerates as $|v| \rightarrow \infty$, it is convenient to use a change of variables developed in [7, Lemma 4.1] that makes \bar{a}^f uniformly elliptic (see also [23, Lemma 3.1] for the extension to the case $\gamma < -2$). This makes it possible to apply the local regularity estimates from [18, 23] and understand precisely how the constants degenerate for large $|v|$. We define this transformation here.

Fix $z_0 \in \mathbb{R}_+ \times \mathbb{R}^6$, and let S be the linear transformation such that

$$Se = \begin{cases} \langle v_0 \rangle^{1+\gamma/2} e, & e \cdot v_0 = 0 \\ \langle v_0 \rangle^{\gamma/2} e, & e \cdot v_0 = |v_0|, \end{cases}$$

and

$$r_1 = \langle v_0 \rangle^{-(1+\gamma/2)+} \min \left(1, \sqrt{t_0/2} \right). \quad (2.4)$$

Next, define

$$\mathcal{S}_{z_0}(t, x, v) = (t_0 + t, x_0 + Sx + tv_0, v_0 + Sv) \quad \text{and} \quad \delta_{r_1} z = (r_1^2 t, r_1^3 x, r_1 v). \quad (2.5)$$

There are two important, elementary features of these transformations that we require throughout the proof:

$$\begin{aligned} \rho(\delta_{r_1} z, \delta_{r_1} z') &= r_1 \rho(z, z') \quad \text{and} \\ \min \left(1, \sqrt{t_0/2} \right) \langle v_0 \rangle^{-(1+\gamma/2)+\gamma/2} \rho(z, z') &\leq \rho(\mathcal{S}_{z_0}(\delta_{r_1} z), \mathcal{S}_{z_0}(\delta_{r_1} z')) \\ &\leq \min \left(1, \sqrt{t_0/2} \right) \rho(z, z'). \end{aligned} \quad (2.6)$$

Given any function g and any point $z = (t, x, v) \in Q_1$, we define

$$g_{z_0}(z) = g(\mathcal{S}_{z_0}(\delta_{r_1} z)). \quad (2.7)$$

First we prove an L^∞ -based bound on the coefficients. This is necessary to obtain the Hölder regularity of f .

Lemma 2.6. Fix $z_0 = (t_0, x_0, v_0) \in \mathbb{R}_+ \times \mathbb{R}^3 \times \mathbb{R}^3$ and $\ell \in \mathbb{R}$. Let \bar{a} be a matrix and g be a scalar- or vector-valued function, and assume that, for all $z \in [t_0/2, t_0] \times \mathbb{R}^3 \times \mathbb{R}^3$ and $e \in \mathbb{S}^2$,

$$|g(z)| \lesssim \langle v \rangle^\ell \quad \text{and} \quad \bar{a}_{ij}(z) e_i e_j \approx \begin{cases} \langle v \rangle^\gamma & \text{if } |e \cdot v| = |v|, \\ \langle v \rangle^{2+\gamma} & \text{if } e \cdot v = 0. \end{cases} \quad (2.8)$$

Let $\bar{A} = S^{-1} \bar{a}_{z_0} S^{-1}$. Then there exist $0 < \lambda < \Lambda$, depending only on the implied constants in (2.8), such that, in Q_1 ,

$$\lambda I \leq \bar{A}(z) \leq \Lambda I \quad \text{and} \quad |g_{z_0}(z)| \leq \Lambda \langle v_0 \rangle^{-\ell} \|g\|_{L^{\infty, \ell}([t_0/2, t_0] \times \mathbb{R}^6)}.$$

In particular, λ and Λ are independent of z_0 .

Proof. The proof of the bounds of \bar{A} is exactly as in [23, Lemma 3.1]. The bound on g_{z_0} follows directly from the fact that if $z \in Q_1$, then $\langle r_1 S v + v_0 \rangle \approx \langle v_0 \rangle$. \square

In order to obtain bounds on the $C_{\text{kin}}^{2, \alpha}$ norm of f , we require Hölder regularity of the coefficients of (1.1) after applying our transformation. The following lemma is similar to [23, Lemma 3.3].

Lemma 2.7. Fix $z_0 \in \mathbb{R}_+ \times \mathbb{R}^6$, $\alpha \in (0, 1)$, and $m > \max\{5, 5 + \gamma + \alpha/3\}$. Suppose that $\langle v \rangle^m f \in C_{\text{kin}}^\alpha([t_0/2, t_0] \times \mathbb{R}^6)$. Let $\bar{A}(z) = S^{-1} \bar{a}_{z_0}^f S^{-1}$ and $\bar{C} = r_1^2 \bar{c}_{z_0}^f$. Then we have

$$\begin{aligned} [\bar{A}]_{C_{\text{kin}}^{2\alpha/3}(Q_1)} &\lesssim \langle v_0 \rangle^{2+\alpha/3} [\langle v \rangle^m f]_{C_{\text{kin}}^\alpha([t_0/2, t_0] \times \mathbb{R}^6)}, \\ [\bar{C}]_{C_{\text{kin}}^{2\alpha/3}(Q_1)} &\lesssim \langle v_0 \rangle^{-(2+\gamma)+\gamma+\alpha/3} [\langle v \rangle^m f]_{C_{\text{kin}}^\alpha([t_0/2, t_0] \times \mathbb{R}^6)}, \\ [f_{z_0}]_{C_{\text{kin}}^{2\alpha/3}(Q_1)} &\lesssim \min\{1, t_0^{\alpha/3}\} \langle v_0 \rangle^{2+\gamma+\alpha/3} [\langle v \rangle^m f]_{C_{\text{kin}}^\alpha(Q_1(z_0) \cap ([t_0/2, t_0] \times \mathbb{R}^6))}. \end{aligned}$$

Proof. We prove only the first inequality; that is, the inequality for \bar{A} . The bound for \bar{C} is exactly analogous, and the bound for f_{z_0} is straightforward after using (2.6).

The following calculation is similar to [23, Lemma 3.3]. For $z, z' \in Q_1$, let $\tilde{z} = \mathcal{S}_{z_0}(\delta_{r_1} z)$ and $\tilde{z}' = \mathcal{S}_{z_0}(\delta_{r_1} z')$. For $\bar{A}(z) = S^{-1} \bar{a}^f(\mathcal{S}_{z_0}(\delta_{r_1}(z))) S^{-1}$, we have, using $|S^{-1} e| \lesssim \langle v_0 \rangle^{-\gamma/2} |e|$,

$$\begin{aligned} |\bar{A}(z) - \bar{A}(z')| &\lesssim \langle v_0 \rangle^{-\gamma} \int_{\mathbb{R}^3} |w|^{\gamma+2} |f(\tilde{t}, \tilde{x}, \tilde{v} - w) - f(\tilde{t}', \tilde{x}', \tilde{v}' - w)| dw \\ &\lesssim \langle v_0 \rangle^{-\gamma} \int_{\mathbb{R}^3} |w|^{\gamma+2} \left(\rho(\tilde{z}, \tilde{z}')^\alpha + \rho(\tilde{z}, \tilde{z}')^{2\alpha/3} |w|^{\alpha/3} \right) [f]_{C_{\text{kin}}^\alpha([t_0/2, t_0] \times \mathbb{R}^3 \times B_1(v_0 - w))} dw, \end{aligned} \quad (2.9)$$

where we have used that

$$\rho(\tilde{t}, \tilde{x}, \tilde{v} - w), (\tilde{t}', \tilde{x}', \tilde{v}' - w)) \leq \rho(\tilde{z}, \tilde{z}') + \rho(\tilde{z}, \tilde{z}')^{2/3} |w|^{1/3},$$

which can be seen by a direct computation.

The inequality $[gh]_{C_{\text{kin}}^\alpha} \leq \|g\|_{L^\infty} [h]_{C_{\text{kin}}^\alpha} + [g]_{C_{\text{kin}}^\alpha} \|h\|_{L^\infty}$ implies

$$[f]_{C_{\text{kin}}^\alpha([t_0/2, t_0] \times \mathbb{R}^3 \times B_1(v_0 - w))} \lesssim \langle v_0 - w \rangle^{-m} \|\langle v \rangle^m f\|_{C_{\text{kin}}^\alpha}.$$

(Here, and for the remainder of the proof, we write $C_{\text{kin}}^\alpha = C_{\text{kin}}^\alpha([t_0/2, t_0] \times \mathbb{R}^6)$.) Feeding this estimate into (2.9) and using (2.6) and that $m > \max\{3, 5 + \gamma + \alpha/3\}$, we have

$$[\bar{A}]_{C_{\text{kin}}^{2\alpha/3}(Q_1)} = \sup_{z \neq z' \in Q_1} \frac{|\bar{A}(z) - \bar{A}(z')|}{\rho(z, z')^{2\alpha/3}} \lesssim \|\langle v \rangle^m f\|_{C_{\text{kin}}^\alpha} \langle v_0 \rangle^{2+\alpha/3}.$$

A similar calculation, with γ replacing $\gamma + 2$, implies

$$[\bar{C}]_{C_{\text{kin}}^{2\alpha/3}(Q_1)} \lesssim \langle v_0 \rangle^{-(2+\gamma)+\gamma+\alpha/3} \|\langle v \rangle^m f\|_{C_{\text{kin}}^\alpha}. \quad \square$$

2.4. Existence of a solution: Theorem 1.2

We are now ready to prove the existence of solutions:

Proof of Theorem 1.2. (i) Fix any $\varepsilon > 0$ and define the following mollification and cut-off functions. Fix any $\psi \in C_c^\infty(\mathbb{R}^6)$ such that $\psi \geq 0$ and $\int_{\mathbb{R}^6} \psi \, dx \, dv = 1$. Let $\psi_\varepsilon = \varepsilon^{-6} \psi(x/\varepsilon, v/\varepsilon)$. In addition, let $\zeta_\varepsilon \in C^\infty(\mathbb{R}^3)$ be such that $\zeta_\varepsilon(v) = 1$ when $|v| \leq 1/\varepsilon$, $\zeta_\varepsilon(v) = 0$ when $|v| \geq 1/\varepsilon + 1$, and $|\nabla_v \zeta_\varepsilon| \lesssim 1$.

Then let

$$f_{\text{in}}^\varepsilon = \zeta_\varepsilon(v)(f_{\text{in}} * \psi_\varepsilon)(x, v). \quad (2.10)$$

Note that $\|f_{\text{in}}^\varepsilon\|_{L^{\infty,k}} \lesssim \|f_{\text{in}}\|_{L^{\infty,k}}$. The smoothed, cut-off initial condition $f_{\text{in}}^\varepsilon$ is compactly supported in v , smooth, and nonnegative, so $e^{\rho|v|^2} f_{\text{in}}^\varepsilon$ is in $H_{\text{ul}}^4(\mathbb{R}^6)$ for any $\rho > 0$. Hence, we can apply [24, Theorem 1.1] to obtain a solution $f^\varepsilon : [0, T_\varepsilon] \times \mathbb{R}^6 \rightarrow \mathbb{R}_+$. We may assume T_ε is the maximal time of existence of the solution f^ε . By the existence theorem [24, Theorem 1.1], we have $e^{(\rho/2)|v|^2} f^\varepsilon \in L^\infty([0, T_\varepsilon], H_{\text{ul}}^4(\mathbb{R}^6)) \subset L^\infty([0, T_\varepsilon] \times \mathbb{R}^6)$, which clearly implies $f^\varepsilon \in L^{\infty,k}([0, T_\varepsilon] \times \mathbb{R}^6)$. Since $k > 5$, Lemma 2.3 yields

$$\|f^\varepsilon\|_{L^{\infty,k}([0, T_1], \mathbb{R}^6)} \leq C \|f_{\text{in}}^\varepsilon\|_{L^{\infty,k}} \leq C \|f_{\text{in}}\|_{L^{\infty,k}}, \quad (2.11)$$

where $T_1 = \min(T_0, T_\varepsilon)$, $T_0 \lesssim (\|f_{\text{in}}\|_{L^{\infty,k_0}})^{-1}$, and C is independent of ε .

For ε small enough (depending only on δ , R , and r), we have that $f_{\text{in}}^\varepsilon(x, v)$ is well-distributed with parameters $\delta/2$, $r/2$, and R . Therefore, [24, Theorem 1.5] implies f^ε can be extended for as long as

$$\begin{cases} \|f^\varepsilon(t, \cdot, \cdot)\|_{L_x^\infty L_v^{1,2}}, & \gamma \in (-2, 0), \\ \|f^\varepsilon(t, \cdot, \cdot)\|_{L_x^\infty L_v^{1,\ell}} + \|f^\varepsilon(t, \cdot, \cdot)\|_{L_{x,v}^\infty}, & \gamma \in [-3, -2], \end{cases}$$

remains finite, where $\ell > 3|\gamma|/(5 + \gamma)$. This quantity is controlled by $\|f^\varepsilon(t, \cdot, \cdot)\|_{L^{\infty,k}(\mathbb{R}^6)}$ since $k > \max\{5, 15/(5 + \gamma)\}$. Therefore, (2.11) implies the maximal time of existence T_ε must be larger than T_0 , i.e. f^ε exists for $t \in [0, T_0]$ for all ε .

The bound (2.11) holds for f^ε on $[0, T_0]$, so the upper bounds for \bar{a}^{f^ε} , \bar{b}^{f^ε} , and \bar{c}^{f^ε} of Lemma 2.1 and Lemma 2.2 hold independently of ε , since all the relevant norms of f are controlled by $\|f\|_{L^{\infty,k}}$. Since $f_{\text{in}}^\varepsilon$ is well-distributed, the smoothing theorem [24, Theorem 1.3] implies f^ε is a C^∞ classical solution of (1.1), with regularity estimates that may depend on ε . The decay estimate (2.11) and Lemma 2.5 with $K \lesssim \|f_{\text{in}}\|_{L^{\infty,k}}$ imply lower bounds for f^ε and \bar{a}^{f^ε} that depend only on t , x , δ , r , and $\|f_{\text{in}}\|_{L^{\infty,k}}$, but not on ε , for $t \in [0, T_0]$.

Next, we want to apply local regularity estimates at any point $z_0 \in \mathbb{R}_+ \times \mathbb{R}^6$. To track the dependence of these estimates on v , we must use the change of variables defined in (2.5). Recall the definition of f_{z_0} via (2.7). By Lemma 2.6

(which relies on our ε -independent bounds for \bar{a}^{f^ε} , \bar{b}^{f^ε} , and \bar{c}^{f^ε} , most crucially the anisotropic bounds (2.1) and (2.3)), we have that $f_{z_0}^\varepsilon$ satisfies

$$\partial_t f_{z_0}^\varepsilon + v \cdot \nabla_x f_{z_0}^\varepsilon = \nabla_v \cdot (\bar{A}(z) \nabla_v f_{z_0}^\varepsilon) + \bar{B}(z) \cdot \nabla_v f_{z_0}^\varepsilon + \bar{C}(z) f_{z_0}^\varepsilon, \quad (2.12)$$

which comes from (1.1), and

$$\partial_t f_{z_0}^\varepsilon + v \cdot \nabla_x f_{z_0}^\varepsilon = \operatorname{tr}(\bar{A}(z) D_v^2 f_{z_0}^\varepsilon) + \bar{C}(z) f_{z_0}^\varepsilon, \quad (2.13)$$

which comes from (1.3), in Q_1 , with the coefficients

$$\begin{aligned} \bar{A}(z) &= S^{-1} \bar{a}^{f^\varepsilon}(\mathcal{S}_{z_0}(\delta_{r_1}(z))) S^{-1}, & \bar{B}(z) &= r_1 S^{-1} \bar{b}^{f^\varepsilon}(\mathcal{S}_{z_0}(\delta_{r_1}(z))), \\ \text{and} & & \bar{C}(z) &= r_1^2 \bar{c}^{f^\varepsilon}(\mathcal{S}_{z_0}(\delta_{r_1}(z))) \end{aligned} \quad (2.14)$$

satisfying

$$\lambda_{t_0} I \leq \bar{A}(z) \leq \Lambda I, \quad |\bar{B}(z)| + |\bar{C}(z)| \leq \Lambda, \quad (2.15)$$

with Λ depending only on $\|f^\varepsilon\|_{L^{\infty,k}([0,T_0] \times \mathbb{R}^6)} \lesssim K$, and λ_{t_0, x_0} depending on K , t_0 , δ , r , and R . The dependence on t_0 comes from $c(t)$ in Lemma 2.5, which is uniformly positive on any compact subset of $(0, T]$.

The divergence-form equation (2.12) allows us to apply [18, Theorem 3] to $f_{z_0}^\varepsilon$:

$$\|f_{z_0}^\varepsilon\|_{C_{\text{kin}}^\alpha(Q_{1/2})} \leq C(\|f_{z_0}^\varepsilon\|_{L^2(Q_1)} + \|\bar{C} f_{z_0}^\varepsilon\|_{L^\infty(Q_1)}) \lesssim \langle v_0 \rangle^{-k}, \quad (2.16)$$

with implied constant depending on λ_{t_0} and K . The Hölder exponent $\alpha \in (0, 1)$ also depends on λ_{t_0} and Λ , and therefore on K . Undoing this change of variables and using (2.6), we find that

$$\begin{aligned} \|f^\varepsilon\|_{C_{\text{kin}}^\alpha(Q_{r_1/2}(z_0))} &\leq \min\{1, t_0\}^{-\alpha/2} \langle v_0 \rangle^{\alpha((1+\gamma/2)+-\gamma/2)} \|f_{z_0}^\varepsilon\|_{C_{\text{kin}}^\alpha(Q_{1/2})} \\ &\lesssim \min\{1, t_0\}^{-\alpha/2} \langle v_0 \rangle^{-k+\alpha((1+\gamma/2)+-\gamma/2)}, \end{aligned}$$

where r_1 is defined in (2.4). Applying the straightforward interpolation

$$\|g\|_{C_{\text{kin}}^\alpha(Q_1(z_1))} \leq r^{-\alpha} \|g\|_{L^\infty(Q_1(z_1))} + \sup_{z_2 \in Q_1(z_1)} \|g\|_{C_{\text{kin}}^\alpha(Q_r(z_2))}$$

for any g , z_1 , z_2 , and r , we deduce

$$\|f^\varepsilon\|_{C_{\text{kin}}^\alpha(Q_1(z_0) \cap ([0,T] \times \mathbb{R}^6))} \lesssim \min\{1, t_0\}^{-\alpha/2} \langle v_0 \rangle^{-k+\alpha((1+\gamma/2)+-\gamma/2)}. \quad (2.17)$$

Since it is not an important point in this proof, we absorb all dependence on t_0 into the implied constant for the remainder of this section.

Next, we pass this regularity to \bar{A} and \bar{C} via Lemma 2.7, which requires $[\langle v \rangle^m f^\varepsilon]_{C_{\text{kin}}^\alpha([t_0/2, t_0] \times \mathbb{R}^6)} \leq C_{t_0}$, where $m > \max\{5, 5 + \gamma + \alpha/3\}$. By assumption $k > 5$ so this holds with $m = k - \alpha((1 + \gamma/2)_+ - \gamma/2)$, up to decreasing α . Thus,

$$[\bar{A}]_{C_{\text{kin}}^{2\alpha/3}(Q_1)} \lesssim \langle v_0 \rangle^{2+\alpha/3}, \quad [\bar{C}]_{C_{\text{kin}}^{2\alpha/3}(Q_1)} \lesssim \langle v_0 \rangle^{-(2+\gamma)_++\gamma+\alpha/3},$$

with constants depending on t_0 and K . It is then straightforward to show that

$$\begin{aligned} \|\bar{A}\|_{C_{\text{kin}}^{2\alpha/3}(Q_1)} &\lesssim \langle v_0 \rangle^{2+\alpha/3}, \quad \text{and} \\ [\bar{C} f_{z_0}^\varepsilon]_{C_{\text{kin}}^{2\alpha/3}(Q_1)} &\lesssim \langle v_0 \rangle^{-k-(2+\gamma)_++\gamma+\alpha/3}. \end{aligned}$$

Now, using the non-divergence form equation (2.13), we can apply the Schauder-type estimate [23, Theorem 2.9] to $f_{z_0}^\varepsilon$:

$$\begin{aligned} [f_{z_0}^\varepsilon]_{C_{\text{kin}}^{2,2\alpha/3}(Q_{1/2})} &\lesssim [\bar{C} f_{z_0}^\varepsilon]_{C_{\text{kin}}^{2\alpha/3}(Q_1)} + \|\bar{A}\|_{C_{\text{kin}}^{2\alpha/3}(Q_1)}^{3+\frac{2\alpha}{3}+\frac{3}{\alpha}} \|f_{z_0}^\varepsilon\|_{L^\infty(Q_1)} \\ &\lesssim \langle v_0 \rangle^{-k-(2+\gamma)_++\gamma+\alpha/3} + \langle v_0 \rangle^p \langle v_0 \rangle^{-k}, \end{aligned} \quad (2.18)$$

where $p > 0$ depends on α , which in turn depends on K and t_0 . The implied constant in (2.18) depends on the same quantities.

Translating from $f_{z_0}^\varepsilon$ back to f^ε , we clearly see that f^ε is $C_{\text{kin}}^{2,2\alpha/3}$ away from $t = 0$ and that

$$\| \langle v \rangle^{k-p} f^\varepsilon \|_{C_{\text{kin}}^{2,2\alpha/3}([t_0/2, t_0] \times \mathbb{R}^6)} \lesssim 1 \quad (2.19)$$

for some p depending on K and t_0 . We, again, stress that the implied constant in (2.19) depends on t_0 and may degenerate faster than t_0^{-1} to any power since λ_{t_0} may be exponentially small in t_0^{-1} .

For any $z_0 \in (0, T] \times \mathbb{R}^6$ and $\alpha' \in (0, 2\alpha/3)$, since $C_{\text{kin}}^{2,\alpha'}(Q_{r_1}(z_0))$ is precompact in $C_{\text{kin}}^{2,2\alpha/3}(Q_{r_1}(z_0))$, a subsequence of f^ε converges to a limit f in $C_{\text{kin}}^{2,\alpha'}(Q_{r_1}(z_0))$. Since $z_0 \in (0, T] \times \mathbb{R}^6$ is arbitrary, we have that $f \in C_{\text{kin,loc}}^{2,2\alpha/3}((0, T] \times \mathbb{R}^6)$ and satisfies a bound such as (2.19). Since $f^\varepsilon \rightarrow f$ pointwise, the bound (2.11) extends to f . By (2.11) and the Dominated Convergence Theorem, we conclude $\bar{a}^{f^\varepsilon} \rightarrow \bar{a}^f$, $\bar{b}^{f^\varepsilon} \rightarrow \bar{b}^f$, and $\bar{c}^{f^\varepsilon} \rightarrow \bar{c}^f$ as $\varepsilon \rightarrow 0$. This establishes the existence part of the theorem.

(ii) The proof of (1.5) relies on the following lemma:

Lemma 2.8. Fix any compact set $K_x \subset \mathbb{R}^3$. With f^ε as above, we have

- (i) $f^\varepsilon \rightarrow f$ in $L^{2,p}([0, T] \times K_x \times \mathbb{R}^3)$ for any $p < 7/2$,
- (ii) $\bar{a}^{f^\varepsilon} \nabla_v f^\varepsilon \rightarrow \bar{a}^f \nabla_v f$ weakly in $L^{2, -(1+\gamma/2)}([0, T] \times K_x \times \mathbb{R}^3)$, and
- (iii) $f^\varepsilon \bar{b}^{f^\varepsilon}$ converges weakly to $f \bar{b}^f$ in $L^{2,p}([0, T] \times K_x \times \mathbb{R}^3)$ for any $p < \max\{5, 15/(5+\gamma)\} - 5/2 - \gamma$.

We postpone the proof of this lemma momentarily and proceed with the proof of Theorem 1.2.(ii). Recall that our test functions ϕ satisfy

$$\begin{aligned} \phi &\in C([0, T]; L^{2, -7/2+\eta}(K_x \times \mathbb{R}^3)), \quad (\partial_t + v \cdot \nabla_x) \phi \in L^{2, -7/2+\eta}([0, T] \times K_x \times \mathbb{R}^3) \\ \nabla_v \phi &\in L^{2, 1+\gamma/2}([0, T] \times K_x \times \mathbb{R}^3) \quad \text{and} \quad \text{supp}(\phi) \subset [0, T] \times K_x \times \mathbb{R}^3, \end{aligned}$$

for some small $\eta > 0$. Note that the collision term $Q(f, f)$ in (1.1) may be written $Q(f, f) = \nabla_v \cdot (\bar{a}^f \nabla_v f + \bar{b}^f f)$. Since f^ε is smooth and satisfies (1.1), we find

$$\int_{\mathbb{R}^6} f_{\text{in}}^\varepsilon \phi(0) \, dx \, dv = \int_{[0, T] \times \mathbb{R}^6} \left(f^\varepsilon (\partial_t + v \cdot \nabla_x) \phi - \nabla_v \phi \cdot (\bar{a}^{f^\varepsilon} \nabla_v f^\varepsilon) - f^\varepsilon \bar{b}^{f^\varepsilon} \cdot \nabla_v \phi \right) \, dx \, dv \, dt, \quad (2.20)$$

where $\phi(0) = \phi|_{t=0}$. It is straightforward to show from the definition (2.10) that $f_{\text{in}}^\varepsilon \rightarrow f_{\text{in}}$ in L_{loc}^2 as $\varepsilon \rightarrow 0$. On the other hand, by Lemma 2.8, the right hand side of (2.20) converges to

$$\int_{[0, T] \times \mathbb{R}^6} \left(f (\partial_t + v \cdot \nabla_x) \phi - \nabla_v \phi \cdot (\bar{a}^f \nabla_v f) - f \bar{b}^f \cdot \nabla_v \phi \right) \, dx \, dv \, dt$$

as $\varepsilon \rightarrow 0$. Thus, we recover (1.5) as claimed.

(iii) For higher regularity, we return to the sequence $f_{z_0}^\varepsilon$ and apply the argument of [23]: pass the regularity provided by (2.19) to the coefficients of the equation (if the pointwise decay in v of f^ε is sufficiently strong, i.e. if k is large enough compared to p), apply local Schauder estimates, and repeat, differentiating the equation to estimate higher derivatives in $C_{\text{kin,loc}}^\alpha$. As $\varepsilon \rightarrow 0$, this implies the same local regularity for f , with α' replacing α . The number of iterations allowed is limited by the decay of f^ε in v (which, by (2.11), is determined by the decay of f_{in}). If f_{in} decays faster than any polynomial, then the solution $f_{z_0}^\varepsilon$ is C^∞ , which implies f is C^∞ as in [23]. The details are omitted. \square

We now prove the lemma.

Proof of Lemma 2.8. (i) We first recall that $f^\varepsilon \rightarrow f$ locally uniformly in $(0, T] \times \mathbb{R}^6$. Thus, for any $\delta > 0$, we have

$$\begin{aligned} & \limsup_{\varepsilon \rightarrow 0} \int_0^T \int_{K_x \times \mathbb{R}^3} \langle v \rangle^{2p} |f^\varepsilon - f|^2 \, dx \, dv \, dt \\ & \leq \limsup_{\varepsilon \rightarrow 0} \int_0^\delta \int_{K_x \times \mathbb{R}^3} \langle v \rangle^{2p} |f^\varepsilon - f|^2 \, dx \, dv \, dt + \limsup_{\varepsilon \rightarrow 0} \int_\delta^T \int_{K_x \times B_{1/\delta}(0)} \langle v \rangle^{2p} |f^\varepsilon - f|^2 \, dx \, dv \, dt \\ & \quad + \limsup_{\varepsilon \rightarrow 0} \int_\delta^T \int_{K_x \times B_{1/\delta}(0)^c} \langle v \rangle^{2p} |f^\varepsilon - f|^2 \, dx \, dv \, dt \\ & \leq C\delta |K_x| (\limsup_{\varepsilon \rightarrow 0} \|f^\varepsilon\|_{L^{\infty,5}} + \|f\|_{L^{\infty,5}})^2 + 0 + C(T - \delta) |K_x| (\limsup_{\varepsilon \rightarrow 0} \|f^\varepsilon\|_{L^{\infty,5}} + \|f\|_{L^{\infty,5}})^2 \delta^{7-2p}. \end{aligned}$$

Recall that $2p < 7$. Taking $\delta \rightarrow 0$ establishes the result.

(ii) We begin by showing that $|\bar{a}^{f^\varepsilon} \nabla_v f^\varepsilon|$ is bounded in $L^{2, -(1+\gamma/2)}([0, T] \times K_x \times \mathbb{R}^3)$. This guarantees that $\bar{a}^{f^\varepsilon} \nabla_v f^\varepsilon$ has a weak subsequential limit. First, note that

$$\begin{aligned} \|\bar{a}^{f^\varepsilon} \nabla_v f^\varepsilon\|_{L^{2, -(1+\gamma/2)}}^2 &= \int_0^T \int_{K_x \times \mathbb{R}^3} \langle v \rangle^{-(2+\gamma)} |(\bar{a}^{f^\varepsilon})^{1/2} (\bar{a}^{f^\varepsilon})^{1/2} \nabla_v f^\varepsilon|^2 \, dv \, dx \, dt \\ &\lesssim \int_0^T \int_{K_x \times \mathbb{R}^3} \langle v \rangle^{-(2+\gamma)} \left(\langle v \rangle^{2+\gamma} \|f^\varepsilon\|_{L^{\infty,k}} \right) |(\bar{a}^{f^\varepsilon})^{1/2} \nabla_v f^\varepsilon|^2 \, dv \, dx \, dt \\ &= \|f^\varepsilon\|_{L^{\infty,k}} \int_0^T \int_{K_x \times \mathbb{R}^3} \nabla_v f^\varepsilon \cdot (\bar{a}^{f^\varepsilon} \nabla_v f^\varepsilon) \, dv \, dx \, dt, \end{aligned}$$

since \bar{a}^{f^ε} is symmetric. Therefore, it is enough to show that $(\bar{a}^{f^\varepsilon})^{1/2} \nabla_v f^\varepsilon$ is bounded in $L^2([0, T] \times K_x \times \mathbb{R}^3)$ uniformly in ε . To this end, fix any non-negative $\psi \in C_c^\infty(\mathbb{R}^3)$ that equals 1 on K_x . Using that f^ε satisfies (1.1) and that \bar{a}^{f^ε} is non-negative definite, we find

$$\begin{aligned} 0 &\leq \int_0^T \int_{K_x} \nabla_v f^\varepsilon \cdot (\bar{a}^{f^\varepsilon} \nabla_v f^\varepsilon) \, dx \, dv \, dt \leq \int_0^T \int_{\mathbb{R}^6} \psi \nabla_v f^\varepsilon \cdot (\bar{a}^{f^\varepsilon} \nabla_v f^\varepsilon) \, dx \, dv \, dt \\ &\leq \frac{1}{2} \int_{\mathbb{R}^6} \psi |f_{\text{in}}^\varepsilon|^2 \, dx \, dv + \frac{1}{2} \int_0^T \int_{\mathbb{R}^6} |f^\varepsilon|^2 (v \cdot \nabla_x \psi + \frac{1}{2} \bar{c}^f \psi) \, dx \, dv \, dt. \end{aligned} \tag{2.21}$$

It is clear that the right hand side is bounded uniformly in ε , due to the uniform $L^{\infty,k}(\mathbb{R}^6)$ bound that holds on $f_{\text{in}}^\varepsilon$ and f^ε .

We next show that any subsequence of $\bar{a}^{f^\varepsilon} \nabla_v f^\varepsilon$ has a subsequence that converges weakly to $\bar{a}^f \nabla_v f$. It is an elementary fact that this is equivalent to the weak convergence of $\bar{a}^{f^\varepsilon} \nabla_v f^\varepsilon$ to $\bar{a}^f \nabla_v f$. Fix any subsequence $\varepsilon_n \rightarrow 0$. From the uniform bound above, we find $g \in L^{2, -(1+\gamma/2)}([0, T] \times K_x \times \mathbb{R}^3)^3$ and a further subsequence $\varepsilon_{n_j} \rightarrow 0$ such that $\bar{a}^{f^{\varepsilon_{n_j}}} \nabla_v f^{\varepsilon_{n_j}}$ converges weakly to g in $L^2([0, T] \times K_x \times \mathbb{R}^3)^3$.

On the other hand, from the Schauder estimates of the proof of part (i), and the uniform bounds on $f^\varepsilon \in L^{\infty,5}$, we find that $f^{\varepsilon_{n_j}} \in C_{\text{kin}}^{2, 2\alpha/3}(K)$ and $\bar{a}^{f^{\varepsilon_{n_j}}} \in C_{\text{kin}}^{2\alpha/3}(K)$ with uniform-in- ε bounds for any compact $K \subset (0, T] \times \mathbb{R}^6$. Thus, up to passing to a further subsequence, $\nabla_v f^{\varepsilon_{n_j}} \rightarrow \nabla_v f$ and $\bar{a}^{f^{\varepsilon_{n_j}}} \rightarrow \bar{a}^f$ uniformly on K . It follows that

$g = (\bar{a}^f)^{1/2} \nabla_v f$ on K . Since this holds for any K , we find $g = \bar{a}^f \nabla_v f$ almost everywhere in $[0, T] \times K_x \times \mathbb{R}^3$, which concludes the proof.

(iii) We omit this proof as it is similar to and easier than the proof of (ii). \square

2.5. Continuation of solutions: Theorem 1.3

Now, we prove our continuation criterion for the solutions of Theorem 1.2:

Proof of Theorem 1.3. Suppose first that $\gamma \in [-3, -2]$. If, for some $T' > 0$, the quantity $\Psi(T') = \|f\|_{L_{t,x}^\infty L_v^p([0, T'] \times \mathbb{R}^6)} + \|f\|_{L_{t,x}^\infty L_v^1([0, T'] \times \mathbb{R}^6)}$ is finite, Lemma 2.3 yields

$$\|f(t, \cdot, \cdot)\|_{L^{\infty,k}(\mathbb{R}^6)} \leq \|f_{\text{in}}\|_{L^{\infty,k}(\mathbb{R}^6)} e^{CT'}, \quad 0 \leq t \leq T',$$

with C depending on $\Psi(T')$. Since f_{in} is well-distributed, Lemma 2.5 with $K = \Psi(T')$ implies $f(T', \cdot, \cdot)$ is well-distributed with parameters that can only degenerate if $T' \rightarrow \infty$. Therefore, $f(T', \cdot, \cdot)$ satisfies the hypotheses of Theorem 1.2, and we can continue the solution past $t = T'$.

If $\gamma \in (-2, 0)$, we can bound $\|f\|_{L_{t,x}^\infty L_v^p([T'/2, T'] \times \mathbb{R}^6)}$ by applying the argument of [7, Theorem 1.1]. This gives an upper bound for $\|f\|_{L^\infty([T'/2, T'] \times \mathbb{R}^6)}$ depending only on $\|f\|_{L_{t,x}^\infty L_v^{1,2}}$ and the lower ellipticity bound for \bar{a}^f on $[T'/2, T']$ given by Lemma 2.5 (which can be bounded in terms of $\Psi(T')$ and the initial data). This gives a bound for f in $L_x^\infty L_v^p$ for $t \in [T'/2, T']$, which, combined with the reasoning of the previous paragraph, lets us continue the solution past $t = T'$. \square

3. Pointwise matching of initial data

In this section we show that, under the additional assumption that f_{in} is continuous, we have $f_{\text{in}}(x, v) = \lim_{t \rightarrow 0+} f(t, x, v)$. The proof uses a simple barrier argument.

Proposition 3.1. *Let f be the solution to (1.1) constructed in Theorem 1.2. If $f_{\text{in}} = f(0, \cdot, \cdot) \in L^{\infty,k}(\mathbb{R}^6)$ is continuous, then $f(t, x, v) \rightarrow f_{\text{in}}(x, v)$ as $t \rightarrow 0+$, uniformly on compact sets of \mathbb{R}^6 . If f_{in} is uniformly continuous in x , then the convergence as $t \rightarrow 0+$ is uniform.*

Proof. We work with the approximating solutions f^ε from Theorem 1.2, obtaining a uniform bound in ε . Thus, we obtain the result in the limit $\varepsilon \rightarrow 0$. Importantly, these f^ε are smooth on $[0, T] \times \mathbb{R}^6$, so we may use the classical comparison principle. In an abuse of notation, we denote f^ε simply by f for the remainder of the proof.

We show that continuity of f_{in} at a fixed $(x_0, v_0) \in \mathbb{R}^6$ implies $f(t, x_0, v_0) \rightarrow f_{\text{in}}(x_0, v_0)$ as $t \rightarrow 0$. Fix $\eta > 0$. If $|v_0|$ is sufficiently large, then the finiteness of $\|f\|_{L^{\infty,k}([0, T] \times \mathbb{R}^6)}$ guarantees that $f(t, x, v_0) < \eta$ for all x and for all t sufficiently small. Hence, we need only consider $v_0 \in B_{R_\eta}(0)$ for some large $R_\eta > 0$ depending only on η and $\|f\|_{L^{\infty,k}([0, T] \times \mathbb{R}^6)}$.

Let $\delta > 0$ be such that

$$|f_{\text{in}}(x, v) - f_{\text{in}}(x_0, v_0)| < \eta, \quad \text{if } |x - x_0|^2 + |v - v_0|^2 \leq \delta^2.$$

With $M, \beta, \rho > 0$ to be determined, define

$$\bar{h}(t, x, v) = e^{\beta t} \left[M(|x - x_0 - vt|^2 + |v - v_0|^2) + \eta + f_{\text{in}}(x_0, v_0) + \rho t \right].$$

Recall the linear operator $Lg = \partial_t g + v \cdot \nabla_x g - \text{tr}(\bar{a}^f D_v^2 g) - \bar{c}^f g$ from the proof of Lemma 2.3. We have $D_v^2 \bar{h} = 2Me^{\beta t}(1 + t^2)\text{Id}$, and

$$L\bar{h} = \beta \bar{h} + \rho e^{\beta t} - 2Me^{\beta t}(1 + t^2)\text{tr}(\bar{a}^f) - \bar{c}^f \bar{h}.$$

Since $\bar{c}^f \lesssim \|f\|_{L^\infty([0, T], L^{\infty,k}(\mathbb{R}^6))}$ and $\bar{a}^f \lesssim \langle v \rangle^{(\gamma+2)+} \|f\|_{L^\infty([0, T], L^{\infty,k}(\mathbb{R}^6))}$ (by Lemma 2.1), one has $L\bar{h} \geq 0$ in $[0, 1] \times \mathbb{R}^6$, provided β and ρ are large enough. By our choice of δ , we have

$$f_{\text{in}}(x, v) \leq f_{\text{in}}(x_0, v_0) + \eta \leq \bar{h}(0, x, v), \quad \text{for } |x - x_0|^2 + |v - v_0|^2 \leq \delta^2.$$

Next, choose $M > 0$ large enough so that, for $t \leq \delta/(4(R_\eta + \delta))$ and $|x - x_0|^2 + |v - v_0|^2 = \delta^2$, there holds

$$\begin{aligned} \bar{h}(t, x, v) &\geq M(|x - x_0|^2 + t^2|v|^2 - 2t(x - x_0) \cdot v + |v - v_0|^2) \\ &\geq M(\delta^2 - 2t(x - x_0) \cdot v) \\ &\geq M\delta^2/2 \geq \|f\|_{L^\infty([0, T] \times \mathbb{R}^6)}. \end{aligned}$$

We may now apply the maximum principle to obtain $f(t, x, v) \leq \bar{h}(t, x, v)$ in $[0, \delta/(4(R_\eta + \delta))] \times B_\delta(x_0, v_0)$.

Let $\delta' = \sqrt{\eta/M}$ and

$$t_0 = \min \left\{ \frac{1}{\beta} \log(1 + \eta), \frac{\delta'}{(|v_0| + \delta')}, \frac{\eta}{\rho} \right\}.$$

If $(t, x, v) \in [0, t_0) \times B_{\delta'}(x_0, v_0)$, we have

$$|x - x_0 - tv|^2 + |v - v_0|^2 \leq 2|x - x_0|^2 + |v - v_0|^2 + 2t^2|v|^2 < 4(\delta')^2,$$

and, thus,

$$f(t, x, v) \leq \bar{h}(t, x, v) < e^{\beta t} \left[4M(\delta')^2 + \eta + f_{\text{in}}(x_0, v_0) + \rho t \right] < (1 + \eta) [4\eta + \eta + f_{\text{in}}(x_0, v_0) + \eta].$$

Since η was arbitrary, we conclude $\limsup_{t \rightarrow 0} f(t, x_0, v_0) \leq f_{\text{in}}(x_0, v_0)$.

For the reverse inequality, define

$$\underline{h}(t, x, v) = e^{-\beta t} \left[f_{\text{in}} - M(|x - x_0 - vt|^2 - |v - v_0|^2) - \eta - \rho t \right].$$

By a similar calculation, we have $L\underline{h} \leq 0$ for $t \leq 1$, and $f_{\text{in}}(t, x, v) \geq \underline{h}(t, x, v)$ on the parabolic boundary of $[0, \delta/(2(|v_0| + \delta))] \times B_\delta(x_0, v_0)$, if M, β, ρ are chosen large enough. The rest of the proof follows similarly to establish that $f_{\text{in}}(x_0, v_0) = \lim_{t \rightarrow 0+} f(t, x_0, v_0)$.

Clearly, δ (and therefore M, δ' , and t_0) can be chosen uniformly on any compact set of \mathbb{R}^6 . If δ is independent of x_0 , then t_0 can be chosen depending only on η and $\|f\|_{L^{\infty, k}([0, T] \times \mathbb{R}^6)}$ (recall that R_η depends only on η and $\|f\|_{L^{\infty, k}([0, T] \times \mathbb{R}^6)}$), which yields the uniform convergence. Thus, the proof is complete. \square

4. Propagation of Hölder regularity and higher regularity estimates

In this section, we prove that if $f_{\text{in}}(x, v)$ is Hölder continuous, with Hölder norm decaying appropriately for large velocities, then our solution f is Hölder continuous in (t, x, v) up to some short time T_H . This is a necessary ingredient of our proof of uniqueness (see Section 5).

In this section, we are not interested in the dependence of constants on $\|f\|_{L^{\infty, k}}$. Thus, we allow the implied constant in any “ \lesssim ” to depend on $\|f\|_{L^{\infty, k}([0, T] \times \mathbb{R}^6)}$.

Following the proof outline given in the introduction, the first step is to revisit the Schauder estimates from the proof of Theorem 1.2.(i) under stronger assumptions on f_{in} that ensure good lower bounds for f as $t \rightarrow 0$. The non-scale-invariance of this estimate reflects the dependence of the coefficients in (1.3) on f .

Lemma 4.1. *Let $f \in L^{\infty, k}([0, T] \times \mathbb{R}^6)$ be a solution to (1.1) with f_{in} satisfying the hypotheses of Theorem 1.2. Assume in addition that for all x , there is some $|v_m| < R$ with $f_{\text{in}}(\cdot, \cdot) \geq \delta \mathbb{1}_{B_r(x, v_m)}$. For any $t_0 \in (0, T]$, we have*

$$\|\langle v \rangle^{m-q(\gamma, \alpha, k, m)} D_v^2 f\|_{L^\infty([t_0/2, t_0] \times \mathbb{R}^6)} \lesssim t_0^{-1+\alpha^2/(6-\alpha)} (1 + \|\langle v \rangle^m f\|_{C_{\text{kin}}^\alpha([t_0/2, t_0] \times \mathbb{R}^6)})^{p(\alpha)},$$

for any $m \in (\max\{3, 5 + \gamma + \alpha/3\}, k]$ such that the right-hand side is finite, with

$$\begin{aligned} p(\alpha) &= 3 + 2\alpha/3 + 3/\alpha, \\ q(\gamma, \alpha, k, m) &= (2 + \gamma)_+ - \gamma \\ &\quad + \left(1 - \frac{\alpha^2}{6 - \alpha}\right) \max\{-(2 + \gamma)_+ + \gamma - (k - m)/3, (2 + \alpha/3)p(\alpha) - k + m\} \end{aligned}$$

and the implied constant depending only on $m, \gamma, \alpha, T, \delta, r, R, k$, and $\|f\|_{L^{\infty, k}([t_0/2, t_0] \times \mathbb{R}^6)}$.

Proof. Let $z_0 \in (0, T] \times \mathbb{R}^6$ be fixed, and let $f_{z_0}(z) = f(\mathcal{S}_{z_0}(\delta_{r_1} z))$, with r_1 defined by (2.4) and \mathcal{S}_{z_0} defined by (2.5) as in the proof of Theorem 1.2. The function f_{z_0} satisfies (2.13) in Q_1 . By Lemma 2.5, Lemma 2.6, and our extra assumption on f_{in} , the diffusion matrix $\bar{A}(z)$, given by (2.14), satisfies upper and lower ellipticity estimates that are independent of z_0 . In other words, λ_{t_0} in (2.15) depends only on the initial data and T , not on t_0 .

Our goal is to apply the Schauder estimate [23, Theorem 2.12(a)]. Using the bounds in Lemma 2.7, along with the fact that $[f_{z_0}]_{C_{\text{kin}}^{2\alpha/3}(Q_1)} \leq [f_{z_0}]_{C_{\text{kin}}^\alpha(Q_1)}$, we find

$$\begin{aligned} [D_v^2 f_{z_0}]_{C_{\text{kin}}^{2\alpha/3}(Q_{1/2})} &\lesssim [\bar{C} f_{z_0}]_{C_{\text{kin}}^{2\alpha/3}(Q_1)} + \|\bar{A}\|_{C_{\text{kin}}^{2\alpha/3}(Q_1)}^{p(\alpha)} \|f_{z_0}\|_{L^\infty(Q_1)} \\ &\lesssim [\bar{C}]_{C_{\text{kin}}^{2\alpha/3}(Q_1)} \|f_{z_0}\|_{L^\infty(Q_1)} + \|\bar{C}\|_{L^\infty(Q_1)} \|f_{z_0}\|_{C_{\text{kin}}^{2\alpha/3}(Q_1)} \\ &\quad + \langle v_0 \rangle^{(2+\gamma)p(\alpha)} \|\langle v \rangle^m f\|_{C_{\text{kin}}^\alpha}^{p(\alpha)} \langle v_0 \rangle^{-k} \\ &\lesssim \left(\langle v_0 \rangle^{-(2+\gamma)_+ + \gamma + \alpha/3} \langle v_0 \rangle^{-k} + \langle v_0 \rangle^{-(2+\gamma)_+ + \gamma - k/3 - 2m/3} \right) \|\langle v \rangle^m f\|_{C_{\text{kin}}^\alpha} \\ &\quad + \langle v_0 \rangle^{(2+\gamma)p(\alpha) - k} \|\langle v \rangle^m f\|_{C_{\text{kin}}^\alpha}^{p(\alpha)} \\ &\lesssim \langle v_0 \rangle^{-m + \tilde{q}(\alpha, \gamma, k, m)} \left(1 + \|\langle v \rangle^m f\|_{C_{\text{kin}}^\alpha} \right)^{p(\alpha)}, \end{aligned} \quad (4.1)$$

where, as above, we use the shorthand $C_{\text{kin}}^\alpha = C_{\text{kin}}^\alpha([t_0/2, t_0] \times \mathbb{R}^6)$, and we have defined

$$\tilde{q}(\alpha, \gamma, k, m) = \max \{ -(2+\gamma)_+ + \gamma - (k-m)/3, (2+\alpha/3)p(\alpha) - (k-m) \}.$$

The reduction to the maximum of these two values is due to the fact that $-(2+\gamma)_+ + \gamma + \alpha/3 - (k-m) \leq (2+\alpha/3)p(\alpha) - (k-m)$. In the third line of (4.1), we also used an interpolation between C_{kin}^α and $L^{\infty, k}$ (see Lemma B.2) to write

$$[f_{z_0}]_{C_{\text{kin}}^{2\alpha/3}(Q_1)} \lesssim \|f_{z_0}\|_{L^\infty(Q_1)}^{1/3} [f_{z_0}]_{C_{\text{kin}}^\alpha(Q_1)}^{2/3} \lesssim \min\{1, t_0^{\alpha/3}\} \langle v_0 \rangle^{-k/3 - 2m/3} [\langle v \rangle^m f]_{C_{\text{kin}}^\alpha}^{2/3}, \quad (4.2)$$

and used $t_0 \lesssim 1$.

Using (4.2) again, and an interpolation between C_v^α and $C_v^{2, \alpha}$ (see Lemma B.1), we find

$$\begin{aligned} \|D_v^2 f_{z_0}\|_{L^\infty(Q_{1/4})} &\lesssim [f_{z_0}]_{C_{\text{kin}}^\alpha(Q_{1/2})} + [f_{z_0}]_{C_v^{2\alpha/3}(Q_{1/2})}^{\frac{2\alpha/3}{2+(2\alpha/3)-\alpha}} [D_v^2 f_{z_0}]_{C_v^\alpha(Q_{1/2})}^{1-\frac{2\alpha/3}{2+(2\alpha/3)-\alpha}} \\ &\lesssim \left(\min\{1, t_0^{\alpha/2}\} \langle v_0 \rangle^{-m} + \min\{1, t_0^{\frac{\alpha}{2} \frac{2\alpha}{6-\alpha}}\} \langle v_0 \rangle^{-m \frac{2\alpha}{6-\alpha}} \langle v_0 \rangle^{(1-\frac{2\alpha}{6-\alpha})(-m+\tilde{q}(\alpha, \gamma, k, m))} \right) \\ &\quad (1 + \|\langle v \rangle^m f\|_{C_{\text{kin}}^\alpha})^{p(\alpha)} \\ &\lesssim \min\{1, t_0^{\frac{\alpha^2}{6-\alpha}}\} \langle v_0 \rangle^{-m + (1-\frac{2\alpha}{6-\alpha})\tilde{q}(\alpha, \gamma, k, m)} (1 + \|\langle v \rangle^m f\|_{C_{\text{kin}}^\alpha})^{p(\alpha)}. \end{aligned}$$

Undoing the change of variables, we get that

$$|D_v^2 f(z_0)| \lesssim |r_1^{-2} S^{-2} D_v^2 f_{z_0}(0)| \lesssim \left(1 + t_0^{-1 + \frac{\alpha^2}{6-\alpha}} \right) \langle v_0 \rangle^{((2+\gamma)_+ - \gamma) - m + (1 - \frac{\alpha^2}{6-\alpha})\tilde{q}(\alpha, \gamma, k, m)} \|\langle v \rangle^m f\|_{C_{\text{kin}}^\alpha}.$$

Since $q(\alpha, \gamma, k, m) = ((2+\gamma)_+ - \gamma) + (1 - \frac{\alpha^2}{6-\alpha})\tilde{q}(\alpha, \gamma, k, m)$ and since z_0 was arbitrary, the claim is proved. \square

The purpose of the next lemma is passing from Hölder regularity in (x, v) to Hölder regularity in (t, x, v) . This lemma is proven in Appendix A, in a more general form. Since the reverse implication is immediate, we see that bounding our solution f in $L_t^\infty C_{\text{kin}, x, v}^\alpha$ is equivalent to bounding f in C_{kin}^α , up to velocity weights.

Lemma 4.2. For any locally Hölder continuous solution to (1.1) with $f \in L^{\infty, k}([0, T] \times \mathbb{R}^6)$ and k as in Theorem 1.2, there holds

$$\|f\|_{C_{\text{kin}}^\alpha(Q_1(z_0) \cap [0, T] \times \mathbb{R}^6)} \lesssim \langle v_0 \rangle^{\alpha(1+\gamma/2)+} \left(\|f\|_{L^\infty([0, T] \times \mathbb{R}^6)} + \|f\|_{L_t^\infty C_{\text{kin}, x, v}^\alpha(Q_2(z_0) \cap [0, T] \times \mathbb{R}^6)} \right),$$

for any $z_0 \in [0, T] \times \mathbb{R}^6$, where, for any $A \subset \mathbb{R}^6$,

$$[f]_{C_{\text{kin},x,v}^\alpha(A)} = \sup_{(x_1,v_1),(x_2,v_2) \in A} \frac{|f(x_1, v_1) - f(x_2, v_2)|}{(|x_1 - x_2|^{1/3} + |v_1 - v_2|)^\alpha}.$$

The implied constant depends only on $\|f\|_{L^{\infty,k}}$.

By combining Lemma 4.2 and Lemma 4.1, we deduce the following, whose proof is omitted.

Lemma 4.3. *Let the assumptions of Lemma 4.1 hold and additionally assume that $m > \max\{3, 5 + \gamma + \alpha/3\} + \alpha(1 + \gamma/2)_+$. Let*

$$q'(\alpha, \gamma, k, m) = q(\alpha, \gamma, k, m - \alpha(1 + \gamma/2)_+) + \alpha(1 + \gamma/2)_+.$$

Then

$$\|\langle v \rangle^{m-q'(\gamma,\alpha,k,m)} D_v^2 f\|_{L^\infty} \lesssim t_0^{-1+\alpha^2/(6-\alpha)} (1 + \|\langle v \rangle^m f\|_{L_t^\infty C_{\text{kin},x,v}^\alpha([t_0/2, t_0] \times \mathbb{R}^6)})^{p(\alpha)}$$

Now we are ready to show that Hölder continuity at $t = 0$ implies Hölder continuity for positive time. The proof requires us to work with the Euclidean Hölder norm $\|\cdot\|_{C^\alpha(\mathbb{R}^6)}$. This norm is always only in x, v variables.

Proposition 4.4. *Let f be the solution constructed in Theorem 1.2. Suppose that $\langle v \rangle^m f_{\text{in}} \in C^\alpha(\mathbb{R}^6)$ and $f_{\text{in}} \in L^{\infty,k}(\mathbb{R}^6)$ and that α, m , and k satisfy*

$$m > \max\{3, 5 + \gamma + \alpha/3\} + \alpha(1 + \gamma/2)_+ \quad \text{and} \quad q'(\alpha, \gamma, k, m) \leq (2 + \gamma)_+.$$

Then there exists $T_H \in (0, T]$ such that

$$\|\langle v \rangle^m f\|_{L_t^\infty C^\alpha([0, T_H] \times \mathbb{R}^6)} \lesssim \|\langle v \rangle^m f_{\text{in}}\|_{C^\alpha(\mathbb{R}^6)}.$$

The implied constant above and T_H depend only on $m, k, \alpha, \gamma, \|f\|_{L^{\infty,k}([0, T] \times \mathbb{R}^6)}, \delta, r$, and R .

Proof. Let f^ε be the regularizing approximation from the proof of Theorem 1.2. We show that the conclusion of the proposition holds for f^ε , with T_H independent of ε , so the same conclusion for f follows. The smoothness and decay of f^ε is used to obtain a first touching point with a supersolution, and to ensure the right-hand side of Lemma 4.1 is finite, but none of the estimates depend quantitatively on ε . To keep the notation clean, we denote $f = f^\varepsilon$ for this proof.

Step 1: Defining g and deriving its equation. As discussed in the introduction, for $(t, x, v, \chi, v) \in \mathbb{R}_+ \times \mathbb{R}^6 \times B_1(0)^2$ and $m > 0$, we define

$$\begin{aligned} \tau f(t, x, v, \chi, v) &:= f(t, x + \chi, v + v), \\ \delta f(t, x, v, \chi, v) &= \tau f(t, x, v, \chi, v) - f(t, x, v), \\ g(t, x, v, \chi, v) &= \frac{|\delta f(t, x, v, \chi, v)|^2}{(|\chi|^2 + |v|^2)^\alpha} \langle v \rangle^{2m}. \end{aligned} \tag{4.3}$$

Note that g encodes the Euclidean Hölder norm of f , as stated in the following elementary lemma.

Lemma 4.5. *Let $m \geq 0$. Fix any $f : \mathbb{R}^6 \rightarrow \mathbb{R}$ and let $g : \mathbb{R}^6 \times B_1(0)^2 \rightarrow \mathbb{R}$ be defined by $g(x, v, \chi, v) = |\delta f(x, v, \chi, v)|^2 \langle v \rangle^{2m} / (|\chi|^2 + |v|^2)^\alpha$. There holds*

$$\|g\|_{L^\infty(\mathbb{R}^6 \times B_1(0)^2)} + \|\langle v \rangle^m f\|_{L^\infty(\mathbb{R}^6)}^2 \approx \|\langle v \rangle^m f\|_{C^\alpha(\mathbb{R}^6)}^2 \approx \sup_{x,v} \langle v \rangle^{2m} \|f\|_{C^\alpha(B_1(x,v))}^2,$$

where the implied constants depend only on m and α .

Hence, deriving a time-dependent upper bound on g suffices to prove the proposition. We obtain a bound on g by showing that it satisfies an equation where all terms are either bounded nicely or respect a maximum principle. By a short computation, g satisfies

$$\begin{aligned} \partial_t g + v \cdot \nabla_x g + v \cdot \nabla_\chi g + \frac{2\alpha v \cdot \chi}{|\chi|^2 + |v|^2} g \\ = 2 \frac{(\operatorname{tr}(\bar{a}^{\delta f} D_v^2 \tau f + \bar{a}^f D_v^2 \delta f) + \bar{c}^{\delta f} \tau f + \bar{c}^f \delta f) \delta f}{(|\chi|^2 + |v|^2)^\alpha} \langle v \rangle^m. \end{aligned} \quad (4.4)$$

The first three terms on the left enjoy a maximum principle. The last term on the left is clearly bounded by g . The terms on the right can be shown to be bounded, which is the main thrust of the argument below.

Before continuing with the proof, we discuss why the Euclidean Hölder norm occurs naturally here. In the definition of g , one might expect to see a denominator of $(|\chi|^2 + |v|^6)^{\alpha/3}$ or another term with this balance of powers that respects the kinetic scaling that is natural to the equation. However, this would replace the last term on the left hand side of (4.4) by

$$\frac{2\alpha}{3} \frac{v \cdot \chi}{|\chi|^2 + |v|^6} g$$

which is not bounded. This forces the choice of $(|\chi|^2 + |v|^2)^\alpha$ for the denominator of g , which is why this proposition is stated in terms of the Euclidean Hölder norm.

To find a time-dependent upper bound of g that remains finite for some time interval, we use (4.4) and construct a super-solution of g .

Step 2: A super-solution and the maximum principle argument. With $N > 0$ to be chosen later, define \bar{G} to be the unique solution to

$$\begin{cases} \frac{d}{dt} \bar{G}(t) = N t^{-1 + \frac{\alpha^2}{6-\alpha}} (1 + \bar{G}(t))^{\frac{p(\alpha)+1}{2}}, \\ \bar{G}(0) = 1 + \|g(0, \cdot)\|_{L^\infty(\mathbb{R}^6 \times B_1(0)^2)} + N \|f\|_{L^{\infty,m}([0,T] \times \mathbb{R}^6)}. \end{cases} \quad (4.5)$$

This solution \bar{G} exists on a maximal time interval $[0, T_G)$ with T_G depending on $N, \beta, \alpha, \|g(0, \cdot)\|_{L^\infty(\mathbb{R}^6 \times B_1(0)^2)}$, and $\|f\|_{L^\infty([0,T] \times \mathbb{R}^6)}$. Our goal is to show that $g(t, x, v, \chi, v) < \bar{G}(t)$ for all $t \in [0, \min\{T, T_G\})$. By Lemma 4.5, this implies the existence of T_H as in the statement of the proposition. Let t_0 be the first time that $\|g\|_{L^\infty([0,t_0] \times \mathbb{R}^6 \times B_1(0)^2)} = \bar{G}(t_0)$. It is clear, by construction, that $t_0 > 0$. We seek a contradiction at $t = t_0$.

First, we claim that we may assume there exists $(x_0, v_0, \chi_0, v_0) \in \mathbb{R}^6 \times \bar{B}_1(0)^2$ such that $g(t_0, x_0, v_0, \chi_0, v_0) = \bar{G}(t_0)$. If there is no such point, then fix any sequence $z_n \in \mathbb{R}^6 \times B_1(0)^2$ such that $g(t_0, z_n) \rightarrow \bar{G}(t_0)$. Recall that f (which is actually the regularization f^ε) is C^∞ and satisfies pointwise Gaussian decay in v . Because of this Gaussian decay, we can take $z_n \in \mathbb{R}^3 \times B_R(0) \times B_1(0)^2$ for some $R > 0$. Since g does not decay as $|x| \rightarrow \infty$, we need to re-center as follows: let $g_n(t, x, v, \chi, v) = g(t, x + x_n, v, \chi, v)$. Up to passing to a subsequence, it is clear that there exists \bar{g} and $(v_0, \chi_0, v_0) \in \bar{B}_R(0) \times \bar{B}_1(0)^2$ such that $g_n \rightarrow \bar{g}$ locally uniformly and $\bar{g}(t_0, 0, v_0, \chi_0, v_0) = \bar{G}(t_0)$. Further, $\|\bar{g}\|_{L^\infty([0,t] \times \mathbb{R}^6 \times B_1(0)^2)} < \bar{G}(t)$ for all $t < t_0$. The smoothness of g implies that, again up to passing to a subsequence, $g_n \rightarrow \bar{g}$ in C_{loc}^ℓ for any $\ell \in \mathbb{N}$, so that \bar{g} satisfies the same equation as g , i.e. (4.4). The proof may then proceed using \bar{g} in the place of g . Hence, we may assume the existence of (x_0, v_0, χ_0, v_0) .

Next, we notice that $(\chi_0, v_0) \in B_1(0)^2$. Indeed, if χ_0 or v_0 were in $\partial B_1(0)$, then

$$\begin{aligned} g(t_0, x_0, v_0, \chi_0, v_0) &\leq \frac{|\delta f(t_0, x_0, v_0, \chi_0, v_0)|^2}{1} \langle v_0 \rangle^{2m} \\ &\lesssim f(t_0, x_0 + \chi_0, v_0 + v_0)^2 \langle v_0 \rangle^{2m} + f(t_0, x_0, v_0)^2 \langle v_0 \rangle^{2m} \lesssim \|f\|_{L^{\infty,m}}^2. \end{aligned}$$

Then, choosing N sufficiently large, we find $g(t_0, x_0, v_0, \chi_0, v_0) \leq N \|f\|_{L^{\infty,m}}^2$. On the other hand, by (4.5), \bar{G} is increasing. Hence, we have $\bar{G}(t_0) \geq \bar{G}(0) > N \|f\|_{L^{\infty,m}}^2$, which contradicts the fact that $\bar{G}(t_0) = g(t_0, x_0, v_0, \chi_0, v_0)$. Thus, we conclude $(\chi_0, v_0) \in B_1(0)^2$.

In order to conclude the proof, we establish that, at $(t_0, x_0, v_0, \chi_0, v_0)$,

$$\partial_t g < \frac{N}{t^{1-\frac{\alpha^2}{6-\alpha}}} (1+g)^{\frac{p(\alpha)+1}{2}} \quad (4.6)$$

as long as N is chosen sufficiently large, depending on α, m, k , and the constant in Lemma 4.1. Since, at this location, there holds $g = \bar{G}$ and $\partial_t g \geq (d/dt)\bar{G}$, this yields a contradiction in view of (4.5).

Step 3: Re-writing (4.4) at $(t_0, x_0, v_0, \chi_0, v_0)$. In order to prove that (4.6) holds, we return to (4.4) and examine the right hand side. Notice that

$$\begin{aligned} & 2 \frac{\text{tr}(\bar{a}^{\delta f} D_v^2 \tau f + \bar{a}^f D_v^2 \delta f)}{(|\chi|^2 + |v|^2)^\alpha} \delta f \langle v \rangle^{2m} \\ &= \text{tr}(\bar{a}^f D_v^2 g) + 2 \frac{\text{tr}(\bar{a}^{\delta f} D_v^2 (\tau f))}{(|\chi|^2 + |v|^2)^\alpha} \delta f \langle v \rangle^{2m} - 2 \nabla_v (\delta f) \cdot (\bar{a}^f \nabla_v (\delta f)) \frac{\langle v \rangle^{2m}}{(|\chi|^2 + |v|^2)^\alpha} \\ & \quad - \frac{2m \langle v \rangle^{2m-2}}{(|\chi|^2 + |v|^2)^\alpha} \left(4v \cdot (\bar{a}^f \nabla_v (\delta f)) \delta f + \text{tr}(\bar{a}^f) |\delta f|^2 + \frac{(2m-2)v \cdot (\bar{a}^f v)}{\langle v \rangle^2} |\delta f|^2 \right). \end{aligned} \quad (4.7)$$

On the other hand (x_0, v_0, χ_0, v_0) is the location of a global maximum of g . Hence, at (x_0, v_0, χ_0, v_0) , we obtain the identity

$$0 = \nabla_v g = \langle v_0 \rangle^{2m} \frac{2\delta f \nabla_v (\delta f)}{(|\chi_0|^2 + |v_0|^2)^\alpha} + 2m v_0 \langle v_0 \rangle^{2m-2} \frac{|\delta f|^2}{(|\chi_0|^2 + |v_0|^2)^\alpha},$$

or $\nabla_v (\delta f) = -m \langle v_0 \rangle^{-2} \delta f v_0$. Then (4.7) becomes, at (x_0, v_0, χ_0, v_0) ,

$$\begin{aligned} & 2 \frac{\text{tr}(\bar{a}^{\delta f} D_v^2 \tau f + \bar{a}^f D_v^2 \delta f)}{(|\chi_0|^2 + |v_0|^2)^\alpha} \delta f \langle v_0 \rangle^{2m} \\ &= \text{tr}(\bar{a}^f D_v^2 g) + 2 \frac{\text{tr}(\bar{a}^{\delta f} D_v^2 (\tau f))}{(|\chi_0|^2 + |v_0|^2)^\alpha} \delta f \langle v_0 \rangle^{2m} + \frac{2mg}{\langle v_0 \rangle^4} \left((m+2) v_0 \cdot (\bar{a}^f v_0) - \langle v_0 \rangle^2 \text{tr}(\bar{a}^f) \right). \end{aligned}$$

Using again that (x_0, v_0, χ_0, v_0) is the location of a maximum, we have $\bar{a}^f D_v^2 g \leq 0$ and $\nabla_x g = \nabla_\chi g = 0$. Therefore, from (4.4) we obtain, at (x_0, v_0, χ_0, v_0) ,

$$\begin{aligned} \partial_t g &\leq -2\alpha \frac{v_0 \cdot \chi_0}{|\chi_0|^2 + |v_0|^2} g + 2 \frac{\text{tr}(\bar{a}^{\delta f} D_v^2 (\tau f))}{(|\chi_0|^2 + |v_0|^2)^\alpha} \delta f \langle v_0 \rangle^{2m} + \frac{2mg}{\langle v_0 \rangle^4} \left((m+2) v_0 \cdot (\bar{a}^f v_0) - \langle v_0 \rangle^2 \text{tr}(\bar{a}^f) \right) \\ & \quad + 2 \frac{\bar{c}^{\delta f} \tau f \delta f}{(|\chi_0|^2 + |v_0|^2)^\alpha} \langle v_0 \rangle^{2m} + 2\bar{c}^f g \\ &=: J_1 + J_2 + J_3 + J_4 + J_5. \end{aligned} \quad (4.8)$$

It is clear that $J_1, J_3, J_5 \lesssim g$ (see Lemma 2.1 for J_3 and J_5). We now bound the remaining two terms.

Step 4: Bounding J_2 and J_4 . Re-writing J_4 and taking the absolute value, we find

$$|J_4| \leq \frac{2|\bar{c}^{\delta f}| |\tau f| \sqrt{g}}{(|\chi_0|^2 + |v_0|^2)^{\alpha/2}} \langle v_0 \rangle^m.$$

It is clear that we must bound the Hölder modulus of $\bar{c}^{\delta f}$. If $\gamma \in (-3, 0)$, then using Lemma 4.5, we have

$$\begin{aligned} \frac{\bar{c}^{\delta f}}{(|\chi_0|^2 + |v_0|^2)^{\alpha/2}} &= c_\gamma \int_{\mathbb{R}^3} |w|^\gamma \frac{\delta f(t_0, x_0, v_0 - w, \chi_0, v_0)}{(|\chi_0|^2 + |v_0|^2)^{\alpha/2}} dw \\ &\leq c_\gamma \int_{\mathbb{R}^3} \frac{|w|^\gamma}{\langle v_0 - w \rangle^m} \|g(t_0, \cdot)\|_{L^\infty(\mathbb{R}^6 \times B_1(0)^2)}^{\frac{1}{2}} dw \leq C \|g(t_0, \cdot)\|_{L^\infty(\mathbb{R}^6 \times B_1(0)^2)}^{\frac{1}{2}}, \end{aligned}$$

because $m > 3 + \gamma$. On the other hand, if $\gamma = -3$, we have (up to a constant) $\bar{c}^{\delta f} = \delta f$, and Lemma 4.5 directly implies the same upper bound for $\bar{c}^{\delta f}(|\chi_0|^2 + |\nu_0|^2)^{-\alpha/2}$. In addition, it is clear that $|\tau f| \langle v \rangle^m \lesssim \|f\|_{L^{\infty,k}}$ because $k \geq m$. By construction, we have $\|g(t_0, \cdot)\|_{L^\infty(\mathbb{R}^6 \times B_1(0)^2)} = g(t_0, x_0, \nu_0, \chi_0, \nu_0)$. Thus, at $(t_0, x_0, \nu_0, \chi_0, \nu_0)$, we have

$$J_4 \leq Cg.$$

For J_2 , we begin with a similar approach; observe that

$$\begin{aligned} \frac{\bar{a}^{\delta f}}{(|\chi_0|^2 + |\nu_0|^2)^{\alpha/2}} &\leq a_\gamma \int_{\mathbb{R}^3} |w|^{2+\gamma} \frac{\delta f(t_0, x_0, \nu_0 - w, \chi_0, \nu_0)}{(|\chi_0|^2 + |\nu_0|^2)^{\alpha/2}} dw \\ &\leq a_\gamma \int_{\mathbb{R}^3} \frac{|w|^{2+\gamma}}{\langle \nu_0 - w \rangle^m} \|g(t_0, \cdot)\|_{L^\infty(\mathbb{R}^6 \times B_1(0)^2)}^{\frac{1}{2}} dw \\ &\leq C \|g(t_0, \cdot)\|_{L^\infty(\mathbb{R}^6 \times B_1(0)^2)}^{1/2} \langle \nu_0 \rangle^{(2+\gamma)_+}. \end{aligned} \quad (4.9)$$

This holds since $m > 5 + \gamma$ by assumption. Thus

$$J_2 \lesssim |D_v^2(\tau f)| \langle \nu_0 \rangle^{m+(2+\gamma)_+} \|g(t_0, \cdot)\|_{L^\infty(\mathbb{R}^6 \times B_1(0)^2)}^{1/2}. \quad (4.10)$$

To close the estimate, we require an upper bound on $\|\langle v \rangle^{m+(2+\gamma)_+} D_v^2 f\|_{L^\infty}$, which is provided by Lemma 4.1; however, the results in Lemma 4.1 require working with the kinetic Hölder norms. In this case, it suffices to notice that

$$\|\langle v \rangle^m f\|_{C_{\text{kin},x,v}^\alpha([t_0/2, t_0] \times \mathbb{R}^6)} \lesssim \|\langle v \rangle^m f\|_{C^\alpha([t_0/2, t_0] \times \mathbb{R}^6)}.$$

This inequality follows from the easy-to-establish fact that $|z - z'| \lesssim \rho(z, z')$ when $t = t'$ and $|z - z'| \leq 1$. Putting this together with Lemma 4.1 and using the fact that $q(\alpha, \gamma, k, m) \leq -(2 + \gamma)_+$, we have

$$\|\langle v \rangle^{m+(2+\gamma)_+} D_v^2 f\|_{L^\infty} \leq \|\langle v \rangle^{m-q(\alpha, \gamma, k, m)} D_v^2 f\|_{L^\infty} \lesssim \frac{1}{t_0^{\frac{1-\frac{\alpha^2}{6-\alpha}}{2}}} (1 + \|\langle v \rangle^m f\|_{C^\alpha([t_0/2, t_0] \times \mathbb{R}^6)})^{p(\alpha)}.$$

Thus,

$$J_2 \lesssim \frac{1}{t_0^{\frac{1-\frac{\alpha^2}{6-\alpha}}{2}}} \left(\|\langle v \rangle^m f\|_{L_t^\infty C_{x,v}^\alpha([t_0/2, t_0] \times \mathbb{R}^6)}^{p(\alpha)} + 1 \right) \|g(t_0, \cdot)\|_{L^\infty(\mathbb{R}^6 \times B_1(0)^2)}^{1/2}.$$

Then, using Lemma 4.5 and the fact that $\|g(t_0, \cdot)\|_{L^\infty(\mathbb{R}^6 \times B_1(0)^2)} = \|g\|_{L^\infty([0, t_0] \times \mathbb{R}^6 \times B_1(0)^2)}$, since \bar{G} is increasing, we find

$$J_2 \lesssim \frac{1}{t_0^{\frac{1-\frac{\alpha^2}{6-\alpha}}{2}}} (1 + g)^{\frac{p(\alpha)+1}{2}},$$

which concludes the bound of J_2 .

Step 5: Establishing (4.6) and concluding the proof. Putting together the bounds on J_1, \dots, J_5 with (4.8), we find, at $(t_0, x_0, \nu_0, \chi_0, \nu_0)$,

$$\partial_t g \lesssim g + \frac{1}{t_0^{\frac{1-\frac{\alpha^2}{6-\alpha}}{2}}} (1 + g)^{\frac{p(\alpha)+1}{2}} \lesssim \frac{1}{t_0^{\frac{1-\frac{\alpha^2}{6-\alpha}}{2}}} (1 + g)^{\frac{p(\alpha)+1}{2}}. \quad (4.11)$$

The second inequality follows by using Young's inequality to show that $g \lesssim 1 + g^{P(\beta)}$. Choosing N to be larger than the implied constant in (4.11), we obtain (4.6). This yields a contradiction, as outlined above. Thus, our proof is concluded. \square

Finally, combining Lemma 4.3, Lemma 4.2, and Proposition 4.4, we obtain a time-integrable bound on $D_v^2 f$ for our solution:

Lemma 4.6. *Let the assumptions of Proposition 4.4 hold, and let f be the solution constructed in Theorem 1.2 corresponding to f_{in} . Then for $t \in [0, T_H]$,*

$$\| \langle v \rangle^{m+(\gamma+2)+} D_v^2 f(t, \cdot, \cdot) \|_{L^\infty(\mathbb{R}^6)} \lesssim \frac{1}{t^{1-\frac{\alpha^2}{6-\alpha}}}.$$

The implied constant depends on $\gamma, k, m, \alpha, \|f_{\text{in}}\|_{L^{\infty,k}}$, and $\|\langle v \rangle^m f_{\text{in}}\|_{C^\alpha}$.

5. Uniqueness

We are now ready to prove our solutions are unique. In this section, we allow all implied constants in the \lesssim notation to additionally depend on $\|f\|_{L^{\infty,k}}$ and $\|g\|_{L^{\infty,5+\gamma+\eta}}$, where k, g , and η are given below.

In order to state our result, we clarify the type of weak solution we work with. To use the results of [18,23], we require g to be in the kinetic Sobolev space required in these works; namely, for $\Omega \subset [0, \infty) \times \mathbb{R}^6$, let

$$H_{\text{kin}}^1(\Omega) = \{\phi \in L^2(\Omega) : \nabla_v \phi \in L^2(\Omega), (\partial_t + v \cdot \nabla_x) \phi \in L_{t,x}^2 H_v^{-1}(\Omega)\},$$

and let $H_{\text{kin,loc}}^1$ be defined in the obvious way. By a weak solution of (1.1), we mean a solution in the sense of integration against test functions in $H_{\text{kin}}^1([0, T_0] \times \mathbb{R}^6)$ with compact support.

First, we show uniformly continuous weak solutions have pointwise regularity:

Lemma 5.1. *Suppose that $g \in H_{\text{kin,loc}}^1((0, T_0] \times \mathbb{R}^6) \cap L^{\infty,5+\eta}([0, T_0] \times \mathbb{R}^6)$, with $\eta > 0$, solves (1.1) weakly and is uniformly continuous on $[0, T_0] \times \mathbb{R}^6$ and $g(0, \cdot, \cdot) = f_{\text{in}}$. Then $g \in C_{\text{kin,loc}}^2((0, T_0] \times \mathbb{R}^6)$.*

Proof. In view of the arguments used in Theorem 1.2, namely an application, up to rescaling, of [18, Theorem 3] and [23, Theorem 2.12(a)], it is enough to verify that M_g^{-1} , M_g , E_g , and H_g (recall the notation from Section 1.2) are bounded uniformly. The upper bounds on M_g , E_g , and H_g follow directly from the $L^{\infty,5+\eta}$ bound on g . Therefore, the proof is completed after establishing a uniform positive lower bound on M_g on $[0, T_0] \times \mathbb{R}^6$.

First, we set some notation. Let

$$A = 2\|\text{tr}(\bar{a}^g)\|_{L^\infty([0, T_0] \times \mathbb{R}^6)} + 1 \quad \text{and} \quad R = 2\sqrt{AT_0}.$$

Next, notice that since $g \in L^{\infty,5+\eta}([0, T_0] \times \mathbb{R}^6)$, g is uniformly continuous, and $M_{f_{\text{in}}} > (4\pi/3)\delta r^3$, then there exists $T_1 \in (0, T_0]$ such that, for all $(t, x) \in [0, T_1] \times \mathbb{R}^3$,

$$M_g(t, x) \geq \frac{1}{T_1}. \tag{5.1}$$

This concludes the proof on $[0, T_1] \times \mathbb{R}^3$, establishing that $g \in C_{\text{kin,loc}}^2((0, T_1] \times \mathbb{R}^6)$. Now that g is sufficiently regular, we note that a classical comparison principle argument yields $g \geq 0$ on $[0, T_1] \times \mathbb{R}^6$.

We now obtain a lower bound on M_g on $[T_1, T_0] \times \mathbb{R}^3$. Since we have bounds on M_g^{-1} , M_g , E_g , and H_g on $[0, T_1]$, we may iteratively apply the Harnack inequality for the Landau equation [18, Theorem 4] in order to find $\varepsilon > 0$ such that

$$\delta R^2 < g \quad \text{on } \{T_1\} \times \mathbb{R}^3 \times B_R(0).$$

Let

$$\underline{g}(t, x, v) = \varepsilon(R^2 - |v|^2 - A(t - T_1))_+,$$

and notice that, when $|v|^2 \leq R^2 - A(t - T_1)$,

$$(\partial_t + v \cdot \nabla_x) \underline{g} - \text{tr}(\bar{a}^g D_v^2 \underline{g}) - \bar{c}^g \underline{g} < 0. \tag{5.2}$$

If $\underline{g} \leq g$ on $[T_1, T_0] \times \mathbb{R}^6$, it is clear that we are finished since the choice of R and A imply that, for any $(t, x) \in [T_1, T_0] \times \mathbb{R}^3$,

$$\int_{B_{R/\sqrt{2}}} \underline{g}(t, x, v) dv \geq \frac{\varepsilon \pi R^5}{6\sqrt{2}} \geq \varepsilon R^5/10,$$

and, hence, $M_g \geq \varepsilon R^5/10$ on $[T_1, T_0] \times \mathbb{R}^3$. This, combined with (5.1) yields the desired lower bound on M_g and would complete the proof.

Hence, we assume that $\underline{g} \not\leq g$ on $[T_1, T_0] \times \mathbb{R}^6$. Let

$$T_h = \sup\{t \in [T_1, T_0] : \underline{g}(s, x, v) \leq g(s, x, v) \text{ for all } (s, x, v) \in [T_1, t] \times \mathbb{R}^6\}.$$

We claim that $T_h = T_0$. If this were true, our proof would be finished; hence, we suppose it is not true. At time T_h , we have that $g \geq \underline{g}$, by continuity. Hence $M_g(T_h, x) \geq \varepsilon R^5/10$ for all x . As above, we find $\mu > 0$ such that $M_g(t, x) > \mu$ for all $(t, x) \in [T_1, T_1 + \mu] \times \mathbb{R}^3$, and, hence, a $g \in C_{\text{kin,loc}}^2$ and a classical comparison principle argument shows that $g \geq 0$ on $[T_1, T_1 + \mu] \times \mathbb{R}^6$.

From above, we have established that M_g is positive on $[0, T_h + \mu] \times \mathbb{R}^6$ and $g \geq 0$ on the same set. Thus, we may apply the Harnack inequality [18, Theorem 4] in order to conclude that $g > 0$ on $[0, T_h + \mu] \times \mathbb{R}^6$.

By definition and since $T_h < T_0$, we find $z_0 \in [T_h, T_h + \mu] \times \mathbb{R}^6$ such that $g(z_0) < \underline{g}(z_0)$. By the positivity of g and up to recentering, we have that there exists $z_h \in [T_h, T_h + \mu] \times B_R \times \mathbb{R}^3$ such that $g(z_h) = \underline{g}(z_h)$, while $g \geq \underline{g}$ on $[0, t_h] \times \mathbb{R}^6$.

Let $\phi = g - \underline{g}$. From the work above, we have that z_h is the location of a minimum of ϕ where $\phi(z_h) = 0$. Hence

$$(\partial_t + v \cdot \nabla_x)\phi - \text{tr}(\bar{a}^g D_v^2 \phi) - \bar{c}^g \phi \leq 0$$

On the other hand, using (1.1) and (5.2), we find

$$(\partial_t + v \cdot \nabla_x)\phi - \text{tr}(\bar{a}^g D_v^2 \phi) - \bar{c}^g \phi > 0,$$

which contradicts the previous inequality. We conclude that $T_h = T_0$, and the proof is finished. \square

Proposition 5.2. Suppose that $\alpha \in (0, 1)$, k and m satisfy the conditions in Proposition 4.4 and $m \geq 5$, $0 \leq f_{\text{in}} \in L^{\infty,k}(\mathbb{R}^6)$, and $\langle v \rangle^m f_{\text{in}} \in C^\alpha(\mathbb{R}^6)$. Let f be the solution of (1.1) with initial data f_{in} constructed in Theorem 1.2, with $T_0 > 0$ such that $\langle v \rangle^m f \in C^\alpha([0, T_0] \times \mathbb{R}^6)$.

Let g be a weak solution of (1.1) satisfying the hypotheses of Lemma 5.1. Then $f = g$.

Before proceeding we comment briefly on the assumptions. The above is a form of weak-strong uniqueness; that is, the uniqueness holds in a weaker class than the constructed solution. We believe that this is not the weakest class in which uniqueness holds and, at the expense of more technical arguments, one may remove the added assumption that g is uniformly continuous. Indeed, if the lower bounds of Lemma 2.5 were extended to weak solutions, one would get uniform continuity “for free” from the Hölder estimate of [18].

Proof. We assume that $f_{\text{in}} \not\equiv 0$. Indeed if $f_{\text{in}} \equiv 0$, uniqueness holds via [24, Theorem 1.1]. By Lemma 5.1, $g \in C_{\text{kin,loc}}^2([0, T_0] \times \mathbb{R}^6)$.

By Proposition 3.1 and our assumptions, f and g are both continuous up to $t = 0$. For a positive function $r \in C(0, T_0] \cap L^1(0, T_0]$ to be determined, let

$$w = e^{-\int_0^t r(s) ds} (g - f) \quad \text{and} \quad W = \frac{1}{2} \langle v \rangle^{10} w^2.$$

Then a straightforward computation yields that, whenever $W \neq 0$,

$$\begin{aligned} \partial_t W + v \cdot \nabla_x W &= \text{tr}(\bar{a}^g D_v^2 W) - W^{-1} \nabla_v W \cdot (\bar{a}^g \nabla_v W) + 10 \langle v \rangle^{-2} v \cdot (\bar{a}^g \nabla_v W) \\ &+ \left(-35 \langle v \rangle^{-4} v \cdot (\bar{a}^g v) + 5 \langle v \rangle^{-2} \text{tr}(\bar{a}^g) + \bar{c}^g \right) W + \langle v \rangle^{10} w \text{tr}(\bar{a}^w D_v^2 f) + \langle v \rangle^{10} w \bar{c}^w f - r W. \end{aligned} \quad (5.3)$$

Fix $\varepsilon > 0$, and assume by contradiction that $\sup_{[0, T_0] \times \mathbb{R}^6} W(t, x, v) \geq \varepsilon$. Up to re-centering the equation as in Step 2 of the proof of Proposition 4.4, we may assume there exists $z_\varepsilon = (t_\varepsilon, x_\varepsilon, v_\varepsilon) \in (0, T_0] \times \mathbb{R}^6$ such that $W(z_\varepsilon) = \varepsilon$

and $W(t, x, v) < \varepsilon$ for all $t < t_\varepsilon$ and $(x, v) \in \mathbb{R}^6$. Note that this reduction uses strongly that the f and g are uniformly continuous for two reasons: (i) the re-centering requires an Arzela-Ascoli-based compactness argument on the x -translates of W and (2) to conclude that $t_\varepsilon > 0$ after the re-centering.

We immediately have that, at z_ε , $\nabla_v W = 0$ and $D_v^2 W \leq 0$. Derivatives in t and x may not exist pointwise, but we have $(\partial_t + v \cdot \nabla_x)W \geq 0$. (This follows by considering the directional derivative of $W(t, x, v)$ at z_ε in the direction $(1, v, 0)$, since z_ε is a maximum point in $[0, t_\varepsilon] \times \mathbb{R}^6$.) It is also clear that $W(z_\varepsilon) = \varepsilon = \|W\|_{L^\infty([0, t_\varepsilon] \times \mathbb{R}^6)}$. Finally, since $g \in L^{\infty, 5+\eta}$, we have that \bar{a}^g and \bar{c}^g are bounded according to Lemma 2.1. Using the above in (5.3) and moving the rW term to the left hand side, we find

$$rW \lesssim W + \langle v \rangle^{10} |w| |\bar{a}^w| |D_v^2 f| + \langle v \rangle^5 |w| |\bar{c}^w|. \quad (5.4)$$

Next, we notice that, at z_ε ,

$$|\bar{a}^w| \lesssim \langle v \rangle^{7+\gamma} \|w(t_\varepsilon, \cdot, \cdot)\|_{L^\infty(\mathbb{R}^6)} = \langle v \rangle^{7+\gamma} w \quad \text{and, similarly,} \quad |\bar{c}^w| \lesssim \langle v \rangle^5 w. \quad (5.5)$$

Recall that $m \geq 5$, and, thus, $m + (2 + \gamma)_+ \geq 7 + \gamma$. We can also apply Lemma 4.6 to obtain

$$\langle v_\varepsilon \rangle^{7+\gamma} |D_v^2 f| \leq \|D_v^2 f(t_\varepsilon, \cdot, \cdot)\|_{L^{\infty, 7+\gamma}(\mathbb{R}^6)} \lesssim t_\varepsilon^{-1+\frac{\alpha^2}{6-\alpha}}. \quad (5.6)$$

Using (5.5) and (5.6) in (5.4) and recalling the relationship between w and W , we find a constant C , such that

$$rW \leq C \left(1 + t_\varepsilon^{-1+\frac{\alpha^2}{6-\alpha}}\right) W.$$

With this value of C , we obtain a contradiction by defining

$$r(t) = C \left(2 + t^{-1+\frac{\alpha^2}{6-\alpha}}\right).$$

We therefore conclude that z_ε cannot exist. Since this is true for all ε , we find that $W = 0$, which implies that $g = f$. \square

Finally, it is straightforward to prove Theorem 1.4 from Proposition 5.2:

Proof of Theorem 1.4. By Proposition 4.4, there exists $T_H > 0$ such that $\langle v \rangle^m f \in C_{\text{kin}}^\alpha([0, T_H] \times \mathbb{R}^6)$. Thus, we may apply Proposition 5.2 to conclude the first part of Theorem 1.4.

We now consider the case where $f_{\text{in}} \in L^{\infty, k'}(\mathbb{R}^6)$ for all k' . Combining Theorem 1.2.(iii) and Theorem 1.3, we see that f is smooth and all its derivatives lie in $L^{\infty, k'}([0, T] \times \mathbb{R}^6)$ for all k' . Thus, Proposition 5.2 applies with $T_H = T$, which concludes the proof. \square

Declaration of competing interest

There are no competing interests.

Appendix A. Regularity in x and v implies regularity in t

We consider solutions f to the linear equation

$$\partial_t f + v \cdot \nabla_x f = \text{tr}(a D_v^2 f) + cf, \quad (A.1)$$

where a is a non-negative definite matrix that grows at most like $\langle v \rangle^{2+\gamma}$, c is uniformly bounded, and (A.1) enjoys a maximum principle. All implied constants depend only on the upper bounds of a and c , but *do not* depend on any lower bound of a .

In this appendix, we show the following:

Proposition A.1. Suppose that $f \in C_{\text{kin},\text{loc}}^\alpha([0, T] \times \mathbb{R}^6) \cap L^\infty([0, T] \times \mathbb{R}^6)$ and solves (A.1). Then, for any z_0 , we have

$$\|f\|_{C_{\text{kin}}^\alpha(Q_1(z_0) \cap [0, T] \times \mathbb{R}^6)} \lesssim \langle v_0 \rangle^{\alpha(1+\gamma/2)+} \left(\|f\|_{L^\infty([0, T] \times \mathbb{R}^6)} + \sup_{t \in [\max\{0, t_0-1\}, t_0]} [f(t, \cdot, \cdot)]_{C_{\text{kin},x,v}^\alpha(B_2(x_0, v_0))} \right).$$

The implied constant depends only on the upper bounds of a and c .

A useful transformation here is given by

$$\mathcal{T}_{z_0}(z) = (t_0 + t, x_0 + x + tv_0, v_0 + v).$$

Recall also δ_r from (2.6). The main lemma is the following.

Lemma A.2. For any $z_0 \in [0, T] \times \mathbb{R}^6$, $r \in (0, 1]$, $t_1 \in [0, \langle v_0 \rangle^{-(2+\gamma)+}]$ such that $r^2 t_1 + t_0 \in [0, T]$ and $|x_1|, |v_1| < 1$, and for $z_1 = (t_1, x_1, v_1)$, we have

$$|f(\mathcal{T}_{z_0}(\delta_r(z_1))) - f(z_0)| \lesssim |r|^\alpha \left(\|f\|_{L^\infty([0, T] \times \mathbb{R}^6)} + [f(t_0, \cdot, \cdot)]_{C_{\text{kin},x,v}^\alpha(B_2(x_0, v_0))} \right).$$

A.1. Concluding Proposition A.1 from Lemma A.2

Proof of Proposition A.1. Fix any z_1 and z_2 in $Q_1(z_0)$. Recall that $\rho(z_1, z_2) \approx \rho(z_2, z_1)$ so we may assume that $t_2 \geq t_1$.

We consider first the case when $\rho(z_2, z_1) \geq \langle v_1 \rangle^{-(1+\gamma/2)+}$. Then we have

$$\frac{|f(z_2) - f(z_1)|}{\rho(z_2, z_1)^\alpha} \leq \langle v_1 \rangle^{(1+\gamma/2)+\alpha} \|f\|_{L^\infty([0, T] \times \mathbb{R}^6)} \lesssim \langle v_0 \rangle^{(1+\gamma/2)+\alpha} \|f\|_{L^\infty([0, T] \times \mathbb{R}^6)},$$

since $\langle v_1 \rangle^{(1+\gamma/2)+} \approx \langle v_0 \rangle^{(1+\gamma/2)+}$.

Next we consider the case when $\rho(z_2, z_1) < \langle v_1 \rangle^{-(1+\gamma/2)+}$. Clearly, then, $t_2 - t_1 < \langle v_0 \rangle^{-(2+\gamma)+}$. Let

$$s_2 = \langle v_1 \rangle^{-(2+\gamma)+}, \quad r^2 = \frac{t_2 - t_1}{s_2}, \quad w_2 = \frac{v_2 - v_1}{r}, \quad \text{and} \quad y_2 = \frac{x_2 - x_1 - r^2 s_2 v_1}{r^3}.$$

Notice that $z_2 = \mathcal{T}_{z_1}(s_2, y_2, w_2)$, $|s_2| \leq \langle v_1 \rangle^{-(2+\gamma)+}$, and $r \in (0, 1)$. Hence, we may apply Lemma A.2 to find

$$|f(z_2) - f(z_1)| \lesssim r^\alpha \left(\|f\|_{L^\infty([0, T] \times \mathbb{R}^6)} + [f(t_1, \cdot, \cdot)]_{C_{\text{kin},x,v}^\alpha(B_2(x_0, v_0))} \right).$$

Recalling the definition of r , we have

$$\begin{aligned} \frac{|f(z_2) - f(z_1)|}{\rho(z_2, z_1)^\alpha} &\lesssim \frac{r^\alpha}{\rho(z_2, z_1)^\alpha} \left(\|f\|_{L^\infty([0, T] \times \mathbb{R}^6)} + [f(t_1, \cdot, \cdot)]_{C_{\text{kin},x,v}^\alpha(B_2(x_0, v_0))} \right) \\ &= \frac{|t_2 - t_1|^{\alpha/2}}{s_1^{\alpha/2} \rho(z_2, z_1)^\alpha} \left(\|f\|_{L^\infty([0, T] \times \mathbb{R}^6)} + [f(t_1, \cdot, \cdot)]_{C_{\text{kin},x,v}^\alpha(B_2(x_0, v_0))} \right) \end{aligned}$$

The proof is finished after noting that $s_1^{-\alpha/2} \lesssim \langle v_0 \rangle^{\alpha(1+\gamma/2)+}$ and $|t_2 - t_1|^{\alpha/2} \leq \rho(z_2, z_1)^\alpha$. \square

A.2. The proof of Lemma A.2

Proof of Lemma A.2. The proof is based on a maximum principle argument. Let z_0 be as in the statement of the lemma. Without loss of generality, we may assume that $t_0 = 0$ and $x_0 = 0$. Then $z_1 \in [0, \min\{\langle v_0 \rangle^{(2+\gamma)+}, T\}] \times B_1(0) \times B_1(v_0)$.

Step 1: A cut-off function. Let $\phi \in C_c^\infty(\mathbb{R}^6)$ be a cut-off function such that

$$\begin{aligned} 0 &\leq \phi \leq 1, \quad \phi \equiv 1 \text{ on } B_1(0), \\ \phi &\lesssim \langle v \rangle^{-3} \langle x \rangle^{-3}, \\ |\partial_{v_i} \phi| &\lesssim \langle v \rangle^{-4} \langle x \rangle^{-3} \quad \text{for all } i = 1, 2, 3, \\ |\partial_{x_i} \phi| &\lesssim \langle v \rangle^{-3} \langle x \rangle^{-4} \quad \text{for all } i = 1, 2, 3, \quad \text{and} \\ |\partial_{v_i v_j} \phi| &\lesssim \langle v \rangle^{-5} \langle x \rangle^{-3} \quad \text{for all } i, j = 1, 2, 3. \end{aligned} \tag{A.2}$$

We also set some useful notation. For any $r \geq 0$ and any function g , let

$$g_r(z) = g(\mathcal{T}_{z_0}(\delta_r(z))).$$

Step 2: An auxiliary function and its equation. Then, let

$$\tilde{F}(z) = f_r(z) - f(0, 0, v_0).$$

It is straightforward to check that

$$\tilde{F}_t + v \cdot \nabla_x \tilde{F} - \text{tr}(a_r D_v^2 \tilde{F}) = r^2 c_r f_r.$$

Let

$$\psi(t, x, v) = \phi(x - tv, v) \quad \text{and} \quad F = \psi \tilde{F}.$$

Again, after a straightforward computation, we find

$$\begin{aligned} F_t + v \cdot \nabla_x F - \text{tr}(a_r D_v^2 F) + 2(a_r \nabla_v \log(\psi)) \cdot \nabla_v F \\ = F \left(2\nabla_v \log(\psi) \cdot (a_r \nabla_v \log(\psi)) - \text{tr}(a_r \psi^{-1} D_v^2 \psi) \right) + r^2 \psi c_r f_r. \end{aligned}$$

Fix $R = 2\langle v_0 \rangle^3 / r$. We claim that there exists a constant $C > 0$ such that, on $[0, t_1] \times \mathbb{R}^3 \times B_R(0)$,

$$2|\nabla_v \log(\psi) \cdot (a_r \nabla_v \log(\psi))| + |\text{tr}(a_r \psi^{-1} D_v^2 \psi)| < C \langle v_0 \rangle^{(2+\gamma)+}. \tag{A.3}$$

This is established at the end of this proof.

Assuming (A.3), let

$$\overline{F}(t) = e^{tC\langle v_0 \rangle^{(2+\gamma)+}} \left(\|F(0, \cdot, \cdot)\|_{L^\infty(\mathbb{R}^6)} + \sup_{s \in [0, t_1], x \in \mathbb{R}^3, |v|=R} F(s, x, v)_+ + r^2 t \|c\|_{L^\infty} \|f\|_{L^\infty} \right).$$

Then, since $\overline{F} \geq 0$,

$$\begin{aligned} \overline{F}_t + v \cdot \nabla_x \overline{F} - \text{tr}(a_r D_v^2 \overline{F}) + 2(a_r \nabla_v \log(\psi)) \cdot \nabla_v \overline{F} \\ = C \langle v_0 \rangle^{(2+\gamma)+} \overline{F} + r^2 \|c\|_{L^\infty} \|f\|_{L^\infty} e^{C\langle v_0 \rangle^{(2+\gamma)+} t} \\ > \overline{F} \left(2\nabla_v \log(\psi) \cdot (a_r \nabla_v \log(\psi)) - \text{tr}(a \psi^{-1} D_v^2 \psi) \right) + r^2 \psi c_r f_r. \end{aligned}$$

Above, we used (A.3) and that $\|f_r\|_{L^\infty} = \|f\|_{L^\infty}$. Note that ψ is compactly supported in x . Thus, by the maximum principle, we find $F(t, x, v) \leq \overline{F}(t)$ on $[0, t_1] \times \mathbb{R}^3 \times B_R(0)$. In particular, we have

$$F(t_1, x_1, v_1) \leq e^{t_1 C \langle v_0 \rangle^{(2+\gamma)+}} \left(\|F(0, \cdot, \cdot)\|_{L^\infty(\mathbb{R}^6)} + \sup_{s \in [0, t_1], x \in \mathbb{R}^3, |v|=R} F(s, x, v)_+ + t_1 r^2 \|c\|_{L^\infty} \|f\|_{L^\infty} \right).$$

For the lower bound of $F(t_1, x_1, v_1)$, consider

$$\underline{F}(t, x, v) = -e^{tC\langle v_0 \rangle^{(2+\gamma)+}} \left(\|F(0)\|_{L^\infty} + \sup_{s \in [0, t_1], x \in \mathbb{R}^3, |v|=R} F(s, x, v)_- \right).$$

By an argument similar to the one for \overline{F} , but simpler because one can use $0 \leq r^2 \psi c_r f_r$, we have $F(t, x, v) \geq \underline{F}(t)$ on $[0, t_1] \times \mathbb{R}^3 \times B_R(0)$. Altogether, we have

$$|F(t_1, x_1, v_1)| \leq e^{t_1 C \langle v_0 \rangle^{(2+\gamma)+}} \left(\|F(0, \cdot, \cdot)\|_{L^\infty(\mathbb{R}^6)} + \sup_{s \in [0, t_1], x \in \mathbb{R}^3, |v|=R} |F(s, x, v)| + r^2 \|c\|_{L^\infty} \|f\|_{L^\infty} \right). \quad (\text{A.4})$$

Step 3: Quantitative bounds on the right hand side of (A.4). Unpacking the coordinate transformations and using the decay of ϕ , it is easy to verify that

$$\|F(0, \cdot, \cdot)\|_{L^\infty(\mathbb{R}^6)} \lesssim r^\alpha \left([f(0, \cdot, \cdot)]_{C_{\text{kin}}^\alpha(B_2(0, v_0))} + \|f(0, \cdot, \cdot)\|_{L^\infty(\mathbb{R}^6)} \right). \quad (\text{A.5})$$

Indeed, fix any $(x, v) \in \mathbb{R}^6$. We consider two cases. First, if $r^3|x|, r|v| \leq 2$, then

$$\begin{aligned} |\psi(0, x, v)(f_r(0, x, v) - f(0, 0, v_0))| &\lesssim \langle x \rangle^{-3} \langle v \rangle^{-3} |f(0, r^3 x, r v + v_0) - f(0, 0, v_0)| \\ &\leq \langle x \rangle^{-3} \langle v \rangle^{-3} \rho((0, r^3 x, r v + v_0), (0, 0, v_0))^\alpha [f]_{C_{\text{kin}}^\alpha(Q_2(0, 0, v_0))} \\ &= \langle x \rangle^{-3} \langle v \rangle^{-3} r^\alpha (|x|^{1/3} + |v|)^\alpha [f]_{C_{\text{kin}}^\alpha(Q_2(0, 0, v_0))} \leq r^\alpha [f]_{C_{\text{kin}}^\alpha(Q_2(0, 0, v_0))}. \end{aligned}$$

Second, if $r^3|x|$ or $r|v|$ is greater than 2, then

$$|\psi(0, x, v)(f_r(0, x, v) - f(0, 0, v_0))| \lesssim \frac{\|f(0, \cdot, \cdot)\|_{L^\infty(\mathbb{R}^6)}}{\langle x \rangle^3 \langle v \rangle^3} \lesssim r^3 \|f(0, \cdot, \cdot)\|_{L^\infty(\mathbb{R}^6)} \leq r^\alpha \|f(0, \cdot, \cdot)\|_{L^\infty(\mathbb{R}^6)},$$

since $r \leq 1$. Hence (A.5) holds.

Next, we check the boundary term. Indeed, if $(s, x, v) \in [0, t_1] \times \mathbb{R}^3 \times \partial B_R(0)$, then

$$|F(s, x, v)| = \psi(s, x, v) |\tilde{F}(s, x, v)| \lesssim \frac{\|f\|_{L^\infty}}{\langle x - vs \rangle^3 \langle v \rangle^3} \lesssim \frac{r^3 \|f\|_{L^\infty}}{\langle v_0 \rangle^9} \leq r^\alpha \|f\|_{L^\infty}.$$

We conclude that

$$\sup_{s \in [0, t_1], x \in \mathbb{R}^3, |v|=R} |F(s, x, v)| \lesssim r^\alpha \|f\|_{L^\infty}. \quad (\text{A.6})$$

Plugging (A.5) and (A.6) into (A.4), and using the fact that $t_1 \leq \langle v_0 \rangle^{-(2+\gamma)+}$, we find

$$|F(t_1, x_1, v_1)| \lesssim e^{C \langle v_0 \rangle^{2+\gamma} t_1} r^\alpha \left([f(0, \cdot, \cdot)]_{C_{\text{kin}}^\alpha(B_2(0, v_0))} + \|f\|_{L^\infty} \right). \quad (\text{A.7})$$

Step 4: Unpacking the transformations to obtain the desired estimate. Since $|t_1|, |x_1|, |v_1| < 1$, then $\psi(t_1, x_1, v_1) = 1$ and $v_1 \in B_R(0)$. We thus conclude, from (A.7), that

$$|f_r(t_1, x_1, v_1) - f(0, 0, v_0)| = F(t_1, x_1, v_1) \lesssim e^{C \langle v_0 \rangle^{2+\gamma} t_1} r^\alpha \left([f(0, \cdot, \cdot)]_{C_{\text{kin}}^\alpha(B_2(0, v_0))} + \|f\|_{L^\infty} \right).$$

Hence, the proof is finished once we establish (A.3), which is the last step.

Step 5: The coefficient bounds (A.3). We show that the first term on the left in (A.3) is bounded by $C \langle v_0 \rangle^{(2+\gamma)+}$. We omit the proof for the second term, which is similar.

Notice that $\nabla_v \psi(t, x, v) = (\nabla_v - t \nabla_x) \phi(x - vt, v)$. Using the non-negative definiteness of a , we obtain

$$|\nabla_v \log(\psi) \cdot (a_r \nabla_v \log(\psi))| \leq 2 \left(\nabla_v \log(\phi) \cdot (a_r \nabla_v \log(\phi)) + t^2 \nabla_x \log(\phi) \cdot (a_r \nabla_x \log(\phi)) \right). \quad (\text{A.8})$$

In the above, we have abused notation in the following way: a_r is evaluated at (t, x, v) while ϕ is evaluated at $(x - vt, v)$.

For the first term on the right in (A.8), we find from (A.2) that

$$\nabla_v \log(\phi) \cdot (a_r \nabla_v \log(\phi)) \leq |a_r| |\nabla_v \log(\phi)|^2 \lesssim \frac{\langle v_0 + rv \rangle^{2+\gamma}}{\langle v \rangle^2}. \quad (\text{A.9})$$

It is straightforward to establish, by considering the four cases when $|rv|$ is comparable or not to $|v_0|$ and when $\gamma \geq -2$ or $\gamma < -2$, that the right hand side is bounded by $\langle v_0 \rangle^{2+\gamma}$.

The second term in (A.8) can be handled as follows. When $\gamma < -2$, since all terms are bounded, we find

$$t^2 \nabla_x \log(\phi) \cdot (a_r \nabla_x \log(\phi)) \lesssim 1 \leq \langle v_0 \rangle^{(2+\gamma)_+}.$$

Next, consider the case when $2 + \gamma \geq 0$. Again using (A.2), we find

$$t^2 \nabla_x \log(\phi) \cdot (a_r \nabla_x \log(\phi)) \leq t^2 |a_r| |\nabla_x \log(\phi)|^2 \lesssim t^2 \frac{\langle v_0 + rv \rangle^{2+\gamma}}{\langle x - vt \rangle^2}.$$

This case is more subtle than (A.9) because the $x - vt$ term in the denominator may not be large, even if v is large. Therefore, we must use the smallness of t to balance the fact that $\langle v_0 + rv \rangle$ may be large. Recall that $|v| \leq R = 2\langle v_0 \rangle^3/r$ and $t \leq \langle v_0 \rangle^{-(2+\gamma)_+}$. Since $\langle x - vt \rangle \geq 1$, we have

$$t^2 \frac{\langle v_0 + rv \rangle^{2+\gamma}}{\langle x - vt \rangle^2} \lesssim \langle v_0 \rangle^{-2(2+\gamma)} \langle v_0 \rangle^{3(2+\gamma)} = \langle v_0 \rangle^{2+\gamma} = \langle v_0 \rangle^{(2+\gamma)_+},$$

as desired. This completes the proof of (A.3), which completes the proof of the lemma. \square

Appendix B. Interpolation lemmas

Here, we collect the technical interpolation lemmas used in Section 4. The first lemma is for functions on \mathbb{R}^3 with the standard Euclidean Hölder metric:

Lemma B.1. Suppose that $d \in \mathbb{N}$ and $\phi : \mathbb{R}^d \rightarrow \mathbb{R}$ is a bounded $C_{\text{loc}}^{2,\beta}$ function, with $\beta \in (0, 1)$. For any $\alpha \in (0, 1)$,

$$\|D^2\phi\|_{L^\infty(B_1(z))} \lesssim [\phi]_{C^\alpha(B_2(z))} + [\phi]_{C^\alpha(B_2(z))}^{\frac{\beta}{2+\beta-\alpha}} [D^2\phi]_{C^\beta(B_2(z))}^{1-\frac{\beta}{2+\beta-\alpha}}. \quad (\text{B.1})$$

Proof. Let $z \in \mathbb{R}^d$ be a given point. Taking $x \in B_1(z)$ and $h \in \mathbb{R}^d$ sufficiently small, we see that

$$\phi(x+h) - \phi(x) = \nabla\phi(\xi_1)h = \nabla\phi(x)h + (\nabla\phi(\xi_1) - \nabla\phi(x))h,$$

where $\xi_1 = x + \theta_1 h$ for some $\theta_1 \in (0, 1)$. Let $\sigma > 0$ be a constant, to be chosen later. If $|h| \leq \sigma$, then

$$|\nabla\phi(\xi_1) - \nabla\phi(x)| \leq \sigma \|D^2\phi\|_{L^\infty(B_\sigma(x))}.$$

If we let $h = \sigma \nabla\phi(x)/|\nabla\phi(x)|$, we then get

$$|\nabla\phi(x)| \leq \sigma^{\alpha-1} [\phi]_{C^\alpha(B_\sigma(x))} + \sigma \|D^2\phi\|_{L^\infty(B_\sigma(x))}. \quad (\text{B.2})$$

Now, taking the Taylor expansion of ϕ to first order about a point $x \in B_1(z)$ gives

$$\phi(x+h) = \phi(x) + \nabla\phi(x)h + \frac{1}{2}hD^2\phi(\xi_2)h,$$

where $\xi_2 = x + \theta_2 h$ for some $\theta_2 \in (0, 1)$. We rewrite this as

$$hD^2\phi(x)h = 2(\phi(x+h) - \phi(x)) - 2\nabla\phi(x)h + h(D^2\phi(x) - D^2\phi(\xi_2))h.$$

Since $D^2\phi(x)$ is a real symmetric matrix, its matrix norm is given by its largest eigenvalue, which is achieved by some unit eigenvector \hat{v} . With $r > 0$ to be chosen later, setting $h = r\hat{v}$ yields

$$\begin{aligned} r^2 |D^2\phi(x)| &= 2(\phi(x+h) - \phi(x)) - 2\nabla\phi(x)h + h(D^2\phi(x) - D^2\phi(\xi_2))h \\ &\leq 2r^\alpha [\phi]_{C^\alpha(B_r(x))} + 2r \|\nabla\phi\|_{L^\infty(B_r(x))} + r^{2+\beta} [D^2\phi]_{C^\beta(B_r(x))}. \end{aligned}$$

Taking the supremum in x over $B_1(z)$, we get that

$$\|D^2\phi\|_{L^\infty(B_1(z))} \lesssim r^{\alpha-2} [\phi]_{C^\alpha(B_{1+r}(z))} + r^{-1} \|\nabla\phi\|_{L^\infty(B_{1+r}(z))} + r^\beta [D^2\phi]_{C^\beta(B_{1+r}(z))}.$$

Using (B.2) on the middle term yields

$$\begin{aligned} \|D^2\phi\|_{L^\infty(B_1(z))} &\lesssim \left(r^{\alpha-2} + r^{-1}\sigma^{\alpha-1}\right) [\phi]_{C^\alpha(B_{1+r+\sigma}(z))} \\ &\quad + \sigma r^{-1} \|D^2\phi\|_{L^\infty(B_{1+r+\sigma}(x))} + r^\beta [D^2\phi]_{C^\beta(B_{1+r}(z))}. \end{aligned} \quad (\text{B.3})$$

We can then take $\sigma = cr$ for $c < 1$ small enough that we can absorb the middle term into the norm on the left hand side. As long as $r < 1/2$, this yields

$$\|D^2\phi\|_{L^\infty} \lesssim r^{\alpha-2} [\phi]_{C^\alpha(B_2(z))} + r^\beta [D^2\phi]_{C^\beta(B_2(z))}.$$

Thus, choosing

$$r = \min \left\{ 1/2, \left([\phi]_{C^\alpha(B_2(z))} / [D^2\phi]_{C^\beta(B_2(z))} \right)^{\frac{1}{2+\beta-\alpha}} \right\},$$

then we obtain (B.1). \square

Our next, somewhat elementary lemma allows us to trade regularity (in the form of a larger Hölder exponent) for pointwise decay. It is stated for the (time-independent) kinetic Hölder spaces, however, it is straightforward to prove it for the time-dependent kinetic and standard Hölder spaces as well.

Lemma B.2. *Suppose that $\phi : \mathbb{R}^6 \rightarrow \mathbb{R}$ is such that $\phi \in L^{\infty, k_1}(\mathbb{R}^6)$ and $\langle v \rangle^{k_2} \phi \in C_{\text{kin}}^\alpha(\mathbb{R}^6)$, for some $\alpha \in (0, 1)$ and $k_1 \geq k_2 \geq 0$. If $\beta \in (0, \alpha)$ and $\ell \in [k_2, k_1]$ are such that*

$$\ell \leq k_1 \left(1 - \frac{\beta}{\alpha} \right) + k_2 \frac{\beta}{\alpha},$$

then

$$\|\langle v \rangle^\ell \phi\|_{C_{\text{kin}}^\beta(\mathbb{R}^6)} \lesssim [\langle v \rangle^{k_2} \phi]_{C_{\text{kin}}^\alpha(\mathbb{R}^6)}^{\frac{\beta}{\alpha}} \|\phi\|_{L^{\infty, k_1}(\mathbb{R}^6)}^{1 - \frac{\beta}{\alpha}}.$$

The same result is true with the standard Hölder spaces replacing the kinetic ones.

Proof. We omit the proof with the standard Hölder spaces since it is exactly the same. Let

$$R = \left(\langle v \rangle^{k_2 - k_1} \|\phi\|_{L^{\infty, k_1}(\mathbb{R}^6)} [\langle v \rangle^{k_2} \phi]_{C_{\text{kin}}^\alpha(\mathbb{R}^6)}^{-1} \right)^{1/\alpha}.$$

Let $(x_1, v_1), (x_2, v_2) \in \mathbb{R}^6$. If $|x_1 - x_2|^{1/3} + |v_1 - v_2| \geq R$, then

$$\frac{|\phi(x_1, v_1) - \phi(x_2, v_2)|}{(|x_1 - x_2|^{1/3} + |v_1 - v_2|)^\beta} \lesssim \langle v \rangle^{-k_1} \frac{2\|\phi\|_{L^{\infty, k_1}(\mathbb{R}^6)}}{R^\beta} = \frac{[\langle v \rangle^{k_2} \phi]_{C_{\text{kin}}^\alpha(B)}^{\frac{\beta}{\alpha}} \|\phi\|_{L^{\infty, k_1}(\mathbb{R}^6)}^{1 - \frac{\beta}{\alpha}}}{\langle v \rangle^{k_1 \left(1 - \frac{\beta}{\alpha} \right) + k_2 \frac{\beta}{\alpha}}}.$$

On the other hand, suppose that $|x_1 - x_2|^{1/3} + |v_1 - v_2| < R$. Let $\theta = |x_1 - x_2|^{1/3} + |v_1 - v_2|$. We have

$$\frac{|\phi(x_1, v_1) - \phi(x_2, v_2)|}{\theta^\beta} = \frac{|\phi(x_1, v_1) - \phi(x_2, v_2)|}{\theta^\alpha} \frac{\theta^\alpha}{\theta^\beta} \lesssim \langle v \rangle^{-k_2} [\langle v \rangle^{k_2} \phi]_{C_{\text{kin}}^\alpha(B)} R^{\alpha - \beta}.$$

From the definition of R , we obtain

$$\frac{|\phi(x_1, v_1) - \phi(x_2, v_2)|}{\theta^\beta} \leq \frac{[\langle v \rangle^{k_2} \phi]_{C_{\text{kin}}^\alpha(B)}^{\frac{\beta}{\alpha}} \|\phi\|_{L^{\infty, k_1}(\mathbb{R}^6)}^{1 - \frac{\beta}{\alpha}}}{\langle v \rangle^{k_1 \left(1 - \frac{\beta}{\alpha} \right) + k_2 \frac{\beta}{\alpha}}}. \quad \square$$

References

- [1] R. Alexandre, J. Liao, C. Lin, Some a priori estimates for the homogeneous Landau equation with soft potentials, *Kinet. Relat. Models* 8 (4) (2015) 617–650.
- [2] R. Alexandre, Y. Morimoto, S. Ukai, C.-J. Xu, T. Yang, Regularizing effect and local existence for the non-cutoff Boltzmann equation, *Arch. Ration. Mech. Anal.* 198 (1) (2010) 39–123.
- [3] R. Alexandre, Y. Morimoto, S. Ukai, C.-J. Xu, T. Yang, Bounded solutions of the Boltzmann equation in the whole space, *Kinet. Relat. Models* 4 (1) (2011) 17–40.
- [4] R. Alexandre, Y. Morimoto, S. Ukai, C.-J. Xu, T. Yang, Local existence with mild regularity for the Boltzmann equation, *Kinet. Relat. Models* 6 (4) (2013) 1011–1041.
- [5] R. Alexandre, C. Villani, On the Landau approximation in plasma physics, *Ann. Inst. Henri Poincaré, Anal. Non Linéaire* 21 (1) (2004) 61–95.
- [6] A.A. Arsen'ev, N.V. Peskov, The existence of a generalized solution of Landau's equation, *Ž. Vyčisl. Mat. Mat. Fiz.* 17 (4) (1977) 1063–1068, 1096.
- [7] S. Cameron, L. Silvestre, S. Snelson, Global a priori estimates for the inhomogeneous Landau equation with moderately soft potentials, *Ann. Inst. Henri Poincaré, Anal. Non Linéaire* 35 (3) (2018) 625–642.
- [8] K. Carrapatoso, S. Mischler, Landau equation for very soft and Coulomb potentials near Maxwellians, *Ann. PDE* 3 (1) (Jan 2017) 1.
- [9] K. Carrapatoso, I. Tristani, K.-C. Wu, Cauchy problem and exponential stability for the inhomogeneous Landau equation, *Arch. Ration. Mech. Anal.* 221 (1) (2016) 363–418.
- [10] S. Chapman, T.G. Cowling, *The Mathematical Theory of Non-uniform Gases: An Account of the Kinetic Theory of Viscosity, Thermal Conduction and Diffusion in Gases*, 3rd edition, Cambridge University Press, 1970.
- [11] S. Chaturvedi, Local existence for the Landau equation with hard potentials, Preprint, arXiv:1910.11866, 2019.
- [12] Y. Chen, L. Desvillettes, L. He, Smoothing effects for classical solutions of the full Landau equation, *Arch. Ration. Mech. Anal.* 193 (1) (2009) 21–55.
- [13] P. Constantin, A. Tarfulea, V. Vicol, Long time dynamics of forced critical SQG, *Commun. Math. Phys.* 335 (1) (2015) 93–141.
- [14] L. Desvillettes, Entropy dissipation estimates for the Landau equation in the Coulomb case and applications, *J. Funct. Anal.* 269 (5) (2015) 1359–1403.
- [15] L. Desvillettes, C. Villani, On the spatially homogeneous Landau equation for hard potentials part I: existence, uniqueness and smoothness, *Commun. Partial Differ. Equ.* 25 (1–2) (2000) 179–259.
- [16] R. Duan, S. Liu, S. Sakamoto, R.M. Strain, Global mild solutions of the Landau and non-cutoff Boltzmann equations, Preprint, arXiv:1904.12086, 2019.
- [17] F. Golse, M. Gualdani, C. Imbert, A. Vasseur, Partial regularity in time for the space homogeneous Landau equation with Coulomb potential, Preprint, arXiv:1906.02841, 2019.
- [18] F. Golse, C. Imbert, C. Mouhot, A. Vasseur, Harnack inequality for kinetic Fokker-Planck equations with rough coefficients and application to the Landau equation, *Ann. Sc. Norm. Super. Pisa, Cl. Sci.* XIX (1) (2019) 253–295.
- [19] M. Gualdani, N. Guillen, Estimates for radial solutions of the homogeneous Landau equation with Coulomb potential, *Anal. PDE* 9 (8) (2016) 1773–1810.
- [20] M. Gualdani, N. Guillen, On A_p weights and the Landau equation, *Calc. Var. Partial Differ. Equ.* 58 (1) (2018) 17.
- [21] Y. Guo, The Landau equation in a periodic box, *Commun. Math. Phys.* 231 (3) (2002) 391–434.
- [22] L. He, X. Yang, Well-posedness and asymptotics of grazing collisions limit of Boltzmann equation with Coulomb interaction, *SIAM J. Math. Anal.* 46 (6) (2014) 4104–4165.
- [23] C. Henderson, S. Snelson, C^∞ smoothing for weak solutions of the inhomogeneous Landau equation, *Arch. Ration. Mech. Anal.* 236 (1) (2020) 113–143, <https://doi.org/10.1007/s00205-019-01465-7>.
- [24] C. Henderson, S. Snelson, A. Tarfulea, Local existence, lower mass bounds, and a new continuation criterion for the Landau equation, *J. Differ. Equ.* 266 (2–3) (2019) 1536–1577.
- [25] C. Imbert, C. Mouhot, L. Silvestre, Decay estimates for large velocities in the Boltzmann equation without cut-off, *J. Éc. Polytech.* 7 (2020) 143–184.
- [26] C. Imbert, L. Silvestre, The Schauder estimate for kinetic integral equations, Preprint, arXiv:1812.11870, 2018.
- [27] C. Imbert, L. Silvestre, Global regularity estimates for the Boltzmann equation without cut-off, Preprint, arXiv:1909.12729, 2019.
- [28] C. Imbert, L. Silvestre, Weak Harnack inequality for the Boltzmann equation without cut-off, *J. Eur. Math. Soc.* 2 (2) (2020) 507–592, <https://doi.org/10.4171/JEMS/928>.
- [29] A. Kiselev, F. Nazarov, R. Shterenberg, Blow up and regularity for fractal Burgers equation, *Dyn. Partial Differ. Equ.* 5 (3) (2008) 211–240.
- [30] E.M. Lifshitz, L.P. Pitaevskii, *Physical Kinetics*, 1st edition, Course of Theoretical Physics, vol. 10, Butterworth-Heinemann, 1981.
- [31] P.-L. Lions, On Boltzmann and Landau equations, *Philos. Trans. R. Soc. Lond., Ser. A* 346 (1679) (1994) 191–204.
- [32] J. Luk, Stability of vacuum for the Landau equation with moderately soft potentials, *Ann. PDE* 5 (1) (2019) 11.
- [33] Y. Morimoto, T. Yang, Local existence of polynomial decay solutions to the Boltzmann equation for soft potentials, *Anal. Appl. (Singap.)* 13 (6) (2015) 663–683.
- [34] C. Mouhot, De Giorgi-Nash-Moser and Hörmander theories: new interplay, in: *Proc. ICM 2018, Vol. III, 2020*, pp. 2467–2493, <https://doi.org/10.1142/11060>.
- [35] C. Mouhot, L. Neumann, Quantitative perturbative study of convergence to equilibrium for collisional kinetic models in the torus, *Nonlinearity* 19 (4) (2006) 969.
- [36] A. Pascucci, S. Polidoro, The Moser's iterative method for a class of ultraparabolic equations, *Commun. Contemp. Math.* 06 (03) (2004) 395–417.

- [37] L. Silvestre, A new regularization mechanism for the Boltzmann equation without cut-off, *Commun. Math. Phys.* 348 (1) (2016) 69–100.
- [38] L. Silvestre, Upper bounds for parabolic equations and the Landau equation, *J. Differ. Equ.* 262 (3) (2017) 3034–3055.
- [39] S. Snelson, Gaussian bounds for the inhomogeneous Landau equation with hard potentials, *SIAM J. Math. Anal.* 52 (2) (2020) 2081–2097, <https://doi.org/10.1137/19M1244275>.
- [40] R.M. Strain, Y. Guo, Almost exponential decay near Maxwellian, *Commun. Partial Differ. Equ.* 31 (3) (2006) 417–429.
- [41] R.M. Strain, Y. Guo, Exponential decay for soft potentials near Maxwellian, *Arch. Ration. Mech. Anal.* 187 (2) (2008) 287–339.
- [42] C. Villani, On the Cauchy problem for Landau equation: sequential stability, global existence, *Adv. Differ. Equ.* 1 (5) (1996) 793–816.
- [43] C. Villani, On the spatially homogeneous Landau equation for Maxwellian molecules, *Math. Models Methods Appl. Sci.* 08 (06) (1998) 957–983.
- [44] W. Wang, L. Zhang, The C^α regularity of weak solutions of ultraparabolic equations, *Discrete Contin. Dyn. Syst.* 29 (3) (2011) 1261–1275.
- [45] K.-C. Wu, Global in time estimates for the spatially homogeneous Landau equation with soft potentials, *J. Funct. Anal.* 266 (5) (2014) 3134–3155.