

OPTIMAL DISTRIBUTED AND TANGENTIAL BOUNDARY CONTROL FOR THE UNSTEADY STOCHASTIC STOKES EQUATIONS

PETER BENNER AND CHRISTOPH TRAUTWEIN*

Abstract. We consider a control problem constrained by the unsteady stochastic Stokes equations with nonhomogeneous boundary conditions in connected and bounded domains. In this paper, controls are defined inside the domain as well as on the boundary. Using a stochastic maximum principle, we derive necessary and sufficient optimality conditions such that explicit formulas for the optimal controls are derived. As a consequence, we are able to control the stochastic Stokes equations using distributed controls as well as boundary controls in a desired way.

Mathematics Subject Classification. 76D07, 93E20.

Received August 22, 2018. Accepted July 6, 2019.

1. INTRODUCTION

In this paper, we consider a linear quadratic control problem for the unsteady stochastic Stokes equations with linear multiplicative noise. Here, controls appear as distributed controls inside the domain as well as tangential controls on the boundary. Concerning fluid dynamics, noise enters the system due to structural vibration and other environmental effects, see [46] and the references therein. The aim is to find controls such that the velocity field is as close as possible to a given desired velocity field.

In the last decades, optimal control problems constrained by the Stokes equations have been studied extensively. Simultaneous distributed and boundary controls can be found in [21]. In [36, 43], discretization schemes for control problems are considered. For stochastic distributed controls, we refer to [5]. In [32], an optimal control problem for the Stokes equations is presented, where the viscosity satisfies a transport equation. A control problem motivated by Stokes flow in an artificial heart is considered in [13]. We extend this setting by allowing additional noise terms arising from random environmental effects. We address this complication of the problem by decomposing the external force into a control term and a noise term. Moreover, control problems are mainly considered for the case of distributed controls. Therefore, we include nonhomogeneous Dirichlet boundary conditions to involve tangential boundary controls.

Using stochastic processes, one can model structural vibration and other environmental effects affecting flow fields. This leads us immediately to the formulation of a stochastic partial differential equation, which belongs to the modern research areas of infinite dimensional stochastic analysis. Such equations can be interpreted as stochastic evolution equations and the solutions are defined in a generalized sense. There exist different

Keywords and phrases: Stochastic control, Stokes equations, Q-Wiener process, boundary control, maximum principle.

Max Planck Institute for Dynamics of Complex Technical Systems, Sandtorstraße 1, 39106 Magdeburg, Germany.

* Corresponding author: trautwein@mpi-magdeburg.mpg.de

approaches on how to deal with these solutions. In [8, 10, 22, 40], the concept of weak solutions is introduced, where the construction is mainly based on inner products. Using Gelfand triples, another approach is given by variational solutions, see [40, 44]. For problems containing a linear operator as the generator of a semigroup on a Hilbert space, one can use mild solutions, see [8, 10, 22]. Mild solutions are considered as solutions to integral equations of Itô-Volterra type containing a stochastic convolution. All of these concepts are based on a given probability space and they are called (probabilistic) strong solutions. Solutions constructing the probability space are called (probabilistic) weak solutions or martingale solutions, see [8, 10].

For deterministic linear systems, the theory of mild solutions is a well-established concept in order to involve nonhomogeneous boundary conditions, see, *e.g.*, [3, 7]. For that reason, we use this theory such that boundary control problems can be considered, which have an important meaning for applications in engineering. The construction of the solution is mainly based on an approach for the deterministic Stokes equations, see [42]. Although this approach is applicable for a broad class of boundary conditions, we restrict to the case of tangential boundary conditions. Therefore, we can reformulate the Stokes equations as an evolution equation in a suitable Hilbert space. Since we assume that the external force can be decomposed into a control term and a noise term, we obtain immediately a linear stochastic partial differential equation with distributed and Dirichlet boundary controls. We prove the existence and uniqueness of a mild solution being square integrable with respect to the time variable. In order to get a well defined solution, we need the definition of stochastic integrals with respect to adapted processes, see [22].

Due to the presence of a noise term, it might be the case that the velocity field of the fluid flow reveals an undesired behavior. Thus, it is reasonable to control the velocity field in a certain desired way, which leads us to a stochastic control problem in infinite dimensions defined by an optimization problem of a given cost functional constrained by the stochastic Stokes equations. The minimizer of the cost functional is then called an optimal control.

In general, there exist mainly two approaches to solve stochastic control problems:

- (i) stochastic maximum principle;
- (ii) dynamic programming.

Based on existence and uniqueness results for the solution of the stochastic system, one can often reformulate the control problem as a minimization problem on a set of admissible controls. For that reason, the idea of the stochastic maximum principle is to state necessary and sufficient optimality conditions the optimal control has to satisfy. Using the necessary optimality condition, one can derive an explicit formula of the optimal control based on the adjoint equation. For stochastic control problems in finite dimensions and a comparison to the corresponding deterministic setting, we refer to [39, 53]. The main difference to the deterministic setting is that the adjoint equation becomes a backward stochastic differential equation. For application to stochastic control problems in infinite dimensions, see [15, 26, 35]. In contrast to this methods, the dynamic programming principle considers the control problem at different initial times and initial states through the so called value function. This value function is the solution of a nonlinear partial differential equation given by the Hamilton-Jacobi-Bellman equation. The finite dimensional approach is presented in [14, 53]. For the infinite dimensional setting, we refer to [9, 11, 12]. Here, we are interested to provide a theory for solutions to a specific stochastic control problem such that it can be treated numerically. Since sufficient optimality conditions are useful to obtain the convergence to an optimal control, we use a stochastic maximum principle. This fact is already known for deterministic problems, see [28].

The concrete control problem considered in this paper is formulated as a tracking problem motivated by [2, 5, 31, 41, 43]. We derive a stochastic maximum principle to obtain first order optimality conditions, which are necessary and sufficient. To utilize these optimality conditions, a duality principle is required. In general, a duality principle gives a relation between forward and backward stochastic partial differential equations using an Itô product formula, which is not applicable for mild solutions. Hence, we approximate the mild solutions by strong solutions using an approach based on the resolvent operator, see [25, 30]. As a consequence, we obtain the duality principle for the approximating strong solutions and due to convergence results, the duality

principle holds also for the mild solutions. Based on the optimality conditions and the duality principle, we deduce formulas the optimal distributed control and the optimal boundary control have to satisfy.

The main contribution of this paper is to provide a mild solution to the stochastic Stokes equations with nonhomogeneous tangential boundary conditions. Moreover, we solve a control problem using a stochastic maximum principle such that optimal distributed controls and optimal boundary controls are derived.

The paper is organized as follows. In Section 3, we introduce common spaces and operators concerning the Stokes equations. Moreover, we discuss the deterministic Stokes equations with nonhomogeneous boundary conditions and we give an introduction to stochastic integrals with respect to adapted processes. In Section 4, we provide an existence and uniqueness result for the stochastic Stokes equations with nonhomogeneous boundary conditions. Section 5 addresses the control problem. We derive optimality conditions and a duality principle such that formulas for the optimal distributed control as well as the optimal boundary control are derived.

2. PRELIMINARIES

2.1. Functional analysis background

Throughout the paper, let $\mathcal{D} \subset \mathbb{R}^n$, $n \geq 2$, be a connected and bounded domain with C^2 boundary $\partial\mathcal{D}$. For $s \geq 0$, let $H^s(\mathcal{D})$ denote the usual Sobolev space and for $s > \frac{1}{2}$, let $H_0^s(\mathcal{D}) = \{y \in H^s(\mathcal{D}) : y = 0 \text{ on } \partial\mathcal{D}\}$. We introduce the following common spaces:

$$\begin{aligned} H &= \text{Completion of } \{y \in (C_0^\infty(\mathcal{D}))^n : \operatorname{div} y = 0 \text{ in } \mathcal{D}\} \text{ in } (L^2(\mathcal{D}))^n \\ &= \{y \in (L^2(\mathcal{D}))^n : \operatorname{div} y = 0 \text{ in } \mathcal{D}, y \cdot \eta = 0 \text{ on } \partial\mathcal{D}\}, \\ V &= \text{Completion of } \{y \in (C_0^\infty(\mathcal{D}))^n : \operatorname{div} y = 0 \text{ in } \mathcal{D}\} \text{ in } (H^1(\mathcal{D}))^n \\ &= \{y \in (H_0^1(\mathcal{D}))^n : \operatorname{div} y = 0 \text{ in } \mathcal{D}\}, \end{aligned}$$

where η denotes the unit outward normal to $\partial\mathcal{D}$. The spaces H and V equipped with the inner product of $(L^2(\mathcal{D}))^n$ and $(H_0^1(\mathcal{D}))^n$, respectively, become Hilbert spaces. Furthermore, we get the orthogonal Helmholtz decomposition

$$(L^2(\mathcal{D}))^n = H \oplus \{\nabla y : y \in H^1(\mathcal{D})\},$$

where \oplus denotes the direct sum. Then there exists an orthogonal projection $\Pi: (L^2(\mathcal{D}))^n \rightarrow H$, see [18]. Next, we define the Stokes Operator $A: D(A) \subset H \rightarrow H$ by $Ay = -\Pi\Delta y$ for every $y \in D(A)$ with domain $D(A) = (H^2(\mathcal{D}))^n \cap V$. The Stokes operator A is positive, self adjoint and has a bounded inverse. Moreover, the operator $-A$ is the infinitesimal generator of an analytic semigroup $(e^{-At})_{t \geq 0}$ such that $\|e^{-At}\|_{\mathcal{L}(H)} \leq 1$ for all $t \geq 0$. For more details, see [16, 23, 24, 50]. Hence, we can introduce fractional powers of the Stokes operator, see [38, 49, 50]. For $\alpha > 0$, we define

$$A^{-\alpha} = \frac{1}{\Gamma(\alpha)} \int_0^\infty t^{\alpha-1} e^{-At} dt, \quad (2.1)$$

where $\Gamma(\cdot)$ denotes the gamma function. The operator $A^{-\alpha}$ is linear, bounded and injective in H . Hence, we define for all $\alpha > 0$

$$A^\alpha = (A^{-\alpha})^{-1}.$$

Moreover, we set $A^0 = I$, where I is the identity operator in H . For $\alpha > 0$, the operator A^α is linear and closed in H with dense domain $D(A^\alpha) = R(A^{-\alpha})$, where $R(A^{-\alpha})$ denotes the range of $A^{-\alpha}$. Next, we provide some useful properties of fractional powers of the Stokes operator.

Lemma 2.1 (cf. [38], Sect. 2.6). *Let $A: D(A) \subset H \rightarrow H$ be the Stokes operator. Then*

- (i) for $\alpha, \beta \in \mathbb{R}$, we have $A^{\alpha+\beta}y = A^\alpha A^\beta y$ for every $y \in D(A^\gamma)$, where $\gamma = \max\{\alpha, \beta, \alpha + \beta\}$,
- (ii) $e^{-At}: H \rightarrow D(A^\alpha)$ for all $t > 0$ and $\alpha \geq 0$,
- (iii) we have $A^\alpha e^{-At}y = e^{-At}A^\alpha y$ for every $y \in D(A^\alpha)$ with $\alpha \in \mathbb{R}$,
- (iv) the operator $A^\alpha e^{-At}$ is bounded for all $t > 0$ and there exist constants $M_\alpha, \theta > 0$ such that

$$\|A^\alpha e^{-At}\|_{\mathcal{L}(H)} \leq M_\alpha t^{-\alpha} e^{-\theta t},$$

- (v) $0 \leq \beta \leq \alpha \leq 1$ implies $D(A^\alpha) \subset D(A^\beta)$ and there exists a constant $C > 0$ such that for every $y \in D(A^\alpha)$

$$\|A^\beta y\|_H \leq C \|A^\alpha y\|_H.$$

As a consequence of the previous lemma, we obtain that the space $D(A^\alpha)$ for all $\alpha \geq 0$ equipped with the inner product

$$\langle y, z \rangle_{D(A^\alpha)} = \langle A^\alpha y, A^\alpha z \rangle_H$$

for every $y, z \in D(A^\alpha)$ becomes a Hilbert space. Furthermore, we get the following result.

Lemma 2.2. *Let $A: D(A) \subset H \rightarrow H$ be the Stokes operator. Then the operator A^α is self adjoint for all $\alpha \in \mathbb{R}$.*

Proof. First, we show the claim for negative exponents. Recall the Stokes operator $-A$ is self adjoint. Hence, the semigroup $(e^{-At})_{t \geq 0}$ is self adjoint as well. By equation (2.1), we get for every $y, z \in \mathcal{H}$ and all $\alpha > 0$

$$\begin{aligned} \langle A^{-\alpha} y, z \rangle_{\mathcal{H}} &= \left\langle \frac{1}{\Gamma(\alpha)} \int_0^\infty t^{\alpha-1} S(t) y dt, z \right\rangle_{\mathcal{H}} = \frac{1}{\Gamma(\alpha)} \int_0^\infty t^{\alpha-1} \langle S(t) y, z \rangle_{\mathcal{H}} dt \\ &= \left\langle y, \frac{1}{\Gamma(\alpha)} \int_0^\infty t^{\alpha-1} S(t) z dt \right\rangle_{\mathcal{H}} = \langle y, A^{-\alpha} z \rangle_{\mathcal{H}}. \end{aligned} \quad (2.2)$$

Next, we show the claim for positive exponents. Using Lemma 2.1 (iv) and equation (2.2), we obtain for every $y, z \in D(A^\alpha)$ and all $\alpha > 0$

$$\langle A^\alpha y, z \rangle_{\mathcal{H}} = \langle A^\alpha y, A^{-\alpha} A^\alpha z \rangle_{\mathcal{H}} = \langle A^{-\alpha} A^\alpha y, A^\alpha z \rangle_{\mathcal{H}} = \langle y, A^\alpha z \rangle_{\mathcal{H}}.$$

For $\alpha = 0$, the claim is obvious. □

Next, we introduce the resolvent operator of $-A$ and we state some of its basic properties. For more details, see [38]. Let $\lambda \in \mathbb{R}$ be such that $\lambda I + A$ is invertible, i.e. $(\lambda I + A)^{-1}$ is a linear and bounded operator in the space H . Then the operator $R(\lambda; -A) = (\lambda I + A)^{-1}$ is called the resolvent operator. The operator $R(\lambda; -A)$ maps H into $D(A)$ and using the closed graph theorem, we can conclude that the operator $AR(\lambda; -A): H \rightarrow H$ is linear and bounded. Moreover, we have the following representation:

$$R(\lambda; -A) = \int_0^\infty e^{-\lambda r} e^{-Ar} dr. \quad (2.3)$$

For all $\lambda > 0$, we get $\|R(\lambda; -A)\|_{\mathcal{L}(H)} \leq 1/\lambda$ and since the semigroup $(e^{-At})_{t \geq 0}$ is self adjoint, the operator $R(\lambda; -A)$ is self adjoint as well. Let the operator $R(\lambda): H \rightarrow D(A)$ be defined by $R(\lambda) = \lambda R(\lambda; -A)$. Hence, we get for all $\lambda > 0$

$$\|R(\lambda)\|_{\mathcal{L}(H)} \leq 1. \quad (2.4)$$

By Lemma 2.1 (iii) and equation (2.3), we obtain for every $y \in D(A^\alpha)$ with $\alpha \in \mathbb{R}$

$$A^\alpha R(\lambda)y = R(\lambda)A^\alpha y. \quad (2.5)$$

Moreover, we have for every $y \in H$

$$\lim_{\lambda \rightarrow \infty} \|R(\lambda)y - y\|_H = 0. \quad (2.6)$$

If the domain \mathcal{D} is connected and bounded with C^∞ boundary $\partial\mathcal{D}$, then we can specify the domain of the operator A^α for $\alpha \in (0, 1)$ explicitly. Let $A_D: D(A_D) \subset (L^2(\mathcal{D}))^n \rightarrow (L^2(\mathcal{D}))^n$ be the Laplace operator with homogeneous Dirichlet boundary condition defined by $A_D y = -\Delta y$ for all $y \in D(A_D)$. The domain is given by $D(A_D) = (H_0^1(\mathcal{D}))^n \cap (H^2(\mathcal{D}))^n$. Then A_D is a positive and self adjoint operator and $-A_D$ is the infinitesimal generator of an analytic semigroup $(e^{-A_D t})_{t \geq 0}$ such that $\|e^{-A_D t}\|_{\mathcal{L}(H)} \leq 1$ for all $t \geq 0$. Hence, we can define fractional powers of the Laplace operator denoted by A_D^α for $\alpha \in \mathbb{R}$. We get the following result.

Proposition 2.3 ([16], Thm. 1.1). *Let the operator $A: D(A) \subset H \rightarrow H$ be the Stokes Operator and let the operator $A_D: D(A_D) \subset (L^2(\mathcal{D}))^n \rightarrow (L^2(\mathcal{D}))^n$ be the Laplace operator with homogeneous Dirichlet boundary condition. Then we have for any $\alpha \in (0, 1)$*

$$D(A^\alpha) = D(A_D^\alpha) \cap H.$$

The domain of the operator A_D^α can be determined explicitly for $\alpha \in (0, 1)$.

Proposition 2.4 (cf. [17], Thm. 1). *Let $A_D: D(A_D) \subset (L^2(\mathcal{D}))^n \rightarrow (L^2(\mathcal{D}))^n$ be the Laplace operator with homogeneous Dirichlet boundary condition. Then we have*

- (i) $D(A_D^\alpha) = (H^{2\alpha}(\mathcal{D}))^n$ for $\alpha \in (0, \frac{1}{4})$,
- (ii) $D(A_D^{1/4}) \subset (H^{1/2}(\mathcal{D}))^n$,
- (iii) $D(A_D^\alpha) = (H_0^{2\alpha}(\mathcal{D}))^n$ for $\alpha \in (\frac{1}{4}, \frac{3}{4})$,
- (iv) $D(A_D^{3/4}) \subset (H_0^{3/2}(\mathcal{D}))^n$,
- (v) $D(A_D^\alpha) = (H_0^{2\alpha}(\mathcal{D}))^n$ for $\alpha \in (\frac{3}{4}, 1)$.

2.2. The Stokes equations

In this section, we consider the deterministic Stokes equations with nonhomogeneous boundary conditions. Here, we restrict the problem to tangential boundary conditions. A general formulation can be found in [42].

Throughout the paper, let $T > 0$. We introduce the Stokes equations with nonhomogeneous boundary conditions:

$$\left\{ \begin{array}{ll} \frac{\partial}{\partial t} y(t, x) - \Delta y(t, x) + \nabla p(t, x) = f(t, x) & \text{in } (0, T) \times \mathcal{D}, \\ \operatorname{div} y(t, x) = 0 & \text{in } (0, T) \times \mathcal{D}, \\ y(t, x) = g(t, x) & \text{on } (0, T) \times \partial\mathcal{D}, \\ y(0, x) = \xi(x) & \text{in } \mathcal{D}, \end{array} \right. \quad (2.7)$$

where $y(t, x) \in \mathbb{R}^n$ denotes the velocity field with initial value $\xi(x) \in \mathbb{R}^n$, $p(t, x) \in \mathbb{R}$ describes the pressure of the fluid, and $f(t, x) \in \mathbb{R}^n$ is the external force. The boundary condition $g(t, x) \in \mathbb{R}^n$ is assumed to be tangential, *i.e.*

$$g(t, x) \cdot \eta(x) = 0 \quad \text{on } (0, T) \times \partial\mathcal{D},$$

where η denotes the unit outward normal to $\partial\mathcal{D}$. The goal is to reformulate system (2.7) as an evolution equation. We define the following spaces for $s \geq 0$:

$$\begin{aligned} V^s(\mathcal{D}) &= \{y \in (H^s(\mathcal{D}))^n : \operatorname{div} y = 0 \text{ in } \mathcal{D}, y \cdot \eta = 0 \text{ on } \partial\mathcal{D}\}, \\ V^s(\partial\mathcal{D}) &= \{y \in (H^s(\partial\mathcal{D}))^n : y \cdot \eta = 0 \text{ on } \partial\mathcal{D}\}. \end{aligned}$$

For $s < 0$, the space $V^s(\partial\mathcal{D})$ is the dual space of $V^{-s}(\partial\mathcal{D})$ with $V^0(\partial\mathcal{D})$ as pivot space. Moreover, let $H^s(\mathcal{D})/\mathbb{R}$ with $s \geq 0$ be the quotient space of $H^s(\mathcal{D})$ by \mathbb{R} , *i.e.* $H^s(\mathcal{D})/\mathbb{R} = \{y + c : y \in H^s(\mathcal{D}), c \in \mathbb{R}\}$. We set $\|y\|_{H^s(\mathcal{D})/\mathbb{R}} = \inf_{c \in \mathbb{R}} \|y + c\|_{H^s(\mathcal{D})}$ for every $y \in H^s(\mathcal{D})/\mathbb{R}$. The dual space is denoted by $(H^s(\mathcal{D})/\mathbb{R})'$ with $H^0(\mathcal{D})/\mathbb{R}$ as pivot space.

Next, let us consider the system

$$\begin{cases} -\Delta w + \nabla \pi = 0 & \text{and } \operatorname{div} w = 0 & \text{in } \mathcal{D}, \\ w = g & & \text{on } \partial\mathcal{D}. \end{cases} \quad (2.8)$$

In [19, 20, 42], existence and uniqueness results of a solution to system (2.8) are provided. This allows us to introduce the Dirichlet operators D and D_p defined by

$$Dg = w \quad \text{and} \quad D_p g = \pi,$$

where (w, π) is the solution of system (2.8). We get the following properties of these operators.

Proposition 2.5 (*cf.* [42], Cor. A.1). *The operator D is linear and continuous from $V^s(\partial\mathcal{D})$ into $V^{s+1/2}(\mathcal{D})$ for all $-\frac{1}{2} \leq s \leq \frac{3}{2}$. If $-\frac{1}{2} \leq s < \frac{1}{2}$, then the operator D_p is linear and continuous from $V^s(\partial\mathcal{D})$ into $(H^{1/2-s}(\mathcal{D})/\mathbb{R})'$, and if $\frac{1}{2} \leq s \leq \frac{3}{2}$, then D_p is a linear and continuous operator from $V^s(\partial\mathcal{D})$ into $H^{s-1/2}(\mathcal{D})/\mathbb{R}$.*

As a consequence of Proposition 2.3, Proposition 2.4 and Proposition 2.5, we get $D \in \mathcal{L}(V^0(\partial\mathcal{D}); D(A^\beta))$ for $\beta \in (0, \frac{1}{4})$. By the closed graph theorem, we have $A^\beta D \in \mathcal{L}(V^0(\partial\mathcal{D}); V^0(\mathcal{D}))$. Note that $V^0(\mathcal{D}) = H$. Furthermore, system (2.7) can be rewritten in the following form:

$$\begin{cases} \frac{d}{dt} y(t) = -Ay(t) + ADg(t) + \Pi f(t), \\ y(0) = \Pi \xi, \end{cases} \quad (2.9)$$

where the operators A and Π are introduced in Section 2.1. For the sake of simplicity, we assume $f(t), \xi \in H$ for $t \in [0, T]$. Hence, we obtain a linear evolution equation and the solution is given by

$$y(t) = e^{-At} \xi + \int_0^t A e^{-A(t-s)} Dg(s) ds + \int_0^t e^{-A(t-s)} f(s) ds.$$

For more details about linear evolution equations, see [3]. The following existence and uniqueness result is stated in [42] for more general boundary conditions and $f = 0$.

Theorem 2.6. *Let $g \in L^2([0, T]; V^0(\partial\mathcal{D}))$ and $f \in L^2([0, T]; H)$. If $\alpha \in [0, \frac{1}{4})$, then for any $\xi \in D(A^\alpha)$, there exists a unique solution $y \in L^2([0, T]; D(A^\alpha))$ of system (2.9) and the following estimate holds:*

$$\|y\|_{L^2([0, T]; D(A^\alpha))} \leq C^* (\|\xi\|_{D(A^\alpha)} + \|g\|_{L^2([0, T]; V^0(\partial\mathcal{D}))} + \|f\|_{L^2([0, T]; H)}),$$

where $C^* > 0$ is a constant.

2.3. Stochastic processes and the stochastic integral

In this section, we give a brief introduction to stochastic integrals, where the noise term is defined as a Hilbert space valued Wiener process. For more details, see [8].

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a complete probability space endowed with a filtration $(\mathcal{F}_t)_{t \in [0, T]}$ satisfying $\mathcal{F}_t = \bigcap_{s > t} \mathcal{F}_s$ for all $t \in [0, T]$. We denote by $\mathcal{L}(E)$ the space of linear and bounded operators defined on a separable Hilbert space E . Assume that $(W(t))_{t \in [0, T]}$ is an E -valued Q-Wiener process with covariance operator $Q \in \mathcal{L}(E)$, see ([8], Def. 4.2). There exists a unique operator $Q^{1/2} \in \mathcal{L}(E)$ such that $Q^{1/2} \circ Q^{1/2} = Q$. We denote by $\mathcal{L}_{(HS)}(Q^{1/2}(E); \mathcal{H})$ the space of Hilbert-Schmidt operators mapping from $Q^{1/2}(E)$ into another Hilbert space \mathcal{H} . Let $(\Phi(t))_{t \in [0, T]}$ be a predictable process with values in $\mathcal{L}_{(HS)}(Q^{1/2}(E); \mathcal{H})$ such that \mathbb{P} -a.s. $\int_0^T \|\Phi(t)\|_{\mathcal{L}_{(HS)}(Q^{1/2}(E); \mathcal{H})}^2 dt < \infty$. Then one can define the stochastic integral

$$\psi(t) = \int_0^t \Phi(s) dW(s)$$

for all $t \in [0, T]$ and we have

$$\mathbb{E} \|\psi(t)\|_{\mathcal{H}}^2 = \mathbb{E} \int_0^t \|\Phi(s)\|_{\mathcal{L}_{(HS)}(Q^{1/2}(E); \mathcal{H})}^2 ds. \quad (2.10)$$

The following proposition is useful when dealing with a closed operator $\mathcal{A}: D(\mathcal{A}) \subset \mathcal{H} \rightarrow \mathcal{H}$.

Proposition 2.7 (cf. [8], Prop. 4.30). *If $\Phi(t)y \in D(\mathcal{A})$ for every $y \in E$, all $t \in [0, T]$ and \mathbb{P} -almost surely,*

$$\mathbb{P} \left(\int_0^T \|\Phi(t)\|_{\mathcal{L}_{(HS)}(Q^{1/2}(E); \mathcal{H})}^2 dt \right) < \infty \quad \text{and} \quad \mathbb{P} \left(\int_0^T \|\mathcal{A}\Phi(t)\|_{\mathcal{L}_{(HS)}(Q^{1/2}(E); \mathcal{H})}^2 dt \right) < \infty,$$

then we have \mathbb{P} -a.s. $\int_0^T \Phi(t) dW(t) \in D(\mathcal{A})$ and

$$\mathcal{A} \int_0^T \Phi(t) dW(t) = \int_0^T \mathcal{A}\Phi(t) dW(t).$$

Next, we state a product formula for infinite dimensional stochastic processes, which we use to obtain a duality principle. The formula is an immediate consequence of the Itô formula, see ([8], Thm. 4.32).

Lemma 2.8. *For $i = 1, 2$, assume that X_i^0 are \mathcal{F}_0 -measurable \mathcal{H} -valued random variables, $(f_i(t))_{t \in [0, T]}$ are \mathcal{H} -valued predictable processes such that \mathbb{P} -a.s. $\int_0^T \|f_i(t)\|_{\mathcal{H}} dt < \infty$, and $(\Phi_i(t))_{t \in [0, T]}$ are $\mathcal{L}_{(HS)}(Q^{1/2}(E); \mathcal{H})$ -valued predictable processes such that \mathbb{P} -a.s. $\int_0^T \|\Phi_i(t)\|_{\mathcal{L}_{(HS)}(Q^{1/2}(E); \mathcal{H})}^2 dt < \infty$. For $i = 1, 2$, let the processes*

$(X_i(t))_{t \in [0, T]}$ satisfy for all $t \in [0, T]$ and \mathbb{P} -a.s.

$$X_i(t) = X_i^0 + \int_0^t f_i(s) ds + \int_0^t \Phi_i(s) dW(s).$$

Then we have for all $t \in [0, T]$ and \mathbb{P} -a.s.

$$\begin{aligned} \langle X_1(t), X_2(t) \rangle_{\mathcal{H}} &= \langle X_1^0, X_2^0 \rangle_{\mathcal{H}} + \int_0^t \left[\langle X_1(s), f_2(s) \rangle_{\mathcal{H}} + \langle X_2(s), f_1(s) \rangle_{\mathcal{H}} + \langle \Phi_1(s), \Phi_2(s) \rangle_{\mathcal{L}_{(HS)}(Q^{1/2}(E); \mathcal{H})} \right] ds \\ &\quad + \int_0^t \langle X_1(s), \Phi_2(s) dW(s) \rangle_{\mathcal{H}} + \int_0^t \langle X_2(s), \Phi_1(s) dW(s) \rangle_{\mathcal{H}}. \end{aligned}$$

Finally, we state a martingale representation theorem for Q-Wiener processes, which we use to construct solutions of backward SPDE's. Let $Q \in \mathcal{L}(E)$ be the covariance operator of a Q-Wiener process $(W(t))_{t \in [0, T]}$. Recall that the operator $Q \in \mathcal{L}(E)$ is a symmetric and nonnegative semidefinite such that $\text{Tr } Q < \infty$. Hence, there exists a complete orthonormal system $(e_k)_{k \in \mathbb{N}}$ in E and a bounded sequence of nonnegative real numbers $(\mu_k)_{k \in \mathbb{N}}$ such that $Qe_k = \mu_k e_k$ for each $k \in \mathbb{N}$. Then for arbitrary $t \in [0, T]$, a Q-Wiener process has the expansion

$$W(t) = \sum_{k=1}^{\infty} \sqrt{\mu_k} w_k(t) e_k,$$

where $(w_k(t))_{t \in [0, T]}$, $k \in \mathbb{N}$, are real valued mutually independent Brownian motions. The convergence is in $L^2(\Omega)$. Furthermore, we assume that the complete probability space $(\Omega, \mathcal{F}, \mathbb{P})$ is endowed with the filtration $\mathcal{F}_t = \sigma\{\bigcup_{k=1}^{\infty} \mathcal{F}_t^k\}$, where $\mathcal{F}_t^k = \sigma\{w_k(s) : 0 \leq s \leq t\}$ for $t \in [0, T]$ and we require that the σ -algebra \mathcal{F} satisfies $\mathcal{F} = \mathcal{F}_T$. Then we have the following martingale representation theorem.

Proposition 2.9 ([22], Thm. 2.5). *Let the process $(M(t))_{t \in [0, T]}$ be a continuous \mathcal{F}_t -martingale with values in \mathcal{H} such that $\mathbb{E}\|M(t)\|_{\mathcal{H}}^2 < \infty$ for all $t \in [0, T]$. Then there exists a unique predictable process $(\Phi(t))_{t \in [0, T]}$ with values in $\mathcal{L}_{(HS)}(Q^{1/2}(E); \mathcal{H})$ such that $\mathbb{E} \int_0^T \|\Phi(t)\|_{\mathcal{L}_{(HS)}(Q^{1/2}(E); \mathcal{H})}^2 dt < \infty$ and we have for all $t \in [0, T]$ and \mathbb{P} -a.s.*

$$M(t) = \mathbb{E}M(0) + \int_0^t \Phi(s) dW(s).$$

3. THE STOCHASTIC STOKES EQUATIONS

In this section, we consider the controlled stochastic Stokes equations. Here, controls appear as distributed controls inside the domain as well as tangential controls on the boundary.

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a complete probability space endowed with a filtration $(\mathcal{F}_t)_{t \in [0, T]}$ satisfying $\mathcal{F}_t = \bigcap_{s > t} \mathcal{F}_s$ for all $t \in [0, T]$. We assume that the external force $f(t)$ in equation (2.9) can be decomposed as the sum of a control term and a noise term dependent on the velocity field $y(t)$. Using the spaces and operators introduced

in Section 2.1 and Section 2.2, we obtain the stochastic Stokes equations:

$$\begin{cases} dy(t) = [-Ay(t) + Bu(t) + ADv(t)] dt + G(y(t))dW(t), \\ y(0) = \xi, \end{cases} \quad (3.1)$$

where the H -valued initial value ξ is assumed to be \mathcal{F}_0 -measurable and the process $(W(t))_{t \in [0, T]}$ is a Q-Wiener process with values in H and covariance operator $Q \in \mathcal{L}(H)$. The set of admissible distributed controls U contains all predictable processes $(u(t))_{t \in [0, T]}$ with values in H such that

$$\mathbb{E} \int_0^T \|u(t)\|_H^2 dt < \infty.$$

The space U equipped with the inner product of $L^2(\Omega; L^2([0, T]; H))$ becomes a Hilbert space. Similarly, the set of admissible boundary controls V contains all predictable processes $(v(t))_{t \in [0, T]}$ with values in $V^0(\partial\mathcal{D})$ such that

$$\mathbb{E} \int_0^T \|v(t)\|_{V^0(\partial\mathcal{D})}^2 dt < \infty.$$

The space V equipped with the inner product of $L^2(\Omega; L^2([0, T]; V^0(\partial\mathcal{D})))$ becomes a Hilbert space. The operators $B: H \rightarrow H$ and $G: H \rightarrow \mathcal{L}_{(HS)}(Q^{1/2}(H); H)$ are linear and bounded. Motivated by Section 2.2, we introduce the definition of a mild solution to system (3.1).

Definition 3.1. A predictable process $(y(t))_{t \in [0, T]}$ with values in $D(A^\alpha)$ is called a **mild solution of system (3.1)** if

$$\mathbb{E} \int_0^T \|y(t)\|_{D(A^\alpha)}^2 dt < \infty, \quad (3.2)$$

and we have for $t \in [0, T]$ and \mathbb{P} -a.s.

$$y(t) = e^{-At}\xi + \int_0^t e^{-A(t-s)}Bu(s)ds + \int_0^t Ae^{-A(t-s)}Dv(s)ds + \int_0^t e^{-A(t-s)}G(y(s))dW(s).$$

We get the following existence and uniqueness result.

Theorem 3.2. *Let the controls $u \in U$ and $v \in V$ be fixed. If $\alpha \in [0, \frac{1}{4})$, then for any $\xi \in L^2(\Omega; D(A^\alpha))$, there exists a unique mild solution $(y(t))_{t \in [0, T]}$ of system (3.1).*

Proof. For all $t_0, t_1 \in [0, T]$ with $t_0 < t_1$, let the space $\mathcal{Z}_{[t_0, t_1]}$ contain all predictable processes $(\tilde{y}(t))_{t \in [t_0, t_1]}$ with values in $D(A^\alpha)$ such that $\mathbb{E} \int_{t_0}^{t_1} \|\tilde{y}(t)\|_{D(A^\alpha)}^2 dt < \infty$. The space $\mathcal{Z}_{[t_0, t_1]}$ equipped with the inner product

$$\langle \tilde{y}_1, \tilde{y}_2 \rangle_{\mathcal{Z}_{[t_0, t_1]}}^2 = \mathbb{E} \int_{t_0}^{t_1} \langle \tilde{y}_1(t), \tilde{y}_2(t) \rangle_{D(A^\alpha)}^2 dt$$

for every $\tilde{y}_1, \tilde{y}_2 \in \mathcal{Z}_{[t_0, t_1]}$ becomes a Hilbert space. We define for $t \in [0, T]$ and \mathbb{P} -a.s.

$$\mathcal{J}(\tilde{y})(t) = e^{-At}\xi + \int_0^t e^{-A(t-s)}Bu(s)ds + \int_0^t Ae^{-A(t-s)}Dv(s)ds + \int_0^t e^{-A(t-s)}G(\tilde{y}(s))dW(s).$$

Let $T_1 \in (0, T]$ and let us denote by \mathcal{Z}_{T_1} the space $\mathcal{Z}_{[0, T_1]}$. First, we prove that \mathcal{J} maps \mathcal{Z}_{T_1} into itself. We define for $t \in [0, T_1]$ and \mathbb{P} -a.s.

$$\psi_1(t) = e^{-At}\xi + \int_0^t e^{-A(t-s)}Bu(s)ds, \quad \psi_2(t) = \int_0^t Ae^{-A(t-s)}Dv(s)ds, \quad \psi_3(\tilde{y})(t) = \int_0^t e^{-A(t-s)}G(\tilde{y}(s))dW(s).$$

Recall that $\|e^{-At}\|_{\mathcal{L}(H)} \leq 1$ for all $t \in [0, T]$ and $B: H \rightarrow H$ is bounded. Using Lemma 2.1 and the Cauchy-Schwarz inequality, the process $(\psi_1(t))_{t \in [0, T_1]}$ takes values in $D(A^\alpha)$ and there exists a constant $C_1 > 0$ such that

$$\begin{aligned} \mathbb{E} \int_0^{T_1} \|\psi_1(t)\|_{D(A^\alpha)}^2 dt &\leq 2\mathbb{E} \int_0^{T_1} \|e^{-At}A^\alpha\xi\|_H^2 dt + 2\mathbb{E} \int_0^{T_1} \left(\int_0^t \|A^\alpha e^{-A(t-s)}Bu(s)\|_H ds \right)^2 dt \\ &\leq 2T_1 \mathbb{E} \|\xi\|_{D(A^\alpha)}^2 + 2M_\alpha^2 \mathbb{E} \int_0^{T_1} \left(\int_0^t (t-s)^{-\alpha} \|Bu(s)\|_H ds \right)^2 dt \\ &\leq C_1 \left[\mathbb{E} \|\xi\|_{D(A^\alpha)}^2 + \mathbb{E} \int_0^{T_1} \|u(t)\|_H^2 dt \right]. \end{aligned}$$

Recall that $A^\beta D: V^0(\partial\mathcal{D}) \rightarrow H$ is bounded for all $\beta \in (0, \frac{1}{4})$. We choose β such that $\alpha < \beta$. By Lemma 2.1 and Young's inequality for convolutions, the process $(\psi_2(t))_{t \in [0, T_1]}$ takes values in $D(A^\alpha)$ and there exists a constant $C_2 > 0$ such that

$$\begin{aligned} \mathbb{E} \int_0^{T_1} \|\psi_2(t)\|_{D(A^\alpha)}^2 dt &\leq \mathbb{E} \int_0^{T_1} \left(\int_0^t \|A^{1+\alpha-\beta}e^{-A(t-s)}A^\beta Dv(s)\|_H ds \right)^2 dt \\ &\leq M_{1+\alpha-\beta}^2 \mathbb{E} \int_0^{T_1} \left(\int_0^t (t-s)^{-1-\alpha+\beta} \|A^\beta Dv(s)\|_H ds \right)^2 dt \\ &\leq M_{1+\alpha-\beta}^2 \left(\int_0^{T_1} t^{-1-\alpha+\beta} dt \right)^2 \mathbb{E} \int_0^{T_1} \|A^\beta Dv(t)\|_H^2 dt \\ &\leq C_2 \mathbb{E} \int_0^{T_1} \|v(t)\|_{V^0(\partial\mathcal{D})}^2 dt. \end{aligned}$$

Due to Lemma 2.1 and the fact that the operator $G: H \rightarrow \mathcal{L}_{(HS)}(Q^{1/2}(H); H)$ is bounded, one can verify the assumptions of Proposition 2.7 with $\mathcal{A} = A^\alpha$ and hence, the process $(\psi_3(\tilde{y})(t))_{t \in [0, T_1]}$ takes values in $D(A^\alpha)$.

Using Lemma 2.1, Fubini's theorem, the Itô isometry (2.10) and Young's inequality for convolutions, there exists a constant $C_3 > 0$ such that

$$\begin{aligned} \mathbb{E} \int_0^{T_1} \|\psi_3(\tilde{y})(t)\|_{D(A^\alpha)}^2 dt &= \int_0^{T_1} \mathbb{E} \int_0^t \left\| A^\alpha e^{-A(t-s)} G(\tilde{y}(s)) \right\|_{\mathcal{L}_{(HS)}(Q^{1/2}(H); H)}^2 ds dt \\ &\leq M_\alpha^2 \mathbb{E} \int_0^{T_1} \int_0^t (t-s)^{-2\alpha} \|G(\tilde{y}(s))\|_{\mathcal{L}_{(HS)}(Q^{1/2}(H); H)}^2 ds dt \\ &\leq C_3 T_1^{1-2\alpha} \mathbb{E} \int_0^{T_1} \|\tilde{y}(t)\|_{D(A^\alpha)}^2 dt. \end{aligned} \quad (3.3)$$

Hence, we can conclude that for fixed $\tilde{y} \in \mathcal{Z}_{T_1}$, the process $(\mathcal{J}(\tilde{y})(t))_{t \in [0, T_1]}$ takes values in $D(A^\alpha)$ such that $\mathbb{E} \int_0^{T_1} \|\mathcal{J}(\tilde{y})(t)\|_{D(A^\alpha)}^2 dt < \infty$. Obviously, the process $(\mathcal{J}(\tilde{y})(t))_{t \in [0, T_1]}$ is predictable. We conclude that \mathcal{J} maps \mathcal{Z}_{T_1} into itself.

Next, we show that \mathcal{J} is a contraction on \mathcal{Z}_{T_1} . Recall that the operator $G: H \rightarrow \mathcal{L}_{(HS)}(Q^{1/2}(H); H)$ is linear. Using inequality (3.3), we get for every $\tilde{y}_1, \tilde{y}_2 \in \mathcal{Z}_{T_1}$

$$\begin{aligned} \mathbb{E} \int_0^{T_1} \|\mathcal{J}(\tilde{y}_1)(t) - \mathcal{J}(\tilde{y}_2)(t)\|_{D(A^\alpha)}^2 dt &= \mathbb{E} \int_0^{T_1} \|\psi_3(\tilde{y}_1 - \tilde{y}_2)(t)\|_{D(A^\alpha)}^2 dt \\ &\leq C_3 T_1^{1-2\alpha} \mathbb{E} \int_0^{T_1} \|\tilde{y}_1(t) - \tilde{y}_2(t)\|_{D(A^\alpha)}^2 dt. \end{aligned}$$

We choose $T_1 \in (0, T]$ such that $C_3 T_1^{1-2\alpha} < 1$. Applying Banach fixed point theorem, we get a unique element $y \in \mathcal{Z}_{T_1}$ such that for $t \in [0, T_1]$ and \mathbb{P} -a.s. $y(t) = \mathcal{J}(y)(t)$.

Next, we consider for $t \in [T_1, T]$ and \mathbb{P} -a.s.

$$\mathcal{J}(\tilde{y})(t) = e^{-A(t-T_1)} y(T_1) + \int_{T_1}^t e^{-A(t-s)} B u(s) ds + \int_{T_1}^t A e^{-A(t-s)} D v(s) ds + \int_{T_1}^t e^{-A(t-s)} G(\tilde{y}(s)) dW(s).$$

Again, for a certain $T_2 \in [T_1, T]$, there exists a unique fixed point of \mathcal{J} on $\mathcal{Z}_{[T_1, T_2]}$. By continuing the method, we get the existence and uniqueness of a predictable process $(y(t))_{t \in [0, T]}$ satisfying for $t \in [0, T]$ and \mathbb{P} -a.s. $y(t) = \mathcal{J}(y)(t)$. \square

For the rest of the paper, we assume that $(y(t))_{t \in [0, T]}$ satisfies condition (3.2) with $\alpha = 0$ and we assume that the initial value $\xi \in L^2(\Omega; H)$ is fixed. To illustrate the dependence on the controls $u \in U$ and $v \in V$, let us denote by $(y(t; u, v))_{t \in [0, T]}$ the mild solution of system (3.1). Whenever the process is considered for fixed controls, we omit the dependency.

Next, we show some useful properties. Therefore, we need the following formulation of Gronwall's inequality for integrable functions. The result might be deduced from more general formulations, see [6, 51, 52].

Lemma 3.3. *Let $a, x: [0, T] \rightarrow [0, \infty)$ be integrable functions and let $b \geq 0$. If*

$$x(t) \leq a(t) + b \int_0^t x(s) ds$$

for all $t \in [0, T]$, then

$$x(t) \leq a(t) + b \int_0^t e^{b(t-s)} a(s) ds.$$

for all $t \in [0, T]$. If $a(t)$ is nondecreasing on $[0, T]$, then for $t \in [0, T]$

$$x(t) \leq a(t)e^{bt}.$$

Corollary 3.4. *Let $(y(t; u, v))_{t \in [0, T]}$ be the mild solution of system (3.1) corresponding to the controls $u \in U$ and $v \in V$. Then the process $(y(t; u, v))_{t \in [0, T]}$ is affine linear with respect to u and v , and we have for every $u_1, u_2 \in U$ and every $v_1, v_2 \in V$*

$$\mathbb{E} \int_0^T \|y(t; u_1, v_1) - y(t; u_2, v_2)\|_H^2 dt \leq \widehat{C} \left[\mathbb{E} \int_0^T \|u_1(t) - u_2(t)\|_H^2 dt + \mathbb{E} \int_0^T \|v_1(t) - v_2(t)\|_{V_0(\partial\mathcal{D})}^2 dt \right], \quad (3.4)$$

where $\widehat{C} > 0$ is a constant.

Proof. First, we show that $(y(t; u, v))_{t \in [0, T]}$ is affine linear with respect to $u \in U$. Let $\xi = 0$ and $v = 0$. Moreover, let $a, b \in \mathbb{R}$ and $u_1, u_2 \in U$. Recall that the operators $B: H \rightarrow H$ and $G: H \rightarrow \mathcal{L}_{(HS)}(Q^{1/2}(H); H)$ are linear and bounded. Moreover, we have $\|e^{-At}\|_{\mathcal{L}(H)} \leq 1$ for all $t \in [0, T]$. By definition, we get for $t \in [0, T]$ and \mathbb{P} -a.s.

$$y(t; a u_1 + b u_2, 0) - a y(t; u_1, 0) - b y(t; u_2, 0) = \int_0^t e^{-A(t-s)} G(y(s; a u_1 + b u_2, 0) - a y(s; u_1, 0) - b y(s; u_2, 0)) dW(s).$$

Using the Itô isometry (2.10) and Fubini's theorem, there exists a constant $C^* > 0$ such that for $t \in [0, T]$

$$\begin{aligned} & \mathbb{E} \|y(t; a u_1 + b u_2, 0) - a y(t; u_1, 0) - b y(t; u_2, 0)\|_H^2 \\ & \leq C^* \int_0^t \mathbb{E} \|y(s; a u_1 + b u_2, 0) - a y(s; u_1, 0) - b y(s; u_2, 0)\|_H^2 ds. \end{aligned}$$

By Lemma 3.3 and Fubini's theorem, we get $\mathbb{E} \int_0^T \|y(t; a u_1 + b u_2, 0) - a y(t; u_1, 0) - b y(t; u_2, 0)\|_H^2 dt = 0$. We obtain that $(y(t; u, 0))_{t \in [0, T]}$ with initial value $\xi = 0$ is linear with respect to $u \in U$. For arbitrary $\xi \in L^2(\Omega; H)$ and $v \in V$, we can conclude that $(y(t; u, v))_{t \in [0, T]}$ is affine linear with respect to $u \in U$. Similarly, we obtain that $(y(t; u, v))_{t \in [0, T]}$ is affine linear with respect to $v \in V$.

Next, we show that inequality (3.4) holds. Let $u_1, u_2 \in U$ and $v_1, v_2 \in V$. Recall that $A^\alpha D: V^0(\partial\mathcal{D}) \rightarrow H$ is linear and bounded for all $\alpha \in (0, \frac{1}{4})$. By definition and Lemma 2.1, we get for $t \in [0, T]$ and \mathbb{P} -a.s.

$$\begin{aligned} y(t; u_1, v_1) - y(t; u_2, v_2) &= \int_0^t e^{-A(t-s)} B[u_1(s) - u_2(s)] ds + \int_0^t A^{1-\alpha} e^{-A(t-s)} A^\alpha D[v_1(s) - v_2(s)] ds \\ &\quad + \int_0^t e^{-A(t-s)} G(y(s; u_1, v_1) - y(s; u_2, v_2)) dW(s). \end{aligned}$$

Due to the Itô isometry (2.10), Lemma 2.1 and Fubini's theorem, there exist constants $C_1, C_2, C_3 > 0$ such that for $t \in [0, T]$

$$\begin{aligned} \mathbb{E} \|y(t; u_1, v_1) - y(t; u_2, v_2)\|_H^2 &\leq C_1 \mathbb{E} \int_0^t \|u_1(s) - u_2(s)\|_H^2 ds + C_2 \mathbb{E} \left(\int_0^t (t-s)^{\alpha-1} \|v_1(s) - v_2(s)\|_{V^0(\partial\mathcal{D})} ds \right)^2 \\ &\quad + C_3 \int_0^t \mathbb{E} \|y(s; u_1, v_1) - y(s; u_2, v_2)\|_H^2 ds. \end{aligned}$$

Using Lemma 3.3, Fubini's theorem and Young's inequality for convolutions, we get for $t \in [0, T]$

$$\begin{aligned} &\mathbb{E} \|y(t; u_1, v_1) - y(t; u_2, v_2)\|_H^2 \\ &\leq C_1 \mathbb{E} \int_0^t \|u_1(s) - u_2(s)\|_H^2 ds + C_2 \mathbb{E} \left(\int_0^t (t-s)^{\alpha-1} \|v_1(s) - v_2(s)\|_{V^0(\partial\mathcal{D})} ds \right)^2 \\ &\quad + C_3 \int_0^t e^{C_3(t-s)} \left[C_1 \mathbb{E} \int_0^s \|u_1(r) - u_2(r)\|_H^2 dr + C_2 \mathbb{E} \left(\int_0^s (s-r)^{\alpha-1} \|v_1(r) - v_2(r)\|_{V^0(\partial\mathcal{D})} dr \right)^2 \right] ds \\ &\leq C_1 (1 + C_3 e^{C_3 t}) \mathbb{E} \int_0^t \|u_1(s) - u_2(s)\|_H^2 ds + C_2 \mathbb{E} \left(\int_0^t (t-s)^{\alpha-1} \|v_1(s) - v_2(s)\|_{V^0(\partial\mathcal{D})} ds \right)^2 \\ &\quad + \frac{C_2 C_3 e^{C_3 t} t^{2\alpha}}{\alpha^2} \mathbb{E} \int_0^t \|v_1(s) - v_2(s)\|_{V^0(\partial\mathcal{D})}^2 ds. \end{aligned}$$

By Fubini's theorem and Young's inequality for convolutions, there exists a constant $\widehat{C} > 0$ such that

$$\begin{aligned} &\mathbb{E} \int_0^T \|y(t; u_1, v_1) - y(t; u_2, v_2)\|_H^2 dt \\ &\leq \int_0^T \left[C_1 (1 + C_3 e^{C_3 t}) \mathbb{E} \int_0^t \|u_1(s) - u_2(s)\|_H^2 ds + C_3 e^{C_3 t} \frac{C_2 t^{2\alpha}}{\alpha^2} \mathbb{E} \int_0^t \|v_1(s) - v_2(s)\|_{V^0(\partial\mathcal{D})}^2 ds \right] dt \end{aligned}$$

$$\begin{aligned}
& + C_2 \mathbb{E} \int_0^T \left(\int_0^t (t-s)^{\alpha-1} \|v_1(s) - v_2(s)\|_{V^0(\partial\mathcal{D})} ds \right)^2 dt \\
& \leq \widehat{C} \left[\mathbb{E} \int_0^T \|u_1(t) - u_2(t)\|_H^2 dt + \mathbb{E} \int_0^T \|v_1(t) - v_2(t)\|_{V^0(\partial\mathcal{D})}^2 dt \right].
\end{aligned}$$

□

4. THE CONTROL PROBLEM

Due to the presence of the noise term in system (3.1), it might be the case that the velocity field of the fluid flow reveals an undesired behavior. Thus, it is reasonable to control the velocity field in a desired way. Motivated by [2, 5, 31, 41, 43], the control problem considered in this paper is formulated as a tracking-type problem arising typically in data assimilation. For general fluid flows, we also refer to [27, 34, 37, 48].

Let us introduce the following cost functional:

$$J(u, v) = \frac{1}{2} \mathbb{E} \int_0^T \|y(t; u, v) - y_d(t)\|_H^2 dt + \frac{\kappa_1}{2} \mathbb{E} \int_0^T \|u(t)\|_H^2 dt + \frac{\kappa_2}{2} \mathbb{E} \int_0^T \|v(t)\|_{V^0(\partial\mathcal{D})}^2 dt, \quad (4.1)$$

where $(y(t; u, v))_{t \in [0, T]}$ is the mild solution of system (3.1) corresponding to the controls $u \in U$ and $v \in V$. The function $y_d \in L^2([0, T]; H)$ is a given desired velocity field and $\kappa_1, \kappa_2 > 0$ are weights. The task is to find controls $\bar{u} \in U$ and $\bar{v} \in V$ such that

$$J(\bar{u}, \bar{v}) = \inf_{u \in U, v \in V} J(u, v).$$

The controls $\bar{u} \in U$ and $\bar{v} \in V$ are called optimal controls. Note that the control problem is formulated as an unbounded optimization problem constrained by a SPDE. The functional $J: U \times V \rightarrow \mathbb{R}$ given by equation (4.1) is continuous, coercive and strictly convex, which is a consequence of Corollary 3.4. Hence, we get the existence and uniqueness of optimal controls. For more details, we refer to [33, 54].

4.1. Necessary and sufficient optimality conditions

First, let us introduce the following systems:

$$\begin{cases} dz_1(t) = [-Az_1(t) + Bu(t)] dt + G(z_1(t))dW(t), \\ z_1(0) = 0, \end{cases} \quad (4.2)$$

$$\begin{cases} dz_2(t) = [-Az_2(t) + ADv(t)] dt + G(z_2(t))dW(t), \\ z_2(0) = 0, \end{cases} \quad (4.3)$$

where $u \in U$, $v \in V$ and $(W(t))_{t \in [0, T]}$ is a Q-Wiener process with values in H and covariance operator $Q \in \mathcal{L}(H)$. The operators A, B, D, G and the spaces U, V are introduced in Section 2 and Section 3, respectively. We consider the solutions of system (4.2) and system (4.3) in a mild sense according to Definition 3.1. Hence, existence and uniqueness results can be obtained similarly to Theorem 3.2. To illustrate the dependence on the controls $u \in U$ and $v \in V$, let us denote by $(z_1(t; u))_{t \in [0, T]}$ and $(z_2(t; v))_{t \in [0, T]}$ the mild solutions of system (4.2) and system

(4.3), respectively. Whenever these processes are considered for fixed controls, we omit the dependency. Similarly to Corollary 3.4, we get the following result.

Lemma 4.1. *Let $(z_1(t; u))_{t \in [0, T]}$ and $(z_2(t; v))_{t \in [0, T]}$ be the mild solutions of system (4.2) and system (4.3) corresponding to the controls $u \in U$ and $v \in V$, respectively. Then the process $(z_1(t; u))_{t \in [0, T]}$ is linear with respect to u and the process $(z_2(t; v))_{t \in [0, T]}$ is linear with respect to v . Moreover, we have for every $u_1, u_2 \in U$ and every $v_1, v_2 \in V$*

$$\begin{aligned} \mathbb{E} \int_0^T \|z_1(t; u_1) - z_1(t; u_2)\|_H^2 dt &\leq \widehat{C} \mathbb{E} \int_0^T \|u_1(t) - u_2(t)\|_H^2 dt, \\ \mathbb{E} \int_0^T \|z_2(t; v_1) - z_2(t; v_2)\|_H^2 dt &\leq \widehat{C} \mathbb{E} \int_0^T \|v_1(t) - v_2(t)\|_{V^0(\partial\mathcal{D})}^2 dt, \end{aligned}$$

where $\widehat{C} > 0$ is a constant.

Next, we calculate the Fréchet derivative of the mild solution to system (3.1). Let X, Y and Z be arbitrary Banach spaces. For a mapping $f: M_X \times M_Y \rightarrow Z$ with $M_X \subset X$, $M_Y \subset Y$ nonempty and open, the (partial) Fréchet derivative at $x \in M_X$ in direction $h \in X$ for fixed $y \in Y$ is denoted by $d_x f(x, y)[h]$. Analogously, the (partial) Fréchet derivative at $y \in M_Y$ in direction $h \in Y$ for fixed $x \in X$ is denoted by $d_y f(x, y)[h]$. We get the following result.

Theorem 4.2. *Let $(y(t; u, v))_{t \in [0, T]}$, $(z_1(t; u))_{t \in [0, T]}$ and $(z_2(t; v))_{t \in [0, T]}$ be the mild solutions of systems (3.1), (4.2) and (4.3) corresponding to the controls $u \in U$ and $v \in V$, respectively. Then the Fréchet derivative of $y(t; u, v)$ at $u \in U$ in direction $\tilde{u} \in U$ satisfies for fixed $v \in V$, $t \in [0, T]$ and \mathbb{P} -a.s.*

$$d_u y(t; u, v)[\tilde{u}] = z_1(t; \tilde{u}).$$

The Fréchet derivative of $y(t; u, v)$ at $v \in V$ in direction $\tilde{v} \in V$ satisfies for fixed $u \in U$, $t \in [0, T]$ and \mathbb{P} -a.s.

$$d_v y(t; u, v)[\tilde{v}] = z_2(t; \tilde{v}).$$

Proof. First, we calculate the Fréchet derivative of $y(t; u, v)$ at $u \in U$ in direction $\tilde{u} \in U$. Let $v \in V$ be fixed. Recall that the operators $B: H \rightarrow H$ and $G: H \rightarrow \mathcal{L}_{(HS)}(Q^{1/2}(H); H)$ are linear and bounded. Moreover, we have $\|e^{-At}\|_{\mathcal{L}(H)} \leq 1$ for all $t \in [0, T]$. Using the Itô isometry (2.10) and Fubini's theorem, there exists a constant $C^* > 0$ such that for $t \in [0, T]$

$$\begin{aligned} \mathbb{E} \|y(t; u + \tilde{u}, v) - y(t; u, v) - z_1(t; \tilde{u})\|_H^2 &= \mathbb{E} \left\| \int_0^t e^{-A(t-s)} G(y(s; u + \tilde{u}, v) - y(s; u, v) - z_1(s; \tilde{u})) dW(s) \right\|_H^2 \\ &\leq C^* \int_0^t \mathbb{E} \|y(s; u + \tilde{u}, v) - y(s; u, v) - z_1(s; \tilde{u})\|_H^2 ds. \end{aligned}$$

By Lemma 3.3 and Fubini's theorem, we get

$$\mathbb{E} \int_0^T \|y(t; u + \tilde{u}, v) - y(t; u, v) - z_1(t; \tilde{u})\|_H^2 dt = 0.$$

Hence, the Fréchet derivative of $y(t; u, v)$ at $u \in U$ in direction $\tilde{u} \in U$ satisfies for every $v \in V$, $t \in [0, T]$ and \mathbb{P} -a.s.

$$d_u y(t; u, v)[\tilde{u}] = z_1(t; \tilde{u}).$$

Due to Lemma 4.1, the operator $d_u y(t; u, v)$ is linear and bounded on U . Similarly, we obtain the Fréchet derivative of $y(t; u, v)$ at $v \in V$ in direction $\tilde{v} \in V$. \square

As a direct consequence of the previous theorem and the chain rule for Fréchet derivatives, we get the following result.

Theorem 4.3. *Let the functional $J: U \times V \rightarrow \mathbb{R}$ be defined by (4.1). Then the Fréchet derivative at $u \in U$ in direction $\tilde{u} \in U$ for fixed $v \in V$ satisfies*

$$d_u J(u, v)[\tilde{u}] = \mathbb{E} \int_0^T \langle y(t; u, v) - y_d(t), z_1(t; \tilde{u}) \rangle_H dt + \kappa_1 \mathbb{E} \int_0^T \langle u(t), \tilde{u}(t) \rangle_H dt,$$

where $(z_1(t; \tilde{u}))_{t \in [0, T]}$ is the mild solution of system (4.2) corresponding to the control $\tilde{u} \in U$. The Fréchet derivative at $v \in V$ in direction $\tilde{v} \in V$ for fixed $u \in U$ satisfies

$$d_v J(u, v)[\tilde{v}] = \mathbb{E} \int_0^T \langle y(t; u, v) - y_d(t), z_2(t; \tilde{v}) \rangle_H dt + \kappa_2 \mathbb{E} \int_0^T \langle v(t), \tilde{v}(t) \rangle_{V^0(\partial\mathcal{D})} dt,$$

where $(z_2(t; \tilde{v}))_{t \in [0, T]}$ is the mild solution of system (4.3) corresponding to the control $\tilde{v} \in V$.

As a result of the previous theorem and the fact that the cost functional $J: U \times V \rightarrow \mathbb{R}$ given by (4.1) is strictly convex, the optimal controls $\bar{u} \in U$ and $\bar{v} \in V$ satisfy the following necessary and sufficient optimality conditions:

$$d_u J(\bar{u}, \bar{v})[\tilde{u}] = 0, \tag{4.4}$$

$$d_v J(\bar{u}, \bar{v})[\tilde{v}] = 0 \tag{4.5}$$

for every $\tilde{u} \in U$ and every $\tilde{v} \in V$. For more details about optimality conditions of convex differentiable functionals, we refer to [33, 54]. Next, we use the optimality conditions (4.4) and (4.5) to derive explicit formulas for the optimal controls $\bar{u} \in U$ and $\bar{v} \in V$. Therefor, we need a duality principle, which gives us a relation between the Fréchet derivatives of the mild solution to system (3.1) and the adjoint equation, which is given by a backward SPDE.

4.2. The adjoint equation

We introduce the following backward SPDE:

$$\begin{cases} dz^*(t) = -[-Az^*(t) + G^*(\Phi(t)) + y(t) - y_d(t)]dt + \Phi(t)dW(t), \\ z^*(T) = 0, \end{cases} \tag{4.6}$$

where the process $(y(t))_{t \in [0, T]}$ is the mild solution of system (3.1) and $y_d \in L^2([0, T]; H)$ is the desired velocity field. The process $(W(t))_{t \in [0, T]}$ is a Q-Wiener process with values in H and covariance operator $Q \in \mathcal{L}(H)$ and the operator $G^*: \mathcal{L}_{(HS)}(Q^{1/2}(H); H) \rightarrow H$ is linear and bounded. A precise meaning is given in the following remark.

Remark 4.4. Since the operator $G: H \rightarrow \mathcal{L}_{(HS)}(Q^{1/2}(H); H)$ is linear and bounded, there exists a linear and bounded operator $G^*: \mathcal{L}_{(HS)}(Q^{1/2}(H); H) \rightarrow H$ satisfying for every $h \in H$ and every $\Phi \in \mathcal{L}_{(HS)}(Q^{1/2}(H); H)$

$$\langle G(h), \Phi \rangle_{\mathcal{L}_{(HS)}(Q^{1/2}(H); H)} = \langle h, G^*(\Phi) \rangle_H. \quad (4.7)$$

Definition 4.5. A pair of predictable processes $(z^*(t), \Phi(t))_{t \in [0, T]}$ with values in $H \times \mathcal{L}_{(HS)}(Q^{1/2}(H); H)$ is called a **mild solution of system (4.6)** if

$$\sup_{t \in [0, T]} \mathbb{E} \|z^*(t)\|_H^2 < \infty, \quad \mathbb{E} \int_0^T \|\Phi(t)\|_{\mathcal{L}_{(HS)}(Q^{1/2}(H); H)}^2 dt < \infty,$$

and we have for all $t \in [0, T]$ and \mathbb{P} -a.s.

$$z^*(t) = \int_t^T e^{-A(s-t)} G^*(\Phi(s)) ds + \int_t^T e^{-A(s-t)} (y(s) - y_d(s)) ds - \int_t^T e^{-A(s-t)} \Phi(s) dW(s).$$

An existence and uniqueness result is mainly based on the following lemma.

Lemma 4.6 ([29], Lem. 2.1). *Let $z \in L^2(\Omega; H)$ be \mathcal{F}_T -measurable and let $(f(t))_{t \in [0, T]}$ be a predictable process with values in H such that $\mathbb{E} \int_0^T \|f(t)\|_H^2 dt < \infty$. Then there exists a unique pair of predictable processes $(\varphi(t), \phi(t))_{t \in [0, T]}$ with values in $H \times \mathcal{L}_{(HS)}(Q^{1/2}(H); H)$ such that for all $t \in [0, T]$ and \mathbb{P} -a.s.*

$$\varphi(t) = e^{-A(T-t)} z + \int_t^T e^{-A(s-t)} f(s) ds - \int_t^T e^{-A(s-t)} \phi(s) dW(s).$$

Moreover, there exists a constant $c > 0$ such that for all $t \in [0, T]$

$$\mathbb{E} \|\varphi(t)\|_H^2 \leq c \left[\mathbb{E} \|z\|_H^2 + (T-t) \mathbb{E} \int_t^T \|f(s)\|_H^2 ds \right], \quad (4.8)$$

$$\mathbb{E} \int_t^T \|\phi(s)\|_{\mathcal{L}_{(HS)}(Q^{1/2}(H); H)}^2 ds \leq c \left[\mathbb{E} \|z\|_H^2 + (T-t) \mathbb{E} \int_t^T \|f(s)\|_H^2 ds \right]. \quad (4.9)$$

Existence and uniqueness results of mild solutions to backward SPDE's with cylindrical Wiener processes can be found in [29]. Similarly, we get the existence of a unique mild solution to system (4.6). Furthermore, note that the mild solution of system (3.1) depends on the controls $u \in U$ and $v \in V$. Thus, we get this property for the mild solution of system (4.6) as well. To illustrate the dependence on the controls $u \in U$ and $v \in V$, let us denote by $(z^*(t; u, v), \Phi(t; u, v))_{t \in [0, T]}$ the mild solution of system (4.6). Whenever these processes are considered for fixed controls, we omit the dependency. For the process $(z^*(t; u, v))_{t \in [0, T]}$, one can show another important regularity property. Therefore, we need a modification of Young's inequality for convolutions.

Lemma 4.7. *Let $f \in L^p([0, T])$ and $g \in L^q([0, T])$ be arbitrary. We set for $t \in [0, T]$*

$$h(t) = \int_t^T f(s-t)g(s)ds.$$

If $p, q, r \geq 1$ satisfy $\frac{1}{p} + \frac{1}{q} = \frac{1}{r} + 1$, then $h \in L^r([0, T])$ and

$$\|h\|_{L^r([0, T])} \leq \|f\|_{L^p([0, T])} \|g\|_{L^q([0, T])}.$$

Proof. The proof can be obtained similarly to the classical version of Young's inequality for convolutions, see ([4], Thm. 3.9.4). \square

Proposition 4.8. *Let $(z^*(t; u, v), \Phi(t; u, v))_{t \in [0, T]}$ be the mild solution of system (4.6) corresponding to the controls $u \in U$ and $v \in V$. Then $(z^*(t; u, v))_{t \in [0, T]}$ takes values in $D(A^\varepsilon)$ with $\varepsilon \in [0, 1)$ such that*

$$\mathbb{E} \int_0^T \|z^*(t; u, v)\|_{D(A^\varepsilon)}^2 dt < \infty.$$

Proof. Since $(z^*(t; u, v))_{t \in [0, T]}$ is predictable, we get for $t \in [0, T]$ and \mathbb{P} -a.s.

$$z^*(t; u, v) = \mathbb{E} \left[\int_t^T e^{-A(s-t)} G^*(\Phi(s; u, v)) ds + \int_t^T e^{-A(s-t)} (y(s; u, v) - y_d(s)) ds \middle| \mathcal{F}_t \right].$$

Recall that the operator $G^*: \mathcal{L}_{(HS)}(Q^{1/2}(H); H) \rightarrow H$ is bounded. Using Lemma 2.1 and Lemma 4.7, the process $(z^*(t))_{t \in [0, T]}$ takes values in $D(A^\varepsilon)$ with $\varepsilon \in [0, 1)$ and there exists a constant $C^* > 0$ such that

$$\begin{aligned} & \mathbb{E} \int_0^T \|z^*(t; u, v)\|_{D(A^\varepsilon)}^2 dt \\ & \leq 2 \mathbb{E} \int_0^T \left(\int_t^T \|A^\varepsilon e^{-A(s-t)} G^*(\Phi(s; u, v))\|_H ds \right)^2 dt + 2 \mathbb{E} \int_0^T \left(\int_t^T \|A^\varepsilon e^{-A(s-t)} (y(s; u, v) - y_d(s))\|_H^2 ds \right)^2 dt \\ & \leq 2M_\varepsilon^2 \mathbb{E} \int_0^T \left(\int_t^T (s-t)^{-\varepsilon} \|G^*(\Phi(s; u, v))\|_H ds \right)^2 dt + 2M_\varepsilon^2 \mathbb{E} \int_0^T \left(\int_t^T (s-t)^{-\varepsilon} \|y(s; u, v) - y_d(s)\|_H ds \right)^2 dt \\ & \leq C^* \left[\mathbb{E} \int_0^T \|\Phi(t; u, v)\|_{\mathcal{L}_{(HS)}(Q^{1/2}(H); H)}^2 dt + \mathbb{E} \int_0^T \|y(t; u, v)\|_H^2 dt + \int_0^T \|y_d(t)\|_H^2 dt \right]. \end{aligned}$$

\square

4.3. Approximation by a strong formulation

In general, a duality principle of solutions to forward and backward SPDE's can be obtained by applying an Itô product formula. This formula is not applicable to solutions in a mild sense. Hence, we need to approximate

the mild solutions of systems (4.2), (4.3) and (4.6) by strong formulations. One method is given by introducing the Yosida approximation of the operator A , see [8]. For applications regarding duality principles, see [15, 47]. However, we apply the method introduced in [25, 30]. The basic idea is to formulate a mild solution with values in $D(A)$ by using the resolvent operator $R(\lambda)$ introduced in Section 2.1. Thus, we get the required convergence results and the mild solutions coincide with the strong solutions. In this section, we omit the dependence on the controls for the sake of simplicity.

4.3.1. The forward equations

Here, we provide approximations of the mild solutions to system (4.2) and system (4.3). We introduce the following systems:

$$\begin{cases} dz_1(t, \lambda) = [-Az_1(t, \lambda) + R(\lambda)Bu(t)] dt + R(\lambda)G(R(\lambda)z_1(t, \lambda))dW(t), \\ z_1(0, \lambda) = 0, \end{cases} \quad (4.10)$$

$$\begin{cases} dz_2(t, \lambda) = [-Az_2(t, \lambda) + AR(\lambda)Dv(t)] dt + R(\lambda)G(R(\lambda)z_2(t, \lambda))dW(t), \\ z_2(0, \lambda) = 0, \end{cases} \quad (4.11)$$

where $\lambda > 0$, $u \in U$ and $v \in V$. The process $(W(t))_{t \in [0, T]}$ is a Q-Wiener process with values in H and covariance operator $Q \in \mathcal{L}(H)$. The operators $A, R(\lambda), B, D, G$ and the spaces U, V are introduced in Section 2 and Section 3, respectively.

Remark 4.9. Note that the approximation scheme provided in [25, 30] differs to the approximation scheme introduced by system (4.10) or system (4.11). Here, the additional operator $R(\lambda)$ arises in order to obtain a duality principle.

We consider the solutions of system (4.10) and system (4.11) in a mild sense according to Definition 3.1. Recall that the operators $R(\lambda)$ and $AR(\lambda)$ are linear and bounded on H . Hence, existence and uniqueness results of mild solutions to system (4.10) and system (4.11) can be obtained similarly to Theorem 3.2 for fixed $\lambda > 0$ such that the processes $(z_1(t, \lambda))_{t \in [0, T]}$ and $(z_2(t, \lambda))_{t \in [0, T]}$ take values in $D(A)$. In the following lemma, we state that the mild solutions of system (4.10) and system (4.11) also satisfy a strong formulation, which is an immediate consequence of ([30], Prop. 2.3).

Lemma 4.10. *Let $(z_1(t, \lambda))_{t \in [0, T]}$ and $(z_2(t, \lambda))_{t \in [0, T]}$ be the mild solutions of system (4.10) and system (4.11), respectively. Then we have for fixed $\lambda > 0$, $t \in [0, T]$ and \mathbb{P} -a.s.*

$$\begin{aligned} z_1(t, \lambda) &= \int_0^t (-A)z_1(s, \lambda) + R(\lambda)Bu(s)ds + \int_0^t R(\lambda)G(R(\lambda)z_1(s, \lambda))dW(s), \\ z_2(t, \lambda) &= \int_0^t (-A)z_2(s, \lambda) + AR(\lambda)Dv(s)ds + \int_0^t R(\lambda)G(R(\lambda)z_2(s, \lambda))dW(s). \end{aligned}$$

We have the following convergence results.

Lemma 4.11. (i) Let $(z_1(t))_{t \in [0, T]}$ and $(z_1(t, \lambda))_{t \in [0, T]}$ be the mild solutions of system (4.2) and system (4.10), respectively. Then we have

$$\lim_{\lambda \rightarrow \infty} \mathbb{E} \int_0^T \|z_1(t) - z_1(t, \lambda)\|_H^2 dt = 0.$$

(ii) Let $(z_2(t))_{t \in [0, T]}$ and $(z_2(t, \lambda))_{t \in [0, T]}$ be the mild solutions of system (4.3) and system (4.11), respectively. Then we have

$$\lim_{\lambda \rightarrow \infty} \mathbb{E} \int_0^T \|z_2(t) - z_2(t, \lambda)\|_H^2 dt = 0.$$

Proof. First, we show part (i). Let I be the identity operator in H . Recall that $G: H \rightarrow \mathcal{L}(H_S)(Q^{1/2}(H); H)$ is linear and bounded. By definition, we have for all $\lambda > 0$, $t \in [0, T]$ and \mathbb{P} -a.s.

$$\begin{aligned} & z_1(t) - z_1(t, \lambda) \\ &= \int_0^t e^{-A(t-s)} [I - R(\lambda)] Bu(s) ds + \int_0^t e^{-A(t-s)} G([I - R(\lambda)] z_1(s)) dW(s) \\ & \quad + \int_0^t e^{-A(t-s)} [I - R(\lambda)] G(R(\lambda) z_1(s)) dW(s) + \int_0^t e^{-A(t-s)} R(\lambda) G(R(\lambda) [z_1(s) - z_1(s, \lambda)]) dW(s). \end{aligned}$$

The remaining part of the proof can be obtained similarly to ([30], Lem. 3.1) using Lemma 3.3.

Next, we prove part (ii). By definition, we obtain for all $\lambda > 0$, $t \in [0, T]$ and \mathbb{P} -a.s.

$$\begin{aligned} & z_2(t) - z_2(t, \lambda) \\ &= \int_0^t A e^{-A(t-s)} [I - R(\lambda)] Dv(s) ds + \int_0^t e^{-A(t-s)} G([I - R(\lambda)] z_2(s)) dW(s) \\ & \quad + \int_0^t e^{-A(t-s)} [I - R(\lambda)] G(R(\lambda) z_2(s)) dW(s) + \int_0^t e^{-A(t-s)} R(\lambda) G(R(\lambda) [z_2(s) - z_2(s, \lambda)]) dW(s). \end{aligned}$$

Thus, we get for all $\lambda > 0$ and $t \in [0, T]$

$$\mathbb{E} \|z_2(t) - z_2(t, \lambda)\|_H^2 \leq 4\mathcal{I}_1(t, \lambda) + 4\mathcal{I}_2(t, \lambda) + 4\mathcal{I}_3(t, \lambda), \quad (4.12)$$

where

$$\begin{aligned} \mathcal{I}_1(t, \lambda) &= \mathbb{E} \left\| \int_0^t A e^{-A(t-s)} [I - R(\lambda)] Dv(s) ds \right\|_H^2, \\ \mathcal{I}_2(t, \lambda) &= \mathbb{E} \left\| \int_0^t e^{-A(t-s)} G([I - R(\lambda)] z_2(s)) dW(s) \right\|_H^2 + \mathbb{E} \left\| \int_0^t e^{-A(t-s)} [I - R(\lambda)] G(R(\lambda) z_2(s)) dW(s) \right\|_H^2, \end{aligned}$$

$$\mathcal{I}_3(t, \lambda) = \mathbb{E} \left\| \int_0^t e^{-A(t-s)} R(\lambda) G(R(\lambda)) [z_2(s) - z_2(s, \lambda)] dW(s) \right\|_H^2.$$

Recall that $D: V^0(\partial\mathcal{D}) \rightarrow D(A^\alpha)$ for all $\alpha \in (0, \frac{1}{4})$. Using Lemma 2.1, equation (2.5), Fubini's theorem and Young's inequality for convolutions, there exists a constant $C_1 > 0$ such that for all $\lambda > 0$ and all $t \in [0, T]$

$$\begin{aligned} \int_0^t \mathcal{I}_1(s, \lambda) ds &\leq \mathbb{E} \int_0^t \left(\int_0^s \left\| A^{1-\alpha} e^{-A(s-r)} [I - R(\lambda)] A^\alpha Dv(r) \right\|_H dr \right)^2 ds \\ &\leq C_1 \mathbb{E} \int_0^T \|[I - R(\lambda)] A^\alpha Dv(t)\|_H^2 dt. \end{aligned} \quad (4.13)$$

Recall that $\|e^{-At}\|_{\mathcal{L}(H)} \leq 1$ for all $t \in [0, T]$. Due to the Itô isometry (2.10) and Fubini's theorem, there exists a constant $C_2 > 0$ such that for all $\lambda > 0$ and all $t \in [0, T]$

$$\begin{aligned} \int_0^t \mathcal{I}_2(s, \lambda) ds &\leq \int_0^t \mathbb{E} \int_0^s \left\| e^{-A(s-r)} G([I - R(\lambda)] z_2(r)) \right\|_{\mathcal{L}(HS)(Q^{1/2}(H); H)}^2 dr ds \\ &\quad + \int_0^t \mathbb{E} \int_0^s \left\| e^{-A(s-r)} [I - R(\lambda)] G(R(\lambda) z_2(r)) \right\|_{\mathcal{L}(HS)(Q^{1/2}(H); H)}^2 dr ds \\ &\leq C_2 \left[\mathbb{E} \int_0^T \|[I - R(\lambda)] z_2(t)\|_H^2 dt + \mathbb{E} \int_0^T \|[I - R(\lambda)] G(R(\lambda) z_2(t))\|_{\mathcal{L}(HS)(Q^{1/2}(H); H)}^2 dt \right]. \end{aligned} \quad (4.14)$$

By the Itô isometry (2.10), inequality (2.4) and Fubini's theorem, there exists a constant $C_3 > 0$ such that for all $\lambda > 0$ and all $t \in [0, T]$

$$\mathcal{I}_3(t, \lambda) \leq C_3 \int_0^t \mathbb{E} \|z_2(s) - z_2(s, \lambda)\|_H^2 ds.$$

Due to inequality (4.12), we get for all $\lambda > 0$ and $t \in [0, T]$

$$\mathbb{E} \|z_2(t) - z_2(t, \lambda)\|_H^2 \leq 4 \mathcal{I}_1(t, \lambda) + 4 \mathcal{I}_2(t, \lambda) + 4C_3 \int_0^t \mathbb{E} \|z_2(s) - z_2(s, \lambda)\|_H^2 ds.$$

Applying Lemma 3.3, we obtain for all $\lambda > 0$ and $t \in [0, T]$

$$\mathbb{E} \|z_2(t) - z_2(t, \lambda)\|_H^2 \leq 4 \mathcal{I}_1(t, \lambda) + 4 \mathcal{I}_2(t, \lambda) + 16C_3 e^{4C_3 t} \int_0^t [\mathcal{I}_1(s, \lambda) + \mathcal{I}_2(s, \lambda)] ds.$$

Using Fubini's theorem, inequality (4.13) and inequality (4.14), there exists a constant $C^* > 0$ such that for all $\lambda > 0$

$$\begin{aligned} \mathbb{E} \int_0^T \|z_2(t) - z_2(t, \lambda)\|_H^2 dt &\leq C^* \mathbb{E} \int_0^T \|[I - R(\lambda)]A^\alpha Dv(t)\|_H^2 dt + C^* \mathbb{E} \int_0^T \|[I - R(\lambda)]z_2(t)\|_H^2 dt \\ &\quad + C^* \mathbb{E} \int_0^T \|[I - R(\lambda)]G(R(\lambda)z_2(t))\|_{\mathcal{L}(HS)(Q^{1/2}(H);H)}^2 dt. \end{aligned}$$

By equation (2.6) and Lebesgue's dominated convergence theorem ([4], Thm. 2.8.1), we can infer

$$\lim_{\lambda \rightarrow \infty} \mathbb{E} \int_0^T \|z_2(t) - z_2(t, \lambda)\|_H^2 dt = 0.$$

□

4.3.2. The backward equation

Here we provide an approximation of the mild solution to system (4.6). We introduce the following backward SPDE:

$$\begin{cases} dz^*(t, \lambda) = -[-Az^*(t, \lambda) + R(\lambda)G^*(R(\lambda)\Phi(t, \lambda)) + R(\lambda)(y(t) - y_d(t))]dt + \Phi(t, \lambda)dW(t), \\ z^*(T, \lambda) = 0, \end{cases} \quad (4.15)$$

where $\lambda > 0$. The process $(y(t))_{t \in [0, T]}$ is the mild solution of system (3.1) and $(W(t))_{t \in [0, T]}$ is a Q-Wiener process with values in H and covariance operator $Q \in \mathcal{L}(H)$. The function $y_d \in L^2([0, T]; H)$ is the desired velocity field. The operators $A, R(\lambda), G^*$ are introduced in Section 2.1 and Section 4.2, respectively. We consider the solution of system (4.15) in a mild sense according to Definition 3.1. Recall that the operators $R(\lambda)$ and $AR(\lambda)$ are linear and bounded in H . Hence, existence and uniqueness results of the mild solution to system (4.15) can be obtained similarly to [29] such that the pair of processes $(z^*(t, \lambda), \Phi(t, \lambda))_{t \in [0, T]}$ takes values in $D(A) \times \mathcal{L}(HS)(Q^{1/2}(H); H)$. In the following lemma, we state that the mild solution of system (4.15) also satisfies a strong formulation, which is an immediate consequence of Theorem 4.2 from [1].

Lemma 4.12. *Let the pair of stochastic processes $(z^*(t, \lambda), \Phi(t, \lambda))_{t \in [0, T]}$ be the mild solution of system (4.15). Then we have for fixed $\lambda > 0$, all $t \in [0, T]$ and \mathbb{P} -a.s.*

$$z^*(t, \lambda) = \int_t^T (-A)z^*(s, \lambda) + R(\lambda)G^*(R(\lambda)\Phi(s, \lambda)) + R(\lambda)(y(s) - y_d(s)) ds - \int_t^T \Phi(s, \lambda)dW(s).$$

We have the following convergence results.

Lemma 4.13. *Let $(z^*(t), \Phi(t))_{t \in [0, T]}$ and $(z^*(t, \lambda), \Phi(t, \lambda))_{t \in [0, T]}$ be the mild solutions of system (4.6) and system (4.15), respectively. Then we have*

$$\lim_{\lambda \rightarrow \infty} \sup_{t \in [0, T]} \mathbb{E} \|z^*(t) - z^*(t, \lambda)\|_H^2 = 0,$$

$$\lim_{\lambda \rightarrow \infty} \mathbb{E} \int_0^T \|\Phi(t) - \Phi(t, \lambda)\|_{\mathcal{L}(HS)(Q^{1/2}(H); H)}^2 dt = 0.$$

Proof. Recall that the operator $G^* : \mathcal{L}(HS)(Q^{1/2}(H); H) \rightarrow H$ is linear and bounded. By definition, we get for all $\lambda > 0$, all $t \in [0, T]$ and \mathbb{P} -a.s.

$$\begin{aligned} z^*(t) - z^*(t, \lambda) &= \int_t^T e^{-A(s-t)} G^*([I - R(\lambda)]\Phi(s)) ds + \int_t^T e^{-A(s-t)} [I - R(\lambda)] G^*(R(\lambda)\Phi(s)) ds \\ &+ \int_t^T e^{-A(s-t)} R(\lambda) G^*(R(\lambda)[\Phi(s) - \Phi(s, \lambda)]) ds + \int_t^T e^{-A(s-t)} [I - R(\lambda)] (y(s) - y_d(s)) ds \\ &- \int_t^T e^{-A(s-t)} [\Phi(s) - \Phi(s, \lambda)] dW(s), \end{aligned}$$

where I is the identity operator on H . Note that the assumptions of Lemma 4.6 are fulfilled. Thus, inequalities (4.8) and (4.9) hold. Let $T_1 \in [0, T]$. We obtain for all $\lambda > 0$

$$\sup_{t \in [T_1, T]} \mathbb{E} \|z^*(t) - z^*(t, \lambda)\|_H^2 \leq 4c(T - T_1) [\mathcal{I}_1(\lambda) + \mathcal{I}_2(\lambda)], \quad (4.16)$$

$$\mathbb{E} \int_{T_1}^T \|\Phi(t) - \Phi(t, \lambda)\|_{\mathcal{L}(HS)(Q^{1/2}(H); H)}^2 dt \leq 4c(T - T_1) [\mathcal{I}_1(\lambda) + \mathcal{I}_2(\lambda)], \quad (4.17)$$

where

$$\begin{aligned} \mathcal{I}_1(\lambda) &= \mathbb{E} \int_{T_1}^T [\|G^*([I - R(\lambda)]\Phi(t))\|_H^2 + \|[I - R(\lambda)]G^*(R(\lambda)\Phi(t))\|_H^2 + \|[I - R(\lambda)](y(t) - y_d(t))\|_H^2] dt, \\ \mathcal{I}_2(\lambda) &= \mathbb{E} \int_{T_1}^T \|R(\lambda)G^*(R(\lambda)[\Phi(t) - \Phi(t, \lambda)])\|_H^2 dt. \end{aligned}$$

Using equation (2.6) and the Lebesgue's dominated convergence theorem ([4], Thm. 2.8.1), we can conclude

$$\lim_{\lambda \rightarrow \infty} \mathcal{I}_1(\lambda) = 0. \quad (4.18)$$

By inequality (2.4), there exists a constant $C^* > 0$ such that for all $\lambda > 0$

$$\mathcal{I}_2(\lambda) \leq C^* \mathbb{E} \int_{T_1}^T \|\Phi(t) - \Phi(t, \lambda)\|_{\mathcal{L}(HS)(Q^{1/2}(H); H)}^2 dt. \quad (4.19)$$

Due to inequality (4.17) and inequality (4.19), we get for all $\lambda > 0$

$$\mathbb{E} \int_{T_1}^T \|\Phi(t) - \Phi(t, \lambda)\|_{\mathcal{L}_{(HS)}(Q^{1/2}(H); H)}^2 dt \leq 4c(T - T_1) \mathcal{I}_1(\lambda) + 4cC^*(T - T_1) \mathbb{E} \int_{T_1}^T \|\Phi(t) - \Phi(t, \lambda)\|_{\mathcal{L}_{(HS)}(Q^{1/2}(H); H)}^2 dt.$$

We chose $T_1 \in [0, T)$ such that $4cC^*(T - T_1) < 1$. Thus, we have for all $\lambda > 0$

$$\mathbb{E} \int_{T_1}^T \|\Phi(t) - \Phi(t, \lambda)\|_{\mathcal{L}_{(HS)}(Q^{1/2}(H); H)}^2 dt \leq \frac{4c(T - T_1) \mathcal{I}_1(\lambda)}{1 - 4cC^*(T - T_1)}.$$

Due to equation (4.18), we can conclude

$$\lim_{\lambda \rightarrow \infty} \mathbb{E} \int_{T_1}^T \|\Phi(t) - \Phi(t, \lambda)\|_{\mathcal{L}_{(HS)}(Q^{1/2}(H); H)}^2 dt = 0. \quad (4.20)$$

Using inequality (4.16), inequality (4.19), equation (4.18) and equation (4.20), we have

$$\lim_{\lambda \rightarrow \infty} \sup_{t \in [T_1, T]} \mathbb{E} \|z^*(t) - z^*(t, \lambda)\|_H^2 = 0.$$

By definition, we get for all $\lambda > 0$, all $t \in [0, T_1]$ and \mathbb{P} -a.s.

$$\begin{aligned} z^*(t) - z^*(t, \lambda) &= e^{-A(T_1-t)} [z^*(T_1) - z^*(T_1, \lambda)] + \int_t^{T_1} e^{-A(s-t)} [G^*(\Phi(s)) - R(\lambda)G^*(R(\lambda)\Phi(s, \lambda))] ds \\ &\quad + \int_t^{T_1} e^{-A(s-t)} [I - R(\lambda)] (y(s) - y_d(s)) ds - \int_t^{T_1} e^{-A(s-t)} [\Phi(s) - \Phi(s, \lambda)] dW(s). \end{aligned}$$

Again, we find $T_2 \in [0, T_1]$ such that

$$\begin{aligned} \lim_{\lambda \rightarrow \infty} \sup_{t \in [T_2, T_1]} \mathbb{E} \|z^*(t) - z^*(t, \lambda)\|_H^2 &= 0, \\ \lim_{\lambda \rightarrow \infty} \mathbb{E} \int_{T_2}^{T_1} \|\Phi(t) - \Phi(t, \lambda)\|_{\mathcal{L}_{(HS)}(Q^{1/2}(H); H)}^2 dt &= 0. \end{aligned}$$

By continuing the method, we obtain the result. \square

5. MAIN RESULTS

In this section, we derive a duality principle. Since we formulated a control problem with simultaneous distributed controls and boundary controls, we obtain two equations. The first equation gives us a relation between the mild solution of system (4.2) and the mild solution of the adjoint equation (4.6). The second equation provides a relation between the mild solution of system (4.3) and the mild solution of the adjoint

equation (4.6). As a consequence, we can utilize the necessary and sufficient optimality conditions (4.4) and (4.5) to deduce explicit formulas for the optimal controls.

5.1. Duality principle

Based on the results provided in the previous sections, we are able to show the following duality principle.

Theorem 5.1. *Let $(y(t; u, v))_{t \in [0, T]}$ and $(z^*(t; u, v), \Phi(t; u, v))_{t \in [0, T]}$ be the mild solutions of system (3.1) and system (4.6) corresponding to the distributed control $u \in U$ and the boundary control $v \in V$, respectively. Moreover, let $(z_1(t; \tilde{u}))_{t \in [0, T]}$ and $(z_2(t; \tilde{v}))_{t \in [0, T]}$ be the mild solutions of system (4.2) and system (4.3) corresponding to the controls $\tilde{u} \in U$ and $\tilde{v} \in V$, respectively. Then we have for all $\alpha \in (0, \frac{1}{4})$*

$$\mathbb{E} \int_0^T \langle y(t; u, v) - y_d(t), z_1(t; \tilde{u}) \rangle_H dt = \mathbb{E} \int_0^T \langle z^*(t; u, v), B\tilde{u}(t) \rangle_H dt, \quad (5.1)$$

$$\mathbb{E} \int_0^T \langle y(t; u, v) - y_d(t), z_2(t; \tilde{v}) \rangle_H dt = \mathbb{E} \int_0^T \langle A^{1-\alpha} z^*(t; u, v), A^\alpha D\tilde{v}(t) \rangle_H dt. \quad (5.2)$$

Proof. For the sake of simplicity, we omit the dependence on the controls. First, we prove the result for the approximations derived in Section 4.3. Let $(z_1(t, \lambda))_{t \in [0, T]}$ and $(z_2(t, \lambda))_{t \in [0, T]}$ be the mild solutions of system (4.10) and system (4.11), respectively. Using Lemma 4.10, we have for all $\lambda > 0$, $t \in [0, T]$ and \mathbb{P} -a.s.

$$z_1(t, \lambda) = \int_0^t (-A)z_1(s, \lambda) + R(\lambda)B\tilde{u}(s)ds + \int_0^t R(\lambda)G(R(\lambda)z_1(s, \lambda))dW(s), \quad (5.3)$$

$$z_2(t, \lambda) = \int_0^t (-A)z_2(s, \lambda) + AR(\lambda)D\tilde{v}(s)ds + \int_0^t R(\lambda)G(R(\lambda)z_2(s, \lambda))dW(s). \quad (5.4)$$

Next, let the pair of stochastic processes $(z^*(t, \lambda), \Phi(t, \lambda))_{t \in [0, T]}$ be the mild solution of system (4.15). Due to Lemma 4.12, we get for all $\lambda > 0$, all $t \in [0, T]$ and \mathbb{P} -a.s.

$$z^*(t, \lambda) = \int_t^T (-A)z^*(s, \lambda) + R(\lambda)G^*(R(\lambda)\Phi(s, \lambda)) + R(\lambda)(y(s) - y_d(s)) ds - \int_t^T \Phi(s, \lambda)dW(s). \quad (5.5)$$

By definition, the process $(z^*(t, \lambda))_{t \in [0, T]}$ is predictable. Hence, we have for all $\lambda > 0$, all $t \in [0, T]$ and \mathbb{P} -a.s.

$$z^*(t, \lambda) = \mathbb{E} \left[\int_0^T (-A)z^*(s, \lambda) + R(\lambda)G^*(R(\lambda)\Phi(s, \lambda)) + R(\lambda)(y(s) - y_d(s)) ds \middle| \mathcal{F}_t \right] - \int_0^t (-A)z^*(s, \lambda) + R(\lambda)G^*(R(\lambda)\Phi(s, \lambda)) + R(\lambda)(y(s) - y_d(s)) ds.$$

By the martingale representation theorem given by Proposition 2.9 with $(M(t))_{t \in [0, T]}$ satisfying for all $t \in [0, T]$ and \mathbb{P} -a.s.

$$M(t) = \mathbb{E} \left[\int_0^T (-A)z^*(s, \lambda) + R(\lambda)G^*(R(\lambda)\Phi(s, \lambda)) + R(\lambda)(y(s) - y_d(s)) ds \middle| \mathcal{F}_t \right],$$

there exists a unique predictable process $(\Psi(t, \lambda))_{t \in [0, T]}$ with values in $\mathcal{L}_{(HS)}(Q^{1/2}(H); H)$ such that for all $\lambda > 0$, all $t \in [0, T]$ and \mathbb{P} -a.s.

$$\begin{aligned} z^*(t, \lambda) = & \mathbb{E} \left[\int_0^T (-A)z^*(s, \lambda) + R(\lambda)G^*(R(\lambda)\Phi(s, \lambda)) + R(\lambda)(y(s) - y_d(s)) ds \right] \\ & - \int_0^t (-A)z^*(s, \lambda) + R(\lambda)G^*(R(\lambda)\Phi(s, \lambda)) + R(\lambda)(y(s) - y_d(s)) ds + \int_0^t \Psi(s, \lambda) dW(s). \end{aligned} \quad (5.6)$$

Since the pair $(z^*(t, \lambda), \Phi(t, \lambda))_{t \in [0, T]}$ satisfies equation (5.5) uniquely, we can conclude $\Psi(t, \lambda) = \Phi(t, \lambda)$ for all $\lambda > 0$, almost all $t \in [0, T]$ and \mathbb{P} -almost surely. Applying the Itô product formula given by Lemma 2.8 to equation (5.3) and equation (5.6), we get for all $\lambda > 0$, all $t \in [0, T]$ and \mathbb{P} -a.s.

$$\langle z_1(t, \lambda), z^*(t, \lambda) \rangle_H = \mathcal{I}_1(t, \lambda) + \mathcal{I}_2(t, \lambda) + \mathcal{I}_3(t, \lambda) + \mathcal{I}_4(t, \lambda),$$

where

$$\begin{aligned} \mathcal{I}_1(t, \lambda) &= \int_0^t [\langle z_1(s, \lambda), Az^*(s, \lambda) \rangle_H - \langle z^*(s, \lambda), Az_1(s, \lambda) \rangle_H] ds, \\ \mathcal{I}_2(t, \lambda) &= \int_0^t [\langle R(\lambda)G(R(\lambda)z_1(s, \lambda)), \Phi(s, \lambda) \rangle_{\mathcal{L}_{(HS)}(Q^{1/2}(H), H)} - \langle z_1(s, \lambda), R(\lambda)G^*(R(\lambda)\Phi(s, \lambda)) \rangle_H] ds, \\ \mathcal{I}_3(t, \lambda) &= \int_0^t \langle z^*(s, \lambda), R(\lambda)B\tilde{u}(s) \rangle_H ds - \int_0^t \langle z_1(s, \lambda), R(\lambda)(y(s) - y_d(s)) \rangle_H ds, \\ \mathcal{I}_4(t, \lambda) &= \int_0^t \langle z_1(s, \lambda), \Phi(s, \lambda) dW(s) \rangle_H + \int_0^t \langle z^*(s, \lambda), R(\lambda)G(R(\lambda)z_1(s, \lambda)) dW(s) \rangle_H. \end{aligned}$$

By definition, we have $z^*(T, \lambda) = 0$ for all $\lambda > 0$ and \mathbb{P} -almost surely. Hence, we obtain for all $\lambda > 0$ and \mathbb{P} -a.s.

$$0 = \mathcal{I}_1(T, \lambda) + \mathcal{I}_2(T, \lambda) + \mathcal{I}_3(T, \lambda) + \mathcal{I}_4(T, \lambda).$$

Since the operator A is self adjoint, we have for all $\lambda > 0$ and \mathbb{P} -a.s. $\mathcal{I}_1(T, \lambda) = 0$. Recall that the operator $R(\lambda)$ is self adjoint on H . Using equation (4.7), we obtain for all $\lambda > 0$ and \mathbb{P} -a.s. $\mathcal{I}_2(T, \lambda) = 0$. Note that

$\mathbb{E} \mathcal{I}_4(T, \lambda) = 0$ for all $\lambda > 0$. We can conclude $0 = \mathbb{E} \mathcal{I}_3(T, \lambda)$ for all $\lambda > 0$ and hence, we have for all $\lambda > 0$

$$\mathbb{E} \int_0^T \langle R(\lambda) z_1(t, \lambda), y(t) - y_d(t) \rangle_H dt = \mathbb{E} \int_0^T \langle R(\lambda) z^*(t, \lambda), B\tilde{u}(t) \rangle_H dt. \quad (5.7)$$

Next, we show that the left hand side and the right hand side of equation (5.7) converge as $\lambda \rightarrow \infty$. By the Cauchy-Schwarz inequality and inequality (2.4), we have for all $\lambda > 0$

$$\begin{aligned} & \left| \mathbb{E} \int_0^T \langle z_1(t), y(t) - y_d(t) \rangle_H dt - \mathbb{E} \int_0^T \langle R(\lambda) z_1(t, \lambda), y(t) - y_d(t) \rangle_H dt \right|^2 \\ & \leq 2 \left| \mathbb{E} \int_0^T \langle [I - R(\lambda)] z_1(t), y(t) - y_d(t) \rangle_H dt \right|^2 + 2 \left| \mathbb{E} \int_0^T \langle R(\lambda) (z_1(t) - z_1(t, \lambda)), y(t) - y_d(t) \rangle_H dt \right|^2 \\ & \leq 4 \left(\mathbb{E} \int_0^T \|y(t)\|_H^2 dt + \int_0^T \|y_d(t)\|_H^2 dt \right) \left(\mathbb{E} \int_0^T \|[I - R(\lambda)] z_1(t)\|_H^2 dt + \mathbb{E} \int_0^T \|z_1(t) - z_1(t, \lambda)\|_H^2 dt \right). \end{aligned}$$

Using equation (2.6), Lebesgue's dominated convergence theorem ([4], Thm. 2.8.1) and Lemma 4.11, we can conclude

$$\lim_{\lambda \rightarrow \infty} \mathbb{E} \int_0^T \langle R(\lambda) z_1(t, \lambda), y(t) - y_d(t) \rangle_H dt = \mathbb{E} \int_0^T \langle z_1(t), y(t) - y_d(t) \rangle_H dt. \quad (5.8)$$

Recall that the operator $B: H \rightarrow H$ is bounded. Similarly as above, there exists a constant $C^* > 0$ such that for all $\lambda > 0$

$$\begin{aligned} & \left| \mathbb{E} \int_0^T \langle z^*(t), B\tilde{u}(t) \rangle_H dt - \mathbb{E} \int_0^T \langle R(\lambda) z^*(t, \lambda), B\tilde{u}(t) \rangle_H dt \right|^2 \\ & \leq 2 \left| \mathbb{E} \int_0^T \langle [I - R(\lambda)] z^*(t), B\tilde{u}(t) \rangle_H dt \right|^2 + 2 \left| \mathbb{E} \int_0^T \langle R(\lambda) (z^*(t) - z^*(t, \lambda)), B\tilde{u}(t) \rangle_H dt \right|^2 \\ & \leq C^* \left(\mathbb{E} \int_0^T \|\tilde{u}(t)\|_H^2 dt \right) \left(\mathbb{E} \int_0^T \|[I - R(\lambda)] z^*(t)\|_H^2 dt + \sup_{t \in [0, T]} \mathbb{E} \|z^*(t) - z^*(t, \lambda)\|_H^2 \right). \end{aligned}$$

By equation (2.6), Lebesgue's dominated convergence theorem ([4], Thm. 2.8.1) and Lemma 4.13, we can infer

$$\lim_{\lambda \rightarrow \infty} \mathbb{E} \int_0^T \langle R(\lambda) z^*(t, \lambda), B\tilde{u}(t) \rangle_H dt = \mathbb{E} \int_0^T \langle z^*(t), B\tilde{u}(t) \rangle_H dt.$$

We conclude that the left hand side and the right hand side of equation (5.7) converge as $\lambda \rightarrow \infty$ and equation (5.1) holds.

Next, we show that equation (5.2) holds. Again, we apply Lemma 2.8 to equation (5.4) and equation (5.6). Similarly to equation (5.7), we find for all $\lambda > 0$ and all $\alpha \in (0, \frac{1}{4})$

$$\mathbb{E} \int_0^T \langle R(\lambda) z_2(t, \lambda), y(t) - y_d(t) \rangle_H dt = \mathbb{E} \int_0^T \langle R(\lambda) A^{1-\alpha} z^*(t, \lambda), A^\alpha D\tilde{v}(t) \rangle_H dt. \quad (5.9)$$

Similarly to equation (5.8), we can conclude

$$\lim_{\lambda \rightarrow \infty} \mathbb{E} \int_0^T \langle R(\lambda) z_2(t, \lambda), y(t) - y_d(t) \rangle_H dt = \mathbb{E} \int_0^T \langle z_2(t), y(t) - y_d(t) \rangle_H dt.$$

Recall that the operator $A^\alpha D: V^0(\partial\mathcal{D}) \rightarrow H$ is bounded for all $\alpha \in (0, \frac{1}{4})$. Hence, the process $(A^\alpha D\tilde{v}(t))_{t \in [0, T]}$ takes values in H such that $\mathbb{E} \int_0^T \|A^\alpha D\tilde{v}(t)\|_H^2 dt < \infty$. Since $D(A^{1-\alpha})$ is dense in H , there exists a sequence of processes $(v_m(t))_{t \in [0, T]}$, $m \in \mathbb{N}$, taking values in $D(A^{1-\alpha})$ such that

$$\mathbb{E} \int_0^T \|v_m(t)\|_{D(A^{1-\alpha})}^2 dt < \infty$$

for each $m \in \mathbb{N}$ and

$$\lim_{m \rightarrow \infty} \mathbb{E} \int_0^T \|A^\alpha D\tilde{v}(t) - v_m(t)\|_H^2 dt = 0.$$

Due to Proposition 4.8, the process $(z^*(t))_{t \in [0, T]}$ takes values in $D(A^{1-\alpha})$ for all $\alpha \in (0, \frac{1}{4})$. By equation (2.5), Lemma 2.2, the Cauchy-Schwarz inequality, inequality (2.4) and Fubini's theorem, there exists a constant $C^* > 0$ such that for all $\lambda > 0$, all $\alpha \in (0, \frac{1}{4})$ and each $m \in \mathbb{N}$

$$\begin{aligned} & \left| \mathbb{E} \int_0^T \langle A^{1-\alpha} z^*(t), v_m(t) \rangle_H dt - \mathbb{E} \int_0^T \langle R(\lambda) A^{1-\alpha} z^*(t, \lambda), v_m(t) \rangle_H dt \right|^2 \\ & \leq 2 \left| \mathbb{E} \int_0^T \langle [I - R(\lambda)] z^*(t), A^{1-\alpha} v_m(t) \rangle_H dt \right|^2 + 2 \left| \mathbb{E} \int_0^T \langle R(\lambda) (z^*(t) - z^*(t, \lambda)), A^{1-\alpha} v_m(t) \rangle_H dt \right|^2 \\ & \leq C^* \left(\mathbb{E} \int_0^T \|v_m(t)\|_{D(A^{1-\alpha})}^2 dt \right) \left(\mathbb{E} \int_0^T \|[I - R(\lambda)] z^*(t)\|_H^2 dt + \sup_{t \in [0, T]} \mathbb{E} \|z^*(t) - z^*(t, \lambda)\|_H^2 \right). \end{aligned}$$

Using equation (2.6), Lebesgue's dominated convergence theorem ([4], Thm. 2.8.1) and Lemma 4.13, we can infer for each $m \in \mathbb{N}$

$$\lim_{\lambda \rightarrow \infty} \mathbb{E} \int_0^T \langle R(\lambda) A^{1-\alpha} z^*(t, \lambda), v_m(t) \rangle_H dt = \mathbb{E} \int_0^T \langle A^{1-\alpha} z^*(t), v_m(t) \rangle_H dt.$$

Due to the Moore-Osgood theorem ([45], Thm. 7.11), we get

$$\begin{aligned} \lim_{\lambda \rightarrow \infty} \mathbb{E} \int_0^T \langle R(\lambda) A^{1-\alpha} z^*(t, \lambda), A^\alpha D \tilde{v}(t) \rangle_H dt &= \lim_{\lambda \rightarrow \infty} \lim_{m \rightarrow \infty} \mathbb{E} \int_0^T \langle R(\lambda) A^{1-\alpha} z^*(t, \lambda), v_m(t) \rangle_H dt \\ &= \lim_{m \rightarrow \infty} \lim_{\lambda \rightarrow \infty} \mathbb{E} \int_0^T \langle R(\lambda) A^{1-\alpha} z^*(t, \lambda), v_m(t) \rangle_H dt \\ &= \mathbb{E} \int_0^T \langle A^{1-\alpha} z^*(t), A^\alpha D \tilde{v}(t) \rangle_H dt. \end{aligned}$$

We conclude that the left hand side and the right hand side of equation (5.9) converge as $\lambda \rightarrow \infty$ and equation (5.2) holds. \square

5.2. The optimal controls

Based on the optimality conditions given by equation (4.4) and equation (4.5), we deduce formulas of the optimal controls using the duality principle derived in the previous theorem.

Theorem 5.2. *Let $(z^*(t; u, v), \Phi(t; u, v))_{t \in [0, T]}$ be the mild solution of system (4.6) corresponding to the controls $u \in U$ and $v \in V$. Then the optimal controls $\bar{u} \in U$ and $\bar{v} \in V$ satisfy for all $\alpha \in (0, \frac{1}{4})$, almost all $t \in [0, T]$ and \mathbb{P} -a.s.*

$$\bar{u}(t) = -\frac{1}{\kappa_1} B^* z^*(t; \bar{u}, \bar{v}), \quad (5.10)$$

$$\bar{v}(t) = -\frac{1}{\kappa_2} K^* A^{1-\alpha} z^*(t; \bar{u}, \bar{v}), \quad (5.11)$$

where $B^* \in \mathcal{L}(H)$ and $K^* \in \mathcal{L}(H; V^0(\partial D))$ are the adjoint operators of B and $K = A^\alpha D$, respectively.

Proof. Let $(y(t; u, v))_{t \in [0, T]}$ and $(z_1(t; u))_{t \in [0, T]}$ be the mild solutions of system (3.1) and system (4.2) corresponding to the controls $u \in U$ and $v \in V$, respectively. Using equation (4.4) and Theorem 4.3, the optimal control $\bar{u} \in U$ satisfies for every $\tilde{u} \in U$

$$\mathbb{E} \int_0^T \langle y(t; \bar{u}, \bar{v}) - y_d(t), z_1(t; \tilde{u}) \rangle_H dt + \kappa_1 \mathbb{E} \int_0^T \langle \bar{u}(t), \tilde{u}(t) \rangle_H dt = 0.$$

By equation (5.1), we obtain for every $\tilde{u} \in U$

$$\mathbb{E} \int_0^T \langle z^*(t; \bar{u}, \bar{v}), B \tilde{u}(t) \rangle_H dt + \kappa_1 \mathbb{E} \int_0^T \langle \bar{u}(t), \tilde{u}(t) \rangle_H dt = 0.$$

Hence, we get for every $\tilde{u} \in U$

$$\mathbb{E} \int_0^T \langle B^* z^*(t; \bar{u}, \bar{v}) + \kappa_1 \bar{u}(t), \tilde{u}(t) \rangle_H dt = 0.$$

Therefore, the optimal control $\bar{u} \in U$ satisfies equation (5.10) for almost all $t \in [0, T]$ and \mathbb{P} -almost surely.

Let $(z_2(t; v))_{t \in [0, T]}$ be the mild solution of system (4.3) corresponding to the control $v \in V$. Due to equation (4.5) and Theorem 4.3, the optimal control $\bar{v} \in V$ fulfills the following equation for every $\tilde{v} \in V$:

$$\mathbb{E} \int_0^T \langle y(t; \bar{u}, \bar{v}) - y_d(t), z_2(t; \tilde{v}) \rangle_H dt + \kappa_2 \mathbb{E} \int_0^T \langle \bar{v}(t), \tilde{v}(t) \rangle_{V^0(\partial\mathcal{D})} dt = 0.$$

By equation (5.2), we have for all $\alpha \in (0, \frac{1}{4})$ and every $\tilde{v} \in V$

$$\mathbb{E} \int_0^T \langle A^{1-\alpha} z^*(t; \bar{u}, \bar{v}), A^\alpha D\tilde{v}(t) \rangle_H dt + \kappa_2 \mathbb{E} \int_0^T \langle \bar{v}(t), \tilde{v}(t) \rangle_{V^0(\partial\mathcal{D})} dt = 0.$$

Hence, we get for all $\alpha \in (0, \frac{1}{4})$ and every $\tilde{v} \in V$

$$\mathbb{E} \int_0^T \langle K^* A^{1-\alpha} z^*(t; \bar{u}, \bar{v}) + \kappa_2 \bar{v}(t), \tilde{v}(t) \rangle_{V^0(\partial\mathcal{D})} dt = 0.$$

Therefore, the optimal control $\bar{v} \in V$ satisfies equation (5.11) for all $\alpha \in (0, \frac{1}{4})$, almost all $t \in [0, T]$ and \mathbb{P} -almost surely. \square

Remark 5.3. Let us denote by $(\bar{y}(t))_{t \in [0, T]}$ and $(\bar{z}^*(t), \bar{\Phi}(t))_{t \in [0, T]}$ the mild solutions of system (3.1) and system (4.6) corresponding to the optimal controls $\bar{u} \in U$ and $\bar{v} \in V$, respectively. As a consequence of the previous theorem, the optimal state $(\bar{y}(t))_{t \in [0, T]}$ can be computed by solving the stochastic boundary value problem imposed by the following system of coupled forward-backward SPDEs:

$$\begin{cases} d\bar{y}(t) = \left[-A\bar{y}(t) - \frac{1}{\kappa_1} BB^* \bar{z}^*(t) - \frac{1}{\kappa_2} ADK^* A^{1-\alpha} \bar{z}^*(t) \right] dt + G(\bar{y}(t)) dW(t), \\ d\bar{z}^*(t) = - \left[-A\bar{z}^*(t) + G^* (\bar{\Phi}(t)) + \bar{y}(t) - y_d(t) \right] dt + \bar{\Phi}(t) dW(t), \\ \bar{y}(0) = \xi, \quad \bar{z}^*(T) = 0. \end{cases} \quad (5.12)$$

6. CONCLUSION

In this paper, we considered a control problem constrained by the stochastic Stokes equations on connected and bounded domains with linear multiplicative noise, where controls are defined inside the domain as well as on the boundary.

We proved an existence and uniqueness result for the mild solution of the stochastic Stokes equations dependent on inhomogeneous tangential boundary conditions. Based on the Fréchet derivative of the cost functional, we stated necessary and sufficient optimality conditions the optimal distributed control as well as the optimal boundary control have to satisfy. Using the adjoint equation given by a backward SPDE, a duality principle was derived from which we deduced explicit formulas for the optimal controls. As a consequence, the optimal velocity field can be obtained by solving a system of coupled forward-backward SPDEs.

For engineering applications to control problems of fluid dynamics, the inflow is often used as a boundary control, see [42] and the references therein. These boundary controls can not be covered by tangential boundary conditions and thus remain as an open problem.

Acknowledgements. This research is supported by a research grant of the ‘International Max Planck Research School (IMPRS) for Advanced Methods in Process and System Engineering’, Magdeburg. The authors would like to thank Prof. Wilfried Grecksch of the Martin Luther University Halle-Wittenberg for his helpful advice on various technical issues.

REFERENCES

- [1] A. Al-Hussein, Strong, mild and weak solutions of backward stochastic evolution equations. *Random Oper. Stoch. Equ.* **13** (2005) 129–138.
- [2] O. Axelsson, S. Farouq and M. Neytcheva, A preconditioner for optimal control problems, constrained by Stokes equation with a time-harmonic control. *J. Comput. Appl. Math.* **310** (2017) 5–18.
- [3] A. Bensoussan, G. Da Prato, M.C. Delfour and S.K. Mitter, Representation and Control of Infinite Dimensional Systems. Birkhäuser Boston (2007).
- [4] V. Bogachev, Measure Theory. Springer, Berlin (2007).
- [5] P. Chen, A. Quarteroni and G. Rozza, Multilevel and weighted reduced basis method for stochastic optimal control problems constrained by Stokes equations. *Numer. Math.* **130** (2015) 1–36.
- [6] S. Corlay, Partial functional quantization and generalized bridges. *Bernoulli J.* **20** (2014) 716–746.
- [7] R.F. Curtain and H. Zwart, An Introduction to Infinite-Dimensional Linear Systems Theory. Vol. 21 of *Texts in Applied Mathematics*. Springer-Verlag, New York (1995).
- [8] G. Da Prato and J. Zabczyk, Stochastic Equations in Infinite Dimensions. Cambridge University Press (2014).
- [9] A. Debussche, M. Fuhrman and G. Tessitore, Optimal control of a stochastic heat equation with boundary-noise and boundary-control. *ESAIM: COCV* **13** (2007) 178–205.
- [10] J. Duan and W. Wang, Effective Dynamics of Stochastic Partial Differential Equations. Elsevier (2014).
- [11] G. Fabbri and B. Goldys, An LQ problem for the heat equation on the halfline with Dirichlet boundary control and noise. *SIAM J. Control Optim.* **48** (2009) 1473–1488.
- [12] G. Fabbri, F. Gozzi and A. Święch, Stochastic Optimal Control in Infinite Dimension: Dynamic Programming and HJB Equations. Springer, Berlin (2017).
- [13] C. Fang and W. Han, Well-posedness and optimal control of a hemivariational inequality for nonstationary Stokes fluid flow. *Discrete Contin. Dyn. Syst.* **39** (2016) 5369–5386.
- [14] W.H. Fleming and R.W. Rishel, Deterministic and Stochastic Optimal Control. Springer, New York (1975).
- [15] M. Fuhrman and C. Orrieri, Stochastic maximum principle for optimal control of a class of nonlinear SPDEs with dissipative drift. *SIAM J. Control Optim.* **54** (2016) 341–371.
- [16] H. Fujita and H. Morimoto, On fractional powers of the Stokes operator. *Proc. Jpn. Acad.* **46** (1970) 1141–1143.
- [17] D. Fujiwara, Concrete characterization of the domains of fractional powers of some elliptic differential operators of the second order. *Proc. Jpn. Acad.* **43** (1967) 82–86.
- [18] D. Fujiwara and H. Morimoto, An L_r -theorem of the Helmholtz decomposition of vector fields. *J. Fac. Sci. Univ. Tokyo Sec. 1 A* **24** (1977) 685–700.
- [19] G.P. Galdi, An Introduction to the Mathematical Theory of the Navier–Stokes Equations. Springer, New York (2011).
- [20] G.P. Galdi, C.G. Simader and H. Sohr, A class of solutions to the stationary Stokes and Navier–Stokes equations with boundary data in $W^{1/q,q}$. *Math. Ann.* **331** (2005) 41–74.
- [21] C.M. Gariboldi and E.L. Schwindt, Simultaneous Optimal Controls for Unsteady Stokes Systems. In *Conf. of V MACI* (2015).
- [22] L. Gawarecki and V. Mandrekar, Stochastic Differential Equations in Infinite Dimensions. Springer-Verlag Berlin Heidelberg (2011).
- [23] Y. Giga, Analyticity of the Semigroup Generated by the Stokes Operator in L_r Space. *Math. Z.* **178** (1981) 297–329.
- [24] Y. Giga and T. Miyakawa, Solutions in L_r of the Navier-Stokes initial value problem. *Arch. Ration. Mech. Anal.* **89** (1985) 267–281.
- [25] T.E. Govindan, Yosida Approximations of Stochastic Differential Equations in Infinite Dimensions and Applications. Springer (2016).
- [26] G. Guatteri and F. Masiero, On the existence of optimal controls for SPDEs with boundary-noise and boundary-control. *SIAM J. Control Optim.* **51** (2013) 1909–1939.
- [27] M. Hintermüller and M. Hinze, A SQP–Semismooth Newton-type Algorithm applied to Control of the instationary Navier–Stokes System Subject to Control Constraints. *SIAM J. Optim.* **16** (2006) 1177–1200.
- [28] M. Hinze, R. Pinnau, M. Ulbrich and S. Ulbrich, Optimization with PDE Constraints. Springer, New York (2009).
- [29] Y. Hu and S. Peng, Adapted solution of a backward semilinear stochastic evolution equation. *Stoch. Anal. Appl.* **9** (1991) 445–459.
- [30] A. Ichikawa, Stability of semilinear stochastic evolution equations. *J. Math. Anal. Appl.* **90** (1982) 12–44.
- [31] W. Krendl, V. Simoncini and W. Zulehner, Efficient Preconditioning for an Optimal Control Problem with the Time-Periodic Stokes Equations. *Numerical Mathematics and Advanced Applications - ENUMATH 2013*, edited by A. Abdulle, S. Deparis, D. Kressner, F. Nobile, M. Picasso. Vol. 103 of *Lecture Notes in Computational Science and Engineering*. Springer, Cham (2015) 479–487.
- [32] K. Kunisch and X. Lu, Optimal control for multi-phase fluid Stokes problems. *Nonlinear Anal.* **74** (2011) 585–599.

- [33] A.J. Kurdila and M. Zabaranin, *Convex Functional Analysis*. Birkhäuser Basel (2005).
- [34] H. Liang, L. Hou and J. Ming, The velocity tracking problem for Wick-stochastic Navier–Stokes flows using Wiener chaos expansion. *J. Comput. Appl. Math.* **307** (2016) 25–36.
- [35] Q. Lü and X. Zhang, *General Pontryagin–Type Stochastic Maximum Principle and Backward Stochastic Evolution Equations in Infinite Dimensions*. Springer, Cham (2014).
- [36] S. Nicaise and D. Sirch, Optimal control of the Stokes equations: conforming and non–conforming finite element methods under reduced regularity. *Comput. Optim. Appl.* **49** (2011) 567–600.
- [37] Y.R. Ou, Design of Feedback Compensators for Viscous Flow. In: *Optimal Control of Viscous Flow*. SIAM, Philadelphia (1998) 151–180.
- [38] A. Pazy, *Semigroups of Linear Operators and Applications to Partial Differential Equations*. Springer, New York (1983).
- [39] H. Pham, *Continuous-time Stochastic Control and Optimization with Financial Applications*. Springer, Berlin (2009).
- [40] C. Prévôt and M. Röckner, A Concise Course on Stochastic Partial Differential Equations. In vol. 1905 of *Lecture Notes in Mathematics*. Springer, Berlin (2007).
- [41] J.-P. Raymond, Feedback boundary stabilization of the two-dimensional Navier–Stokes equations. *SIAM J. Control Optim.* **45** (2006) 790–828.
- [42] J.-P. Raymond, Stokes and Navier–Stokes equations with nonhomogeneous boundary conditions. *Ann. Inst. Henri Poincaré Anal. Non Linéaire* **24** (2007) 921–951.
- [43] A. Rösch and B. Vexler, Optimal control of the Stokes equations: A priori error analysis for finite element discretization with postprocessing. *SIAM J. Numer. Anal.* **44** (2006) 1903–1920.
- [44] B.L. Rozovskii, *Stochastic Evolution Systems: Linear Theory and Applications to Non-linear Filtering*. Springer, Netherlands (1990).
- [45] W. Rudin, *Principles of Mathematical Analysis*, Vol. 3. McGraw-Hill, Inc. (1964).
- [46] S. S. Sritharan, An Introduction to Deterministic and Stochastic Control of Viscous Flow. In: *Optimal Control of Viscous Flow*. SIAM, Philadelphia (1998) 1–42.
- [47] G. Tessitore, Existence, uniqueness and space regularity of the adapted solutions of a backward spde. *Stochastic Anal. Appl.* **14** (1996) 461–486.
- [48] M. Ulbrich, Constrained optimal control of Navier–Stokes flow by semismooth Newton methods. *Syst. Control Lett.* **48** (2003) 297–311.
- [49] W. von Wahl, *The Equations of Navier-Stokes and Abstract Parabolic Equations*. Vieweg + Teubner Verlag (1985).
- [50] I. Vrabie, *C_0 -Semigroups and Applications*. In vol. 191 of *Mathematics Studies*. Elsevier, Amsterdam (2003).
- [51] Q. Wu, A new type of the Gronwall-Bellman inequality and its application to fractional stochastic differential equations. *Cogent. Math.* **4** (2017) 2331–1835.
- [52] H. Ye, J. Gao and Y. Ding, A generalized Gronwall inequality and its application to a fractional differential equation. *J. Math. Anal. Appl.* **328** (2007) 1075–1081.
- [53] J. Yong and X.Y. Zhou, *Stochastic Controls: Hamiltonian Systems and HJB Equations*. Springer, New York (1999).
- [54] E. Zeidler, *Nonlinear Functional Analysis and its Applications III: Variational Methods and Optimization*. Springer, New York (1985).