

ON THE SUPER CONNECTIVITY OF DIRECT PRODUCT OF GRAPHS

FARNAZ SOLIEMANY¹, MOHSEN GHASEMI^{1,*} AND REZVAN VARMAZYAR²

Abstract. A vertex-cut S is called a *super vertex-cut* if $G - S$ is disconnected and it contains no isolated vertices. The *super-connectivity*, κ' , is the minimum cardinality over all super vertex-cuts. This article provides bounds for the super connectivity of the direct product of an arbitrary graph and the complete graph K_n . Among other results, we show that if G is a non-complete graph with $\text{girth}(G) = 3$ and $\kappa'(G) = \infty$, then $\kappa'(G \times K_n) \leq \min\{mn - 6, m(n - 1) + 5, 5n + m - 8\}$, where $|V(G)| = m$.

Mathematics Subject Classification. 05C40, 05C82, 05C25.

Received December 15, 2021. Accepted May 24, 2022.

1. INTRODUCTION

We follow [1] for graph theoretic terminologies and notations not defined here. Let G be a simple undirected graph, where $V(G)$ and $E(G)$ denote the set of vertices and the set of edges of G , respectively. For two vertices $u, v \in V(G)$, u and v are *neighbors* if u and v are adjacent and we write $u \sim v$. If u and v are not adjacent in G , then we write $uv \notin E(G)$. For each vertex $v \in V(G)$, the *neighborhood* $N_G(v)$ of v is defined as the set of all vertices adjacent to v and $\deg(v) = |N_G(v)|$ is the *degree* of v . The number $\delta(G) = \min\{\deg(v) \mid v \in V(G)\}$ is the minimum degree of G . Also the *girth* of the graph G , $\text{girth}(G)$, is the length of its shortest cycle if G contains cycle, define $\text{girth}(G) = \infty$ otherwise. A *wheel* graph is a graph formed by connecting a single universal vertex to all vertices of a cycle. We use W_n to denote a wheel graph with $n \geq 3$ vertices. For an arbitrary subset $S \subset V(G)$ we use $G - S$ to denote the graph obtained by removing all vertices in S from G . For any connected graph G , if $G - S$ is disconnected, then S is a *vertex-cut*. The *connectivity* of a graph G , denoted by $\kappa(G)$, is the minimum cardinality of a set $S \subset V(G)$ such that $G - S$ is either disconnected or the trivial graph K_1 . It is known that $\kappa(G) \leq \delta(G)$. A vertex-cut S is called a *super vertex-cut* if $G - S$ is disconnected and it contains no isolated vertices. The *super-connectivity* κ' is the minimum cardinality over all super vertex-cuts, that is,

$$\kappa'(G) = \min\{|S| \mid S \subseteq V \text{ is a super vertex-cut of } G\}.$$

Clearly, the super connectivity $\kappa'(G)$ does not always exist for a connected graph G . We write $\kappa'(G) = \infty$ if $\kappa'(G)$ does not exist. For example, $\kappa'(G) = \infty$ if G is the star $K_{1,n}$.

It is well known that when the underlying topology of an interconnection network is modeled by a graph $G = (V, E)$, where V represents the set of processors and E represents the set of communication links in the

Keywords. Direct product, super connectivity, vertex-cut.

¹ Department of Mathematics, Urmia University, Urmia 57135, Iran.

² Department of Mathematics, Khoy Branch, Islamic Azad University, Khoy 58168-44799, Iran.

*Corresponding author: m.ghasemi@urmia.ac.ir, math.ghasemi@gmail.com

network, the connectivity $\kappa(G)$ of G is an important measurement for the fault tolerance of the network. It has been shown that a super connected network is most reliable and has the smallest vertex failure rate among all the networks with the same connectivity (see, e.g. [20, 21]).

The direct product $X \times Y$ of two graphs X and Y is the graph having $V(X \times Y) = V(X) \times V(Y)$ and $E(X \times Y) = \{(x_1, y_1)(x_2, y_2) \mid x_1x_2 \in E(X) \text{ and } y_1y_2 \in E(Y)\}$.

We state two known results of the direct product of graphs that will be used in the proof of our main results.

Proposition 1.1 ([17]). *Let G and H be connected graphs. The graph $G \times H$ is connected if and only if G or H contains an odd cycle.*

Proposition 1.2 ([17]). *Let G be a connected graph. If G has no odd cycle, then $G \times K_2$ has exactly two components isomorphic to G .*

The direct product plays an important role in design and analysis of network [18]. This product has generated a lot of interest mainly due to its various applications. For instance, it is used in complex networks to generate realistic networks [12], in multiprocessor systems to model of concurrency [11] and in automata theory [5]. The connectivity of direct product graphs has been investigated in [15] and [17]. Also the connectivity of direct product of a bipartite graph and a complete graph has been presented by Guji and Vumar (see [6]). Moreover, the super connectivity of $K_{m,r} \times K_n$ is determined by Ekinici and Kirlangic (see [3]). For more results we refer the reader to [2, 4, 7–10, 13, 14, 16, 19, 22]. In this paper we investigate the super connectivity κ' of the direct product of an arbitrary graph and the complete graph K_n . We show that if $\kappa'(G) = t < \infty$ then $\kappa'(G \times K_n) \leq tn$. Also if $\kappa'(G) = \infty$ and $\text{girth}(G) = 3$, then $\kappa'(G \times K_n) \leq \min\{mn - 6, m(n - 1) + 5, 5n + m - 8\}$, where $|V(G)| = m$.

2. SUPER CONNECTIVITY OF $G \times K_n$

Throughout this section, G is a connected non-complete graph.

Let G be a graph with $V(G) = \{x_1, x_2, \dots, x_m\}$ and $V(K_n) = \{v_1, v_2, \dots, v_n\}$. Suppose that $S_i = V(G) \times v_i$ for $i \in \mathbb{Z}_n$, where $\mathbb{Z}_n = \{1, 2, 3, \dots, n\}$. Hence $V(G \times K_n) = S_1 \cup S_2 \cup \dots \cup S_n$ where $\{S_i\}$ is a partition of $G \times K_n$.

Theorem 2.1. *Let G be a graph with $\kappa'(G) = t < \infty$. Then $\kappa'(G \times K_n) \leq tn$.*

Proof. Let $X = \{x_1, x_2, \dots, x_t\}$ be a minimum super vertex-cut of G . Then $S = \{(x_j, v_i) \mid j \in \mathbb{Z}_t, i \in \mathbb{Z}_n\}$ is a super vertex-cut in $G \times K_n$. So $\kappa'(G \times K_n) \leq tn$. \square

Let C_n be a cycle of length $n \geq 6$ and $V(C_n) = \{x_1, \dots, x_n\}$ where $x_1 \sim x_2 \sim x_3 \sim \dots \sim x_n \sim x_1$. Then $S = \{x_1, x_4\}$ is a super vertex-cut in C_n . Hence, for a graph G with $\text{girth}(G) \geq 6$ we have $\kappa'(G) < \infty$ and by Theorem 2.1, $\kappa'(G \times K_n) < \infty$. Thus we may suppose that $\text{girth}(G) \leq 5$. First suppose that $\text{girth}(G) = 5$. If $|E(G)| \geq 6$ then $\kappa'(G) < \infty$ and so again by Theorem 2.1, $\kappa'(G \times K_n) < \infty$. Thus in the following theorem we may suppose that $\text{girth}(G) = 5$ and $|E(G)| = 5$. It is easy to see that $\kappa'(G \times K_2) = 2$.

Theorem 2.2. *Let G be a cycle of length 5. Then $\kappa'(G \times K_n) = \min\{5n - 8, 3n\}$ for $n \geq 3$.*

Proof. Suppose that G is a cycle of length 5 and $V(G) = \{a, b, c, d, e\}$ where $a \sim b \sim c \sim d \sim e \sim a$. Let $j \in \mathbb{Z}_n$ be constant. Then by deleting all vertices $\{(a, v_j), (b, v_i), (c, v_j), (d, v_t), (e, v_t) \mid i \in \mathbb{Z}_n, t \in \mathbb{Z}_n - \{j\}\}$ we obtain a disconnected graph without any isolated vertex. Therefore, $\kappa'(G \times K_n) \leq 1 + n + 1 + 2(n - 1) = 3n$. Now, let S be a super vertex-cut of $G \times K_n$. Hence $(G \times K_n) - S$ has at least two components, say C_1, C_2 . Let $(x, v_r) \in C_1$ and $(y, v_t) \in C_2$ for some $x, y \in V(G)$. We have four cases:

Case 1. Let $x = y$. Hence $v_r \neq v_t$. Without loss of generality, let $x = a$. So $(a, v_r) \in C_1$ and $(a, v_t) \in C_2$.

Since $N_G(a) = \{b, e\}$, for every $j \in \mathbb{Z}_n - \{r, t\}$, $(a, v_r) \sim (b, v_j) \sim (a, v_t)$ and $(a, v_r) \sim (e, v_j) \sim (a, v_t)$ are paths between (a, v_r) and (a, v_t) in $G \times K_n$. Therefore $\{(b, v_j), (e, v_j) \mid j \in \mathbb{Z}_n - \{r, t\}\} \subset S$. Clearly

$(b, v_t), (e, v_t) \in C_1$ and $(b, v_r), (e, v_r) \in C_2$. Also, for every $j \in \mathbb{Z}_n - \{r, t\}$, $(b, v_t) \sim (a, v_j) \sim (b, v_r)$ and $(e, v_r) \sim (a, v_j) \sim (e, v_t)$ are paths in $G \times K_n$. Thus, $\{(a, v_j) \mid j \in \mathbb{Z}_n - \{r, t\}\} \subset S$. Similarly, $\{(d, v_j), (c, v_j) \mid j \in \mathbb{Z}_n - \{r, t\}\} \subset S$. By deleting these vertices of $G \times K_n$ we obtain an 10-gone, say P_{10} , where

$$V(P_{10}) = \{(a, v_t), (b, v_r), (c, v_t), (d, v_r), (e, v_t), (a, v_r), (b, v_t), (c, v_r), (d, v_t), (e, v_r)\}.$$

Hence $\bigcup_{i=1, i \neq r, t}^n S_i$ with two more vertices is S , that is, $|S| = 5(n-2) + 2 = 5n-8$.

Case 2. Let $x \neq y$, $v_r = v_t$ and $x \sim y$. Without loss of generality, let $x = a$ and $y = b$. Hence $S = \{(a, v_j), (b, v_j) \mid j \in \mathbb{Z}_n - \{r\}\} \cup \{(e, v_r), (c, v_r)\} \cup \{(d, v_i) \mid i \in \mathbb{Z}_n\}$. Therefore, in this case $|S| = 2(n-1) + 2 + n = 3n$.

Case 3. Let $x \neq y$, $v_r = v_t$ and $xy \notin E(G)$. Without loss of generality, let $x = a$ and $y = c$. Since $(a, v_r) \sim (b, v_j) \sim (c, v_r)$ is a path in $G \times K_n$, for every $v_j \neq v_r$, the set $\{(b, v_j) \mid j \in \mathbb{Z}_n - \{r\}\}$ lies in S . Now, we choose one element $l \in \mathbb{Z}_n$ with $l \neq r$. Since $(e, v_l) \in C_1$, $(d, v_l) \in C_2$ and $(e, v_i) \sim (d, v_j)$ for $i \neq j$, the set $\{(e, v_j), (d, v_j) \mid j \in \mathbb{Z}_n - \{l\}\}$ lies in S . In the remaining graph the vertex (b, v_r) is adjacent to all vertices of $\{(a, v_j) \in C_1, (c, v_j) \in C_2 \mid j \neq r\}$, so $(b, v_r) \in S$. Finally, (a, v_l) and (c, v_l) are isolated vertices. Thus they belong to S . Therefore, in this case $|S| = n + 2 + 2(n-1) = 3n$.

Case 4. Let $x \neq y$, $v_r \neq v_t$. Clearly $xy \notin E(G)$. Suppose that $x = a$, $y = c$ and $l \neq r, t$. Thus $(e, v_l) \in C_1$ and $(d, v_l) \in C_2$. Now with the similar arguments in Case 3, we get $S = \{(e, v_j), (d, v_j) \mid j \in \mathbb{Z}_n - \{l\}\} \cup \{(b, v_i) \mid i \in \mathbb{Z}_n\} \cup \{(a, v_l), (c, v_l)\}$. Therefore, in this case $|S| = 3n$.

Thus $\kappa'(G \times K_n) \leq \min\{5n-8, 3n\}$. Furthermore, by the process of the proof, in all cases if $|S| < \min\{5n-8, 3n\}$ then $(G \times K_n) - S$ is either connected or has some isolated vertices. Therefore, $\kappa'(G \times K_n) = \min\{5n-8, 3n\}$. \square

By the above theorem, the following result holds.

Corollary 2.3. Let G be a cycle of length 5. Then $\kappa'(G \times K_n) = 3n$ for $n \geq 4$.

Suppose that G is a bipartite graph with $\kappa'(G) = \infty$. Hence $\text{girth}(G) = \infty$ or $\text{girth}(G)$ is even. If $\text{girth}(G) \geq 6$ then $\kappa'(G) < \infty$ and by Theorem 2.1, $\kappa'(G \times K_n) < \infty$. So we have the following result when $\text{girth}(G) = 4$ or $\text{girth}(G) = \infty$.

Theorem 2.4. Let G be a bipartite graph and $\kappa'(G) = \infty$. Then $\kappa'(G \times K_n) \leq m(n-2)$, where $|V(G)| = m$.

Proof. By Proposition 1.2, $(G \times K_n) - (\bigcup_{i=3}^n S_i) \cong G \times K_2$ has two components isomorphic to G . Thus $\kappa'(G \times K_n) \leq m(n-2)$. \square

Finally in Theorem 2.5, we investigate $\kappa'(G \times K_n)$ when $\text{girth}(G) = 3$ and $\kappa'(G) = \infty$.

Theorem 2.5. Let G be a graph with $\text{girth}(G) = 3$, $|V(G)| = m$ and $\kappa'(G) = \infty$. Then $\kappa'(G \times K_n) \leq \min\{mn-6, m(n-1)+5, 5n+m-8\}$.

Proof. First suppose that G has a unique triangle. Let C be the only cycle of G with $V(C) = \{u_1, u_2, u_3\}$. We consider the following cases:

Case 1. Let $\deg(u_j) = 2$ for all $j \in \mathbb{Z}_3$. Then G is K_3 and by Theorem 2.7 of [3] $\kappa'(G \times K_n) = 3n-4$.

Case 2. Let $\deg(u_j) \geq 3$ for all $j \in \mathbb{Z}_3$. Also let $N_{G-\{u_2, u_3\}}(u_1) = \{x_{r'} \mid r' \in \mathbb{Z}_r\}$, $N_{G-\{u_1, u_3\}}(u_2) = \{y_{s'} \mid s' \in \mathbb{Z}_s\}$ and $N_{G-\{u_1, u_2\}}(u_3) = \{w_{t'} \mid t' \in \mathbb{Z}_t\}$ where $r+s+t = m-3$. Now, $\kappa'(G) = \infty$ implies that $\deg(x_{r'}) = \deg(y_{s'}) = \deg(w_{t'}) = 1$, for every $r' \in \mathbb{Z}_r$, $s' \in \mathbb{Z}_s$ and $t' \in \mathbb{Z}_t$. Hence $S = \{(u_j, v_l) \mid j \in \mathbb{Z}_3, l \in \mathbb{Z}_n - \{1\}\} \cup \{(x_{r'}, v_1), (y_{s'}, v_1), (w_{t'}, v_1) \mid r' \in \mathbb{Z}_r, s' \in \mathbb{Z}_s, t' \in \mathbb{Z}_t\}$ is a super vertex-cut in $G \times K_n$ with $|S| = 3(n-1) + (m-3) = 3n+m-6$.

Case 3. Let $\deg(u_1) \geq 3$ and $\deg(u_2) = \deg(u_3) = 2$. In this case $S = \bigcup_{i=3}^n S_i \cup \{(u_3, v_1), (u_3, v_2)\}$ is a super vertex-cut in $G \times K_n$ with $|S| = m(n-2) + 2$.




FIGURE 1.




FIGURE 2.

Case 4. Let $\deg(u_1) \geq 3$, $\deg(u_2) \geq 3$ and $\deg(u_3) = 2$. Let $N_{G-\{u_2, u_3\}}(u_1) = \{x_{r'} \mid r' \in \mathbb{Z}_r\}$ and $N_{G-\{u_1, u_3\}}(u_2) = \{y_{s'} \mid s' \in \mathbb{Z}_s\}$ where $r+s = m-3$. We have two subcases:

Subcase 1. If $\deg(x_{r'}) = \deg(y_{s'}) = 1$ for every $x_{r'} \in N_{G-\{u_2, u_3\}}(u_1)$ and $y_{s'} \in N_{G-\{u_1, u_3\}}(u_2)$ then $S = \{(u_3, v_i) \mid i \in \mathbb{Z}_n\} \cup \{(u_1, v_j), (u_2, v_j) \mid j \in \mathbb{Z}_n - \{1\}\} \cup \{(y_{s'}, v_1), (x_{r'}, v_1) \mid r' \in \mathbb{Z}_r, s' \in \mathbb{Z}_s\}$ is a super vertex-cut in $G \times K_n$ with $|S| = n + 2(n-1) + (m-3) = 3n + m - 5$.

Subcase 2. Let there exist some vertices $x_p \in N_{G-\{u_2, u_3\}}(u_1)$ and $y_q \in N_{G-\{u_1, u_3\}}(u_2)$ such that $x_p \sim y_q$. Since $\kappa'(G) = \infty$ if $x_p \sim y_q$ and $x_{p'} \sim y_{q'}$ then $x_p \sim y_{q'}$ or $x_{p'} \sim y_q$. Let h be the number of vertices $x_{r'}$ with $\deg(x_{r'}) \geq 2$ and l be the number of vertices y_s with $\deg(y_s) \geq 2$. Without loss of generality let $l \leq h$, and $\{y_1, y_2, \dots, y_l\} \subset N_{G-\{u_1, u_3\}}(u_2)$ be such that $\deg(y_q) \geq 2$ for $q \in \mathbb{Z}_l$. Then $S = \{(u_3, v_i) \mid i \in \mathbb{Z}_n\} \cup \{(u_1, v_j), (u_2, v_j) \mid j \in \mathbb{Z}_n - \{1\}\} \cup \{(y_q, v_i) \mid q \in \mathbb{Z}_l, i \in \mathbb{Z}_n - \{1\}\} \cup \{(y_{s'}, v_1) \mid l < s' \leq s\} \cup \{(x_{r'}, v_1) \mid r' \in \mathbb{Z}_r\}$ is a super vertex-cut in $G \times K_n$ with $|S| = n + 2(n-1) + l(n-1) + (s-l) + r = 3n + m - 5 + l(n-2)$. If $l = s$ then (u_2, v_1) is an isolated vertex in $(G \times K_n) - S$. So $l < s$. Now $l \leq m-5$ implies that $|S| \leq n(m-2) - m + 5$.

Now, let $\deg(x_{r'}) \geq 2$ and $\deg(y_{s'}) \geq 2$ for every $x_{r'} \in N_{G-\{u_2, u_3\}}(u_1)$, $y_{s'} \in N_{G-\{u_1, u_3\}}(u_2)$. We choose a constant $l \neq 1$ of \mathbb{Z}_n . Hence $S = \{(u_3, v_i) \mid i \in \mathbb{Z}_n\} \cup \{(u_1, v_i), (u_2, v_i) \mid i \in \mathbb{Z}_n - \{1\}\} \cup \{(x_{r'}, v_i), (y_{s'}, v_i) \mid i \in \mathbb{Z}_n - l\}$ is a super vertex-cut in $G \times K_n$ with $|S| = n + 2(n - 1) + (m - 3)(n - 1) = m(n - 1) + 1$.

Now, suppose that G contains t cycles which have common edges. We consider the following cases:

Case 1. Let G be isomorphic to t triangles which have a common edge $\{a, b\}$. Also, let the other vertices be the set $\{\alpha_1, \alpha_2, \dots, \alpha_{m-2}\}$. Now $S = \{(\alpha_j, v_i), (a, v_i) \mid j \in \mathbb{Z}_{m-2}, i \in \mathbb{Z}_n - \{1, 2\}\} \cup \{(b, v_i) \mid i \in \mathbb{Z}_n\}$ is a super vertex-cut in $G \times K_n$ with $|S| = mn - 2m + 2$. Also let G be isomorphic to t triangles and a square which have a common edge $\{a, b\}$. Let $\{a, b, \alpha_1, \alpha_2\}$ be vertices of square and $\{\alpha_3, \dots, \alpha_{m-2}\}$ be other vertices. Again S is a super vertex-cut in $G \times K_n$ with $|S| = mn - 2m + 2$. Furthermore, if G has more than one square with common edge $\{a, b\}$ then it is clear that $\kappa'(G) < \infty$, a contradiction. Assume that G contains more than one square and there are some edges between the vertices of squares such that $\kappa'(G) = \infty$ then we have two subcases:

Subcase 1. If the vertices of square are such that there is no pentagon then S is as above.

Subcase 2. Let $\{\alpha_1, \alpha_2, \dots, \alpha_s\}$ be the vertices of triangles and $\{\alpha_{s+1}, \alpha_{s+2}, \dots, \alpha_{m-2}\}$ be the vertices of squares. We have to delete some vertices for removing pentagons. Let $\{\alpha_{s+1}, \alpha_{s+2}, \dots, \alpha_{s+l}\}$ be the minimum vertices which deleting them removes pentagons. Hence $S = \{(b, v_i) \mid i \in \mathbb{Z}_n\} \cup \{(a, v_i) \mid i \in \mathbb{Z}_n - \{1, 2\}\} \cup \{(\alpha_j, v_i) \mid j \in \mathbb{Z}_s, i \in \mathbb{Z}_n - \{1, 2\}\} \cup \{(\alpha_j, v_i) \mid s+1 \leq j \leq s+l, i \in \mathbb{Z}_n\} \cup \{(\alpha_j, v_i) \mid s+l+1 \leq j \leq m-2, i \in \mathbb{Z}_n - \{1, 2\}\}$ is a super vertex-cut in $G \times K_n$ with $|S| = n + (n-2) + s(n-2) + ln + (m-2-s-l-1+1)(n-2) = mn - 2m + 2 + 2l$. Thus $l \leq m-5$ implies that $|S| \leq mn - 8$.

Moreover, if G is isomorphic to Figure 1, then $S = \{(a, v_i), (b, v_i) \mid i \in \mathbb{Z}_n - \{1\}\} \cup \{(\beta_j, v_1), (\gamma_r, v_1) \mid j \in \mathbb{Z}_f, r \in \mathbb{Z}_{\tilde{f}}\} \cup \{(\alpha_e, v_i) \mid e \in \mathbb{Z}_f, i \in \mathbb{Z}_n\}$ is a super vertex-cut in $G \times K_n$ with $|S| = 2(n-1) + \tilde{f} + \tilde{f} + fn$. Now, $\tilde{f} + \tilde{f} + f = m-2$ and $f \leq m-4$ implies that $|S| \leq 2(n-1) + m-2 + f(n-1) \leq n(m-2)$.

Also, if G is isomorphic to Figure 2, then $S = \{(a, v_i), (b, v_i), (\alpha_1, v_i) \mid i \in \mathbb{Z}_n - \{1\}\} \cup \{(\beta_j, v_1), (\gamma_r, v_1), (\lambda_w, v_1) \mid j \in \mathbb{Z}_{\tilde{f}}, r \in \mathbb{Z}_{\tilde{f}}, w \in \mathbb{Z}_{\tilde{f}}\} \cup \{(\alpha_e, v_i) \mid 2 \leq e \leq f, i \in \mathbb{Z}_n\}$ is a super vertex-cut in $G \times K_n$. Now $\tilde{f} + \tilde{f} + \tilde{f} + f = m-2$ and $f-1 \leq m-6$ implies that $|S| \leq 3(n-1) + \tilde{f} + \tilde{f} + \tilde{f} + (f-1)n \leq n(m-3)$.

Case 2. Let G contains three cycles C_1, C_2 and C_3 with $V(G) = \{u_1, u_2, u_3, \dots, u_m\}$. Let $V(C_1) = \{u_1, u_2, u_3\}$, $V(C_2) = \{u_2, u_3, u_4\}$ and $V(C_3) = \{u_2, u_4, u_5\}$. If $\deg(u_1) = \deg(u_5) = 2$, $\deg(u_2) = 4$ and $\deg(u_3) = \deg(u_4) = 3$ then $S = \cup_{i=3}^n S_i \cup \{(u_2, v_1), (u_2, v_2)\}$ is a super vertex-cut in $G \times K_n$ with $|S| = m(n-2)+2$. Now suppose that $\deg(u_2) \geq 5$ and either $\deg(u_3) \geq 4$ or $\deg(u_4) \geq 4$. Let $N_{G-\{u_1, u_3, u_4, u_5\}}(u_2) = \{x_{r'} \mid r' \in \mathbb{Z}_r\}$, $N_{G-\{u_1, u_2, u_4\}}(u_3) = \{y_{s'} \mid s' \in \mathbb{Z}_s\}$ and $N_{G-\{u_2, u_3, u_5\}}(u_4) = \{w_{t'} \mid t' \in \mathbb{Z}_t\}$ where $r+s+t = m-5$. Let $\deg(x_{r'}) = \deg(y_{s'}) = \deg(w_{t'}) = 1$, for every $r' \in \mathbb{Z}_r$, $s' \in \mathbb{Z}_s$ and $t' \in \mathbb{Z}_t$. So $S = \{(u_j, v_i) \mid 2 \leq j \leq 4, i \in \mathbb{Z}_n - \{1\}\} \cup \{(u_1, v_1)\} \cup \{(u_5, v_i) \mid i \in \mathbb{Z}_n\} \cup \{(x_{r'}, v_1), (y_{s'}, v_1), (w_{t'}, v_1) \mid r' \in \mathbb{Z}_r, s' \in \mathbb{Z}_s, t' \in \mathbb{Z}_t\}$ is a super vertex-cut in $G \times K_n$ with $|S| = 3(n-1)+1+n+(m-5) = 4n-7+m$. Also if $x_{r_i'} \sim x_{r_j'}$ or $y_{s_i'} \sim y_{s_j'}$ or $w_{t_i'} \sim w_{t_j'}$ or $x_{r_i'} \sim y_{s_j'}$ or $x_{r_i'} \sim w_{t_j'}$ or $y_{s_i'} \sim w_{t_j'}$ for some i, j then $\kappa'(G) < \infty$. Moreover if $u_3 \sim w_{t_i'}$ or $u_4 \sim y_{s_i'}$ then $S = \{(u_j, v_i) \mid 2 \leq j \leq 4, i \in \mathbb{Z}_n - \{1\}\} \cup \{(u_1, v_i), (u_5, v_i) \mid i \in \mathbb{Z}_n\} \cup \{(x_{r'}, v_1), (y_{s'}, v_1), (w_{t'}, v_1) \mid r' \in \mathbb{Z}_r, s' \in \mathbb{Z}_s, t' \in \mathbb{Z}_t\}$ is a super vertex-cut in $G \times K_n$ with $|S| = 3(n-1) + 2n + (m-5) = 5n-8+m$. Now let $x_{r_i'} \sim u_3$ or $x_{r_j'} \sim u_4$ for some i, j . Let $A = \{x \in N_{G-\{u_1, u_3, u_4, u_5\}}(u_2) \mid x \sim u_3 \text{ and } x \sim u_4\}$, $B = \{x \in N_{G-\{u_1, u_3, u_4, u_5\}}(u_2) \mid x \sim u_3 \text{ and } xu_4 \notin E(G)\}$, $C = \{x \in N_{G-\{u_1, u_3, u_4, u_5\}}(u_2) \mid xu_3 \notin E(G) \text{ and } x \sim u_4\}$, $D = \{x \in N_{G-\{u_1, u_3, u_4, u_5\}}(u_2) \mid x \sim \dot{x} \text{ for some } \dot{x}; (\dot{x} \in N_{G-\{u_1, u_3, u_4, u_5\}}(u_2) \cap N_{G-\{u_1, u_2, u_4\}}(u_3) \text{ and } \dot{x} \notin N_{G-\{u_1, u_3, u_4, u_5\}}(u_2) \cap N_{G-\{u_2, u_3, u_5\}}(u_4) \text{ or } (\dot{x} \in N_{G-\{u_1, u_3, u_4, u_5\}}(u_2) \cap N_{G-\{u_2, u_3, u_5\}}(u_4) \text{ and } \dot{x} \notin N_{G-\{u_1, u_3, u_4, u_5\}}(u_2) \cap N_{G-\{u_1, u_2, u_4\}}(u_3))\}$ and $F = \{x \in N_{G-\{u_1, u_3, u_4, u_5\}}(u_2) \mid x \notin A \cup B \cup C \cup D\}$ with $|A| = l$, $|B \cup C| = \tilde{l}$, $|D| = \tilde{l}$ and $|F| = \tilde{l}$. Then $S = \{(u_3, v_i), (u_4, v_i) \mid i \in \mathbb{Z}_n - \{1\}\} \cup \{(u_2, v_i) \mid i \in \mathbb{Z}_n\} \cup \{(u_1, v_1), (u_5, v_1)\} \cup \{(x, v_i) \mid x \in A \text{ or } x \in F, i \in \mathbb{Z}_n\} \cup \{(x, v_1) \mid x \in B \cup C\} \cup \{(y_{s'}, v_1), (w_{t'}, v_1)\}$ is a super vertex-cut in $G \times K_n$ with $|S| = 2(n-1) + n + 2 + (l + \tilde{l})n + (\tilde{l}) + (m-5-r)$. Now, $l + \tilde{l} + \tilde{l} = r - \tilde{l}$, $\tilde{l} \leq m-7$ and $l + \tilde{l} \leq m-5$ implies that $|S| \leq n(m-2) - m + 7$.

Moreover, if there are vertices $h \in N_{G-\{u_2, u_3\}}(u_1) - N_{G-\{u_2, u_4\}}(u_5)$ or $\dot{h} \in N_{G-\{u_2, u_4\}}(u_5) - N_{G-\{u_2, u_3\}}(u_1)$ then $\kappa'(G) < \infty$, a contradiction. Let $\{h_1, \dots, h_f\} = N_{G-\{u_2, u_3\}}(u_1) \cap N_{G-\{u_2, u_4\}}(u_5)$. If for some i, j , $h_i \sim h_j$

or $\deg(u_3) \geq 4$ or $\deg(u_4) \geq 4$ then $\kappa'(G) < \infty$, a contradiction. Hence let for every i, j , $h_i h_j \notin E(G)$, $\deg(u_3) = \deg(u_4) = 3$ and $N_{G-\{u_1, u_3, u_4, u_5\}}(u_2) = \{x_{r'} \mid r' \in \mathbb{Z}_r\}$. So $S = \{(u_3, v_i), (u_4, v_i) \mid i \in \mathbb{Z}_n\} \cup \{(u_1, v_i), (u_2, v_i), (u_5, v_i) \mid i \in \mathbb{Z}_n - \{1\}\} \cup \{(h_j, v_1), (x_{r'}, v_1) \mid j \in \mathbb{Z}_f, r' \in \mathbb{Z}_r\}$ is a super vertex-cut in $G \times K_n$ with $|S| = 2n + 3(n-1) + (m-5) = 5n + m - 8$.

Now, let $\deg(u_2) = 4$, $\deg(u_3) = 3$, $\deg(u_4) = 3$ and $u_1 \sim u_5$. The graph is W_4 and has four triangles. In this case, $S = \{(u_2, v_i) \mid i \in \mathbb{Z}_n\} \cup \{(u_j, v_i) \mid j \in \mathbb{Z}_5 - \{2\}, i \in \mathbb{Z}_n - \{1, 2\}\}$ is a super vertex-cut in $G \times K_n$ with $|S| = n + 4(n-2) = 5n - 8$. Furthermore, let $u_1 \sim u_5$, $\deg(u_2) \geq 4$, $\deg(u_3) \geq 3$ and $\deg(u_4) \geq 3$. Let $N_{G-\{u_1, u_3, u_4, u_5\}}(u_2) = \{x_{r'} \mid r' \in \mathbb{Z}_r\}$, $N_{G-\{u_1, u_2, u_4\}}(u_3) = \{y_{s'} \mid s' \in \mathbb{Z}_s\}$ and $N_{G-\{u_2, u_3, u_5\}}(u_4) = \{w_{t'} \mid t' \in \mathbb{Z}_t\}$ where $r+s+t = m-5$, and $\deg(x_{r'}) = \deg(y_{s'}) = \deg(w_{t'}) = 1$ for each r', s', t' . Then $S = \{(u_j, v_i) \mid 2 \leq j \leq 4, i \in \mathbb{Z}_n - \{1\}\} \cup \{(u_5, v_i), (u_1, v_i) \mid i \in \mathbb{Z}_n\} \cup \{(x_{r'}, v_1), (y_{s'}, v_1), (w_{t'}, v_1) \mid r' \in \mathbb{Z}_r, s' \in \mathbb{Z}_s, t' \in \mathbb{Z}_t\}$ is a super vertex-cut in $G \times K_n$ with $|S| = 3(n-1) + 2n + (m-5) = 5n - 8 + m$. Now if $x_{r_i'} \sim u_3$ or $x_{r_j'} \sim u_4$ for some i, j , then by using A, B, C, D and F as above, and putting $\{(u_1, v_i), (u_5, v_i) \mid i \in \mathbb{Z}_n\}$ in S we get S a super vertex-cut in $G \times K_n$ with $|S| \leq n(m-2) - m + 7 + 2n - 2 = m(n-1) + 5$.

Now let G be the graph W_5 with vertices $\{u_i\}_{i \in \mathbb{Z}_6}$ where $u_1 \sim u_3 \sim u_4 \sim u_5 \sim u_6 \sim u_1$ and u_2 is adjacent to all others. Hence by putting $\{(u_2, v_i) \mid i \in \mathbb{Z}_n\}$ in S and using Theorem 2.2, we get $|S| = \min\{4n, 6n-8\}$. Now let $\deg(u_2) \geq 6$, $N_{G-\{u_1, u_3, u_4, u_5, u_6\}}(u_2) = \{x_{r'} \mid r' \in \mathbb{Z}_r\}$. If for some i, j , $x_{r_i'} \sim x_{r_j'}$ or $x_{r_i'} \sim u_s$ for $s \in \mathbb{Z}_n - \{2\}$, then $\kappa'(G) < \infty$, a contradiction. Let for all $r' \in \mathbb{Z}_r$, $\deg(x_{r'}) = 1$. Then $S = \{(u_1, v_i), (u_2, v_i), (u_5, v_i) \mid i \in \mathbb{Z}_n - \{1, 2\}\} \cup \{(u_3, v_i), (u_4, v_i), (u_6, v_i) \mid i \in \mathbb{Z}_n\} \cup \{(x_{r'}, v_i) \mid r' \in \mathbb{Z}_r, i \in \mathbb{Z}_n - \{1, 2\}\}$ is a super vertex-cut in $G \times K_n$ with $|S| = m(n-2) + 6$. Also let $u_1 \sim u_4$ or $u_1 \sim u_5$ or $u_3 \sim u_5$ or $u_3 \sim u_6$ or $u_4 \sim u_6$ where $\deg(u_2) = 5$. Since G is not complete we may assume that $\deg(u_1) = \deg(u_4) = 4$ and $u_1 u_4 \notin E(G)$. Thus $S = \{(u_2, v_i), (u_5, v_i), (u_6, v_i) \mid i \in \mathbb{Z}_n\} \cup \{(u_1, v_i), (u_3, v_i), (u_4, v_i) \mid i \in \mathbb{Z}_n - \{1, 2\}\}$ is a super vertex-cut in $G \times K_n$ with $|S| = 6n-6$. Similarly, for any non complete graph G with vertices $\{x_1, \dots, x_m\}$ and $\text{girth}(G) = 3$, if $x_1 x_3 \notin E(G)$ and $x_1 \sim x_2 \sim x_3$, then $S = \{(x_j, v_i) \mid j \in \mathbb{Z}_m - \{1, 2, 3\}, i \in \mathbb{Z}_n\} \cup \{(x_j, v_i) \mid j \in \mathbb{Z}_3, i \in \mathbb{Z}_n - \{1, 2\}\}$ is a super vertex-cut in $G \times K_n$ with $|S| = (m-3)n + 3(n-2) = mn - 6$. Also, $\kappa'(W_n) < \infty$ where $n \geq 6$. If a graph G includes W_n with more than $2n$ vertices where $\kappa'(G) = \infty$ then by above argument we see that the cardinality of a super vertex-cut is $mn - 6$. Therefore, if $\text{girth}(G) = 3$ and $\kappa'(G) = \infty$ then, $\kappa'(G \times K_n) \leq \min\{mn - 6, m(n-1) + 5, 5n + m - 8\}$. \square

3. CONCLUSION

In this article we provide bounds for the super connectivity of the direct product of an arbitrary graph and the complete graph K_n . Also, we show that if G is a non-complete graph with $\text{girth}(G) = 3$ and $\kappa'(G) = \infty$, then $\kappa'(G \times K_n) \leq \min\{mn - 6, m(n-1) + 5, 5n + m - 8\}$, where $|V(G)| = m$.

Acknowledgements. We thank the anonymous referees for their careful review and very constructive suggestions and comments which helped to improve the content of this article.

REFERENCES

- [1] J.A. Bondy and U.S.R. Murty, Graph Theory with Applications. Elsevier, New York (1967).
- [2] G.B. Ekinci and J.B. Gauci, The super-connectivity of odd graphs and of their kronecker double cover. *RAIRO-Oper. Res.* **55** (2021) 561–566.
- [3] G.B. Ekinci and A. Kırlangıç, Super connectivity of kronecker product of complete bipartite graphs and complete graphs. *Disc. Math.* **339** (2016) 1950–1953.
- [4] G.B. Ekinci and A. Kırlangıç, The super edge connectivity of Kronecker product graphs. *RAIRO-Oper. Res.* **52** (2018) 561–566.
- [5] S. Ghozati, A finite automata approach to modeling the cross product of interconnection networks. *Math. Compute. Model.* **30** (1999) 185–200.
- [6] R. Guji and E. Vumar, A note on the connectivity of kronecker products of graphs. *Appl. Math. Lett.* **22** (2009) 1360–1363.
- [7] L. Guo and X. Guo, Super connectivity of Kronecker products of some graphs. *Ars Combin.* **123** (2015) 65–73.
- [8] L. Guo, C. Qin and X. Guo, Super connectivity of kronecker products of graphs. *Inf. Process. Lett.* **110** (2010) 659–661.
- [9] L. Guo, W. Yang and X. Guo, Super-connectivity of kronecker products of split graphs, powers of cycles, powers of paths and complete graphs. *Appl. Math. Lett.* **26** (2013) 120–123.

- [10] Kh. Kamyab, M. Ghasemi and R. Varmazyar, Super connectivity of lexicographic product graphs. *Ars Combin.* Preprint [arXiv:2009.04831 \[math.GR\]](https://arxiv.org/abs/2009.04831).
- [11] R. Lammprey and B. Barnes, Products of graphs and applications. *Model. Simul.* **5** (1974) 1119–1123.
- [12] J. Leskovec, D. Chakrabarti, J. Kleinberg, C. Faloutsos and Z. Ghahramani, Kronecker graphs: An approach to modeling networks. *J. Mach. Learn. Res.* **11** (2010) 985–1042.
- [13] M. Lü, C. Wu, G.L. Chen and C. Lv, On super connectivity of cartesian product graphs. *Networks* **52** (2008) 78–87.
- [14] M. Ma, G. Liu and J.M. Xu, The super connectivity of augmented cubes. *Inf. Process. Lett.* **106** (2008) 59–63.
- [15] D.J. Miller, The categorical product of graphs. *Can. J. Math.* **20** (1968) 1511–1521.
- [16] F. Soliemany, M. Ghasemi and R. Varmazyar, Super connectivity of a family of direct product graphs. *Int. J. Comput. Math. Comput. Syst.* **7** (2021) 1–5.
- [17] P.M. Weichsel, The kronecker product of graphs. *Proc. Am. Math. Soc.* **13** (1962) 47–52.
- [18] J.M. Xu, *Topological Structure and Analysis of Interconnection Networks*. Kluwer Academic Publishers, Dordrecht (2001).
- [19] J.M. Xu, M. Lü, M. Ma and A. Hellwig, Super connectivity of line graphs. *Inf. Process. Lett.* **94** (2005) 191–195.
- [20] C.S. Yang, J.F. Wang, J.Y. Lee and F.T. Boesch, Graph theoretic reliable analysis for the Boolean n -cube networks. *IEEE Trans. Circuits Syst.* **35** (1988) 1175–1179.
- [21] C.S. Yang, J.F. Wang, J.Y. Lee and F.T. Boesch, The number of spanning trees of the regular networks. *Int. J. Comput. Math.* **23** (1988) 185–200.
- [22] J.X. Zhou, Super connectivity of direct product of graphs. *Ars Math. Contemp.* **8** (2015) 235–244.

Subscribe to Open (S2O)

A fair and sustainable open access model



This journal is currently published in open access under a Subscribe-to-Open model (S2O). S2O is a transformative model that aims to move subscription journals to open access. Open access is the free, immediate, online availability of research articles combined with the rights to use these articles fully in the digital environment. We are thankful to our subscribers and sponsors for making it possible to publish this journal in open access, free of charge for authors.

Please help to maintain this journal in open access!

Check that your library subscribes to the journal, or make a personal donation to the S2O programme, by contacting subscribers@edpsciences.org

More information, including a list of sponsors and a financial transparency report, available at: <https://www.edpsciences.org/en/math-s2o-programme>