BULLETIN DE LA S. M. F.

MICHEL ZINSMEISTER

A distortion theorem for quasiconformal mappings

Bulletin de la S. M. F., tome 114 (1986), p. 123-133

<http://www.numdam.org/item?id=BSMF_1986__114__123_0>

© Bulletin de la S. M. F., 1986, tous droits réservés.

L'accès aux archives de la revue « Bulletin de la S. M. F. » (http://smf.emath.fr/Publications/Bulletin/Presentation.html) implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/conditions). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
http://www.numdam.org/

Bull. Soc. math. France,
114, 1986, p. 123-133

A DISTORTION THEOREM FOR QUASICONFORMAL MAPPINGS

BY

MICHEL ZINSMEISTER (*)

RÉSUMÉ. — Un théorème de distorsion pour les applications quasiconformes. Nous généralisons aux applications quasiconformes dans la boule unité de $\mathbb{R}^n$ un théorème de Pommerenke relatif aux transformations conformes du disque unité du plan.

ABSTRACT. — A distortion theorem for quasiconformal mappings. We extend to quasiconformal mappings in the unit ball of $\mathbb{R}^n$ a theorem of Pommerenke concerning conformal mappings in the unit disk of the plane.

1. Introduction

(*a*) The purpose of this paper is to extend to quasi-conformal mappings the following distortion theorem for conformal mappings in the unit disk B^2 of $\mathbb{R}^2$, a theorem due to Pommerenke. In this statement, $I(z)$ is, for $z \in B^2$, the interval of B^2 centered at $z/|z|$ of length $2\pi(1-|z|)$.

THÉORÈME [9]. — *There is a universal constant $C>0$ such that if $f: B^2 \to \mathbb{R}^2$ is a conformal mapping, then, for every $z \in B^2$, there exists a non euclidean segment γ from z to $I(z)$ such that*

$$\text{Length}(f(\gamma)) \leqq C \,\text{distance}(f(z), \partial f(B^2)).$$

Before giving the precise results, we set some preliminary notations and results.

For $x \in \mathbb{R}^n$, $n \geqq 2$, let $|x|$ be the euclidean norm of x. For $r>0$,

$$B^n(x, r) = \{y \in \mathbb{R}^n;\ |y-x| < r\},$$
$$B^n = B^n(0, 1) \quad \text{and} \quad S^{n-1} = \partial B^n.$$

(*) Texte reçu le 14 mars 1985.

M. ZINSMEISTER, Mathématiques, Université de Rouen, 76130 Mont St-Aignan Cedex.

BULLETIN DE LA SOCIÉTÉ MATHÉMATIQUE DE FRANCE — 0037-9484/1986/01 123 11/$ 3.10

© Gauthier-Villars

If $E \subset \mathbb{R}^n$ and $x \in \mathbb{R}^n$, $d(x, E)$ is the distance from x to E. If $E \subset \mathbb{R}^n$ is measurable, we note $m(E)$ its Lebesgue measure; σ stands for Lebesgue measure on S^{n-1} and $\sigma_{n-1} = \sigma(S^{n-1})$. For a real $n \times n$ matrix A, let

$$|A| = \sup_{x \in S^{n-1}} |Ax|.$$

If $\Omega \subset \mathbb{R}^n$ is open and $f: \Omega \to \mathbb{R}^n$ is in the Sobolev space $W^1_{n\,\mathrm{loc}}(\Omega)$, $Df(x)$ will denote the Jacobian matrix of f, defined almost everywhere, and $J(x, f) = \det Df(x)$.

For $K \geqq 1$, a continuous one-to-one mapping $f: \Omega \to \mathbb{R}^n$ is said to be *K-quasiconformal* if $f \in W^1_{n\,\mathrm{loc}}(\Omega)$ and if

$$|Df(x)|^n \leqq KJ(x, f) \quad \text{a. e. on } \Omega.$$

For $x \in S^{n-1}$, we define the cone with vertex x as

$$\Gamma(x) = \{ y \in B^n; |y - x| < 3(1 - |y|) \}.$$

If $F: B^n \to \mathbb{R}^n$ is any function, the non-tangential maximal function of F is defined as

$$\forall x \in S^{n-1}, \quad F^*(x) = \sup_{y \in \Gamma(x)} |F(y)|.$$

If $z \in B^n$, we define the « cap » $S(z)$ as

$$S(z) = \{ x \in S^{n-1};\ z \in \Gamma(x) \} = S^{n-1} \cap B^n(z,\ 3(1 - |z|)).$$

Let M be the group of Möbius self-maps of B^n. If $z \in B^n$, $z \neq 0$, we define

$$T_z(x) = \frac{(1 - |z|^2)(x - z) - |x - z|^2 z}{|z|^2 |x - (z/|z|^2)|^2};$$

then $T_z \in M$ and $T_z(z) = 0$.

We will need the following elementary results, the proof of which we omit:

(1) $$S(z) = S^{n-1} \quad \text{if} \quad |z| < 1/2.$$

(2) $T_z(S(z))$ always contains an hemisphere,

(3) If $x, y \in S(z)$,

$$(9(1 - |z|))^{-1} |x - y| \leqslant |T_z(x) - T_z(y)| \leqslant 2(1 - |z|)^{-1} |x - y|,$$

(4) $$\forall z \in B^n, \quad B^n(0, 1/7) \subset T_z\left(B^n\left(z, \frac{1}{4}(1 - |z|) \right) \right) \subset B^n(0, 1/2).$$

(*b*) The main result will be the following.

THEOREM 1. — *For $K \geqslant 1$ there exists a constant $C(K, n) > 0$ such that if $f\, B^n \to \mathbb{R}^n$ is K-quasiconformal, then, for any $z \in B^n$, there exists a non-euclidean segment γ joining z to $S(z)$ such that*

$$\text{Length}(f(\gamma)) \leqslant C(K, n)\, d(f(z), \partial f(B^n)). \tag{5}$$

Adapting an idea of B. DAVIS and J. LEWIS [2], we will show in a moment that Theorem 1 is a corollary of the following, of independent interest:

THEOREM 2. — *For f: $B^n \to \mathbb{R}^n$ k-quasiconformal, define, for $x \in S^{n-1}$, $L_f(x) = \text{Length}\,[f([0, x])]$ where $[0, x]$ is the radius $\{tx;\ 0 \leqslant t \leqslant 1\}$. Then there exist $C(n, K) > 0$ and $p(n, K) > 0$ such that*

$$\left(\int_{S^{n-1}} L_f(x)^p \, d\sigma(x) \right)^{1/p} \leqslant C(K, n)\, d(f(0), \partial f(B^n)). \tag{6}$$

Assuming Theorem 2 is true, let us prove Theorem 1. So let $f: B^n \to \mathbb{R}^n$ be a K-quasiconformal mapping and $z \in B^n$; put $g = f \circ T_z^{-1}$: Applying Theorem 2 to g, we see that for every $M > 0$,

$$\sigma(\{x \in S^{n-1};\ \text{Lg}(x) > M\, d(f(z), \partial f(B^n))\}) \leqslant \left(\frac{C(K, n)}{M} \right)^p. \tag{7}$$

Now choose M large enough so that $(C/M)^p \leqslant (\sigma_{n-1})/4$. By (2) there exists then $x \in T^z(S(z))$ such that

$$\text{Lg}(x) \leqslant M\, d(f(z), \partial f(B^n)),$$

and this proves (5) with $\gamma = T_z^{-1}([0, x])$.

The main tools in proving Theorem 2 will be Theorem 3, due to P. Jones, which we discuss in Part 2, and an estimate for the nontangential maximal function f^*, which is proved in part 3.

This paper was written during a stay at the University of Michigan. I would like to thank Professor F. Gehring for his invitation and the constant help he gave me during my stay. I also would like to thank Professor T. Iwaniec for many helpful conversations.

2. Let f be as in Theorem 2. Performing a preliminary translation, we may assume that f does not vanish in B^n and that

$$|f(0)| = d(f(0), \partial f(B^n)).$$

We will say that a function $g : B^n \to \mathbb{R}^n - \{0\}$ satisfies Harmack property if there exists a constant $C(g) > 0$, called the Harnack constant of g, such that:

$$(8) \quad \forall x \in B^n, \quad \forall y, \quad z \in B^n\left(x, \frac{1}{4}(1-|x|)\right), \qquad |g(y)| \leqslant C(g)|g(z)|.$$

LEMMA 1. — *If f: $B^n \mathbb{R}^n - \{0\}$ is K-quasiconformal, then f satisfies Harnack property with a constant depending only on K and n.*

Proof. — By the special distortion theorem for quasiconformal mappings [4], there exists $C(K, n) > 0$ such that

$$\forall x \in B^n, \quad \forall y, \quad z \in B^n\left(x, \frac{1}{4}(1-|x|)\right), \qquad \frac{|f(y)-f(z)|}{d(f(z), \partial f(B^n))} \leqslant C(K, n),$$

and Lemma 1 follows, for $d(f(z), \partial f(B^n)) \leqslant |f(z)|$ since f does not vanish in B^n.

Now let f: $B^n \to \mathbb{R}^n - \{0\}$ be K-quasiconformal. For almost every $x \in S^{n-1}$, we may write

$$L_f(x) \leqslant \int_0^1 |Df(tx)|\, dt \leqslant V_f(x) + 2^{n-1} f^*(x) H_f(x),$$

where

$$V_f(x) = \int_0^{1/2} |Df(tx)|\, dt$$

and

$$H_f(x) = \int_0^1 \frac{|Df(tx)|}{|f(tx)|} t^{n-1}\, dt.$$

LEMMA 2:

$$V_f \in L^1(S^{n-1}) \qquad \text{with} \quad \|V_f\|_1 \leqslant C(K, n)|f(0)|.$$

Proof:

$$\| V_f \|_1 = \int_{B^n(0,\,1/2)} \frac{|Df(y)|}{|y|^{n-1}}\, dm(y):$$

By Gehring's inequality [5], there exists $p=p(K,\,n)>n$ and $C(K,\,n)>0$ such that

$$(9)\quad \left(\int_{B^n(0,\,1/2)} |Df(y)|^p\, dm(y)\right)^{1/p} \leqslant C(K,\,n)\left(\int_{B^n(0,\,1/2)} J(x,\,f)\, dm(x)\right)^{1/n} = C(K,\,n)\, m[f(B^n(0,\,1/2)]^{1/n} \leqslant C(K,\,n)\,|f(0)|,$$

the last inequality being a consequence of Lemma 1. We now apply Hölder's inequality to the expression of $\| V_f \|_1$, to obtain

$$\| V_f \|_1 \leqslant \left(\int_{B^n(0,\,1/2)} |Df(y)|^p\, dm(y)\right)^{1/p} \times \left(\int_{B^n(0,\,1/2)} |y|^{-(n-1)p/(p-1)}\, dm(y)\right)^{(p-1)/p} \leqslant C(K,\,n)\,|f(0)|,$$

by (9) and the fact that $(n-1)p/(p-1)<n$.

An estimate for H_f is given by the following theorem, due to P. Jones:

THEOREM 3 [8]:

$$H_f \in L^1(S^{n-1}) \qquad \textit{with} \quad \| H_f \|_1 \leqslant C(K,\,n).$$

As P. Jones has observed, Theorem 3 implies that

$$\sup_{T \in M} \| H_{f \circ T} \|_1 \leqslant C(K,\,n) \quad \Leftrightarrow$$

$$(10)\qquad \sup_{T \subset M} \int_{B^n} \frac{|Df(x)|}{|f(x)|}\, |DT(x)|^{n-1}\, dm(x) \leqslant C(K,\,n),$$

and (10) exactly says that $|Df|/|f|\,dm$ is a Carleson measure in B^n [3]. Since $|\nabla |f||\leqslant |Df|$, the same is true for $|\nabla u|\,dm$, where $u=\log|f|$. Writing

$$\| u \|_* = \sup_{T \in M} \int_{B^n} |\nabla u(x)|\, |DT(x)|^{n-1}\, dm(x),$$

we can now invoke the following theorem, due to Varopoulos:

THEOREM 4 [10]. — *Let $u \in W^1_{1\,loc}(B^n)$ be a real-valued function having radical limit $\tilde{u}(x)$ a. e. on S^{n-1}. If $|\nabla u| dm$ is a Carleson measure in B^n, then $\tilde{u} \in BMO(S^{n-1})$ with $\|\tilde{u}\|_{BMO} \leqslant C(n)\|u\|_*$.*

(Varopoulos proves Theorem 4 for $n=2$ only, but his argument is easily seen to extend to the general case.)

From Theorems 3 and 4, it follows that if $f: B^n \to \mathbb{R}^n - \{0\}$ is K-quasiconformal, then

$$\| \operatorname{Log} |f(rx)| \|_{BMO(S^{n-1})} \leqslant C(K, n),$$

for $0 \leqslant r \leqslant 1$. If we now apply the John and Nirenverg inequality [7] and Lemma 1, we get

LEMMA 3. — *If $f: B^n \to \mathbb{R}^n - \{0\}$ is K-quasiconformal there exist $C(K, n) > 0$ and $p = p(K, n) > 0$ such that*

$$\sup_{0<r\leqslant 1} \int_{S^{n-1}} |f(rx)|^p \, d\sigma(x) \leqslant C(K, n) |f(0)|^p. \tag{11}$$

In the case $n=2$ and f conformal, (11) implies that f is in the Hardy space $H^p(B^2)$, so that $f^* \in L^p(S^{n-1})$, and Theorem 2 follows in this particular case. In the general case, we need an extra argument, provided by the next section.

3. Non-tangential maximal function

PROPOSITION 1. — *Let $f: B^n \to \mathbb{R}^n - \{0\} \in W^1_{1\,loc}(B^n)$, and $u = \operatorname{Log}|f|$. If $|\nabla u| dm$ is a Carleson measure and if f satisfies Harnack property (8) then, for every $p > 0$,*

$$\int_{S^{n-1}} f^*(x)^p \, d\sigma(x) \leqslant C_p \int_{S^{n-1}} |f(x)|^p \, d\sigma(x),$$

where C_p depends only on n, p, $\|u\|_*$ and $C(f)$.

We first notice that this statement makes sense, since $|f(x)|$ has radial limits a. e. on S^{n-1}, for $|\nabla u| dm$ is a Carleson measure.

Before going into the proof of Proposition 1, let us see why it implies Theorem 2. So let $f: B^n \to \mathbb{R}^n - \{0\}$ be K-quasiconformal with

$|f(0)| = \text{dist}\,(f(0), f(B^n))$. By the results in Part 2 we can apply Proposition 1 to $f(x)$ and, by Lemma 3, we get

$$\|f^*\|_p \leqslant C(K, n)|f(0)|,$$

for some p depending only on K and n. Recalling now that

$$L_f(x) \leqslant V_f(x) + 2^{n-1} f^*(x) H_f(x),$$

Lemma 2, Theorem 3 and (12) imply that $L_f \in L^{p/p+1}(S^{n-1})$ with

$$\|L_f\|_{p/p+1} \leqslant C(K, n)|f(0)|,$$

and Theorem 2 is proved.

To prove Proposition 1, we need 3 lemmas:

LEMMA 4. — *If f is as in Proposition* 1 *and* $N > C(f)^2$,

$$\sigma\left(\left\{x \in S^{n-1};\ |f(x)| \leqslant \frac{|f(0)|}{N}\right\}\right) \leqslant \frac{7^n \|u\|_*}{\operatorname{Log} N}.$$

Proof: Put

$$F_N = \left\{x \in S^{n-1};\ |f(x)| < \frac{|f(0)|}{N}\right\}$$

and

$$G(x) = \int_0^1 |\nabla u(tx)|\, t^{n-1}\, dt.$$

If $x \in F_N$,

$$G(x) \geqslant 7^{1-n} \int_{1/7}^1 |\nabla u(tx)|\, dt$$

$$\geqslant 7^{1-n} \left| \int_{1/7}^1 |f(tx)|^{-1} \frac{\partial}{\partial t}(|f(tx)|)\, dt \right|$$

$$= 7^{1-n} \left| \operatorname{Log} \frac{|f(x)|}{|f(x/7)|} \right|.$$

By Harnack property, $|f(0)| \leqslant C(f)|f(x/7)|$; so, if $x \in F_N$ and $N > C(f)^2$,

$$G(x) \geqslant 7^{1-n} \left| \operatorname{Log} \frac{N}{C(f)} \right| \geqslant 7^{-n} \operatorname{Log} N,$$

and from this it follows that

$$\sigma(F_N) \leqslant \frac{7^n \|G\|_1}{\operatorname{Log} N} \leqslant \frac{7^n \|u\|_*}{\operatorname{Log} N}.$$

LEMMA 5. — *There is an universal constant* $\alpha > 0$ *such that if* f *is as in Proposition* 1 *and* $z \in B^n$,

$$\sigma\left(\left\{x \in S(z); |f(x)| \leqslant \frac{|f(z)|}{N}\right\}\right) \leqslant \frac{\alpha 7^n \|u\|_*}{\operatorname{Log} N} \sigma(S(z)).$$

Proof. — Define $g(x) = f \circ T_z^{-1}(x)$. By (4), if $x \in S^{n-1}$ then $|g(0)| \leqslant C(f)|g(x/7)|$; also, by definition $\|\operatorname{Log}|g|\|_* = \|u\|_*$. By Lemma 4 we then have, for $N \geqslant C(f)^2$,

$$\sigma\left(\left\{x \in S^{n-1}; g(x) \leqslant \frac{|f(z)|}{N}\right\}\right) \leqslant \frac{7^n \|u\|_*}{\operatorname{Log} N},$$

which implies

$$\sigma\left(T_z\left(\left\{y \in S(z); |f(y)| \leqslant \frac{|f(z)|}{N}\right\}\right)\right) \leqslant \frac{7^n \|u\|_*}{\operatorname{Log} N},$$

and the result follows from (3).

LEMMA 6. — *There exist* $0 < C(n) < 1$ *and* $N(n, \|u\|_*, C(f))$ *such that the following inequality holds*:

$$(13) \quad \forall \lambda > 0, \quad \sigma\left(\left\{x \in S^{n-1}; f^*(x) > \lambda, |f(x)| \leqslant \frac{\lambda}{N}\right\}\right) \leqslant C(n)\, \sigma(\{x \in S^{n-1}; f^*(x) \geqslant \lambda\}).$$

Proof. — Let

$$\mathscr{E}(\lambda) = \{z \in B^n, |f(z)| > \lambda\}$$

and

$$\mathscr{U}(\lambda) = \{x \in S^{n-1}; f^*(x) > \lambda\}.$$

Then

$$\mathscr{U}(\lambda) = \cup_{z \in \mathscr{E}(\lambda)} S(z).$$

By Vitali covering lemma, there exists $\alpha(n)\in(0,1)$ and a sequence $\{z_j\}\subset\mathscr{E}(\lambda)$ such that the $S(z_j)$'s are mutually disjoint and

$$\sum_{j\in\mathbf{N}}\sigma(S(z_j))\geqslant\alpha(n)\,\sigma(\mathscr{U}(\lambda)).$$

Now let

$$E_N=\left\{x\in S^{n-1};\ |f(x)|\leqslant\frac{\lambda}{N}\right\}.$$

Then

$$\sigma(E_N\cap\mathscr{U}(\lambda))\leqslant\sum_{j\in\mathbf{N}}\sigma(E_N\cap S(z_j))+(1-\alpha(n))\,\sigma(\mathscr{U}(\lambda)).$$

But

$$E_N\cap S(z_j)\subset\left\{x\in S(z_j);\ |f(x)|\leqslant\frac{|f(z_j)|}{N}\right\}$$

and so

$$\sigma(E_N\cap S(z_j))\leqslant\frac{\alpha 7^n\|u\|_*}{\operatorname{Log}N}\sigma(S(z_j))\qquad\text{if}\quad N\geqslant C(f)^2,$$

by Lemma 5. So, choosing N so large that

$$\frac{\alpha 7^n\|u\|_*}{\operatorname{Log}N}<\frac{\alpha(n)}{2},$$

we get Lemma 6 with $C(n)=1-\alpha(n)/2$. We can now complete the proof of Proposition 1. For $\lambda>0$, let

$$\chi(\lambda)=\sigma(\{x\in S^{n-1};\ f^*(x)>\lambda\})$$

and

$$\theta(\lambda)=\sigma(\{x\in S^{n-1};\ |f(x)|>\lambda\}).$$

By (13),

$$\chi(\lambda)\leqslant C(n)\chi(\lambda)+\theta(\lambda/N)\quad\Leftrightarrow\quad\chi(\lambda)\leqslant\frac{1}{1-C(n)}\theta(\lambda/N).$$

Finally,

$$\|f^*\|_p^p=p\int_0^\infty\lambda^{p-1}\chi(\lambda)\,d\lambda\leqslant\frac{p}{1-C(n)}\int_0^\infty\lambda^{p-1}\theta\left(\frac{\lambda}{N}\right)d\lambda=\frac{N^p}{1-C(n)}\|f\|_p^p,$$

which proves Proposition 1.

Remark. — A mapping $f: B^n \to \mathbb{R}^n$ is said to be K-quasiregular if $f \in W^1_{n\,\mathrm{loc}}(B^n)$ and if

$$|Df(x)|^n \leqslant KJ(x, f) \quad \text{a. e. on } B^n.$$

The above methods imply the following:

PROPOSITION 2. — *Let $f: B^n \to \mathbb{R}^n - \{0\}$ be a K-quasiregular mapping and $u = \mathrm{Log}|f|$. If $|\nabla u|\,dm$ is a Carleson measure in B^n, there exists an exponent $p = p(n, K, \|u\|_*) > 0$ such that $L_f \in L^p(S^{n-1})$, where L_f has the same meaning as in Theorem* 1.

Sketch of Proof. — Using the same notations as in 2, one may write

$$L_f(x) \leqslant V_f(x) + 2^{n-1} K f^*(x) \tilde{H}_f(x),$$

where

$$\tilde{H}_f(x) = \int_0^1 |\nabla u(tx)|\, t^{n-1}\, dt.$$

From results in [1], Lemma 2 is valid for V_f; also, $\tilde{H}_f \in L^1$ by hypothesis. To prove Proposition 2, it suffices then to show that $f^* \in L^p$ for some $p > 0$. Using Proposition 1 and Theorem 4, this reduces to proving that f satisfies Harnack property. To do this, we use a recent result of IWANIEC and NOLDER [6]: They proved that u is actually in $W^1_{q\,\mathrm{loc}}(B^n)$ for some $q(K, n) > n$ and that

$$\forall x \in B^n, \quad \left(\int_{B^n(x,\, 1/2\,(1-|x|))} |\nabla u(y)|^q\, dm(y) \right)^{1/q} \leqslant C(K, n)(1-|x|)^{n(1/q-1)} \times \int_{B^n(x,\, 3/4\,(1-|x|))} |\nabla u(y)|\, dm(y),$$

and the result follows from the Sobolev embedding theorem and the fact that $|\nabla u|\,dm$ is a Carleson measure.

REFERENCES

[1] BOJARSKI (B.) and IWANIEC (T.). – Analytical foundations of the theory of quasiconformal mappings in $\mathbb{R}^n$, *Ann. Acad. Sc. Fenn.*8 (1983) p. 257-324.

[2] DAVIS (B.) and LEWIS (J.). – Paths for subharmonic functions, *Proc. London Math. Soc.*, Vol. XLVIII, 1984, p. 401-428.

[3] GARNETT (J.), *Bounded analytic functions*, Academic Press, 1981.

[4] GEHRING (F.), Rings and quasiconformal mappings in space, *Trans. AMS*, Vol. 103, 1962, p. 353-393.

[5] GEHRING (F.). – The L^p integrability of the partial derivatives of a quasiconformal mapping, *Acta Math.*, Vol. 130, 1973, pp. 265-277.

[6] IWANIEC (T.) and NOLDER (C.). – Hardy-Littlewood inequality for quasiregular mappings in certain domains in $\mathbb{R}^n$, *Ann. Acad. Sc. Fenn.* (to appear).

[7] JOHN (F.) and NIRENBERG (L.). – On functions of bounded mean oscillation, *Comm. Pure Appl. Math.*, Vol. 14, 1961, pp. 415-426.

[8] JONES (P.). – Extension theorems for BMO, *Indiana J. Math.*, Vol. 29, 1980, pp. 41-66.

[9] POMMERENKE (C.). – *Univalent functions*, Vandenhoeck and Ruprecht, Göttingen, 1975.

[10] VAROPOULOS (N.). – *BMO* functions and the $\bar{\partial}$ equation, *Pacific J. Math.*, Vol. 71, 1977, pp. 221-273.