

Bulletin

de la SOCIÉTÉ MATHÉMATIQUE DE FRANCE

ON CUBIC BIRATIONAL MAPS OF $\mathbb{P}_{\mathbb{C}}^3$

Julie Déserti & Frédéric Han

Tome 144
Fascicule 2

2016

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Publié avec le concours du Centre national de la recherche scientifique

pages 217-249

Le *Bulletin de la Société Mathématique de France* est un
périodique trimestriel de la Société Mathématique de France.

Fascicule 2, tome 144, juin 2016

Comité de rédaction

Valérie BERTHÉ	Marc HERZLICH
Gérard BESSON	O'Grady KIERAN
Emmanuel BREUILLARD	Julien MARCHÉ
Yann BUGEAUD	Emmanuel RUSS
Jean-François DAT	Christophe SABOT
Charles FAVRE	Wilhelm SCHLAG
Raphaël KRIKORIAN (dir.)	

Diffusion

Maison de la SMF	Hindustan Book Agency	AMS
Case 916 - Luminy	O-131, The Shopping Mall	P.O. Box 6248
13288 Marseille Cedex 9	Arjun Marg, DLF Phase 1	Providence RI 02940
France	Gurgaon 122002, Haryana	USA
smf@smf.univ-mrs.fr	Inde	www.ams.org

Tarifs

Vente au numéro : 43 € (\$ 64)

Abonnement Europe : 178 €, hors Europe : 194 € (\$ 291)

Des conditions spéciales sont accordées aux membres de la SMF.

Secrétariat : Nathalie Christiaën

Bulletin de la Société Mathématique de France

Société Mathématique de France

Institut Henri Poincaré, 11, rue Pierre et Marie Curie

75231 Paris Cedex 05, France

Tél : (33) 01 44 27 67 99 • Fax : (33) 01 40 46 90 96

revues@smf.ens.fr • <http://smf.emath.fr/>

© Société Mathématique de France 2016

Tous droits réservés (article L 122-4 du Code de la propriété intellectuelle). Toute représentation ou reproduction intégrale ou partielle faite sans le consentement de l'éditeur est illicite. Cette représentation ou reproduction par quelque procédé que ce soit constituerait une contrefaçon sanctionnée par les articles L 335-2 et suivants du CPI.

ISSN 0037-9484

Directeur de la publication : Marc PEIGNÉ

ON CUBIC BIRATIONAL MAPS OF $\mathbb{P}_{\mathbb{C}}^3$

BY JULIE DÉSERTI & FRÉDÉRIC HAN

ABSTRACT. — We study the birational maps of $\mathbb{P}_{\mathbb{C}}^3$. More precisely we describe the irreducible components of the set of birational maps of bidegree $(3, 3)$ (resp. $(3, 4)$, resp. $(3, 5)$).

RÉSUMÉ (*Sur les transformations birationnelles cubiques de $\mathbb{P}_{\mathbb{C}}^3$*)

Nous étudions les transformations birationnelles de $\mathbb{P}_{\mathbb{C}}^3$. Plus précisément nous décrivons les composantes irréductibles de l'ensemble des transformations birationnelles de $\mathbb{P}_{\mathbb{C}}^3$ de bidegré $(3, 3)$ (resp. $(3, 4)$, resp. $(3, 5)$).

1. Introduction

The Cremona group, denoted $\text{Bir}(\mathbb{P}_{\mathbb{C}}^n)$, is the group of birational maps of $\mathbb{P}_{\mathbb{C}}^n$ into itself. If $n = 2$ a lot of properties have been established (see [4, 9] for example). As far as we know the situation is much more different for $n \geq 3$ (see [14, 5] for example). If ψ is an element of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$ then $\deg \psi = \deg \psi^{-1}$. It is not the case in higher dimensions; if ψ belongs to $\text{Bir}(\mathbb{P}_{\mathbb{C}}^3)$ we only have the inequality $\deg \psi^{-1} \leq (\deg \psi)^2$ so one introduces the bidegree of ψ as the pair $(\deg \psi, \deg \psi^{-1})$. For $n = 2$, $\mathfrak{Bir}_d(\mathbb{P}_{\mathbb{C}}^2)$ is the set of birational maps of the complex projective plane of degree d ; for $n \geq 3$ denote by $\text{Bir}_{d,d'}(\mathbb{P}_{\mathbb{C}}^n)$

Texte reçu le 26 novembre 2014, révisé le 9 février 2015, accepté le 17 juin 2015.

JULIE DÉSERTI • *E-mail* : deserti@math.univ-paris-diderot.fr

FRÉDÉRIC HAN • *E-mail* : frederic.han@imj-prg.fr, Université Paris Diderot, Sorbonne Paris Cité, Institut de Mathématiques de Jussieu-Paris Rive Gauche, UMR 7586, CNRS, Sorbonne Universités, UPMC Univ Paris 06, F-75013 Paris, France.

2010 Mathematics Subject Classification. — 14E05, 14E07.

the set of elements of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^n)$ of bidegree (d, d') , and by $\mathfrak{Bir}_d(\mathbb{P}_{\mathbb{C}}^n)$ the union $\bigcup_{d'} \text{Bir}_{d,d'}(\mathbb{P}_{\mathbb{C}}^n)$. The set $\mathfrak{Bir}_d(\mathbb{P}_{\mathbb{C}}^n)$ inherits a structure of algebraic variety as a locally closed subspace a projective space ([3, Lemma 2.4, Proposition 2.15]), and we will always consider it with the Zariski topology ([8, 17]).

The varieties $\mathfrak{Bir}_2(\mathbb{P}_{\mathbb{C}}^2)$ and $\mathfrak{Bir}_3(\mathbb{P}_{\mathbb{C}}^2)$ are described in [6]: $\mathfrak{Bir}_2(\mathbb{P}_{\mathbb{C}}^2)$ is smooth, and irreducible in the space of quadratic rational maps of the complex projective plane whereas $\mathfrak{Bir}_3(\mathbb{P}_{\mathbb{C}}^2)$ is irreducible, and rationally connected. Besides, $\mathfrak{Bir}_d(\mathbb{P}_{\mathbb{C}}^2)$ is not irreducible as soon as $d > 3$ (see [2]). In [7] Cremona studies three types of generic elements of $\mathfrak{Bir}_2(\mathbb{P}_{\mathbb{C}}^3)$. Then there were some articles on the subject, and finally a precise description of $\mathfrak{Bir}_2(\mathbb{P}_{\mathbb{C}}^3)$; the left-right conjugacy is the following one

$$\text{PGL}(4; \mathbb{C}) \times \text{Bir}(\mathbb{P}_{\mathbb{C}}^3) \times \text{PGL}(4; \mathbb{C}) \rightarrow \text{Bir}(\mathbb{P}_{\mathbb{C}}^3), \quad (A, \psi, B) \mapsto A\psi B^{-1}.$$

Pan, Ronga and Vust give quadratic birational maps of $\mathbb{P}_{\mathbb{C}}^3$ up to left-right conjugacy, and show that there are only finitely many biclasses ([15, Theorems 3.1.1, 3.2.1, 3.2.2, 3.3.1]). In particular they show that $\mathfrak{Bir}_2(\mathbb{P}_{\mathbb{C}}^3)$ has three irreducible components of dimension 26, 28, 29; the component of dimension 26 (resp. 28, resp. 29) corresponds to birational maps of bidegree $(2, 4)$ (resp. $(2, 3)$, resp. $(2, 2)$). We will see that the situation is slightly different for $\mathfrak{Bir}_3(\mathbb{P}_{\mathbb{C}}^3)$; in particular we cannot expect such an explicit list of biclasses because there are infinitely many of biclasses (already the dimension of the family \mathcal{E}_2 of the classic cubo-cubic example is 39 that is strictly larger than $\dim(\text{PGL}(4; \mathbb{C}) \times \text{PGL}(4; \mathbb{C})) = 30$). That's why the approach is different.

We do not have such a precise description of $\mathfrak{Bir}_d(\mathbb{P}_{\mathbb{C}}^3)$ for $d \geq 4$. Nevertheless we can find a very fine and classical contribution for $\mathfrak{Bir}_3(\mathbb{P}_{\mathbb{C}}^3)$ due to Hudson ([11]); in the appendix we reproduce Table VI of [11]. Hudson introduces there some invariants to establish her classification. But it gives rise to many cases, and we also find examples where invariants take values that do not appear in her table. We do not know references explaining how her families fall into irreducible components of $\text{Bir}_{3,d}(\mathbb{P}_{\mathbb{C}}^3)$ so we focus on this natural question.

DEFINITION. — An element ψ of $\text{Bir}_{3,d}(\mathbb{P}_{\mathbb{C}}^3)$ is *ruled* if the strict transform of a generic plane under ψ^{-1} is a ruled cubic surface.

Denote by $\text{ruled}_{3,d}$ the set of $(3, d)$ ruled maps; we detail it in Lemma 2.3. Let us remark that there are no ruled birational maps of bidegree $(3, d)$ with $d \geq 6$.

We describe the irreducible components of $\text{Bir}_{3,d}(\mathbb{P}_{\mathbb{C}}^3)$ for $3 \leq d \leq 5$. Let us recall that the inverse of an element of $\text{Bir}_{3,2}(\mathbb{P}_{\mathbb{C}}^3)$ is quadratic and so treated in [15].

THEOREM A. — Assume that $2 \leq d \leq 5$. The set $\text{ruled}_{3,d}$ is an irreducible component of $\text{Bir}_{3,d}(\mathbb{P}_{\mathbb{C}}^3)$.

In bidegree $(3, 3)$ (resp. $(3, 4)$) there is only an other irreducible component; in bidegree $(3, 5)$ there are three others.

The set $\overline{\text{ruled}_{3,3}}$ intersects the closure of any irreducible component of $\text{Bir}_{3,4}(\mathbb{P}_{\mathbb{C}}^3)$ (the closures being taken in $\mathfrak{Bir}_3(\mathbb{P}_{\mathbb{C}}^3)$).

NOTATIONS 1.1. — Consider a dominant rational map ψ from $\mathbb{P}_{\mathbb{C}}^3$ into itself. For a generic line ℓ , the preimage of ℓ by ψ is a complete intersection Γ_{ℓ} ; define the scheme \mathcal{C}_2 to be the union of the irreducible components of Γ_{ℓ} supported in the base locus of ψ . Define \mathcal{C}_1 by liaison from \mathcal{C}_2 in Γ_{ℓ} . Remark that if ψ is birational, then $\mathcal{C}_1 = \psi_*^{-1}(\ell)$. Let us denote by $\mathbf{p}_a(\mathcal{C}_i)$ the arithmetic genus of \mathcal{C}_i .

It is difficult to find a uniform approach to classify elements of $\mathfrak{Bir}_3(\mathbb{P}_{\mathbb{C}}^3)$. Nevertheless in small genus we succeed to obtain some common detailed results; before stating them, let us introduce some notations.

Let us remark that the inequality $\deg \psi^{-1} \leq (\deg \psi)^2$ mentioned previously directly follows from

$$(\deg \psi)^2 = \deg \psi^{-1} + \deg \mathcal{C}_2.$$

PROPOSITION B. — Let ψ be a $(3, d)$ birational map of $\mathbb{P}_{\mathbb{C}}^3$.

Assume that ψ is not ruled, and $\mathbf{p}_a(\mathcal{C}_1) = 0$, i.e., \mathcal{C}_1 is smooth. Then

- $d \leq 6$;
- and \mathcal{C}_2 is a curve of degree $9 - d$, and arithmetic genus $9 - 2d$.

Suppose $\mathbf{p}_a(\mathcal{C}_1) = 1$, and $2 \leq d \leq 6$. Then

- there exists a singular point p of \mathcal{C}_1 independent of the choice of \mathcal{C}_1 ;
- if $d \leq 4$, all the cubic surfaces of the linear system Λ_{ψ} are singular at p ;
- the curve \mathcal{C}_2 is of degree $9 - d$, of arithmetic genus $10 - 2d$, and lies on a unique quadric Q ; more precisely $\mathcal{I}_{\mathcal{C}_2} = (Q, S_1, \dots, S_{d-2})$ where the S_i 's are independent cubics modulo Q .

We denote by $\text{Bir}_{3,d,\mathbf{p}_2}(\mathbb{P}_{\mathbb{C}}^3)$ the subset of non-ruled $(3, d)$ birational maps such that \mathcal{C}_2 is of degree $9 - d$, and arithmetic genus \mathbf{p}_2 . One has the following statement:

THEOREM C. — If $\mathbf{p}_2 \in \{3, 4\}$, then $\text{Bir}_{3,3,\mathbf{p}_2}(\mathbb{P}_{\mathbb{C}}^3)$ is non-empty, and irreducible; $\text{Bir}_{3,3,\mathbf{p}_2}(\mathbb{P}_{\mathbb{C}}^3)$ is empty as soon as $\mathbf{p}_2 \notin \{3, 4\}$.

If $\mathbf{p}_2 \in \{1, 2\}$, then $\text{Bir}_{3,4,\mathbf{p}_2}(\mathbb{P}_{\mathbb{C}}^3)$ is non-empty, and irreducible; $\text{Bir}_{3,4,\mathbf{p}_2}(\mathbb{P}_{\mathbb{C}}^3)$ is empty as soon as $\mathbf{p}_2 \notin \{1, 2\}$.

The set $\text{Bir}_{3,5,\mathbf{p}_2}(\mathbb{P}_{\mathbb{C}}^3)$ is empty as soon as $\mathbf{p}_2 \notin \{-1, 0, 1\}$ and

- if $\mathbf{p}_2 = -1$, then $\text{Bir}_{3,5,\mathbf{p}_2}(\mathbb{P}_{\mathbb{C}}^3)$ is non-empty, and irreducible;
- if $\mathbf{p}_2 = 0$, then $\text{Bir}_{3,5,\mathbf{p}_2}(\mathbb{P}_{\mathbb{C}}^3)$ is non-empty, and has two irreducible components;

- if $\mathfrak{p}_2 = 1$, then $\text{Bir}_{3,5,\mathfrak{p}_2}(\mathbb{P}_{\mathbb{C}}^3)$ is non-empty, and has three irreducible components.

Organization of the article. — In §2 we explain the particular case of ruled birational maps and set some notations. Then §3 is devoted to liaison theory that plays a big role in the description of the irreducible components of $\text{Bir}_{3,3}(\mathbb{P}_{\mathbb{C}}^3)$ (see §4), $\text{Bir}_{3,4}(\mathbb{P}_{\mathbb{C}}^3)$ (see §5) and $\text{Bir}_{3,5}(\mathbb{P}_{\mathbb{C}}^3)$ (see §6). In the last section we give some illustrations of invariants considered by Hudson, especially concerning the local study of the preimage of a line. Since Hudson's book is very old, let us recall her classification in the first part of the appendix.

Acknowledgment. — The authors would like to thank Jérémy Blanc and the referee for their helpful comments.

2. Definitions, notations and first properties

2.1. Definitions and notations. — Let $\psi: \mathbb{P}_{\mathbb{C}}^3 \dashrightarrow \mathbb{P}_{\mathbb{C}}^3$ be a rational map given, for some choice of coordinates, by

$$(z_0 : z_1 : z_2 : z_3) \dashrightarrow (\psi_0(z_0, z_1, z_2, z_3) : \psi_1(z_0, z_1, z_2, z_3) : \psi_2(z_0, z_1, z_2, z_3) : \psi_3(z_0, z_1, z_2, z_3))$$

where the ψ_i 's are homogeneous polynomials of the same degree d , and without common factors. The map ψ is called a *Cremona transformation* or a *birational map* of $\mathbb{P}_{\mathbb{C}}^3$ if it has a rational inverse ψ^{-1} . The *degree* of ψ , denoted $\deg \psi$, is d . The pair $(\deg \psi, \deg \psi^{-1})$ is the *bidegree* of ψ , we say that ψ is a $(\deg \psi, \deg \psi^{-1})$ birational map. The *indeterminacy set* of ψ is the set of the common zeros of the ψ_i 's. Denote by \mathcal{I}_{ψ} the ideal generated by the ψ_i 's, and by $\Lambda_{\psi} \subset H^0(\mathcal{O}_{\mathbb{P}_{\mathbb{C}}^3}(d))$ the subspace of dimension 4 generated by the ψ_i , and by $\deg \mathcal{I}_{\psi}$ the degree of the scheme defined by the ideal \mathcal{I}_{ψ} . The scheme whose ideal is \mathcal{I}_{ψ} is denoted F_{ψ} . It is called *base locus* of ψ . If $\dim F_{\psi} = 0$ then $F_{\psi}^1 = \emptyset$, otherwise F_{ψ}^1 is the maximal subscheme of F_{ψ} of dimension 1 without isolated point, and without embedded point. Furthermore if \mathcal{C} is a curve we will denote by $\omega_{\mathcal{C}}$ its dualizing sheaf.

REMARK 2.1. — We will sometimes identify a divisor of $\mathbb{P}_{\mathbb{C}}^3$ and its equation; for instance a surface Q of $\mathbb{P}_{\mathbb{C}}^3$ will have ideal (Q) .

Let us give a few comments about Table VI of [11]. For any subscheme X of $\mathbb{P}_{\mathbb{C}}^3$ denote by \mathcal{I}_X the ideal of X in $\mathbb{P}_{\mathbb{C}}^3$. Let ψ be a $(3, d)$ birational map. A point p is a *double point* if all the cubic surfaces of Λ_{ψ} are singular at p . A point p is a *binode* if all the cubic surfaces of Λ_{ψ} are singular at p with order 2 approximation at p a quadratic form of rank ≤ 2 (but this quadratic form

is allowed to vary in Λ_ψ). In other words p is a binode if there is a degree 1 element h of \mathcal{I}_p such that all the cubics belong to $(h \cdot \mathcal{I}_p) + \mathcal{I}_p^3$. A point p is a *double point of contact* if the general element of Λ_ψ is singular at p with order 2 approximation at p a quadratic form generically constant on Λ_ψ . In other words p is a double point of contact if all the cubics belong to $\mathcal{I}_p^3 + (Q)$ with Q of degree 2 and singular at p . A point p is a *point of contact* if all the cubics belong to $\mathcal{I}_p^2 + (S)$ where S is a cubic smooth at p . A point p is a *point of osculation* if all the cubics belong to $\mathcal{I}_p^3 + (S)$ where S is a cubic smooth at p .

NOTATIONS 2.2. — We will denote by \mathcal{E}_i the i -th family of Table VI and by $\mathbb{C}[z_0, z_1, \dots, z_n]_d$ the set of homogeneous polynomials of degree d in the variables z_0, z_1, \dots, z_n .

2.2. First properties. — Let us now focus on particular birational maps that cannot be dealt as the others: the ruled birational maps of $\mathbb{P}_{\mathbb{C}}^3$. Recall that there are two projective models of irreducible ruled cubic surfaces; they both have the same normalization: $\mathbb{P}_{\mathbb{C}}^2$ blown up at one point which can be realized as a cubic surface in $\mathbb{P}_{\mathbb{C}}^4$ (see [10, Chapter 10, introduction of § 4.4], [10, Chapter 9, § 2.1]).

LEMMA 2.3. — Assume that $2 \leq d \leq 5$.

- The set $\text{ruled}_{3,d}$ is irreducible.
- Let ψ be a general element of $\text{ruled}_{3,d}$, and let δ be the common line to all elements of $\{\text{Sing } S \mid S \in \Lambda_\psi\}$; then

$$\mathcal{I}_\psi = \mathcal{I}_\delta^2 \cap \mathcal{I}_{\Delta_1} \cap \mathcal{I}_{\Delta_2} \cap \dots \cap \mathcal{I}_{\Delta_{5-d}} \cap \mathcal{I}_K$$

where Δ_i are disjoint lines that intersect δ at a unique point, and K is a general reduced scheme of length $2d - 4$.

Proof. — Let ψ be an element of $\text{ruled}_{3,d}$. Recall that F_ψ^1 is the maximal subscheme of F_ψ of dimension 1 without isolated point, and without embedded point, i.e., F_ψ^1 is a locally Cohen-Macaulay curve.

An irreducible element S of Λ_ψ is a ruled surface; it is the projection of a smooth cubic ruled surface \tilde{S} of $\mathbb{P}_{\mathbb{C}}^4$. Recall that \tilde{S} is also the blow-up $\widehat{\mathbb{P}_{\mathbb{C}}^2}(p)$ of $\mathbb{P}_{\mathbb{C}}^2$ at a point p . The embedding of \tilde{S} in $\mathbb{P}_{\mathbb{C}}^4$ is given by the linear system $|\mathcal{I}_p(2h)|$, where h is the class of an hyperplane in $\mathbb{P}_{\mathbb{C}}^2$. Let us denote by π the projection $\tilde{S} \rightarrow S$, by H the class of the restriction of an hyperplane of $\mathbb{P}_{\mathbb{C}}^4$ to \tilde{S} , and by E_p the exceptional divisor associated to the blow-up of p . Set $\tilde{\delta} = \pi^{-1}\delta$, $\widetilde{\mathcal{C}}_1 = \pi^{-1}(\mathcal{C}_1)$, and $\widetilde{F}_\psi^1 = \pi^{-1}(F_\psi^1)$. One has

$$\tilde{\delta} \sim \pi^*h, \quad H = 2\pi^*h - E_p, \quad f = \pi^*h - E_p, \quad \widetilde{F}_\psi^1 = 2\tilde{\delta} + D$$

where D is an effective divisor. As $\widetilde{\mathcal{C}}_1 + \widetilde{F}_\psi^1 = 3H$, $\widetilde{\mathcal{C}}_1 \cdot f = 1$ and $\widetilde{\mathcal{C}}_1 \cdot H = d$ one gets $D \cdot f = 0$, and $D \cdot H = 5 - d$; therefore $D = (5 - d)f$. And we conclude that ψ has a residual base scheme of length $2d - 4$ from $\widetilde{\mathcal{C}}_1^2 = 2d - 3$.

Conversely, take a ruled cubic surface \widetilde{S} in $\mathbb{P}_{\mathbb{C}}^4$, choose a general projection π to $\mathbb{P}_{\mathbb{C}}^3$, take a general element Δ of $|\mathcal{O}_{\widetilde{S}}((5 - d)f)|$ and take a set \widetilde{K} of $2d - 4$ general points on \widetilde{S} of ideal $\mathcal{J}_{\widetilde{K}}$. We have $h^0(\mathcal{J}_{\widetilde{K}}(\widetilde{\mathcal{C}}_1)) = 3$, and thanks to the equality $\widetilde{\mathcal{C}}_1^2 = 2d - 3$ and the surjection $H^0\mathcal{O}_{\mathbb{P}_{\mathbb{C}}^3}(3) \twoheadrightarrow H^0\mathcal{O}_S(3)$ we get an element of $\text{ruled}_{3,d}$. Therefore the family of such $(\widetilde{S}, \pi, \Delta, \mathcal{J}_{\widetilde{K}})$ dominates $\text{ruled}_{3,d}$, and $\text{ruled}_{3,d}$ is irreducible.

To prove the claimed decomposition of \mathcal{J}_ψ for the general element $\psi \in \text{ruled}_{3,d}$, let us denote by \mathcal{J} the right part of the equality.

$$\mathcal{J} = \mathcal{J}_\delta^2 \cap \mathcal{J}_{\Delta_1} \cap \mathcal{J}_{\Delta_2} \cap \dots \cap \mathcal{J}_{\Delta_{5-d}} \cap \mathcal{J}_K.$$

We already have $\mathcal{J}_\psi \subset \mathcal{J}$. Furthermore, thanks to the computations on \widetilde{S} , one has $h^0(\mathcal{J}(3)) = \dim \Lambda_\psi = 4$, and $h^0(\mathcal{J}(2)) = 0$; hence to prove $\mathcal{J}_\psi = \mathcal{J}$ we just have to prove that \mathcal{J} is generated by polynomials of degree ≤ 3 . As ψ is general, one can assume that the line defined by (z_2, z_3) is trisecant to $\Delta_1 \cup \Delta_2 \cup \Delta_3$, so up to a change of coordinates we have

$$\begin{aligned}\mathcal{J}_\delta &= (z_0, z_1), \\ \mathcal{J}_{\Delta_1} &= (z_0, z_2), \\ \mathcal{J}_{\Delta_2} &= (z_1, z_3), \\ \mathcal{J}_{\Delta_3} &= (z_0 + z_1, z_2 + z_3)\end{aligned}$$

then

$$\begin{aligned}\mathcal{J}_\delta^2 &= (z_0^2, z_0 z_1, z_1^2), \\ \mathcal{J}_\delta^2 \cap \mathcal{J}_{\Delta_1} &= (z_0^2, z_0 z_1, z_1^2 z_2), \\ \mathcal{J}_\delta^2 \cap \mathcal{J}_{\Delta_1} \cap \mathcal{J}_{\Delta_2} &= (z_0 z_1, z_0^2 z_3, z_1^2 z_2), \\ \mathcal{J}_\delta^2 \cap \mathcal{J}_{\Delta_1} \cap \mathcal{J}_{\Delta_2} \cap \mathcal{J}_{\Delta_3} &= (z_0 z_1(z_0 + z_1), z_0 z_1(z_2 + z_3), z_1 z_2(z_0 + z_1), \\ &\quad z_0 z_3(z_0 + z_1))\end{aligned}$$

so we have the equality $\mathcal{J}_\psi = \mathcal{J}$ when $d = 2$. For $d \in \{3, 4, 5\}$, it is now enough to produce examples of $2d - 4$ points such that this equality is true to obtain it to $2d - 4$ general points. So consider the following 3 pairs of points of ideal

$$\begin{aligned}\mathcal{A}_1 &= (z_0 + z_2, z_1 + 2z_2 + z_3, z_0 z_3), \\ \mathcal{A}_2 &= (2z_0 + z_1 + z_2, z_3 - z_1, z_1 z_2), \\ \mathcal{A}_3 &= (z_2, z_0 + z_3, z_1^2 - z_0^2).\end{aligned}$$

Denote by \mathcal{I}_{ψ_3} (resp. $\mathcal{I}_{\psi_4}, \mathcal{I}_{\psi_5}$) the ideal generated by the 4 cubics of $\mathcal{J}_3 = \mathcal{J}_{\delta}^2 \cap \mathcal{I}_{\Delta_1} \cap \mathcal{I}_{\Delta_2} \cap \mathcal{A}_3$ (resp. $\mathcal{J}_4 = \mathcal{J}_{\delta}^2 \cap \mathcal{I}_{\Delta_1} \cap \mathcal{A}_2 \cap \mathcal{A}_3$, $\mathcal{J}_5 = \mathcal{J}_{\delta}^2 \cap \mathcal{A}_1 \cap \mathcal{A}_2 \cap \mathcal{A}_3$). Then we can compute the following equalities for instance with Macaulay2:
 $\mathcal{I}_{\psi_3} = (z_1^2 z_2, z_0 z_1 z_2, z_0 z_1^2 + z_0^2 z_3, z_0^2 z_1 + z_0 z_1 z_3),$
 $\mathcal{I}_{\psi_4} = (z_1^2 z_2, z_0 z_1 z_2, z_0^2 z_1 - z_0 z_1^2 - z_0^2 z_3 + z_0 z_1 z_3, 2z_0^3 + z_0 z_1^2 + z_0^2 z_2 + 3z_0^2 z_3),$
 $\mathcal{I}_{\psi_5} = (2z_0 z_1 z_2 - z_1^2 z_2, 5z_0 z_1^2 + z_1^3 + 7z_1^2 z_2 + 4z_0^2 z_3 + z_0 z_1 z_3 + z_1^2 z_3, 10z_0^2 z_1 + 2z_1^3 + 9z_1^2 z_2 - 2z_0^2 z_3 + 12z_0 z_1 z_3 + 2z_1^2 z_3, 40z_0^3 - 4z_1^3 + 20z_0^2 z_2 - 3z_1^2 z_2 + 44z_0^2 z_3 - 4z_0 z_1 z_3 - 4z_1^2 z_3),$

$$\mathcal{I}_{\psi_3} = \mathcal{J}_3, \quad \mathcal{I}_{\psi_4} = \mathcal{J}_4, \quad (z_0, z_1, z_2, z_3) = (\mathcal{J}_5 : \mathcal{I}_{\psi_5}).$$

So we have obtained the claimed decomposition of \mathcal{I}_{ψ} but when $d = 5$ the ideal \mathcal{I}_K has an irrelevant component. \square

LEMMA 2.4. — *The following inclusions hold:*

$$\text{ruled}_{3,2} \subset \overline{\text{ruled}_{3,3}}, \quad \text{ruled}_{3,3} \subset \overline{\text{ruled}_{3,4}}, \quad \text{ruled}_{3,4} \subset \overline{\text{ruled}_{3,5}}.$$

Proof (with the notations introduced in the proof of Lemma 2.3)

Let us start with an element of $\overline{\text{ruled}_{3,5}}$ with base curve δ^2 and 6 base points p_i in general position as described in Lemma 2.3. Then move two of the p_i , for instance p_1, p_2 until the line $(p_1 p_2)$ intersects δ . The line $(p_1 p_2)$ is now automatically in the base locus of the linear system Λ_{ψ} , and we obtain like this a generic element of $\text{ruled}_{3,4}$.

A similar argument allows to prove the two other inclusions. \square

Let us recall the notion of genus of a birational map ([11, Chapter IX]). The genus \mathfrak{g}_{ψ} of $\psi \in \text{Bir}(\mathbb{P}_{\mathbb{C}}^3)$ is the geometric genus of the curve $h \cap \psi^{-1}(h')$ where h and h' are generic hyperplanes of $\mathbb{P}_{\mathbb{C}}^3$. The equality $\mathfrak{g}_{\psi} = \mathfrak{g}_{\psi^{-1}}$ holds.

REMARK 2.5. — If ψ is a birational map of $\mathbb{P}_{\mathbb{C}}^3$ of degree 1 (resp. 2, resp. 3) then $\mathfrak{g}_{\psi} = 0$ (resp. $\mathfrak{g}_{\psi} = 0$, resp. $\mathfrak{g}_{\psi} \leq 1$).

One can give a characterization of ruled maps of $\text{Bir}_{3,d}(\mathbb{P}_{\mathbb{C}}^3)$ in terms of the genus.

PROPOSITION 2.6. — *Let ψ be in $\text{Bir}_{3,d}(\mathbb{P}_{\mathbb{C}}^3)$, $2 \leq d \leq 5$. The genus of ψ is zero if and only if ψ is ruled.*

Proof. — On the one hand there exists a curve \mathcal{C} such that $\mathcal{I}_{\psi} \subset \mathcal{J}_{\mathcal{C}}^2$ if and only if ψ is ruled; on the other hand $\mathfrak{g}_{\psi} = 0$ if and only if for generic hyperplanes h, h' of $\mathbb{P}_{\mathbb{C}}^3$ the curve $h \cap \psi^{-1}(h')$ is a singular rational cubic. \square

3. Liaison

According to [16] we say that two curves Γ_1 and Γ_2 of $\mathbb{P}_{\mathbb{C}}^3$ are *geometrically linked* if

- $\Gamma_1 \cup \Gamma_2$ is a complete intersection,
- Γ_1 and Γ_2 have no common component.

Let Γ_1 and Γ_2 be two curves geometrically linked. Recall that $\mathcal{I}_{\Gamma_1 \cup \Gamma_2} = \mathcal{I}_{\Gamma_1} \cap \mathcal{I}_{\Gamma_2}$. According to [16, Proposition 1.1] one has $\frac{\mathcal{I}_{\Gamma_1}}{\mathcal{I}_{\Gamma_1 \cup \Gamma_2}} = \text{Hom}(\mathcal{O}_{\Gamma_2}, \mathcal{O}_{\Gamma_1 \cup \Gamma_2})$. Since the kernel of $\mathcal{O}_{\Gamma_1 \cup \Gamma_2} \longrightarrow \mathcal{O}_{\Gamma_2}$ is $\frac{\mathcal{I}_{\Gamma_1}}{\mathcal{I}_{\Gamma_1 \cup \Gamma_2}}$ one gets the following fundamental statement: if Γ_1, Γ_2 are two curves geometrically linked, then

$$0 \longrightarrow \omega_{\Gamma_1} \longrightarrow \omega_{\Gamma_1 \cup \Gamma_2} \longrightarrow \mathcal{O}_{\Gamma_2} \otimes \omega_{\Gamma_1 \cup \Gamma_2} \longrightarrow 0.$$

LEMMA 3.1. — *Let ψ be a rational map of $\mathbb{P}_{\mathbb{C}}^3$ of degree 3. We have*

$$\omega_{\mathcal{C}_1 \cup \mathcal{C}_2} = \mathcal{O}_{\mathcal{C}_1 \cup \mathcal{C}_2}(2h),$$

where h denotes an hyperplane of $\mathbb{P}_{\mathbb{C}}^3$, and for $i \in \{1, 2\}$

$$(3.1) \quad 0 \longrightarrow \omega_{\mathcal{C}_i} \longrightarrow \mathcal{O}_{\mathcal{C}_i \cup \mathcal{C}_{3-i}}(2h) \longrightarrow \mathcal{O}_{\mathcal{C}_{3-i}}(2h) \longrightarrow 0$$

and

$$(3.2) \quad 0 \longrightarrow \mathcal{I}_{\mathcal{C}_1 \cup \mathcal{C}_2}(3h) \longrightarrow \mathcal{I}_{\mathcal{C}_i}(3h) \longrightarrow \omega_{\mathcal{C}_{3-i}}(h) \longrightarrow 0$$

The first exact sequence (3.1) directly implies the following equalities ($i \in \{1, 2\}$)

$$H^0(\omega_{\mathcal{C}_i}(-h)) = H^0(\mathcal{I}_{\mathcal{C}_{3-i}}(h)), \quad H^0(\omega_{\mathcal{C}_i}) = H^0(\mathcal{I}_{\mathcal{C}_{3-i}}(2h)),$$

$$h^0 \omega_{\mathcal{C}_i}(h) + 2 = h^0(\mathcal{I}_{\mathcal{C}_{3-i}}(3h)), \quad H^0(\omega_{\mathcal{C}_i}(h)) = \frac{H^0(\mathcal{I}_{\mathcal{C}_{3-i}}(3h))}{H^0(\mathcal{I}_{\mathcal{C}_1 \cup \mathcal{C}_2}(3h))}.$$

COROLLARY 3.2. — *Let ψ be a rational map of $\mathbb{P}_{\mathbb{C}}^3$ of degree 3. The ideal $\mathcal{I}_{\mathcal{C}_{3-i}}$ is generated by cubics if and only if $\omega_{\mathcal{C}_i}(h)$ is globally generated.*

Proof. — It directly follows from the exact sequence (3.2). □

COROLLARY 3.3. — *Let ψ be a rational map of $\mathbb{P}_{\mathbb{C}}^3$ of degree 3. Then*

$$\deg \mathcal{C}_2 - \deg \mathcal{C}_1 = \mathfrak{p}_a(\mathcal{C}_2) - \mathfrak{p}_a(\mathcal{C}_1).$$

Proof. — Taking the restriction of (3.1) to \mathcal{C}_i for $i = 1, 2$ gives

$$\deg \omega_{\mathcal{C}_i} = 2 \deg \mathcal{C}_i - \deg(\mathcal{C}_1 \cap \mathcal{C}_2),$$

and hence

$$\deg \mathcal{C}_2 - \deg \mathcal{C}_1 = \mathfrak{p}_a(\mathcal{C}_2) - \mathfrak{p}_a(\mathcal{C}_1). \quad \square$$

Furthermore when \mathcal{C}_1 and \mathcal{C}_2 have no common component, and $\omega_{\mathcal{C}_i}$ is locally free, then $\text{length}(\mathcal{C}_1 \cap \mathcal{C}_2) = \deg \omega_{\mathcal{C}_i}^{\vee}(2h)$, i.e.,

$$\sum_{p \in \mathcal{C}_1 \cap \mathcal{C}_2} \text{length}(\mathcal{C}_1 \cap \mathcal{C}_2)_{\{p\}} = 2 \deg \mathcal{C}_i - 2\mathfrak{p}_a(\mathcal{C}_i) + 2.$$

In the preimage of a generic point of $\mathbb{P}_{\mathbb{C}}^3$ by ψ , the number of points that do not lie in the base locus is given by

$$3 \deg \mathcal{C}_1 - \sum_{p \in \mathcal{C}_1 \cap \mathcal{C}_2} \text{length}(S \cap \mathcal{C}_1)_{\{p\}} - \sum_{p \in \Theta} \text{length}(S \cap \mathcal{C}_1)_{\{p\}}$$

where $S \in \Lambda_{\psi}$ is non-zero modulo $H^0(\mathcal{I}_{\mathcal{C}_1 \cup \mathcal{C}_2}(3h))$, and where Θ denotes the set of irreducible components of dimension 0 of the base locus F_{ψ} of ψ .

LEMMA 3.4. — *Let ψ be a rational map of $\mathbb{P}_{\mathbb{C}}^3$ of degree 3. Let Θ denote the set of irreducible components of dimension 0 of F_{ψ} . The map ψ is birational if and only if*

$$1 = 3 \deg \mathcal{C}_1 - \sum_{p \in \mathcal{C}_1 \cap \mathcal{C}_2} \text{length}(S \cap \mathcal{C}_1)_{\{p\}} - \sum_{p \in \Theta} \text{length}(S \cap \mathcal{C}_1)_{\{p\}}.$$

Remark that the computation of $\text{length}(S \cap \mathcal{C}_1)_{\{p\}}$ depends on the nature of the singularity of the cubic surface and on the behavior of \mathcal{C}_2 in that point (see §7).

LEMMA 3.5. — *Let ψ be a $(3, d)$ Cremona map. Assume that $d \geq 4$, then \mathcal{C}_1 is not contained in a plane.*

Proof. — Suppose for instance that $d = 4$; then \mathcal{C}_1 is contained in an irreducible cubic surface S . If \mathcal{C}_1 is contained in a plane \mathcal{P} then all the lines in \mathcal{P} are quadrisecant to S : contradiction with the irreducibility of S . \square

LEMMA 3.6. — *Let ψ be a $(3, d)$ birational map, and let p be a point on \mathcal{C}_1 . Assume that the degree of the tangent cone of \mathcal{C}_1 at p is strictly less than 4. If any S in Λ_{ψ} is singular at p , then p belongs to \mathcal{C}_2 .*

Proof. — If any S in Λ_{ψ} is singular at p , then the degree of the tangent cone of $\mathcal{C}_1 \cup \mathcal{C}_2$ at p is at least 4 because it is the complete intersection of two surfaces singular at p . Hence p has to belong to \mathcal{C}_2 . \square

LEMMA 3.7. — *Let ψ be a non-ruled $(3, d)$ birational map, and let \mathcal{C}_1 be a general element of Λ_{ψ} . The support of $\text{Sing } \mathcal{C}_1$ is independent of the choice of \mathcal{C}_1 .*

Proof. — Let us show that there is a singular point independent of the choice of \mathcal{C}_1 . Let us consider an element S of Λ_ψ with finite singular locus. Let $\pi: \tilde{S} \rightarrow S$ be a minimal desingularization of S , and let $\tilde{\mathcal{C}}_1$ be the strict transform of \mathcal{C}_1 . The elements of Λ_ψ give a linear system in $|\mathcal{O}_{\tilde{S}}(\tilde{\mathcal{C}}_1)|$ whose base locus denoted Ω is finite. According to Bertini's theorem applied on \tilde{S} one has the inclusion $\text{Sing } \mathcal{C}_1 \subset \pi(\Omega) \cup \text{Sing } S$. The assertion thus follows from the fact that $\Omega \cup \text{Sing } S$ is finite. \square

THEOREM 3.8. — *Let ψ be a $(3, d)$ birational map, $2 \leq d \leq 6$, that is not ruled. Assume that $\mathfrak{p}_a(\mathcal{C}_1) = 1$. Then*

- *there exists a singular point p of \mathcal{C}_1 independent of the choice of \mathcal{C}_1 ;*
- *if $d \leq 4$, all the cubic surfaces of the linear system Λ_ψ are singular at p ;*
- *the curve \mathcal{C}_2 is of degree $9 - d$, of arithmetic genus $10 - 2d$, and lies on a unique quadric Q ; more precisely $\mathcal{I}_{\mathcal{C}_2} = (Q, S_1, \dots, S_{d-2})$ where the S_i 's are independent cubics modulo Q .*

REMARK 3.9. — As soon as $d = 5$ the second assertion is not true. Indeed for $d = 5$ we obtain two families: one for which all the elements of Λ_ψ are singular, and another one for which it is not the case (§6).

Proof. — The first assertion directly follows from Lemma 3.7.

Since $\mathfrak{p}_a(\mathcal{C}_1) = 1$, the curve \mathcal{C}_2 lies on a unique quadric Q . The arithmetic genus of \mathcal{C}_2 is obtained from $\deg \mathcal{C}_2 - \deg \mathcal{C}_1 = \mathfrak{p}_a(\mathcal{C}_2) - \mathfrak{p}_a(\mathcal{C}_1)$ (Corollary 3.3).

As $\mathfrak{p}_a(\mathcal{C}_1) = 1$, $\omega_{\mathcal{C}_1}(h)$ has no base point, and $\mathcal{I}_{\mathcal{C}_2}$ is generated by cubics (Corollary 3.2). The number of cubics containing \mathcal{C}_2 independent modulo the multiple of Q is $d - 2$: the liaison sequence (Lemma 3.1) becomes

$$0 \longrightarrow \mathcal{O}_{\mathcal{C}_1}(h) \longrightarrow \mathcal{O}_{\mathcal{C}_1 \cup \mathcal{C}_2}(3h) \longrightarrow \mathcal{O}_{\mathcal{C}_2}(3h) \longrightarrow 0$$

one gets that

$$h^0 \mathcal{O}_{\mathcal{C}_2}(3h) = h^0 \mathcal{O}_{\mathcal{C}_1 \cup \mathcal{C}_2}(3h) - h^0 \mathcal{O}_{\mathcal{C}_1}(h) = 18 - d.$$

This implies that

$$h^0 \mathcal{I}_{\mathcal{C}_2}(3h) = 20 - h^0 \mathcal{O}_{\mathcal{C}_2}(3h) = d + 2.$$

If we remove the four multiples of Q one obtains $d + 2 - 4 = d - 2$ cubics, and finally $\mathcal{I}_{\mathcal{C}_2} = (Q, S_1, \dots, S_{d-2})$. \square

Corollary 3.3 and Theorem 3.8 imply Proposition B.

PROPOSITION 3.10. — *For $2 \leq d \leq 5$ the set $\text{ruled}_{3,d}$ is an irreducible component of $\text{Bir}_{3,d}(\mathbb{P}_{\mathbb{C}}^3)$.*

Proof. — Let us use the notations introduced in Lemma 2.3. Note that $F_{\psi}^1 \subset \mathcal{C}_2$. If $\psi \in \text{Bir}_{3,d}(\mathbb{P}_{\mathbb{C}}^3)$ is not ruled then at a generic point $p \in F_{\psi}^1$ there exists an element of Λ_{ψ} smooth at p . Hence F_{ψ}^1 is locally complete intersection at p and $\deg F_{\psi}^1 = \deg \mathcal{C}_2$. In particular $\deg \mathcal{I}_{\psi} = 9 - d$.

Consider now a general element ψ in $\text{ruled}_{3,d}$. From Lemma 2.3 there is a line ℓ such that $\ell \subset \text{Sing } S$ for any $S \in \Lambda_{\psi}$; the set F_{ψ}^1 has an irreducible component whose ideal is \mathcal{I}_{ℓ}^2 and F_{ψ}^1 is not locally complete intersection. This multiple structure has to be contained in \mathcal{C}_2 but since \mathcal{C}_2 is generically locally complete intersection the inequality $\deg \mathcal{C}_2 > \deg F_{\psi}^1$ holds; it can be rewritten $\deg \mathcal{I}_{\psi} < 9 - d$.

As $d > 9$ for any $\psi \in \text{Bir}_{3,d}(\mathbb{P}_{\mathbb{C}}^3)$, \mathcal{I}_{ψ} defines a 1-dimensional subscheme of $\mathbb{P}_{\mathbb{C}}^3$ so by the semi-continuity theorem $\psi \mapsto \deg \mathcal{I}_{\psi}$ is upper semi-continuous. Hence the number $\deg \mathcal{I}_{\psi}$ cannot decrease by specialization, and $\text{ruled}_{3,d}$ is not included in an irreducible component in $\text{Bir}_{3,d}(\mathbb{P}_{\mathbb{C}}^3)$ whose generic element is not ruled. \square

COROLLARY 3.11. — *Let ψ be a $(3, \cdot)$ birational map of $\mathbb{P}_{\mathbb{C}}^3$; if the general element of Λ_{ψ} is smooth or with isolated singularities, then $\deg F_{\psi}^1 = \deg \mathcal{C}_2$.*

4. $(3, 3)$ Cremona transformations

4.1. Some known results

4.1.1. — In the literature one can find different points of view concerning the classification of $(3, 3)$ birational maps. For example Hudson introduced many invariants related to singularities of families of surfaces and gave four families described in the appendix; nevertheless we do not understand why the family $\mathcal{C}_{3.5}$ defined below does not appear. Pan chose an other point of view and regrouped $(3, 3)$ birational maps into three families. A $(3, 3)$ birational map ψ of $\mathbb{P}_{\mathbb{C}}^3$ is called *determinantal* if there exists a 4×3 matrix M with linear entries such that ψ is given by the four 3×3 minors of the matrix M ; the inverse ψ^{-1} is also determinantal. Let us denote by $\mathbf{T}_{3,3}^{\mathbf{D}}$ the set of determinantal maps. A $(3, 3)$ Cremona transformation is a *de Jonquières* one if and only if the strict transform of a general line under ψ^{-1} is a singular plane rational cubic curve whose singular point is fixed. For such a map there is always a quadric contracted onto a point, the corresponding fixed point for ψ^{-1} which is also a de Jonquières transformation. The de Jonquières transformations form the set $\mathbf{T}_{3,3}^{\mathbf{J}}$. Pan established the following ([13, Theorem 1.2]):

$$\text{Bir}_{3,3}(\mathbb{P}_{\mathbb{C}}^3) = \mathbf{T}_{3,3}^{\mathbf{D}} \cup \mathbf{T}_{3,3}^{\mathbf{J}} \cup \text{ruled}_{3,3};$$

in other words an element of $\text{Bir}_{3,3}(\mathbb{P}_{\mathbb{C}}^3)$ is a determinantal map, or a de Jonquières map, or a ruled map.

REMARK 4.1. — One has $\mathbf{T}_{3,3}^{\mathbf{D}} = \text{Bir}_{3,3,3}(\mathbb{P}_{\mathbb{C}}^3)$ and $\mathbf{T}_{3,3}^{\mathbf{J}} = \text{Bir}_{3,3,4}(\mathbb{P}_{\mathbb{C}}^3)$; hence $\text{Bir}_{3,3,\mathfrak{p}_2}(\mathbb{P}_{\mathbb{C}}^3)$ is irreducible for $\mathfrak{p}_2 \in \{3, 4\}$ (see [13]).

REMARK 4.2. — The birational involution $(z_0 z_1^2 : z_0^2 z_1 : z_0^2 z_2 : z_1^2 z_3)$ is determinantal, the matrix being

$$\begin{bmatrix} z_0 & z_3 & 0 \\ -z_1 & 0 & z_2 \\ 0 & 0 & -z_1 \\ 0 & -z_0 & 0 \end{bmatrix},$$

and also ruled: all the partial derivatives of the components of the map vanish on $z_0 = z_1 = 0$. The Cremona transformation $(z_0^3 : z_0^2 z_1 : z_0^2 z_2 : z_1^2 z_3)$ is a de Jonquières and a ruled one; note that its primary decomposition (obtained with Macaulay2) is

$$(z_0^2, z_1^2) \cap (z_0^2, z_3) \cap (z_0^3, z_1, z_2)$$

and that its primary component (z_0^2, z_1^2) is locally complete intersection contrary to the general case (Lemma 2.3)⁽¹⁾.

One has ([12])

$$\mathbf{T}_{3,3}^{\mathbf{D}} \cap \mathbf{T}_{3,3}^{\mathbf{J}} = \emptyset, \quad \mathbf{T}_{3,3}^{\mathbf{D}} \cap \text{ruled}_{3,3} \neq \emptyset, \quad \mathbf{T}_{3,3}^{\mathbf{J}} \cap \text{ruled}_{3,3} \neq \emptyset.$$

We deal with the natural description of the irreducible components of $\text{Bir}_{3,3}$ which does not coincide with Pan's point of view since one of his family is contained in the closure of another one.

4.2. Irreducible components of the set of $(3, 3)$ birational maps

4.2.1. *General description of $(3, 3)$ birational maps.* — One already describes an irreducible component of $\text{Bir}_{3,3}(\mathbb{P}_{\mathbb{C}}^3)$, the one that contains $(3, 3)$ ruled birational maps (Proposition 3.10). Hence let us consider the case where the linear system Λ_{ψ} associated to $\psi \in \text{Bir}_{3,3}(\mathbb{P}_{\mathbb{C}}^3)$ contains a cubic surface without double line.

- If \mathcal{C}_1 is smooth then it is a twisted cubic, we are in family \mathcal{E}_2 of Table VI (see the appendix). In that case ψ is determinantal; more precisely a $(3, 3)$ birational map is determinantal if and only if its base locus scheme is an arithmetically Cohen-Macaulay curve of degree 6 and (arithmetic) genus 3 (see [1, Proposition 1]).

⁽¹⁾ Thanks to the referee for mentioning it to us.

- Otherwise $\omega_{\mathcal{E}_1} = \mathcal{O}_{\mathcal{E}_1}$, and ψ belongs to the irreducible family $\mathbf{T}_{3,3}^{\mathbf{J}}$ of Jonquières maps (\mathcal{E}_3 in terms of Hudson's classification). The curve \mathcal{C}_2 lies on a quadric described by the quadratic form Q . According to Theorem 3.8 the ideal of \mathcal{C}_2 is (Q, S) , and there exists a point p such that $p \in Q$, and p is a singular point of S . Furthermore $\mathcal{I}_{\psi} = \mathcal{I}_p Q + (S)$. Reciprocally such a triplet (p, Q, S) induces a birational map.

The family $\mathbf{T}_{3,3}^{\mathbf{J}}$ is stratified as follows by Hudson (all the cases belong to $\overline{\mathcal{E}_3}$):

- *Description of \mathcal{E}_3 .* The general element of $\mathcal{I}_p Q + (S)$ has an ordinary quadratic singularity at p (configuration (2, 2) of Table 1 (see §7)), and the generic cubic is singular at p with a quadratic form of rank 3.
- *Description of $\mathcal{E}_{3.5}$.* The point p lies on Q (p is a smooth point or not) and the generic cubic is singular at p with a quadratic form of rank 2. In other words p is a binode and this happens when one of the two biplanes is contained in $T_p Q$, it corresponds to the configuration (2, 3)' of Table 1 (see §7). The generic cubic is singular at p with a quadratic form of rank 2; this case does not appear in Table VI (see the appendix). Let us denote by $\mathcal{E}_{3.5}$ the set of the associated (3, 3) birational maps. The curve \mathcal{C}_2 has degree 6 and a triple point (in Q).
- *Description of \mathcal{E}_4 .* The point p is a double point of contact, it corresponds to configuration (2, 4) of Table 1 (see §7).

PROPOSITION 4.3. — *One has*

$$\dim \mathcal{E}_2 = 39, \quad \dim \mathcal{E}_3 = 38, \quad \dim \mathcal{E}_{3.5} = 35, \quad \dim \mathcal{E}_4 = 35, \quad \dim \mathcal{E}_5 = 31,$$

and

$$\overline{\mathcal{E}_3} = \mathbf{T}_{3,3}^{\mathbf{J}}, \quad \overset{\circ}{\mathcal{E}_{3.5}} \subset \overline{\mathcal{E}_3}, \quad \overset{\circ}{\mathcal{E}_4} \subset \overline{\mathcal{E}_3}, \quad \overset{\circ}{\mathcal{E}_4} \not\subset \overline{\mathcal{E}_{3.5}}, \quad \overset{\circ}{\mathcal{E}_{3.5}} \not\subset \overline{\mathcal{E}_4}.$$

Proof. — Let us justify the equality $\dim \mathcal{E}_3 = 38$. We have to choose a quadric Q and a point p on Q , this gives $9 + 2 = 11$. Then we take a cubic surface singular at p that yields to $19 - 4 = 15$; since we look at this surface modulo pQ one gets $15 - 3 = 12$ so

$$\dim \mathcal{E}_3 = 11 + 12 + 15 = 38.$$

Let us deal with $\dim \mathcal{E}_4$. We take a singular quadric Q this gives 8. Then we take a cubic singular at p , modulo pQ and this yields to $19 - 4 - 3 = 12$, and finally one obtains $12 + 8 + 15 = 35$. \square

4.2.2. Irreducible components

THEOREM 4.4. — *The set $\text{ruled}_{3,3}$ is an irreducible component of $\text{Bir}_{3,3}(\mathbb{P}_{\mathbb{C}}^3)$, and there is only one another irreducible component in $\text{Bir}_{3,3}(\mathbb{P}_{\mathbb{C}}^3)$. More precisely the set of the de Jonquières maps $\overline{\mathcal{E}}_3$ is contained in the closure of determinantal ones $\overline{\mathcal{E}}_2$ whereas $\text{ruled}_{3,3} \not\subset \overline{\mathcal{E}}_2$.*

Proof. — Let us consider the matrix A given by

$$\begin{bmatrix} 0 & 0 & 0 \\ -z_1 & -z_2 & 0 \\ z_0 & 0 & -z_2 \\ 0 & z_0 & z_1 \end{bmatrix}$$

and let A_i denote the matrix A minus the $(i+1)$ -th line. If $i > 0$, the 2×2 minors of A_i are divisible by z_{i-1} .

Consider the 4×3 matrix B given by $[b_{ij}]_{1 \leq i \leq 4, 1 \leq j \leq 3}$ with $b_{ij} \in H^0(\mathcal{O}_{\mathbb{P}_{\mathbb{C}}^3}(1))$; as previously, B_i is the matrix B minus the $(i+1)$ -th line. Denote by $\Delta^{j,k}$ the determinant of the matrix A_0 minus the j -th line and the k -th column; the $\Delta^{j,k}$ generate $\mathbb{C}[z_0, z_1, z_2]_2$. One has

$$\det(A_0 + tB_0) = t \cdot S \quad [t^2]$$

where

$$S = (b_{21} + b_{43})\Delta^{1,1} - (b_{31} - b_{42})\Delta^{2,1} + (b_{33} - b_{22})\Delta^{1,2} + b_{23}\Delta^{1,3} + b_{32}\Delta^{2,2} + b_{41}\Delta^{3,1}$$

is a generic cubic of the ideal $(z_0, z_1, z_2)^2$. For $i > 0$

$$\det(A_i + tB_i) = \det A_i + t \cdot (z_{i+1}Q)(-1)^{i+1} = t \cdot (z_{i+1}Q)(-1)^{i+1} \quad [t^2]$$

where $Q = b_{1,1}z_2 - b_{1,2}z_1 + b_{1,3}z_0$ is the equation of a generic quadric that contains $(0, 0, 0, 1)$. So the map

$$\left[\frac{\det(A_0 + tB_0)}{t} : \frac{\det(A_1 + tB_1)}{t} : \frac{\det(A_2 + tB_2)}{t} : \frac{\det(A_3 + tB_3)}{t} \right]$$

allows to go from $\overline{\mathcal{E}}_2$ to a general element of $\overline{\mathcal{E}}_3$.

Furthermore $\overline{\mathcal{E}}_3$ and $\text{ruled}_{3,3}$ are different components (Proposition 3.10). \square

5. (3, 4) Cremona transformations

5.1. General description of (3, 4) birational maps. — The ruled maps $\text{ruled}_{3,4}$ give rise to an irreducible component (Proposition 3.10). Let us now focus on the case where the linear system Λ_ψ associated to $\psi \in \text{Bir}_{3,4}(\mathbb{P}_{\mathbb{C}}^3)$ contains a cubic surface without double line.

- First case: \mathcal{C}_1 is smooth. From $h^0\omega_{\mathcal{C}_1}(h) = 3$ one gets that \mathcal{C}_2 lies on five cubics. Since $\mathbf{p}_a(\mathcal{C}_1) = 0$ we have $\omega_{\mathcal{C}_2} = \mathcal{O}_{\mathcal{C}_2}$ (Corollary 3.3). The curve \mathcal{C}_1 lies on a quadric (Lemma 3.1). This configuration corresponds to \mathcal{C}_6 .
- Second case: \mathcal{C}_1 is a singular curve of degree 4 not contained in a plane (see Lemma 3.5) so $\omega_{\mathcal{C}_1} = \mathcal{O}_{\mathcal{C}_1}$. The curve \mathcal{C}_1 lies on two quadrics and \mathcal{C}_2 on six cubics ($h^0\omega_{\mathcal{C}_1}(h) = 4$). Let p be the singular point of \mathcal{C}_1 ; all elements of Λ_ψ are singular at p (Theorem 3.8), and p belongs to \mathcal{C}_2 (Lemma 3.6). The curve \mathcal{C}_2 lies on a unique quadric Q (Theorem 3.8), is linked to a line ℓ in a $(2, 3)$ complete intersection $Q \cap S_1$ (with $\deg Q = 2$ and $\deg S_1 = 3$), and $\mathcal{I}_{\mathcal{C}_2} = (Q, S_1, S_2)$ with $\deg S_2 = 3$ (Theorem 3.8).

Since \mathcal{C}_1 is of degree 4 and arithmetic genus 1, one has $H^0(\mathcal{O}_{\mathcal{C}_1}(h)) = H^0(\mathcal{O}_{\mathbb{P}_{\mathbb{C}}^3}(1))$. Let us consider $L = H^0(\mathcal{I}_{\mathcal{C}_1 \cup \mathcal{C}_2}(3h)) \subset \Lambda_\psi$ and the map

$$H^0(\mathcal{O}_{\mathcal{C}_1}(h)) \longrightarrow \frac{H^0(\mathcal{I}_{\mathcal{C}_2}(3h))}{L}, \quad h \mapsto Qh;$$

it is injective. Indeed $\dim(\mathcal{C}_1 \cap Q) = 0$ thus modulo Q the cubics defining \mathcal{C}_1 are independent. Therefore Λ_ψ is contained in $(Q\mathcal{I}_p, S_1, S_2)$. Let $\pi: \widetilde{\mathcal{C}}_1 \rightarrow \mathcal{C}_1$ be the normalization. The linear system induced by $\psi \circ \pi$ is given by $\pi^*\omega_{\mathcal{C}_1}(h) = \pi^*\mathcal{O}_{\mathcal{C}_1}(h)$, and vanishes on $\pi^{-1}(p)$. This linear system has degree 4 and the conductor $\pi^{-1}(p)$ has length 2 because $\mathbf{p}_a(\mathcal{C}_1) = 1$. So it has a residual base point $p_1 \in \widetilde{\mathcal{C}}_1$ because ψ sends birationally \mathcal{C}_1 onto a line. Deforming p_1 to a general point p' of \mathcal{C}_1 we obtain the 4 dimensional vector space

$$\Lambda = H^0(((Q\mathcal{I}_p, S_1, S_2) \cap \mathcal{I}_{p'})(3)).$$

that is a deformation of Λ_ψ . In the following lines we will prove that the linear system given by Λ is birational.

Reciprocally let Q be a quadric, p be a point on Q , S_1 be a cubic singular at p and that contains a line ℓ of Q . If \mathcal{C}_2 is the residual of ℓ in (Q, S_1) , then there exists S_2 singular at p such that $\mathcal{I}_{\mathcal{C}_2} = (Q, S_1, S_2)$. Take $p_1 \in \mathbb{P}_{\mathbb{C}}^3 \setminus Q$, and set

$$\Lambda = H^0((\mathcal{I}_{p_1} \cap (Q\mathcal{I}_p, S_1, S_2))(3)).$$

Let L be a 2-dimensional general element of Λ ; the general linked curve to \mathcal{C}_2 in L , denoted $\mathcal{C}_{1,L}$, is of degree 4, is singular at p , lies on two quadrics; furthermore the linear system induced by Λ on $\mathcal{C}_{1,L}$ has the two following properties:

- its base locus contains p and p_1 ,
- it sends birationally $\mathcal{C}_{1,L}$ onto a line.

In other words, $\Lambda = \Lambda_\psi$ for a $(3, 4)$ birational map ψ .

Let us give some explicit examples, the generic one and the degeneracies considered by Hudson:

- *Description of \mathcal{E}_7 .* The quadric Q is smooth at p , and the rank of Q is maximal. Hence the point p is an ordinary quadratic singularity of the generic element of Λ_ψ , we are in the configuration $(2, 2)$ of Table 1 (see §7).
- *Description of $\mathcal{E}_{7.5}$.* In that case, p is a binode, Q is smooth at p and one of the two biplanes is contained in $T_p Q$; we are in the configuration $(2, 3)'$ of Table 1 (see §7). The set of such maps is denoted $\mathcal{E}_{7.5}$, this case does not appear in Table VI but should appear.
- *Description of \mathcal{E}_8 .* The second way to obtain a binode is the following one: Q is an irreducible cone with vertex p . This corresponds to the configuration $(2, 3)$ of Table 1 (see §7).
- *Description of \mathcal{E}_9 .* The rank of Q is 2, and the point p is a double point of contact; we are in the configuration $(2, 4)$ of Table 1 (see §7).
- *Description of \mathcal{E}_{10} .* The general element of Λ_ψ has a double point of contact and a binode (configurations $(2, 4)$ and $(1, 4)$ of Table 1, see §7). Hudson details this case carefully ([11, Chap. XV]).

PROPOSITION 5.1. — *One has the following properties:*

$$\dim \mathcal{E}_6 = 38, \quad \mathcal{E}_{7.5} \cup \mathcal{E}_8 \subset \overline{\mathcal{E}_7}$$

and

- *a generic element of $\mathcal{E}_{7.5}$ is not a specialization of a generic element of \mathcal{E}_8 ;*
- *a generic element of \mathcal{E}_8 is not a specialization of a generic element of $\mathcal{E}_{7.5}$;*
- *a generic element of \mathcal{E}_9 is a specialization of a generic element of \mathcal{E}_8 .*

Proof. — The arguments to establish $\dim \mathcal{E}_6 = 38$ are similar to those used in the proof of Proposition 4.3.

Let us justify that a generic element of $\mathcal{E}_{7.5}$ is not a specialization of a generic element of \mathcal{E}_8 (we take the notations of §5.1): as we see when $\psi \in \mathcal{E}_8$ the quadric Q is always singular whereas it is not the case when $\psi \in \mathcal{E}_{7.5}$. Conversely if ψ belongs to $\mathcal{E}_{7.5}$ then \mathcal{C}_2 is reducible but if ψ belongs to \mathcal{E}_8 the curve \mathcal{C}_2 can be irreducible and reduced; hence a generic element of \mathcal{E}_8 is not a specialization of a generic element of $\mathcal{E}_{7.5}$. \square

THEOREM 5.2. — *The set $\text{ruled}_{3,4}$ is an irreducible component of $\text{Bir}_{3,4}(\mathbb{P}_{\mathbb{C}}^3)$. There is only one another irreducible component in $\text{Bir}_{3,4}(\mathbb{P}_{\mathbb{C}}^3)$.*

Proof. — According to Proposition 3.10 the set $\text{rule}\mathfrak{d}_{3,4}$ is an irreducible component of $\text{Bir}_{3,4}(\mathbb{P}_{\mathbb{C}}^3)$.

Any element ψ of $\mathcal{E}_7 \cup \mathcal{E}_{7.5} \cup \mathcal{E}_8 \cup \mathcal{E}_9 \cup \mathcal{E}_{10}$ satisfies the following property:

$$\Lambda_{\psi} = H^0(((Q\mathcal{I}_p, S_1, S_2) \cap \mathcal{I}_{p_1})(3))$$

where p belongs to Q , p_1 is an ordinary base point, and

$$Q = \det \begin{bmatrix} L_0 & L_1 \\ L_2 & L_3 \end{bmatrix}, \quad S_1 = L_0Q_1 + L_1Q_2, \quad S_2 = L_2Q_1 + L_3Q_2$$

with $L_i \in \mathbb{C}[z_0, z_1, z_2, z_3]_1$, $Q_i \in \mathbb{C}[z_0, z_1, z_2]_2$. So \mathcal{E}_7 , $\mathcal{E}_{7.5}$, \mathcal{E}_8 , \mathcal{E}_9 and \mathcal{E}_{10} belong to the same irreducible component \mathcal{E} .

It remains to show that $\mathcal{E} = \overline{\mathcal{E}_6}$: let us consider

$$J = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{bmatrix}, \quad N = \begin{bmatrix} 0 & -z_2 & z_3 & L_0 \\ z_2 & 0 & L_1 & L_2 \\ -z_3 & -L_1 & 0 & L_3 \\ -L_0 & -L_2 & -L_3 & 0 \end{bmatrix}, \quad v = \begin{bmatrix} z_2 \\ z_1 \\ z_0 \\ tz_3 \end{bmatrix}$$

with L_i linear forms and

$$M_t = \begin{bmatrix} Jv \\ Nv \end{bmatrix} = \begin{bmatrix} tz_3 & z_0 - z_1 - z_2 \\ tz_3L_0 + Q & q_1 & q_2 & q_3 \end{bmatrix}$$

with

$$\begin{aligned} Q &= z_0z_3 - z_1z_2, & q_1 &= z_2^2 + z_0L_1 + tz_3L_2, \\ q_2 &= -z_2z_3 - z_1L_1 + tz_3L_3, & q_3 &= -z_2L_0 - z_1L_2 - z_0L_3. \end{aligned}$$

For generic L_i 's and $t \neq 0$ the 2×2 minors of M_t generate the ideal of a generic elliptic quintic curve as in \mathcal{E}_6 . For M_0 the 2×2 minors become Qz_0 , Qz_1 , Qz_2 , S_1 , S_2 , and S_3 with

$$S_1 = -z_2Q, \quad S_2 = -z_1q_3 + z_2q_2, \quad S_3 = z_0q_3 + z_2q_1.$$

Therefore the ideal \mathcal{M}_2 generated by these minors is

$$(Qz_0, Qz_1, Qz_2, S_2, S_3).$$

Denote by ℓ the line defined by $\mathcal{I}_{\ell} = (z_1, z_3)$. According to

$$z_3S_3 = -z_2S_2 + Q(q_3 + L_1z_2) \quad \& \quad z_1S_3 = -z_0S_2 - z_2^2Q$$

\mathcal{M}_2 is the ideal of the residual of ℓ in the complete intersection of ideals (Q, S_2) . To prove $\overline{\mathcal{E}_7} \subset \overline{\mathcal{E}_6}$ we just need to show that one can obtain the generic element of $\overline{\mathcal{E}_7}$ with a good choice of the L_i 's (Proposition 5.1); in other words it remains to prove that S_2 is generic among the cubics singular at p that contain ℓ . Modulo

Q one can assume that $q_3 = -z_3a + b$, with a (resp. b) an element of $\mathbb{C}[z_1, z_2]_1$ (resp. $\mathbb{C}[z_0, z_1, z_2]_2$). Then

$$S_2 = -z_3(z_1a + z_2^2) + z_1(b - z_2L_1);$$

in conclusion $S_2 = z_3A + z_2B$ for generic A and B in $\mathbb{C}[z_0, z_1, z_2]_2$. As \mathcal{C}_6 is irreducible $\mathcal{E} = \overline{\mathcal{C}_6}$. \square

5.2. Relations between $\overline{\text{Bir}_{3,3}(\mathbb{P}_{\mathbb{C}}^3)}$ and $\overline{\text{Bir}_{3,4}(\mathbb{P}_{\mathbb{C}}^3)}$. — One can now state the following result:

PROPOSITION 5.3. — *The set $\text{ruled}_{3,3}$ intersects the closure of any irreducible component of $\overline{\text{Bir}_{3,4}(\mathbb{P}_{\mathbb{C}}^3)}$.*

Proof. — According to Lemma 2.4 and Theorem 5.2 it is sufficient to prove that $\text{ruled}_{3,3}$ intersects the closure of $(3, 4)$ birational maps that are non-ruled.

Let us consider an element ψ of $\text{Bir}_{3,4}(\mathbb{P}_{\mathbb{C}}^3)$ whose \mathcal{C}_2 is the union of the lines of ideals

$$\mathcal{J}_\delta = (z_0, z_1^2), \quad (z_0 - \varepsilon z_2, z_1), \quad \mathcal{J}_{\ell_1} = (z_0, z_3), \quad \mathcal{J}_{\ell_2} = (z_1, z_2).$$

Denote by $\mathcal{J}_\varepsilon = (z_0, z_1^2) \cap (z_0 - \varepsilon z_2, z_1) \cap (z_0, z_3) \cap (z_1, z_2)$. One can check that

$$\mathcal{J}_\varepsilon = (z_0 z_1, z_0^2 z_2 + \varepsilon z_0 z_2^2, z_1^2 z_3).$$

Set $\mathcal{J}_\varepsilon = z_0 z_1(z_0, z_1, z_2) + (z_0^2 z_2 + \varepsilon z_0 z_2^2, z_1^2 z_3)$. For a general p_2 the map ψ_ε defined by $\Lambda_{\psi_\varepsilon} = H^0((\mathcal{J}_\varepsilon \cap \mathcal{J}_{p_2})(3))$ is birational; furthermore

- $\psi_\varepsilon \in \text{Bir}_{3,4}(\mathbb{P}_{\mathbb{C}}^3) \setminus \text{ruled}_{3,4}$ for $\varepsilon \neq 0$;
- $\psi_0 \in \text{ruled}_{3,3}$. \square

As in the case of $(3, 3)$ birational maps one has the following statement:

THEOREM 5.4. — *If $p_2 \in \{1, 2\}$, then $\text{Bir}_{3,4,p_2}(\mathbb{P}_{\mathbb{C}}^3)$ is non-empty and irreducible.*

6. $(3, 5)$ Cremona transformations

6.1. General description of $(3, 5)$ birational maps. — We already find an irreducible component of the $(3, 5)$ birational maps: $\text{ruled}_{3,5}$ (Proposition 3.10). Let us now consider a $(3, 5)$ Cremona transformation ψ such that Λ_ψ contains a cubic surface without double line.

Strategy. By Lemma 3.1 the image of \mathcal{C}_1 by ψ is given by a 2-dimensional vector subspace u of $H^0(\omega_{\mathcal{C}_1}(h))$. The restriction of ψ to \mathcal{C}_1 thus factorises by the composition of $\mathcal{C}_1 \dashrightarrow |H^0(\omega_{\mathcal{C}_1}(h))^\vee|$ with the projection $|H^0(\omega_{\mathcal{C}_1}(h))^\vee| \rightarrow |u^\vee|$. In the following cases we will use the equivalence between the birationality of ψ and the birationality of the composition $\mathcal{C}_1 \dashrightarrow |u^\vee|$. It will be useful to

compute the number of base points but also to show that a linear system is birational.

6.1.1. *Case: \mathcal{C}_1 smooth.* — By (3.2) the image of \mathcal{C}_1 by ψ is given by a sub-linear system of $|\omega_{\mathcal{C}_1}(h)|$. In that situation $\deg \omega_{\mathcal{C}_1}(h) = 3$ so as ψ sends birationally \mathcal{C}_1 onto a line of $\mathbb{P}_{\mathbb{C}}^3$ the map ψ has a residual base scheme of length 2. The curve \mathcal{C}_2 has genus -1 and does not lie on a quadric; \mathcal{C}_2 is the disjoint union of a twisted cubic and a line, so this case gives an irreducible family, and the general element belongs to \mathcal{E}_{12} . Indeed suppose that $\psi \notin \overline{\mathcal{E}_{12}}$, then \mathcal{C}_2 is the union of two smooth conics Γ_1 and Γ_2 that do not intersect. Any Γ_i is contained in a plane \mathcal{P}_i . Denote by ℓ the intersection $\mathcal{P}_1 \cap \mathcal{P}_2$. As $\#(\ell \cap (\Gamma_1 \cup \Gamma_2)) = 4$, all the cubic surfaces that contain $\Gamma_1 \cup \Gamma_2$ contain ℓ . So $\ell \subset \mathcal{C}_2$: contradiction.

6.1.2. *Case: \mathcal{C}_1 not smooth.* — So $\mathfrak{p}_a(\mathcal{C}_1) \geq 1$, and by Corollary 3.3

$$\mathfrak{p}_a(\mathcal{C}_2) = \deg \mathcal{C}_2 - \deg \mathcal{C}_1 + \mathfrak{p}_a(\mathcal{C}_1) = -1 + \mathfrak{p}_a(\mathcal{C}_1) \geq 0.$$

Since \mathcal{C}_1 is not in a plane, $\mathfrak{p}_a(\mathcal{C}_1) \leq 2$. Therefore we only have to distinguish the eventualities $\mathfrak{p}_a(\mathcal{C}_1) = 1$ and $\mathfrak{p}_a(\mathcal{C}_1) = 2$. Before looking at any of these eventualities let us introduce the set

$$\mathcal{C} = \{\text{irreducible curves of } \mathbb{P}_{\mathbb{C}}^3 \text{ of degree 5 and geometric genus 0}\}$$

• Assume first that $\mathfrak{p}_a(\mathcal{C}_1) = 1$. Then $\mathcal{O}_{\mathcal{C}_1} = \omega_{\mathcal{C}_1}$. We will denote by $\pi: \mathbb{P}_{\mathbb{C}}^1 \rightarrow \mathcal{C}_1$ the normalization of \mathcal{C}_1 .

a_1 . — Suppose first that all the elements of Λ_{ψ} are singular at $p \in \mathbb{P}_{\mathbb{C}}^3$. Denote by L the 2-dimensional vector space $\Lambda_{\psi} \cap H^0((\mathcal{J}_p^2 \cap \mathcal{J}_{\mathcal{C}_1})(3h))$ defining \mathcal{C}_1 and \mathcal{C}_2 . Let us follow the strategy explained before. By the liaison sequence (3.2) $\frac{\Lambda_{\psi}}{L}$ gives a vector subspace u of $H^0(\omega_{\mathcal{C}_1}(h)) = H^0(\mathcal{O}_{\mathcal{C}_1}(h))$ of dimension 2. It induces a projection from \mathcal{C}_1 to $|u^{\vee}|$ that coincides with the restriction of ψ to \mathcal{C}_1 ; hence this projection has degree 1. Moreover, via the identification $H^0(\mathcal{O}_{\mathcal{C}_1}(h)) = H^0(\pi^* \mathcal{O}_{\mathbb{P}_{\mathbb{C}}^1}(h))$, u is included in the set V_1 of sections of $\mathcal{O}_{\mathbb{P}_{\mathbb{C}}^1}(5)$ whose base locus contains $\pi^{-1}(p)$; as $\mathfrak{p}_a(\mathcal{C}_1) = 1$ the conductor $\pi^{-1}(p)$ has length 2. So there is an other base scheme of length 2 because $\psi|_{\mathcal{C}_1}: \mathcal{C}_1 \dashrightarrow |u^{\vee}|$ is birational.

We would like to show that \mathcal{C}_2 moves in an irreducible family. We will do this by deforming ψ (and \mathcal{C}_2) while \mathcal{C}_1 is fixed. So, $p \in \mathbb{P}_{\mathbb{C}}^3$ being fixed, let us consider

$$\mathcal{R}_{p,1} = \{\mathcal{C} \in \mathcal{C} \mid \text{Sing } \mathcal{C} = \{p\}, \mathfrak{p}_a(\mathcal{C}) = 1\};$$

the set $\mathcal{R}_{p,1}$ is an irreducible one. Remark that

$$h^0 \mathcal{J}_{\mathcal{C}}(3h) = h^0(\mathcal{O}_{\mathbb{P}_{\mathbb{C}}^3}(3h)) - h^0(\mathcal{O}_{\mathcal{C}}(3h)) = 20 - (3 \deg \mathcal{C} + 1 - \mathfrak{p}_a(\mathcal{C})) = 5$$

and $h^0((\mathcal{J}_p^2 \cap \mathcal{J}_{\mathcal{C}})(3h)) = 5 - 1 = 4$ because \mathcal{C} has a double point at p for all \mathcal{C} in $\mathcal{R}_{p,1}$.

Let us denote by F_1 the set of $(\mathcal{C}, L, u) \in \mathcal{R}_{1,p} \times H^0((\mathcal{J}_p^2 \cap \mathcal{J}_{\mathcal{C}})(3h)) \times V_1$ defined by

- $L \subset H^0((\mathcal{J}_p^2 \cap \mathcal{J}_{\mathcal{C}})(3h))$ of dimension 2 such that the residual of \mathcal{C} in the complete intersection defined by L has no common component with \mathcal{C} , and \mathcal{C} is geometrically linked to a curve denoted by $\mathcal{C}_{2,L}$,
- $u \subset V_1$ of dimension 2 such that $\mathbb{P}_{\mathbb{C}}^1 \dashrightarrow |u^\vee|$ has degree 1 (N.B. the general element of this family will then have two ordinary base points in addition to $\pi^{-1}(p)$).

The set F_1 is irreducible since the choice of \mathcal{C} is irreducible, and thus the choices of L and u too.

If (\mathcal{C}, L, u) belongs to F_1 , let us set

$$h_L: H^0(\mathcal{J}_{\mathcal{C}_{2,L}}(3h)) \rightarrow H^0(\omega_{\mathcal{C}}(h))$$

(recall that $\frac{H^0(\mathcal{J}_{\mathcal{C}_{2,L}}(3h))}{L} \simeq H^0(\omega_{\mathcal{C}}(h))$). Consider the map

$$\kappa_1: F_1 \rightarrow \mathbb{G}(4; H^0(\mathcal{O}_{\mathbb{P}^3}(3))), \quad (\mathcal{C}, L, u) \mapsto h_L^{-1}(u).$$

By construction of F_1 if ψ is a birational map such that $\mathfrak{p}_a(\mathcal{C}_1) = 1$ and all the elements of Λ_ψ are singular at p , then Λ_ψ is in the image of κ_1 .

Conversely one has:

LEMMA 6.1. — *The general element of $\text{im } \kappa_1$ coincides with Λ_ψ for some birational map ψ of \mathcal{E}_{14} .*

Proof. — As F_1 is irreducible it is enough to show that $|h_L^{-1}(u)|$ is a birational system when (\mathcal{C}, L, u) is general in F_1 . In that situation $\mathcal{C}_{2,L}$ is a curve of degree 4, arithmetic genus 0, singular at p , lying on a smooth quadric. Therefore $\mathcal{C}_{2,L}$ is reducible; more precisely it is the union of a twisted cubic and a line of this smooth quadric. All the elements of $|h_L^{-1}(u)|$ are cubic surfaces singular at p because $\mathcal{C}_{2,L}$ has a double point at p , and the residual pencil $u \subset H^0(\omega_{\mathcal{C}}(h))$ vanishes at p by definition of F_1 . Let \mathcal{C}_1 be the residual of $\mathcal{C}_{2,L}$ in the intersection of two general cubics of $|h_L^{-1}(u)|$. Hence \mathcal{C}_1 is singular at p . As u is by definition a pencil of sections of $\mathcal{O}_{\mathbb{P}^1}(5)$ vanishing on $\pi^{-1}(p)$ such that $\mathbb{P}_{\mathbb{C}}^1 \dashrightarrow |u^\vee|$ has degree 1 the linear system $|h_L^{-1}(u)|$ sends birationally \mathcal{C}_1 onto the line $|u^\vee|$. Therefore $|h_L^{-1}(u)|$ gives a birational map. \square

Let us remark that the previous irreducibility result asserts that the following example (belonging to family \mathcal{E}_{18}) whose base locus is not on a smooth quadric is nevertheless a deformation of elements of \mathcal{E}_{14} :

EXAMPLE 6.2. — Let \mathcal{C}_2 be the union of a line doubled on a smooth quadric with two other lines, such that all these lines contain a same point p . Set

$$Q = z_0z_3 - z_1z_2, \quad \mathcal{J}_p = (z_0, z_1, z_2);$$

then $\mathcal{J}_{\mathcal{C}_2} = ((z_2, z_0)^2 + (Q)) \cap (z_1, z_2) \cap (z_0 - z_2, z_1 - z_2)$. Now chose a double point of contact (note that the tangent cone must contain the tangent cone of \mathcal{C}_2):

$$\mathcal{J}_{\text{dpc}} = (z_2^2z_3 - z_0z_1z_3) + (z_0, z_1, z_2)^3,$$

and let p_1 and p_2 be two general points. Define \mathcal{J}_{ψ} by $\mathcal{J}_{\mathcal{C}_2} \cap \mathcal{J}_{\text{dpc}} \cap \mathcal{J}_{p_1} \cap \mathcal{J}_{p_2}$. So \mathcal{J}_{ψ} is the intersections of $\mathcal{J}_{p_1} \cap \mathcal{J}_{p_2}$ with

$$\begin{aligned} \mathcal{J}_{\mathcal{C}_2} \cap \mathcal{J}_{\text{dpc}} = & (z_1z_2^2 - z_2^3, z_0z_2^2 - z_2^3, z_1^2z_2 - z_2^3 - z_0z_1z_3 + z_2^2z_3, \\ & z_0z_1z_2 - z_2^3, z_0^2z_2 - z_2^3, z_0^2z_1 - z_2^3). \end{aligned}$$

The tangent cone of \mathcal{C}_2 at p has degree 4 but the tangent cone of $\mathcal{C}_1 \cup \mathcal{C}_2$ at p has degree 6, so \mathcal{C}_1 belongs to $\mathcal{R}_{p,1}$.

b_1 . — Suppose now that Λ_{ψ} contains a smooth element at p . Then p is a point of contact, all the cubic surfaces are tangent at p ; the curve $\mathcal{C}_2 \subset Q$ is linked to a curve of degree 2 and genus -1 so has degree 4 and genus 0. A general curve of degree 4 and genus 0 is in fact rational, smooth on a smooth quadric Q and we will see that such a curve can be "the \mathcal{C}_2 " of a birational map of that type. Hence this family of birational maps will turn out to be an irreducible one. Set $Q = z_0z_3 - z_1z_2$, $\mathcal{J}_{\ell_1} = (z_0, z_1)$, and $\mathcal{J}_{\ell_2} = (z_2, z_3)$; one has

$$\mathcal{J} = \mathcal{J}_{\ell_1 \cup \ell_2} = (z_0z_2, z_0z_3, z_1z_2, z_1z_3).$$

Let S_0 be the element of \mathcal{J} given by

$$az_0z_2 + bz_0z_3 + cz_1z_3 \quad a, b, c \in \mathbb{C}[z_0, z_1, z_2, z_3]_1;$$

one has $\mathcal{J}_{\mathcal{C}_2} = ((S_0, Q) : \mathcal{J}) = (Q, S_0, S_1, S_2)$ with

$$S_1 = z_0^2a + z_0z_1b + z_1^2c, \quad S_2 = z_2^2a + z_2z_3b + z_3^2c.$$

The dimension of $H^0(\mathcal{J}_{\mathcal{C}_2}(3h))$ is 7; indeed one has the following seven cubics:

$$\mathcal{J}_{\mathcal{C}_2} = \langle Qz_0, Qz_1, Qz_2, Qz_3, S_0, S_1, S_2 \rangle.$$

If ψ is a birational map, then ψ has no base point. Indeed $u = \frac{\Lambda_{\psi}}{H^0(\mathcal{J}_{\mathcal{C}_1 \cup \mathcal{C}_2}(3))}$ is contained in the sections of $\mathcal{O}_{\mathbb{P}_{\mathbb{C}}^3}(5)$ whose base locus contains $2\pi^{-1}(p)$; we thus already have an isomorphism between $\mathbb{P}_{\mathbb{C}}^1$ and $|u^{\vee}|$. The map ψ belongs to $\overline{\mathcal{E}_{23}}$. Conversely let L be a general 2-dimensional subspace of $H^0((\mathcal{J}_{\mathcal{C}_2} \cap$

$(Q + \mathcal{J}_p^2)(3h))$, and let \mathcal{C}_1 be the curve linked to \mathcal{C}_2 defined by L . Then the previous arguments show that the image of \mathcal{C}_1 by

$$\left| \frac{H^0((\mathcal{J}_{\mathcal{C}_2} \cap (Q + \mathcal{J}_p^2))(3h))}{L} \right|$$

is a line. So $H^0((\mathcal{J}_{\mathcal{C}_2} \cap (Q + \mathcal{J}_p^2))(3h)) = \Lambda_\psi$ for some birational map ψ of this type.

• Suppose that $\mathfrak{p}_a(\mathcal{C}_1) = 2$. Then $\mathfrak{p}_a(\mathcal{C}_2) = 1$, \mathcal{C}_1 lies on a quadric and $h^0 \mathcal{J}_{\mathcal{C}_1}(3) = 6$. We will still denote by $\pi: \mathbb{P}_{\mathbb{C}}^1 \rightarrow \mathcal{C}_1$ the normalization of \mathcal{C}_1 .

a_2 . — Assume first that \mathcal{C}_1 has a triple point p . The curve \mathcal{C}_1 is linked to a line by a complete intersection (Q, S_0) where Q (resp. S_0) is a cone (resp. a cubic) singular at p . We can write the normalization π as follows $(\alpha^2 A, \alpha\beta A, \beta^2 A, B)$ with $A \in \mathbb{C}[\alpha, \beta]_3$, $B \in \mathbb{C}[\alpha, \beta]_5$, and A, B without common factors. Then $Q = z_1^2 - z_0 z_2$, and $H^0(\omega_{\mathcal{C}_1}(h))$ can be identified with $H^0(\mathcal{J}_\ell(2))$, where $\mathcal{J}_\ell = (z_0, z_1)$. So $H^0(\omega_{\mathcal{C}_1}(h))$ is the 6-dimensional subspace W of $H^0(\mathcal{O}_{\mathbb{P}_{\mathbb{C}}^1}(6))$ spanned by $(\alpha, \beta) \cdot (\alpha^2 A, \alpha\beta A, \beta^2 A, B)$. Let us consider the subspace $V_A = W \cap (A)$ of W . Let L be the 2-dimensional vector space $\Lambda_\psi \cap H^0(\mathcal{J}_{\mathcal{C}_1}(3h))$. Then $\frac{\Lambda_\psi}{L}$ gives a 2-dimensional vector subspace u of V_A . The restriction of ψ to \mathcal{C}_1 gives a birational map $\mathbb{P}_{\mathbb{C}}^1 \dashrightarrow |u^\vee|$ induced by $u \subset V_A \subset H^0(\mathcal{O}_{\mathbb{P}_{\mathbb{C}}^1}(6))$. Furthermore ψ has two ordinary base points. We would like to show that in that case \mathcal{C}_2 moves in an irreducible family whose general element is the complete intersection of two quadrics. We thus fix a point $p \in \mathbb{P}_{\mathbb{C}}^3$ and introduce the irreducible set

$$\mathcal{R}_{p,2} = \{\mathcal{C} \in \mathcal{C} \mid \text{Sing } \mathcal{C} = \{p\}, \mathfrak{p}_a(\mathcal{C}) = 2\}.$$

We define the set F_2 as the $(\mathcal{C}, L, u) \in \mathcal{R}_{p,2} \times H^0(\mathcal{J}_{\mathcal{C}}(3h)) \times V_A$ given by

- $L \subset H^0(\mathcal{J}_{\mathcal{C}}(3h))$ of dimension 2 such that the residual of \mathcal{C} in the complete intersection defined by L has no common component with \mathcal{C} , and \mathcal{C} is geometrically linked to a curve denoted by $\mathcal{C}_{2,L}$,
- $u \subset V_A$ of dimension 2 such that $\mathbb{P}_{\mathbb{C}}^1 \dashrightarrow |u^\vee|$ is birational and whose base locus contains $\pi^{-1}(p)$.

Let us consider the map

$$\kappa_2: F_2 \rightarrow \mathbb{G}(4; H^0(\mathcal{O}_{\mathbb{P}_{\mathbb{C}}^3}(3))), \quad (\mathcal{C}, L, u) \mapsto h_L^{-1}(u).$$

If ψ is birational, if $\mathfrak{p}_a(\mathcal{C}_1) = 2$, and \mathcal{C}_1 has a triple point then ψ belongs to $\text{im } \kappa_2$.

LEMMA 6.3. — *The general element of $\text{im } \kappa_2$ coincides with Λ_ψ for some birational map ψ of \mathcal{C}_{13} .*

Proof. — As F_2 is irreducible one can consider a general element of F_2 , and then $\mathcal{C}_{2,L}$ is a curve of degree 4, genus 1 and is the complete intersection of two smooth quadrics. The map ψ has two ordinary base points p_1, p_2 , and belongs to \mathcal{E}_{13} . More precisely $\Lambda_{\psi} = H^0((\mathcal{I}_{\mathcal{C}_{2,L}} \cdot \mathcal{I}_p \cap \mathcal{I}_{p_1} \cap \mathcal{I}_{p_2})(3))$. \square

Note that this irreducibility result asserts that the following example in which \mathcal{C}_2 is not a complete intersection of two quadrics is nevertheless a deformation of elements of \mathcal{E}_{13} .

EXAMPLE 6.4. — Let \mathcal{C}_2 be the union of a plane cubic \mathcal{C}_3 singular at p and a line ℓ containing p but not in the plane spanned by \mathcal{C}_3 . For instance take

$$\mathcal{I}_p = (z_0, z_1, z_2), \quad \mathcal{I}_{\ell} = (z_1, z_2), \quad \mathcal{I}_{\mathcal{C}_3} = (z_1 - z_0, (z_1 - z_2)z_1z_3 + z_0^3 + z_1^3 + z_2^3).$$

Let \mathcal{I}_{dpc} be a double point of contact at p . (As we have already chosen \mathcal{C}_2 , we must take a quadric cone containing the tangent cone to \mathcal{C}_2). For instance one can take: $\mathcal{I}_{\text{dpc}} = (z_1^2 - z_0z_2) + \mathcal{I}_p^3$, and let

$$\begin{aligned} \mathcal{J} = \mathcal{I}_{\mathcal{C}_2} \cap \mathcal{I}_{\text{dpc}} = & (z_0z_2^2 - z_1z_2^2, z_0z_1z_2 - z_1^2z_2, z_0^2z_2 - z_1^2z_2, \\ & 2z_1^3 + z_2^3 + z_1^2z_3 - z_0z_2z_3, 2z_0z_1^2 + z_2^3 + z_1^2z_3 - z_0z_2z_3, \\ & 2z_0^2z_1 + z_2^3 + z_1^2z_3 - z_0z_2z_3) \end{aligned}$$

choose two general points p_1 and p_2 and define by \mathcal{I}_{ψ} the ideal generated by the 4 cubics of $\mathcal{J} \cap \mathcal{I}_{p_1} \cap \mathcal{I}_{p_2}$. The tangent cone of \mathcal{C}_2 at p has degree 3, the tangent cone of $\mathcal{C}_1 \cup \mathcal{C}_2$ at p has degree 6 (because p is a double point of contact); hence \mathcal{C}_1 has also a triple point at p , and belongs to $\mathcal{R}_{p,2}$.

b_2 . — Suppose now that \mathcal{C}_1 hasn't a triple point; \mathcal{C}_1 has thus two distinct double points. Fix two distinct points p and q in $\mathbb{P}_{\mathbb{C}}^3$, and set

$$\mathcal{R}_{p,q,2} = \{\mathcal{C} \in \mathcal{C} \mid \text{Sing } \mathcal{C} = \{p, q\}, \mathfrak{p}_a(\mathcal{C}) = 2\}.$$

Let V_3 (resp. V_4) be the sections of $\mathcal{O}_{\mathbb{P}_{\mathbb{C}}^3}(7)$ whose base locus contains $\pi^{-1}(p)$ and $\pi^{-1}(q)$ (resp. $\pi^{-1}(p)$ and $2\pi^{-1}(q)$). The set $\mathcal{R}_{p,q,2}$ is irreducible. Remark that for all \mathcal{C} in $\mathcal{R}_{p,q,2}$ one has

$$h^0(\mathcal{I}_{\mathcal{C}}(3h)) = 6, \quad h^0((\mathcal{I}_{\mathcal{C}} \cap \mathcal{I}_p^2)(3h)) = 5, \quad h^0((\mathcal{I}_{\mathcal{C}} \cap \mathcal{I}_p^2 \cap \mathcal{I}_q^2)(3h)) = 4.$$

REMARK 6.5. — One cannot have two distinct points of contact. Assume by contradiction that there are two distinct points of contact p and q . Denote by $\pi: \widetilde{\mathcal{C}_1} \rightarrow \mathcal{C}_1$ the normalization of \mathcal{C}_1 . One would have $\pi^*\omega_{\mathcal{C}_1}(h) = \mathcal{O}_{\mathbb{P}_{\mathbb{C}}^3}(7)$ but the linear system induced by ψ would contain in the base locus $2\pi^{-1}(p) + 2\pi^{-1}(q)$ which is of length 8: contradiction with the fact that ψ sends birationally \mathcal{C}_1 onto a line.

So one has the following alternative:

b_2 i. — Either all the cubics of Λ_ψ are singular at p and q . One can then define the set F_3 of $(\mathcal{C}, L, u) \in \mathcal{R}_{p,q,2} \times H^0((\mathcal{I}_\mathcal{C} \cap \mathcal{I}_p^2 \cap \mathcal{I}_q^2)(3h)) \times V_3$ given by

- $L \subset H^0((\mathcal{I}_\mathcal{C} \cap \mathcal{I}_p^2 \cap \mathcal{I}_q^2)(3h))$ of dimension 2 such that the residual of \mathcal{C} in the complete intersection defined by L has no common component with \mathcal{C} ;
- $u \in V_3$ of dimension 2 such that $\mathcal{C} \dashrightarrow |u^\vee|$ has degree 1.

Let us consider the map

$$\kappa_3: F_3 \rightarrow \mathbb{G}(4; H^0(\mathcal{O}_{\mathbb{P}^3}(3))), \quad (\mathcal{C}, L, u) \mapsto h_L^{-1}(u).$$

If ψ is birational, if $\mathbf{p}_a(\mathcal{C}_1) = 2$, if \mathcal{C}_1 has two distinct double points at p and q and if all the cubics of Λ_ψ are singular at p and q , then Λ_ψ belongs to $\text{im } \kappa_3$.

LEMMA 6.6. — *The general element of $\text{im } \kappa_3$ coincides with Λ_ψ for some birational map ψ of \mathcal{E}_{19} .*

Proof. — As F_3 is irreducible one can consider a general element (\mathcal{C}, L, u) of F_3 and then $\mathcal{C}_{2,L}$ is a curve of degree 4 and genus 1, is singular at p and q , lies on a smooth quadric, and is reducible: $\mathcal{C}_{2,L}$ is the union of a twisted cubic Γ and the line $\ell = (pq)$. Moreover for all the elements of $h_L^{-1}(u)$ the curve \mathcal{C} is singular at p and q (by definition of V_3 and by the fact that $\mathcal{C}_{2,L}$ is singular at p and q). \square

In this situation as all the cubic surfaces are singular at p and q ,

$$h_L^{-1}(u) = H^0((\mathcal{I}_\Gamma \cdot \mathcal{I}_\ell \cap \mathcal{I}_{p_1} \cap \mathcal{I}_{p_2})(3))$$

where p_1, p_2 are two ordinary base points; ψ belongs to \mathcal{E}_{19} .

b_2 ii. — Or one of the cubics of Λ_ψ is smooth at (for instance) q . Let us introduce the set F_4 of pairs $(\mathcal{C}, L) \in \mathcal{R}_{p,q,2} \times H^0((\mathcal{I}_\mathcal{C} \cap \mathcal{I}_p^2)(3h))$ satisfying: $L \subset H^0((\mathcal{I}_\mathcal{C} \cap \mathcal{I}_p^2)(3h))$ of dimension 2 such that the residual of \mathcal{C} in the complete intersection defined by L has no common component with \mathcal{C} .

Let us consider the map

$$\kappa_4: F_4 \rightarrow \mathbb{G}(4; H^0(\mathcal{O}_{\mathbb{P}^3}(3))), \quad (\mathcal{C}, L) \mapsto h_L^{-1}(V_4);$$

note that $\dim V_4 = 2$.

If ψ is birational, if $\mathbf{p}_a(\mathcal{C}_1) = 2$, \mathcal{C}_1 hasn't a triple point and one of the cubic of Λ_ψ is smooth at (for instance) q , then Λ_ψ belongs to $\text{im } \kappa_4$.

LEMMA 6.7. — *The general element of $\text{im } \kappa_4$ coincides with Λ_ψ for some birational map ψ of \mathcal{E}_{24} .*

Proof. — As F_4 is irreducible one can consider a general element of F_4 , and then $\mathcal{C}_{2,L}$ is a curve of degree 4, genus 1, singular at p , and is the complete intersection of two quadrics. The map ψ has no base point and belongs to \mathcal{E}_{24} . \square

6.2. Irreducible components. — The following statement, and Theorems 4.4 and 5.2 imply Theorem A.

THEOREM 6.8. — *One has the inclusions: $\mathcal{E}_{14} \subset \overline{\mathcal{E}_{12}}$, $\mathcal{E}_{24} \subset \overline{\mathcal{E}_{23}}$, and $\mathcal{E}_{19} \subset \overline{\mathcal{E}_{12}}$.*

The set $\text{Bir}_{3,5}(\mathbb{P}_{\mathbb{C}}^3)$ has four irreducible components: \mathcal{E}_{12} , \mathcal{E}_{13} , \mathcal{E}_{23} , and $\mathcal{E}_{27} = \text{ruled}_{3,5}$.

Proof. — Let us first prove that $\mathcal{E}_{14} \subset \overline{\mathcal{E}_{12}}$. If ψ belongs to \mathcal{E}_{12} , or to \mathcal{E}_{14} the curve \mathcal{C}_2 is the union of a line ℓ and a twisted cubic Γ such that $\text{length}(\ell \cap \Gamma) \leq 1$. Let \mathcal{I}_{ℓ} (resp. \mathcal{I}_{Γ}) be the ideal of ℓ (resp. Γ). We have $\mathcal{I}_{\psi} \subset \mathcal{I}_{\ell} \cap \mathcal{I}_{\Gamma}$. If ψ belongs to \mathcal{E}_{12} , then $\ell \cap \Gamma = \emptyset$, and $\mathcal{I}_{\ell} \cap \mathcal{I}_{\Gamma} = \mathcal{I}_{\ell} \cdot \mathcal{I}_{\Gamma}$. And if ψ is in \mathcal{E}_{14} , then all the cubics are singular at $p = \ell \cap \Gamma$ so \mathcal{I}_{ψ} is again in $\mathcal{I}_{\ell} \cdot \mathcal{I}_{\Gamma}$.

Prove now that $\mathcal{E}_{24} \subset \overline{\mathcal{E}_{23}}$. Consider a general element ψ of \mathcal{E}_{24} ; the curve \mathcal{C}_2 is the complete intersection of a quadric $Q' = az_2 + bz_0 + cz_1$ passing through the double point p and a cone $Q_0 = z_1z_2 - z_0^2$. Furthermore all the cubics of \mathcal{I}_{ψ} are singular at p , and $\mathcal{I}_{\psi} \subset \mathcal{I}'_0 = (Q_0, z_0Q', z_1Q', z_2Q')$. Let \mathfrak{ct}_q be the ideal of the point of contact q ; one has $\mathfrak{ct}_q = \mathcal{I}'_q + (H_q)$ where H_q is a plane passing through q . Denote by \mathcal{I}_0 the intersection of \mathcal{I}'_0 and \mathfrak{ct}_q . Set

$$Z_0 = z_0 + tz_3, \quad Z_1 = z_1, \quad Z_2 = z_2, \quad Z_3 = z_0 - tz_3,$$

$$Q_t = Z_1Z_2 - Z_0Z_3, \quad S_0 = aZ_0Z_2 + bZ_0Z_3 + cZ_1Z_3,$$

$$S_1 = aZ_0^2 + bZ_0Z_1 + cZ_1^2, \quad S_2 = aZ_2^2 + bZ_2Z_3 + cZ_3^2.$$

Hence $\mathcal{I}_t = (Q_t, S_0, S_1, S_2)$ is the ideal of a rational quartic if $t \neq 0$ (cf. the equations in §6.1.1 b_1)). The ideal $\mathcal{I}_t = \mathcal{I}_t \cap \mathfrak{ct}_q$ is the ideal \mathcal{I}_{ψ} of $\psi \in \mathcal{E}_{23}$. Remark that if $t = 0$, then

$$\mathcal{I}_0 = (Q_0, z_0Q', az_0^2 + bz_0z_1 + cz_1^2, az_2^2 + bz_0z_2 + cz_3^2)$$

but $az_0^2 + bz_0z_1 + cz_1^2 = z_1Q'$ modulo Q , and $az_2^2 + bz_0z_2 + cz_3^2 = z_2Q'$ modulo Q , that is $\mathcal{I}'_0 = \mathcal{I}_0$. Therefore \mathcal{I}_t tends to \mathcal{I}_0 as t tends to 0.

The inclusion $\mathcal{E}_{19} \subset \overline{\mathcal{E}_{12}}$ follows from $\Lambda_{\psi} = H^0((\mathcal{I}_{\ell} \cdot \mathcal{I}_{\Gamma} \cap \mathcal{I}_{p_1} \cap \mathcal{I}_{p_2})(3))$ found in b_2 i.

Note that $\mathcal{E}_{12} \not\subset \overline{\mathcal{E}_{13}}$ (resp. $\mathcal{E}_{12} \not\subset \overline{\mathcal{E}_{23}}$): if ψ is in \mathcal{E}_{12} then the associated \mathcal{C}_2 does not lie on a quadric whereas if ψ belongs to \mathcal{E}_{13} (resp. \mathcal{E}_{23}) then \mathcal{C}_2 lies on two quadrics (resp. one quadric). Conversely $\mathcal{E}_{13} \not\subset \overline{\mathcal{E}_{12}}$ (resp. $\mathcal{E}_{23} \not\subset \overline{\mathcal{E}_{12}}$): if ψ is an element of \mathcal{E}_{13} (resp. \mathcal{E}_{23}), then \mathcal{C}_2 is smooth and irreducible whereas

the associated \mathcal{C}_2 of a general element of \mathcal{E}_{12} is the disjoint union of a twisted cubic and a line.

Let us now justify that $\mathcal{E}_{23} \not\subset \overline{\mathcal{E}_{13}}$: the linear system of an element of \mathcal{E}_{23} has a smooth surface whereas the linear system of an element of \mathcal{E}_{13} does not. Conversely $\mathcal{E}_{13} \not\subset \overline{\mathcal{E}_{23}}$; indeed $h^0 \mathcal{J}_{\mathcal{E}_2}(3h) = 6$ for a birational map of \mathcal{E}_{13} and $h^0 \mathcal{J}_{\mathcal{E}_2}(3h) = 7$ for a birational map of \mathcal{E}_{23} . \square

In bidegree $(3, 5)$ the description of $\text{Bir}_{3,5,p_2}(\mathbb{P}_{\mathbb{C}}^3)$ is very different from those of smaller bidegrees. Let us now prove Theorem C.

THEOREM 6.9. — *The set $\text{Bir}_{3,5,p_2}(\mathbb{P}_{\mathbb{C}}^3)$ is empty as soon as $p_2 \notin \{-1, 0, 1\}$ and*

- *if $p_2 = -1$, then $\text{Bir}_{3,5,p_2}(\mathbb{P}_{\mathbb{C}}^3)$ is non-empty, and irreducible;*
- *if $p_2 = 0$, then $\text{Bir}_{3,5,p_2}(\mathbb{P}_{\mathbb{C}}^3)$ is non-empty, and has two irreducible components: one formed by the birational maps of \mathcal{E}_{14} , and the other one by the elements of \mathcal{E}_{23} ;*
- *if $p_2 = 1$, then $\text{Bir}_{3,5,p_2}(\mathbb{P}_{\mathbb{C}}^3)$ is non-empty, and has three irreducible components: one formed by the birational maps of \mathcal{E}_{13} , a second one formed by the birational maps of \mathcal{E}_{19} , and a third one by the elements of \mathcal{E}_{24} .*

Proof. — • Assume $p_2 = -1$. In that case only one family appears : \mathcal{E}_{12} (see § 6.1.1), and according to Theorem 6.8 the family \mathcal{E}_{12} is already an irreducible component of $\text{Bir}_{3,5}(\mathbb{P}_{\mathbb{C}}^3)$ so an irreducible component of $\text{Bir}_{3,5,-1}(\mathbb{P}_{\mathbb{C}}^3)$.

• Suppose $p_2 = 0$. We found two families : \mathcal{E}_{14} (case a_1 of § 6.1.1), and \mathcal{E}_{23} (case b_1 of § 6.1.2). Note that for ψ general in \mathcal{E}_{23} the linear system Λ_{ψ} contains smooth cubics whereas all cubics of Λ_{ψ} are singular as soon as ψ belongs to \mathcal{E}_{14} . Hence $\mathcal{E}_{23} \not\subset \overline{\mathcal{E}_{14}}$.

Take a general element of \mathcal{E}_{14} ; it hasn't a base scheme of dimension 0, connected and of length ≥ 3 whereas elements of $\overline{\mathcal{E}_{23}}$ have. Therefore $\mathcal{E}_{14} \not\subset \overline{\mathcal{E}_{23}}$.

• Assume last that $p_2 = 1$. Our study gives three families: \mathcal{E}_{13} , \mathcal{E}_{19} and \mathcal{E}_{24} (cases a_2 , b_2 i and b_2 ii of § 6.1.2). The general element of \mathcal{E}_{19} has two double points whereas a general element of \mathcal{E}_{13} (resp. \mathcal{E}_{24}) has only one; thus $\mathcal{E}_{19} \not\subset \overline{\mathcal{E}_{13}}$ and $\mathcal{E}_{19} \not\subset \overline{\mathcal{E}_{24}}$.

Take a general element in \mathcal{E}_{13} ; its base locus is a smooth curve. On the contrary if ψ belongs to \mathcal{E}_{19} (resp. \mathcal{E}_{24}), then the base locus of ψ is a singular curve. Thus $\mathcal{E}_{13} \not\subset \overline{\mathcal{E}_{19}}$ (resp. $\mathcal{E}_{13} \not\subset \overline{\mathcal{E}_{24}}$).

If ψ is a general element of \mathcal{E}_{24} its base locus is an irreducible curve and this is not the case if $\psi \in \mathcal{E}_{19}$ so $\mathcal{E}_{24} \not\subset \overline{\mathcal{E}_{19}}$.

Let us now consider a general element of \mathcal{E}_{24} , the tangent plane at all cubic surfaces at the point of contact doesn't contain the double point p ; hence if we denote by Q_1 and Q_2 the quadrics containing \mathcal{C}_2 there isn't a plane h passing through p such that $(hQ_1, hQ_2) \subset \Lambda_{\psi}$. But if we take ψ in \mathcal{E}_{13} then

$\Lambda_\psi = H^0((\mathcal{I}_{\mathcal{C}_2} \cdot \mathcal{I}_p \cap \mathcal{I}_{p_1} \cap \mathcal{I}_{p_2})(3))$ with p_1, p_2 two ordinary base points, and p the triple point lying on \mathcal{C}_1 . If h is the plane passing through p, p_1 and p_2 , if $\mathcal{I}_{\mathcal{C}_2} = (Q_1, Q_2)$, then $(hQ_1, hQ_2) \subset \Lambda_\psi$. Thus $\mathcal{C}_{24} \not\subset \overline{\mathcal{C}_{13}}$. \square

7. Relations with Hudson's invariants

To prove the birationality of a linear system of cubics, the local properties of \mathcal{C}_1 and \mathcal{C}_2 are required. For instance to apply Lemma 3.4 one needs to understand the support of $\mathcal{C}_1 \cup \mathcal{C}_2$ and the local intersection of \mathcal{C}_1 with a general element of Λ_ψ at any point of $\mathcal{C}_1 \cup \mathcal{C}_2$. So in the following table we make a schematic picture of the tangent cone of $\mathcal{C}_1 \cup \mathcal{C}_2$ at one of its singular point in the different cases considered by Hudson. Let us note that the degree of the tangent cone of $\mathcal{C}_1 \cup \mathcal{C}_2$ at a point of $\mathcal{C}_1 \cup \mathcal{C}_2$ varies from 1 to 6. In particular if the linear system has a double point (resp. a double point of contact), then it is a complete intersection of two quadric cones (resp. of one quadric cone and one cubic cone). We draw pictures only when the quadric cone is irreducible. If the linear system has a binode, the tangent cone of $\mathcal{C}_1 \cup \mathcal{C}_2$ has degree 5; more precisely for a binode at $p = (z_0, z_1, z_2)$ whose fixed plane is z_0 , i.e., $\mathcal{I}_\psi \subset \mathcal{I}_p \cdot (z_0)$, then the ideal of the tangent cone of $\mathcal{C}_1 \cup \mathcal{C}_2$ at p is (z_0z_1, z_0z_2, P) where P denotes an element of $\mathbb{C}[z_1, z_2]_4$. In our pictures the marked plane of the binode is vertical.

Convention: If the point is black (resp. white) then \mathcal{C}_2 does not pass (resp. passes) through the point. For all cases mentioned in the paper we precise $(\tilde{d}_1, \tilde{d}_2)$ where \tilde{d}_i is the degree of the tangent cone of \mathcal{C}_i at p .

Let us mention that this table in which we propose local illustrations could help the reader to visualize the different examples but the proofs are not based on it.

D.p. of contact						
binode						
D.p.'s						
pt of osculation						
pt of contact						

Table 1

Appendix

Hudson’s Table

In this appendix we give a reproduction of what Hudson called “Cubic Space Transformations”. The first (resp. second, resp. third, resp. fourth) table concerns birational maps of bidegrees (3,2), (3,3) and (3,4) (resp. (3,5), resp. (3,6), resp. (3,7), (3,8) and (3,9)).

number	degrees	D.p. of contact	binode	D. p.'s	pt of osculation	pt of contact	ordinary pts	F -curves	Remarks
1	3-2	l^2, l_1, l_2, l_3	3 generators meet double line
2	3-3	ω_6 (genus 3)	2 generators meet double line
3		.	.	1	.	.	.	$\omega_6 \equiv O^2$ (genus 3)	
4		1	$\omega_6 \equiv O^4$ (rational)	
5		2	l^2, l_1, l_2	
6	3-4	1	ω_5 (genus 1)	(ϕ) touch plane along l generator meets double line
7		.	.	1	.	.	1	$\omega_5 \equiv O_1^2$ (genus 1)	
8		.	1	.	.	.	1	$\omega_5 \equiv O_1^3(2)$ (rational)	
9		1	1	$\omega_3 \equiv O_1^2, l_1 \equiv O_1, l_2 \equiv O_1$	
10		1	1	1	.	.	1	$\omega_2 \equiv O_1O_2, l \equiv O_1O_2$ (osculation)	
11		4	l^2, l_1	
		

number	D.p. of contact	binode	D. p.'s	pt of osculation	pt of contact	ordinary pts	F-curves	Remarks
12	2	ω_3 (rational), l	
13	.	.	1	.	.	2	$\omega_4 \equiv O_1$ (genus 1)	
14	.	.	1	.	.	2	$\omega_3 \equiv O_1$ (rational), $l \equiv O_1$	
15	.	1	.	.	.	2	$\omega_4 \equiv O_1^2(2)$	
16	.	1	.	.	.	2	$\omega_2 \equiv O_1(1)$, $l_1 \equiv O_1(1)$, $l_2 \equiv O_1$	
17	1	2	$\omega_3 \equiv O_1^2$, $l_1 \equiv O_1$	
18	1	2	$l \equiv O_1$ (contact), $l_1 \equiv O_1$, $l_2 \equiv O_1$	(ϕ) touch quadric
19	.	.	2	.	.	2	$\omega_3 \equiv O_1O_2$ (rational), $l \equiv O_1O_2$	
20	.	1	1	.	.	2	$\omega_2 \equiv O_1(1)O_2$, $l_1 \equiv O_1O_2$, $l_2 \equiv O_1(1)$	(ϕ) touch plane
21	1	.	1	.	.	2	$l \equiv O_1O_2$ (contact), $l_1 \equiv O_1$, $l_2 \equiv O_1$	(ϕ) touch plane
22	1	1	.	.	.	2	$l \equiv O_1O_2(1)$ (osculation), $l_1 \equiv O_1$	
23	1	.	ω_4 (rational)	
24	.	.	1	.	1	.	$\omega_4 \equiv O_1^2$	
25	.	1	.	.	1	.	$\omega_3 \equiv O_1^2(1)$, $l \equiv O_1(1)$	
26	1	.	.	.	1	.	$l_1 \equiv O_1$, $l_2 \equiv O_1$, $l_3 \equiv O_1$, $l_4 \equiv O_1$	
27	6	l^2	

Cubic Space Transformations of bidegree (3, 5)

number	D.p. of contact	binode	D. p.'s	pt of osculation	pt of contact	ordinary pts	F -curves	Remarks
28	3	l (contact), l_1	
29	.	.	1	.	.	3	ω_3 (plane, genus 1)	
30	.	.	1	.	.	3	$\omega_2, l \equiv O_1$	
31	.	.	1	.	.	3	$l \equiv O_1$ (contact), l_1	
32	.	.	1	.	.	3	$l \equiv O_1$ (osculation)	(ϕ) touch quadric
33	.	1	.	.	.	3	$\omega_2 \equiv O_1(1), l \equiv O_1$	
34	1	3	$\omega_3 \equiv O_1^2$	(ϕ) touch quadric
35	1	3	$l \equiv O_1$ (contact), $l_1 \equiv O_1$	
36	.	.	2	.	.	3	$\omega_2 \equiv O_1, l \equiv O_1 O_2$	
37	.	1	1	.	.	3	$\omega_2 \equiv O_1(1)O_2, l \equiv O_1 O_2$	
38	.	1	1	.	.	3	$l \equiv O_1 O_2, l_1 \equiv O_1(1), l_2 \equiv O_1(1)$	
39	1	.	1	.	.	3	$l \equiv O_1 O_2$ (contact), $l_1 \equiv O_1$	(ϕ) touch plane
40	1	1	.	.	.	3	$l \equiv O_1 O_2(1)$ osculation	(ϕ) touch plane
41	1	1	l_1, l_2, l_3	
42	.	.	1	.	1	1	$\omega_3 \equiv O_1$ (rational)	
43	.	.	1	.	1	1	$l_1 \equiv O_1, l_2 \equiv O_1, l_3$	
44	.	1	.	.	1	1	$\omega_3 \equiv O_1^2(1)$	
45	.	1	.	.	1	1	$\omega_2 \equiv O_1(1), l \equiv O_1(1)$	
46	.	1	.	.	1	1	$l \equiv O_1(1)$ (contact), $l_1 \equiv O_1$	
47	1	.	.	.	1	1	$l_1 \equiv O_1, l_2 \equiv O_1, l_3 \equiv O_1$	(ϕ) touch quadric
48	.	.	2	.	1	1	$l \equiv O_1 O_2, l_1 \equiv O_1, l_2 \equiv O_2$	
49	.	1	1	.	1	1	$l \equiv O_1(1)O_2$ (contact), $l_2 \equiv O_1$	(ϕ) touch plane O_2 on fixed plane at O_1
50	.	.	3	.	1	1	$l_1 \equiv O_2 O_3, l_2 \equiv O_3 O_1, l_3 \equiv O_1 O_2$	
51	.	.	.	1	.	.	ω_3 (rational)	
52	.	.	1	1	.	.	$\omega_3 \equiv O_1^2$	

Cubic Space Transformations of bidegree (3, 6)

number	degrees	D.p. of contact	binode	D. p.'s	point of osculation	point of contact	ordinary points	F -curves	Remarks
53	3-7	1	4	$l \equiv O_1$ (contact)	(ϕ) touch quadric
54		.	.	2	.	.	4	$l \equiv O_1O_2, l_1$	(ϕ) touch plane
55		.	1	1	.	.	4	$l \equiv O_1O_2, l_1 \equiv O_1(1)$	
56		1	.	1	.	.	4	$l \equiv O_1O_2$ (contact)	
57		.	.	1	.	1	2	ω_2	
58		.	1	.	.	1	2	$\omega_2 \equiv O_1(1)$	(ϕ) touch quadric O_2 on fixed plane at O_1
59		1	.	.	.	1	2	$l_1 \equiv O_1, l_2 \equiv O_1$	
60		.	.	1	.	2	.	$l \equiv O_1, l_1$	
61		.	1	.	.	2	.	$l_1 \equiv O_1(1), l_2 \equiv O_1$	
62		.	1	.	.	2	.	$l \equiv O_1(1)$ (contact)	
63		.	.	2	.	2	.	$l \equiv O_1O_2, l_1 \equiv O_1$	
64		.	1	1	.	2	.	$l_1 \equiv O_1(1)O_2$ (contact)	
65		.	.	1	1	.	1	$\omega_2 \equiv O_1$	
66		.	1	.	1	.	1	$l_1 \equiv O_1(1), l_2 \equiv O_1(1)$	
67	3-8	.	1	1	.	.	5	$l \equiv O_1O_2$	
68		1	.	.	.	1	3	$l \equiv O_1$	
69		.	1	.	.	2	1	$l \equiv O_1$	
70		.	.	1	1	.	2	l	
71		.	1	.	1	.	2	$l \equiv O_1(1)$	
72	3-9	1	.	.	.	1	4		4-point contact at O_2
73		.	1	.	3	.	.		
74		.	1	.	1	.	3		
75		.	.	1	1	.	2		

BIBLIOGRAPHY

- [1] D. AVRITZER, G. GONZALEZ-SPRINBERG & I. PAN – « On singular quadratic complexes, quintic curves and Cremona transformations », *Rend. Circ. Mat. Palermo* **61** (2012), p. 201–240.
- [2] C. BISI, A. CALABRI & M. MELLA – « On plane Cremona transformations of fixed degree », *J. Geom. Anal.* **25** (2015), p. 1108–1131.
- [3] J. BLANC & J.-P. FURTER – « Topologies and structures of the Cremona groups », *Ann. of Math.* **178** (2013), p. 1173–1198.
- [4] S. CANTAT – « The Cremona group in two variables », *Proceedings of the sixth European Congress of Math.* (2013), p. 211–225.
- [5] ———, « Morphisms between Cremona groups, and characterization of rational varieties », *Compos. Math.* **150** (2014), p. 1107–1124.
- [6] D. CERVEAU & J. DÉSERTE – *Transformations birationnelles de petit degré*, Cours Spécialisés, vol. 19, Soc. Math. France, Paris, 2013.
- [7] L. CREMONA – « Sulle trasformazioni razionali nello spazio », *Annali di Mat.* **V** (1871-1873), p. 131–162.
- [8] M. DEMAZURE – « Sous-groupes algébriques de rang maximum du groupe de Cremona », *Ann. Sci. École Norm. Sup.* **3** (1970), p. 507–588.
- [9] J. DÉSERTE – *Some properties of the Cremona group*, Ensaios Matemáticos, vol. 21, Sociedade Brasileira de Matemática, 2012.
- [10] I. V. DOLGACHEV – *Classical algebraic geometry*, Cambridge Univ. Press, Cambridge, 2012.
- [11] H. P. HUDSON – *Cremona Transformations in Plane and Space*, Cambridge University Press, 1927.
- [12] I. PAN – « Sur les transformations de cremona de bidegré (3,3) », thèse de doctorat, Université de Genève, 1996.
- [13] ———, « Sur les transformations de Cremona de bidegré (3,3) », *Enseign. Math.* **43** (1997), p. 285–297.
- [14] ———, « Une remarque sur la génération du groupe de Cremona », *Bol. Soc. Brasil. Mat. (N.S.)* **30** (1999), p. 95–98.
- [15] I. PAN, F. RONGA & T. VUST – « Transformations birationnelles quadratiques de l'espace projectif complexe à trois dimensions », *Ann. Inst. Fourier (Grenoble)* **51** (2001), p. 1153–1187.
- [16] C. PESKINE & L. SZPIRO – « Liaison des variétés algébriques. I », *Invent. math.* **26** (1974), p. 271–302.
- [17] J-P. SERRE – « Le groupe de Cremona et ses sous-groupes finis », Séminaire Bourbaki, volume 2008/2009. exp. n° 1000, *Astérisque* **332** (2010), p. 75–100.

