

THE WEYL CALCULUS WITH RESPECT TO THE GAUSSIAN MEASURE AND RESTRICTED L^p - L^q BOUNDEDNESS OF THE ORNSTEIN-UHLENBECK SEMIGROUP IN COMPLEX TIME

BY JAN VAN NEERVEN & PIERRE PORTAL

ABSTRACT. — In this paper, we introduce a Weyl functional calculus $a \mapsto a(Q, P)$ for the position and momentum operators Q and P associated with the Ornstein-Uhlenbeck operator $L = -\Delta + x \cdot \nabla$, and give a simple criterion for restricted L^p - L^q boundedness of operators in this functional calculus. The analysis of this non-commutative functional calculus is simpler than the analysis of the functional calculus of L . It allows us to recover, unify, and extend old and new results concerning the boundedness of $\exp(-zL)$ as an operator from $L^p(\mathbb{R}^d, \gamma_\alpha)$ to $L^q(\mathbb{R}^d, \gamma_\beta)$ for suitable values of $z \in \mathbb{C}$ with $\operatorname{Re} z > 0$, $p, q \in [1, \infty)$, and $\alpha, \beta > 0$. Here, γ_τ denotes the centered Gaussian measure on \mathbb{R}^d with density $(2\pi\tau)^{-d/2} \exp(-|x|^2/2\tau)$.

Texte reçu le 13 février 2017, modifié le 14 août 2017, accepté le 13 septembre 2017.

JAN VAN NEERVEN, Delft University of Technology Delft Institute of Applied Mathematics P.O. Box 5031 2600 GA Delft The Netherlands • *E-mail* : J.M.A.M.vanNeerven@tudelft.nl

PIERRE PORTAL, Mathematical Sciences Institute, John Dedman Building The Australian National University ACTON ACT 0200 Australia and Laboratoire Paul Painlevé Université Lille 1 F-59655 Villeneuve D'Ascq France • *E-mail* : Pierre.Portal@anu.edu.au

Mathematical subject classification (2010). — 47A60, 47D06, 47G30, 60H07, 81S05.

Key words and phrases. — Weyl functional calculus, canonical commutation relation, Schur estimate, Ornstein-Uhlenbeck operator, Mehler kernel, restricted L^p - L^q -boundedness, restricted Sobolev embedding.

The authors gratefully acknowledge financial support from the Discovery grant DP160100941 of the Australian Research Council. The second author is also supported by the ARC Future Fellowship FT130100607.

RÉSUMÉ (*Calcul fonctionnel de Weyl pour la mesure gaussienne et estimées L^p - L^q restreintes du semigrroupe d'Ornstein-Uhlenbeck en temps complexe*). — Ce papier introduit un calcul fonctionnel de Weyl $a \mapsto a(Q, P)$ adapté aux opérateurs de position et d'impulsion Q et P associés à l'opérateur d'Ornstein-Uhlenbeck $L = -\Delta + x \cdot \nabla$, et fournit un critère simple pour prouver des estimées L^p - L^q restreintes de ce calcul fonctionnel. L'analyse de ce calcul fonctionnel non-commutatif se révèle être plus simple que celle du calcul fonctionnel de L . Ceci nous permet de redémontrer, d'unifier, et d'étendre des résultats anciens et nouveaux sur les propriétés de bornitude de $\exp(-zL)$ de $L^p(\mathbb{R}^d, \gamma_\alpha)$ dans $L^q(\mathbb{R}^d, \gamma_\beta)$ pour les valeurs appropriées de $z \in \mathbb{C}$ (avec $\operatorname{Re} z > 0$, $p, q \in [1, \infty)$, et $\alpha, \beta > 0$). La notation γ_τ est utilisée pour la mesure Gaussienne centrée sur \mathbb{R}^d de densité $(2\pi\tau)^{-d/2} \exp(-|x|^2/2\tau)$.

1. Introduction

In the standard Euclidean situation, pseudo-differential calculus arises as the Weyl joint functional calculus of a non-commuting pair of operators: the position and momentum operators (see, e.g., [9] and [11, Chapter XII]). In this paper, by transferring this calculus to the Gaussian setting, we introduce a Gaussian version of the Weyl pseudo-differential calculus which assigns to suitable functions $a : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{C}$ a bounded operator $a(Q, P)$ acting on $L^2(\mathbb{R}^d, \gamma)$. Here, $Q = (Q_1, \dots, Q_d)$ and $P = (P_1, \dots, P_d)$ are the position and momentum operators associated with the Ornstein-Uhlenbeck operator

$$L = -\Delta + x \cdot \nabla$$

on $L^2(\mathbb{R}^d, \gamma)$, where $d\gamma(x) = (2\pi)^{-d/2} \exp(-\frac{1}{2}|x|^2) dx$ is the standard Gaussian measure on \mathbb{R}^d . We show that the Ornstein-Uhlenbeck semigroup $\exp(-tL)$ can be expressed in terms of this calculus by the formula

$$(1) \quad \exp(-tL) = \left(1 + \frac{1 - e^{-t}}{1 + e^{-t}}\right)^d \exp\left(-\frac{1 - e^{-t}}{1 + e^{-t}}(P^2 + Q^2)\right).$$

With $s := \frac{1 - e^{-t}}{1 + e^{-t}}$, the expression on the right-hand side is defined through the Weyl calculus as $(1 + s)^d a_s(Q, P)$, where $a_s(x, \xi) = \exp(-s(|x|^2 + |\xi|^2))$. The main ingredient in the proof of (1) is the explicit determination of the integral kernel for $a_s(Q, P)$. By applying a Schur-type estimate to this kernel, we are able to prove the following sufficient condition for restricted L^p - L^q -boundedness of $a_s(Q, P)$:

THEOREM 1.1. — *Let $p, q \in [1, \infty)$ and let $\alpha, \beta > 0$. For $s \in \mathbb{C}$ with $\operatorname{Re} s > 0$, define $r_\pm(s) := \frac{1}{2} \operatorname{Re} \left(\frac{1}{s} \pm s\right)$. If s satisfies $1 - \frac{2}{\alpha p} + r_+(s) > 0$, $\frac{2}{\beta q} - 1 + r_+(s) > 0$, and*

$$(r_-(s))^2 \leq \left(1 - \frac{2}{\alpha p} + r_+(s)\right) \left(\frac{2}{\beta q} - 1 + r_+(s)\right),$$

then the operator $\exp(-s(P^2 + Q^2))$ is bounded from $L^p(\mathbb{R}^d, \gamma_\alpha)$ to $L^q(\mathbb{R}^d, \gamma_\beta)$.

Here, γ_τ denotes the centered Gaussian measure on \mathbb{R}^d with density $(2\pi\tau)^{-d/2} \exp(-|x|^2/2\tau)$ (so that $\gamma_1 = \gamma$ is the standard Gaussian measure). The proof of the theorem provides an explicit estimate for the norm of this operator that is of the correct order in the variable s , as subsequent corollaries show.

Taken together, (1) and Theorem 1.1 can then be used to obtain criteria for $L^p(\mathbb{R}^d, \gamma_\alpha)$ - $L^q(\mathbb{R}^d, \gamma_\beta)$ boundedness of $\exp(-zL)$ for suitable values of $z \in \mathbb{C}$ with $\operatorname{Re} z > 0$. Among other things, in Section 5 we show that the operators $\exp(-zL)$ map $L^1(\mathbb{R}^d; \gamma_2)$ to $L^2(\mathbb{R}^d; \gamma)$ for all $\operatorname{Re} z > 0$. We also prove a more precise boundedness result which, for real values $t > 0$ implies the boundedness of $\exp(-tL)$ from $L^1(\mathbb{R}^d, \gamma_{\alpha_t})$ to $L^2(\mathbb{R}^d, \gamma)$, where $\alpha_t = 1 + e^{-2t}$. The boundedness of these operators was proved recently by Bakry, Bolley, and Gentil [1] as a corollary of their work on hypercontractive bounds on Markov kernels for diffusion semigroups. As such, our results may be interpreted as giving an extension to complex time of the Bakry-Bolley-Gentil result for the Ornstein-Uhlenbeck semigroup.

In the final Section 6, we show that Theorem 1.1 also captures the well-known result of Epperson [2] (see also Weissler [12] for the first boundedness result of this type, and part of the contractivity result) for $1 < p \leq q < \infty$, the operator $\exp(-zL)$ is bounded from $L^p(\mathbb{R}^d, \gamma)$ to $L^q(\mathbb{R}^d, \gamma)$ if and only if $\omega := e^{-z}$ satisfies $|\omega|^2 < p/q$ and

$$(q-1)|\omega|^4 + (2-p-q)(\operatorname{Re} \omega)^2 - (2-p-q+pq)(\operatorname{Im} \omega)^2 + p-1 > 0.$$

In particular, for $p = q$, the semigroup $\exp(-tL)$ on $L^p(\mathbb{R}^d, \gamma)$ extends analytically to the set (see Figure 1.1)

$$(2) \quad E_p := \{z = x + iy \in \mathbb{C} : |\sin(y)| < \tan(\theta_p) \sinh(x)\},$$

where

$$(3) \quad \cos \phi_p = \left| \frac{2}{p} - 1 \right|.$$

These results demonstrate the potential of the Gaussian pseudo-differential calculus. Of course, taking (1) for granted, we could forget about the Gaussian pseudo-differential calculus altogether, and reinterpret all the applications given in this paper as consequences of the realization that through the time change $s \mapsto \frac{1-e^{-t}}{1+e^{-t}}$, various algebraic simplifications allow one to derive sharp results for the Ornstein-Uhlenbeck semigroup in a unified manner. In fact, as the referee of this paper pointed out to us, Weissler took exactly this approach in [12], and obtained the most important special case of our Theorem 1.1 in 1979. Besides generalizing this result to the context of weighted Gaussian measures γ_α arising from [1], the point of the new approach given here is to connect results such as Weissler's and other classical hypercontractivity theorems to the underlying Weyl calculus. In doing so, one sees the reason why certain crucial




FIGURE 1.1. The Epperson region E_p (red/orange) and the sector with angle θ_p for $p = 4/3$ (orange).

algebraic simplifications occur, and one develops a far more flexible tool to study other spectral multipliers associated with the Ornstein-Uhlenbeck operator (and, possibly, perturbations thereof). In such applications, the algebraic consequences of the fact that the Weyl calculus involves non-commuting operators may not be as easily unpacked as in (1). The L^p -analysis of operators in the Weyl calculus of the pair (Q, P) , however, is simpler than the direct analysis of operators in the functional calculus of L (or perturbations of L). In future works, we plan to develop this theory and include harmonic analysis substantially more advanced than the Schur-type estimate employed here, along with applications to non-linear stochastic differential equations.

Acknowledgements. — We are grateful to the anonymous referee for her/his useful suggestions, and, in particular, for pointing out to us the paper [12].

2. The Weyl calculus with respect to the Gaussian measure

In this section we introduce the Weyl calculus with respect to the Gaussian measure. To emphasize its Fourier analytic content, our point of departure is the fact that Fourier-Plancherel transform is unitarily equivalent to the second quantization of multiplication by $-i$. The unitary operator implementing this equivalence is used to define the position and momentum operators Q and P

associated with the Ornstein-Uhlenbeck operator L . This approach bypasses the use of creation and annihilation operators altogether and leads to the same expressions.

2.1. The Wiener-Plancherel transform with respect to the Gaussian measure.

— Let $dm(x) = (2\pi)^{-d/2}dx$ denote the normalized Lebesgue measure on \mathbb{R}^d . The mapping $E : f \mapsto ef$, where

$$e(x) := \exp\left(-\frac{1}{4}|x|^2\right),$$

is unitary from $L^2(\mathbb{R}^d, \gamma)$ onto $L^2(\mathbb{R}^d, m)$, and the dilation $\delta : L^2(\mathbb{R}^d, m) \rightarrow L^2(\mathbb{R}^d, m)$,

$$\delta f(x) := (\sqrt{2})^d f(\sqrt{2}x)$$

is unitary on $L^2(\mathbb{R}^d, m)$. Consequently, the operator

$$U := \delta \circ E$$

is unitary from $L^2(\mathbb{R}^d, \gamma)$ onto $L^2(\mathbb{R}^d, m)$. It was shown by Segal [8, Theorem 2] that U establishes a unitary equivalence

$$\mathscr{W} = U^{-1} \circ \mathscr{F} \circ U$$

of the Fourier-Plancherel transform \mathscr{F} as a unitary operator on $L^2(\mathbb{R}^d, m)$,

$$\mathscr{F}f(y) := \hat{f}(y) := \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} f(x) \exp(-ix \cdot y) dx = \int_{\mathbb{R}^d} f(x) \exp(-ix \cdot y) dm(x),$$

with the unitary operator \mathscr{W} on $L^2(\mathbb{R}^d, \gamma)$, defined for polynomials f by

$$\mathscr{W}f(y) := \int_{\mathbb{R}^d} f(-iy + \sqrt{2}x) d\gamma(x).$$

We have the following beautiful representation of this operator, which is sometimes called the *Wiener-Plancherel transform*, in terms of the second quantization functor Γ [8, Corollary 3.2]:

$$\mathscr{W} = \Gamma(-i).$$

This identity is not used in the sequel; it is stated only to demonstrate that both the operator \mathscr{W} and the unitary U are very natural.

2.2. Position and momentum with respect to the Gaussian measure. — Consider classical position and momentum operators

$$X = (x_1, \dots, x_d), \quad D = \left(\frac{1}{i}\partial_1, \dots, \frac{1}{i}\partial_d\right),$$

viewed as densely defined operators mapping from $L^2(\mathbb{R}^d)$ into $L^2(\mathbb{R}^d; \mathbb{C}^d)$. Explicitly, x_j is the densely defined self-adjoint operator on $L^2(\mathbb{R}^d)$ defined by

pointwise multiplication, i.e., $(x_j f)(x) := x_j f(x)$ for $x \in \mathbb{R}^d$, with maximal domain

$$D(x_j) = \{f \in L^2(\mathbb{R}^d) : x_j f \in L^2(\mathbb{R}^d)\},$$

and $\frac{1}{i}\partial_j$ is the self-adjoint operator $f \mapsto \frac{1}{i}\partial_j f$ with maximal domain

$$D\left(\frac{1}{i}\partial_j\right) = \{f \in L^2(\mathbb{R}^d) : \partial_j f \in L^2(\mathbb{R}^d)\},$$

the partial derivative being interpreted in the sense of distributions.

Having motivated our choice of the unitary U , it is now used to introduce the position and momentum operators $Q = (q_1, \dots, q_d)$ and $P = (p_1, \dots, p_d)$ as densely defined closed operators acting from their natural domains in $L^2(\mathbb{R}^d, \gamma)$ into $L^2(\mathbb{R}^d, \gamma; \mathbb{C}^d)$ by unitary equivalence with $X = (x_1, \dots, x_d)$ and $D = (\frac{1}{i}\partial_1, \dots, \frac{1}{i}\partial_d)$:

$$\begin{aligned} q_j &:= U^{-1} \circ x_j \circ U, \\ p_j &:= U^{-1} \circ \frac{1}{i}\partial_j \circ U. \end{aligned}$$

They satisfy the commutation relations

$$(4) \quad [p_j, p_k] = [q_j, q_k] = 0, \quad [q_j, p_k] = \frac{1}{i}\delta_{jk},$$

as well as the identity

$$(5) \quad \frac{1}{2}(P^2 + Q^2) = L + \frac{d}{2}I.$$

Here, L is the *Ornstein-Uhlenbeck operator* which acts on test functions $f \in C_c^2(\mathbb{R}^d)$ by

$$Lf(x) := -\Delta f(x) + x \cdot \nabla f(x) \quad (x \in \mathbb{R}^d).$$

It follows readily from the definition of the Wiener-Plancherel transform \mathscr{W} that

$$\begin{aligned} q_j \circ \mathscr{W} &= \mathscr{W} \circ p_j, \\ p_j \circ \mathscr{W} &= -\mathscr{W} \circ q_j \end{aligned}$$

consistent with the relations $x_j \circ \mathscr{F} = \mathscr{F} \circ (\frac{1}{i}\partial_j)$ and $(\frac{1}{i}\partial_j) \circ \mathscr{F} = -\mathscr{F} \circ x_j$ for position and momentum in the Euclidean setting.

REMARK 2.1. — Our definitions of P and Q coincide with the physicist's definitions in the theory of the quantum harmonic oscillator (cf. [4]). Other texts, such as [7], use different normalizations. The present choice makes the commutation relation between position and momentum as well as the identity relating the Ornstein-Uhlenbeck operator and position and momentum come out right

in the sense that (4) and (5) hold. The former says that position and momentum satisfy the “canonical commutation relations” and the latter says that the Hamiltonian $\frac{1}{2}(P^2 + Q^2)$ of the quantum harmonic oscillator equals the number operator L (physicists would write N) plus the ground-state energy $\frac{d}{2}$.

2.3. The Weyl calculus with respect to the Gaussian measure. — The Weyl calculus for the pair (X, D) is defined, for Schwartz functions $a : \mathbb{R}^{2d} \rightarrow \mathbb{C}$, by

$$a(X, D)f(y) = \int_{\mathbb{R}^{2d}} \widehat{a}(u, v) \exp(i(uX + vD))f(y) dm(u) dm(v).$$

Here, $m(dx) = (2\pi)^{-d/2} dx$ as before, $\widehat{a} := \mathcal{F}a$ is the Fourier-Plancherel transform of a , and the unitary operators $\exp(i(uX + vD))$ on $L^2(\mathbb{R}^d, \gamma)$ are defined through the action

$$(6) \quad \exp(i(uX + vD))f(y) := \exp\left(iuy + \frac{1}{2}iuv\right)f(v + y)$$

(cf. [11, Formula 51, page 550]). This definition can be motivated by a formal application of the Baker-Campbell-Hausdorff formula to the (unbounded) operators X and D ; alternatively, one may look upon it as defining a unitary representation of the Heisenberg group encoding the commutation relations of X and D , the so-called Schrödinger representation.

Motivated by the constructions in the preceding subsection, we formulate the following definition.

DEFINITION 2.2. — For $u, v \in \mathbb{R}^d$, on $L^2(\mathbb{R}^d, \gamma)$ we define the unitary operators $\exp(i(uQ + vP))$ on $L^2(\mathbb{R}^d; \gamma)$ by

$$\exp(i(uQ + vP)) := U^{-1} \circ \exp(i(uX + vD)) \circ U.$$

This allows us to define, for Schwartz functions $a : \mathbb{R}^{2d} \rightarrow \mathbb{C}$, the bounded operator $a(Q, P)$ on $L^2(\mathbb{R}^d, \gamma)$ by

$$(7) \quad a(Q, P) = U^{-1} \circ a(X, D) \circ U = \int_{\mathbb{R}^{2d}} \widehat{a}(u, v) \exp(i(uQ + vP)) dm(u) dm(v),$$

the integral being understood in the strong sense. An explicit expression for $a(Q, P)$ can be obtained as follows. By (6) and a change of variables, one has (cf. [11, Formula (52), page 551])

$$a(X, D)f(y) = \int_{\mathbb{R}^{2d}} a\left(\frac{1}{2}(v + y), \xi\right) \exp(-i\xi(v - y))f(v) dm(v) dm(\xi).$$

By (7) and the definition of U , this gives the following explicit formula for the Gaussian setting:

$$\begin{aligned}
 a(Q, P)f(y) &= \frac{1}{(2\pi)^d} \int_{\mathbb{R}^{2d}} a\left(\frac{1}{2}\left(x + \frac{y}{\sqrt{2}}\right), \xi\right) \exp\left(-i\xi\left(x - \frac{y}{\sqrt{2}}\right)\right) \\
 &\quad \cdot \exp\left(-\frac{1}{2}|x|^2 + \frac{1}{4}|y|^2\right) f(x\sqrt{2}) d\xi dx \\
 (8) \qquad &= \frac{1}{(2\sqrt{2}\pi)^d} \int_{\mathbb{R}^{2d}} a\left(\frac{x+y}{2\sqrt{2}}, \xi\right) \exp\left(-i\xi\left(\frac{x-y}{\sqrt{2}}\right)\right) \\
 &\quad \cdot \exp\left(-\frac{1}{4}|x|^2 + \frac{1}{4}|y|^2\right) f(x) d\xi dx \\
 &= \int_{\mathbb{R}^d} K_a(y, x) f(x) dx,
 \end{aligned}$$

where

$$\begin{aligned}
 (9) \qquad K_a(y, x) &:= \frac{1}{(2\sqrt{2}\pi)^d} \exp\left(-\frac{1}{4}|x|^2 + \frac{1}{4}|y|^2\right) \\
 &\quad \cdot \int_{\mathbb{R}^d} a\left(\frac{x+y}{2\sqrt{2}}, \xi\right) \exp\left(-i\xi\left(\frac{x-y}{\sqrt{2}}\right)\right) d\xi.
 \end{aligned}$$

3. Expressing the Ornstein-Uhlenbeck semigroup in the Weyl calculus

In order to translate results regarding the Weyl functional calculus of (Q, P) into results for the functional calculus of L , we first need to relate these two calculi. This is done in the next theorem. It is the only place where we rely on the concrete expression of the Mehler kernel.

THEOREM 3.1. — *For all $t > 0$ we have, with $s := \frac{1-e^{-t}}{1+e^{-t}}$,*

$$(10) \qquad \exp(-tL) = (1+s)^d a_s(Q, P),$$

where $a_s(x, \xi) := \exp(-s(|x|^2 + |\xi|^2))$.

In the next section we provide restricted L^p - L^q estimates for $a_s(Q, P)$ for complex values of s based purely on the Weyl calculus.

We need an elementary calculus lemma which is proved by writing out the inner product and square norm in terms of coordinates, thus writing the integral as a product of d integrals with respect to a single variable.

LEMMA 3.2. — *For all $A > 0$, $B \in \mathbb{R}$, and $y \in \mathbb{R}^d$,*

$$\int_{\mathbb{R}^d} \exp(-A|y|^2 + Bxy) dx = \left(\frac{\pi}{A}\right)^{d/2} \exp\left(\frac{B^2}{4A}|y|^2\right).$$

Proof of Theorem 3.1. — By (9) we have

$$\begin{aligned}
 K_{a_s}(y, x) &= \frac{1}{(2\sqrt{2}\pi)^d} \exp\left(-\frac{1}{4}|x|^2 + \frac{1}{4}|y|^2\right) \\
 &\quad \cdot \int_{\mathbb{R}^d} \exp\left(-s(|\xi|^2 + \frac{1}{8}|x+y|^2)\right) \exp\left(-i\xi\left(\frac{x-y}{\sqrt{2}}\right)\right) d\xi \\
 &= \frac{1}{(2\sqrt{2}\pi)^d} \exp\left(-\frac{s}{8}|x+y|^2\right) \exp\left(-\frac{1}{4}|x|^2 + \frac{1}{4}|y|^2\right) \\
 &\quad \cdot \int_{\mathbb{R}^d} \exp\left(-s\left(|\xi|^2 + \frac{i}{s}\xi\left(\frac{x-y}{\sqrt{2}}\right)\right)\right) d\xi \\
 &= \frac{1}{(2\sqrt{2}\pi)^d} \exp\left(-\frac{1}{8s}|x-y|^2\right) \exp\left(-\frac{s}{8}|x+y|^2\right) \\
 (11) \quad &\quad \cdot \exp\left(-\frac{1}{4}|x|^2 + \frac{1}{4}|y|^2\right) \int_{\mathbb{R}^d} \exp(-s|\eta|^2) d\eta \\
 &= \frac{1}{2^d(2\pi s)^{d/2}} \exp\left(-\frac{1}{8s}|x-y|^2\right) \exp\left(-\frac{s}{8}|x+y|^2\right) \\
 &\quad \cdot \exp\left(-\frac{1}{4}|x|^2 + \frac{1}{4}|y|^2\right) \\
 &= \frac{1}{2^d(2\pi s)^{d/2}} \exp\left(-\frac{1}{8s}(1-s)^2(|x|^2 + |y|^2) + \frac{1}{4}\left(\frac{1}{s} - s\right)xy\right) \\
 &\quad \cdot \exp\left(-\frac{1}{2}|x|^2\right)
 \end{aligned}$$

and therefore

$$\begin{aligned}
 (12) \quad \exp(-s(P^2 + Q^2))f(y) &= \int_{\mathbb{R}^d} K_{a_s}(y, x)f(x)dx \\
 &= \frac{1}{2^d(2\pi s)^{d/2}} \\
 &\quad \cdot \int_{\mathbb{R}^d} \exp\left(-\frac{1}{8s}(1-s)^2(|x|^2 + |y|^2) + \frac{1}{4}\left(\frac{1}{s} - s\right)xy\right) \\
 &\quad \cdot f(x)e^{-\frac{1}{2}|x|^2} dx.
 \end{aligned}$$

Taking $s := \frac{1-e^{-t}}{1+e^{-t}}$ in this identity we obtain

$$\begin{aligned}
 (13) \quad & \left(1 + \frac{1-e^{-t}}{1+e^{-t}}\right)^d \exp\left(-\frac{1-e^{-t}}{1+e^{-t}}(P^2 + Q^2)\right) f(y) \\
 &= \frac{1}{(2\pi)^{d/2}} \left(\frac{2}{1+e^{-t}}\right)^d \frac{1}{2^d} \left(\frac{1+e^{-t}}{1-e^{-t}}\right)^{d/2} \\
 &\quad \cdot \int_{\mathbb{R}^d} \exp\left(-\frac{1}{2} \frac{e^{-2t}}{1-e^{-2t}}(|x|^2 + |y|^2) + \frac{e^{-t}}{1-e^{-2t}}xy\right) f(x) \exp\left(-\frac{1}{2}|x|^2\right) dx \\
 &= \frac{1}{(2\pi)^{d/2}} \left(\frac{1}{1-e^{-2t}}\right)^{d/2} \int_{\mathbb{R}^d} \exp\left(-\frac{1}{2} \frac{|e^{-t}y - x|^2}{1-e^{-2t}}\right) f(x) dx \\
 &= \int_{\mathbb{R}^d} M_t(y, x) f(x) dx \\
 &= \exp(-tL) f(y),
 \end{aligned}$$

where

$$M_t(y, x) = \frac{1}{(2\pi)^{d/2}} \left(\frac{1}{1-e^{-2t}}\right)^{d/2} \exp\left(-\frac{1}{2} \frac{|e^{-t}y - x|^2}{1-e^{-2t}}\right)$$

denotes the Mehler kernel; the last step of (13) uses the classical Mehler formula for $\exp(-tL)$. \square

For any $z \in \mathbb{C}$ with $\operatorname{Re} z > 0$, the operator $\exp(-zL)$ is well defined and bounded as a linear operator on $L^2(\mathbb{R}^d, \gamma)$, and the same is true for the expression on the right-hand side in (10) by analytically extending the kernel defining it. By uniqueness of analytic extensions, the identity (10) persists for complex time.

The identity (10), extended analytically into the complex plane, allows the following deeper interpretation. The transformation

$$(14) \quad s = \frac{1-e^{-z}}{1+e^{-z}},$$

which is implicit in Theorem 3.1, is bi-holomorphic from

$$\{z \in \mathbb{C} : \operatorname{Re} z > 0, |\operatorname{Im}(z)| < \pi\}$$

onto

$$\{s \in \mathbb{C} : \operatorname{Re} s > 0, s \notin [1, \infty)\}.$$

For $1 < p < \infty$ it maps $E_p \cap \{z \in \mathbb{C} : |\operatorname{Im}(z)| < \pi\}$, where E_p is the Epperson region defined by (2), onto $\Sigma_{\theta_p} \setminus [1, \infty)$, where $\Sigma_{\theta_p} = \{s \in \mathbb{C} : s \neq 0, |\arg(s)| < \theta_p\}$ is the open sector with angle θ_p given by (3) (see Figure 1.1). Using the periodicity modulo $2\pi i$ of the exponential function, the mapping (14) maps E_p onto $\Sigma_{\theta_p} \setminus \{1\}$.

Using this information, the analytic extendibility of the semigroup $\exp(-tL)$ on $L^p(\mathbb{R}^d, \gamma)$ to E_p can now be proved by showing that $\exp(-s(P^2 + Q^2))$ extends analytically to Σ_{θ_p} ; the details are presented in Theorem 6.4. This shows that $\exp(-s(P^2 + Q^2))$ is a much simpler object than $\exp(-zL)$.

REMARK 3.3. — By (8) and (13), the theorem can be interpreted as giving a representation of the Mehler kernel in terms of the variable $\frac{1-e^{-t}}{1+e^{-t}}$. This representation could be taken as the starting point for the results in the next section, without any reference to the Weyl calculus. As we already pointed out in the Introduction, this would obscure the point that the Weyl calculus explains why the ensuing algebraic simplifications occur. What is more, the calculus can be applied to other functions $a(x, \xi)$ beyond the special choice $a_s(\xi, x) = \exp(-s(|x|^2 + |\xi|^2))$ and may serve as a tool to study spectral multipliers associated with the Ornstein-Uhlenbeck operator.

4. Restricted L^p - L^q estimates for $\exp(-s(P^2 + Q^2))$

Restricting the operators $\exp(-s(P^2 + Q^2))$ to $C_c^\infty(\mathbb{R}^d)$, we now take up the problem of determining when these restrictions extend to bounded operators from $L^p(\mathbb{R}^d, \gamma_\alpha)$ into $L^q(\mathbb{R}^d, \gamma_\beta)$. Here, for $\tau > 0$, we set

$$d\gamma_\tau(x) = (2\pi\tau)^{-d/2} \exp(-|x|^2/2\tau) dx$$

(so that $\gamma_1 = \gamma$ is the standard Gaussian measure). Boundedness (or rather, contractivity) from $L^p(\mathbb{R}^d, \gamma)$ to $L^q(\mathbb{R}^d, \gamma)$ corresponds to classical hypercontractivity of the Ornstein-Uhlenbeck semigroup. For other values of $\alpha, \beta > 0$ this includes restricted ultracontractivity of the type obtained in [1].

We begin with a sufficient condition for $L^p(\mathbb{R}^d, \gamma_\alpha)$ - $L^q(\mathbb{R}^d, \gamma_\beta)$ boundedness (Theorem 4.2 below). Recalling that $\exp(-s(P^2 + Q^2))$ equals the integral operator with kernel K_{a_s} given by (11), an immediate sufficient condition for boundedness derives from Hölder's inequality: if $p, q \in [1, \infty)$ and $\frac{1}{p} + \frac{1}{p'} = 1$, and

(15)

$$\int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} |K_a(y, x)|^{p'} \exp\left(\frac{p'}{2\alpha p} |x|^2\right) dx \right)^{q/p'} \exp(-|y|^2/2\beta) dy =: C < \infty$$

(with the obvious change if $p = 1$), then $a(Q, P)$ extends to a bounded operator from $L^p(\mathbb{R}^d, \gamma_\alpha)$ to $L^q(\mathbb{R}^d, \gamma_\beta)$ with norm at most C . A much sharper criterion can be given by using the following Schur-type estimate (which is a straightforward refinement of [10, Theorem 0.3.1]).

LEMMA 4.1. — Let $p, q, r \in [1, \infty)$ be such that $\frac{1}{r} = 1 - (\frac{1}{p} - \frac{1}{q})$. If $K \in L^1_{\text{loc}}(\mathbb{R}^{2d})$ and $\phi, \psi : \mathbb{R}^d \rightarrow (0, \infty)$ are integrable functions such that

$$\sup_{y \in \mathbb{R}^d} \left(\int_{\mathbb{R}^d} |K(y, x)|^r \frac{\psi^{r/q}(y)}{\phi^{r/p}(x)} dx \right)^{1/r} =: C_1 < \infty,$$

and

$$\sup_{x \in \mathbb{R}^d} \left(\int_{\mathbb{R}^d} |K(y, x)|^r \frac{\psi^{r/q}(y)}{\phi^{r/p}(x)} dy \right)^{1/r} =: C_2 < \infty$$

then

$$T_K f(y) := \int_{\mathbb{R}^d} K(y, x) f(x) dx \quad (f \in C_c(\mathbb{R}^d))$$

defines a bounded operator T_K from $L^p(\mathbb{R}^d, \phi(x)dx)$ to $L^q(\mathbb{R}^d, \psi(x)dx)$ with norm

$$\|T_K\|_{L^p(\mathbb{R}^d, \phi(x)dx), L^q(\mathbb{R}^d, \psi(x)dx)} \leq C_1^{1-\frac{r}{q}} C_2^{\frac{r}{q}}.$$

Proof. — First we consider the case $r \in (1, \infty)$.

For strictly positive functions $\eta \in L^1(\mathbb{R}^d)$ denote by $L^s_\eta(\mathbb{R}^d)$ the Banach space of measurable functions g such that $\eta g \in L^s(\mathbb{R}^d)$, identifying two such functions g if they are equal almost everywhere. From

$$\begin{aligned} |T_K f(y)| &\leq \int_{\mathbb{R}^d} |K(y, x)| |f(x)| \frac{1}{\phi^{1/p}(x)} \phi^{1/p}(x) dx \\ &\leq \left(\int_{\mathbb{R}^d} |K(y, x)|^r \frac{1}{\phi^{r/p}(x)} dx \right)^{1/r} \|f\|_{L^{r'}_{\phi^{1/p}}(\mathbb{R}^d)} \\ &= \frac{1}{\psi^{1/q}(y)} \left(\int_{\mathbb{R}^d} |K(y, x)|^r \frac{\psi^{r/q}(y)}{\phi^{r/p}(x)} dx \right)^{1/r} \|f\|_{L^{r'}_{\phi^{1/p}}(\mathbb{R}^d)} \end{aligned}$$

we find that

$$\|T_K f\|_{L^\infty_{\psi^{1/q}}(\mathbb{R}^d)} \leq C_1 \|f\|_{L^{r'}_{\phi^{1/p}}(\mathbb{R}^d)}.$$

This means that

$$T_K : L^{r'}_{\phi^{1/p}}(\mathbb{R}^d) \rightarrow L^\infty_{\psi^{1/q}}(\mathbb{R}^d)$$

is bounded with norm at most C_1 . With $K'(y, x) := \overline{K(x, y)}$, the same argument gives that $T_K^* = T_{K'}$ extends to a bounded operator from $L^{r'}_{(1/\psi)^{1/q}}(\mathbb{R}^d)$ to $L^\infty_{(1/\phi)^{1/p}}(\mathbb{R}^d)$ with norm at most C_2 . Dualizing, this implies that

$$T_K : L^1_{\phi^{1/p}}(\mathbb{R}^d) \rightarrow L^r_{\psi^{1/q}}(\mathbb{R}^d)$$

is bounded with norm at most C_2 .

This puts us into a position to apply the Riesz-Thorin theorem. Choose $0 < \theta < 1$ in such a way that $\frac{1}{p} = \frac{1-\theta}{r'} + \frac{\theta}{1}$, that is, $\frac{\theta}{r} = \frac{1}{p} - (1 - \frac{1}{r}) = \frac{1}{q}$, so $\theta = \frac{r}{q}$. In view of $\frac{1}{q} = \frac{1-\theta}{\infty} + \frac{\theta}{r}$ it follows that

$$T_K : L^p_{\phi^{1/p}}(\mathbb{R}^d) \rightarrow L^q_{\psi^{1/q}}(\mathbb{R}^d)$$

is bounded with norm at most $C = C_1^{1-\theta} C_2^\theta = C_1^{1-\frac{r}{q}} C_2^{\frac{r}{q}}$. But this means that

$$T_K : L^p(\mathbb{R}^d, \phi(x)dx) \rightarrow L^q(\mathbb{R}^d, \psi(x)dx)$$

is bounded with norm at most C .

The same proof works in the case $r = 1$ (which implies $p = q$), provided we check that in the duality argument, the adjoint operator $T_{K'}^*$ maps $L^1_{\phi^{1/p}}(\mathbb{R}^d)$ into $L^1_{\psi^{1/q}}(\mathbb{R}^d)$ (rather than into the bidual of $L^1_{\psi^{1/q}}(\mathbb{R}^d)$). For this, it suffices to check that for functions $f \in C_c(\mathbb{R}^d)$ one has $T_K f \in L^1_{\psi^{1/q}}(\mathbb{R}^d)$. This gives the desired conclusion, for $\langle T_K f, g \rangle = \langle f, T_{K'} g \rangle = \langle g, T_{K'}^* f \rangle$ for all $g \in L^\infty_{(1/\psi)^{1/q}}(\mathbb{R}^d) = (L^1_{(1/\psi)^{1/q}}(\mathbb{R}^d))^*$ implies that $T_{K'}^* f = T_K f \in L^1_{\psi^{1/q}}(\mathbb{R}^d)$.

If the support of f is in the rectangle $[-\rho, \rho]^d$, then by Fubini's theorem

$$\begin{aligned} \int_{\mathbb{R}^d} |T_K f(y)| \psi^{1/q}(y) dy &= \int_{\mathbb{R}^d} \left| \int_{[-\rho, \rho]^d} K(y, x) \frac{\psi^{1/q}(y)}{\phi^{1/p}(x)} \phi^{1/p}(x) f(x) dx \right| dy \\ &\leq \|f\|_\infty \int_{[-\rho, \rho]^d} \left(\int_{\mathbb{R}^d} |K(y, x)| \frac{\psi^{1/q}(y)}{\phi^{1/p}(x)} dy \right) \phi^{1/p}(x) dx \\ &\leq C_2 \|f\|_\infty \int_{[-\rho, \rho]^d} \phi^{1/p}(x) dx, \end{aligned}$$

and the last expression is finite by Hölder's inequality and the local integrability of ϕ . \square

Motivated by (12), for $s \in \mathbb{C}$ with $\operatorname{Re} s > 0$, we define

$$(16) \quad b_s := \frac{1}{8s}(1-s)^2, \quad c_s := \frac{1}{4} \left(\frac{1}{s} - s \right).$$

Setting

$$r_\pm(s) := \frac{1}{2} \operatorname{Re} \left(\frac{1}{s} \pm s \right)$$

we have the identities $\frac{1}{4} + \operatorname{Re} b_s = \frac{1}{4} r_+(s)$ and $\operatorname{Re} c_s = \frac{1}{2} r_-(s)$.

THEOREM 4.2 (Restricted L^p - L^q boundedness). — *Let $p, q \in [1, \infty)$, let $\frac{1}{r} = 1 - (\frac{1}{p} - \frac{1}{q})$, and let $\alpha, \beta > 0$. If $s \in \mathbb{C}$ with $\operatorname{Re} s > 0$ satisfies $1 - \frac{2}{\alpha p} + r_+(s) > 0$, $\frac{2}{\beta q} - 1 + r_+(s) > 0$, and*

$$(17) \quad (r_-(s))^2 \leq \left(1 - \frac{2}{\alpha p} + r_+(s) \right) \left(\frac{2}{\beta q} - 1 + r_+(s) \right),$$

then the operator $\exp(-s(P^2 + Q^2))$ is bounded from $L^p(\mathbb{R}^d, \gamma_\alpha)$ to $L^q(\mathbb{R}^d, \gamma_\beta)$ with norm

$$\begin{aligned} & \|\exp(-s(P^2 + Q^2))\|_{\mathcal{L}(L^p(\mathbb{R}^d, \gamma_\alpha), L^q(\mathbb{R}^d, \gamma_\beta))} \\ & \leq \frac{1}{(2rs)^{d/2}} \frac{\left(\frac{\alpha r}{2}\right)^{d/2p} \left(\frac{\beta r}{2}\right)^{-d/2q}}{\left(1 - \frac{2}{\alpha p} + r_+(s)\right)^{\frac{d}{2}(1 - \frac{1}{p})} \left(\frac{2}{\beta q} - 1 + r_+(s)\right)^{\frac{d}{2}\frac{1}{q}}}. \end{aligned}$$

REMARK 4.3. — We have no reason to believe that the numerical constant $\left(\frac{1}{2r}\right)^{d/2} \left(\frac{\alpha r}{2}\right)^{d/2p} \left(\frac{\beta r}{2}\right)^{-d/2q}$ is sharp, but the examples that we are about to work out indicate that the dependence on s is of the correct order.

REMARK 4.4. — For $s = x + iy \in \mathbb{C}$ with $x > 0$, we have $r_+(s) = \frac{1}{2} \left(\frac{x}{x^2 + y^2} + x \right) > 0$. It follows that the positivity assumptions $1 - \frac{2}{\alpha p} + r_+(s) > 0$ and $\frac{2}{\beta q} - 1 + r_+(s) > 0$ are fulfilled for all $\operatorname{Re} s > 0$ if, respectively, $\alpha p \geq 2$ and $\beta q \leq 2$.

Proof. — Using the notation of (16), the condition (17) is equivalent to

$$(18) \quad (\operatorname{Re} c_s)^2 \leq 4 \left(\frac{1}{2} - \frac{1}{2\alpha p} + \operatorname{Re} b_s \right) \left(\frac{1}{2\beta q} + \operatorname{Re} b_s \right).$$

We prove the theorem by checking the criterion of Lemma 4.1 for $K = K_{a_s}$ with $a_s(x, \xi) = \exp(-s(|x|^2 + |\xi|^2))$, and $\phi(x) = (2\pi\alpha)^{-d/2} \exp(-|x|^2/2\alpha)$, $\psi(x) = (2\pi\beta)^{-d/2} \exp(-|x|^2/2\beta)$.

By (11), for almost all $x, y \in \mathbb{R}^d$, we have

$$K_{a_s}(y, x) = \frac{1}{2^d (2\pi s)^{d/2}} \exp(-b_s(|x|^2 + |y|^2) + c_s xy) \exp\left(-\frac{1}{2}|x|^2\right).$$

Let $r \in [1, \infty)$ be such that $\frac{1}{r} = 1 - \left(\frac{1}{p} - \frac{1}{q}\right)$. Using Lemma 3.2, applied with $A = r \left(\frac{1}{2} - \frac{1}{2\alpha p} + \operatorname{Re} b_s\right)$ and $B = r \operatorname{Re} c_s$, we may estimate

$$\begin{aligned} & \sup_{y \in \mathbb{R}^d} \left(\int_{\mathbb{R}^d} |K_{a_s}(y, x)|^r (2\pi\alpha)^{rd/2p} \exp\left(\frac{1}{p}r|x|^2/2\alpha\right) (2\pi\alpha)^{-rd/2p} \right. \\ & \quad \cdot \exp\left(-\frac{1}{q}r|y|^2/2\beta\right) dx \Big)^{1/r} \\ & = \frac{(2\pi\alpha)^{d/2p} (2\pi\beta)^{-d/2q}}{2^d (2\pi s)^{d/2}} \\ & \quad \cdot \sup_{y \in \mathbb{R}^d} \left(\int_{\mathbb{R}^d} \exp(-r \operatorname{Re} b_s(|x|^2 + |y|^2) + r \operatorname{Re} c_s xy) \right. \\ & \quad \cdot \exp\left(-r \left(\frac{1}{2} - \frac{1}{2\alpha p}\right)|x|^2\right) \exp\left(-\frac{1}{2\beta q}r|y|^2\right) dx \Big)^{1/r} \end{aligned}$$

$$\begin{aligned}
&= \frac{(2\pi\alpha)^{d/2p}(2\pi\beta)^{-d/2q}}{2^d(2\pi s)^{d/2}} \\
&\quad \cdot \sup_{y \in \mathbb{R}^d} \left[\exp \left(- \left(\frac{1}{2\beta q} + \operatorname{Re} b_s \right) |y|^2 \right) \right. \\
&\quad \cdot \left. \left(\int_{\mathbb{R}^d} \exp \left(-r \left(\frac{1}{2} - \frac{1}{2\alpha p} + \operatorname{Re} b_s \right) |x|^2 + r \operatorname{Re} c_s xy \right) dx \right)^{1/r} \right] \\
&= \frac{(2\pi\alpha)^{d/2p}(2\pi\beta)^{-d/2q}}{2^d(2\pi s)^{d/2}} \left(\frac{\pi}{r \left(\frac{1}{2} - \frac{1}{2\alpha p} + \operatorname{Re} b_s \right)} \right)^{d/2r} \\
&\quad \cdot \sup_{y \in \mathbb{R}^d} \left[\exp \left(- \left(\frac{1}{2\beta q} + \operatorname{Re} b_s \right) |y|^2 \right) \exp \left(\frac{(\operatorname{Re} c_s)^2}{4 \left(\frac{1}{2} - \frac{1}{2\alpha p} + \operatorname{Re} b_s \right)} |y|^2 \right) \right] \\
&= \frac{(2\pi\alpha)^{d/2p}(2\pi\beta)^{-d/2q}}{2^d(2\pi)^{d/2}} \left(\frac{\pi}{r} \right)^{d/2r} \frac{1}{s^{d/2}} \left(\frac{1}{\frac{1}{2} - \frac{1}{2\alpha p} + \operatorname{Re} b_s} \right)^{d/2r} \\
&= \frac{(2\alpha)^{d/2p}(2\beta)^{-d/2q}}{2^{3d/2}} \frac{1}{r^{d/2r}} \frac{1}{s^{d/2}} \left(\frac{1}{\frac{1}{2} - \frac{1}{2\alpha p} + \operatorname{Re} b_s} \right)^{d/2r}.
\end{aligned}$$

In the same way, using Lemma 3.2 applied with $A = r(\frac{1}{\beta q} + \operatorname{Re} b_s)$ and $B = r \operatorname{Re} c_s$,

$$\begin{aligned}
&\sup_{x \in \mathbb{R}^d} \left(\int_{\mathbb{R}^d} |K_{a_s}(y, x)|^r \exp \left(\frac{1}{2\alpha p} r |x|^2 \right) \exp \left(- \frac{1}{2\beta q} r |y|^2 \right) dy \right)^{1/r} \\
&= \frac{(2\alpha)^{d/2p}(2\beta)^{-d/2q}}{2^{3d/2}} \frac{1}{r^{d/2r}} \frac{1}{s^{d/2}} \left(\frac{1}{\frac{1}{2\beta q} + \operatorname{Re} b_s} \right)^{d/2r}.
\end{aligned}$$

Denoting these two bounds by C_1 and C_2 , Lemma 4.1 bounds the norm of the operator by $C_1^{1-\frac{r}{q}} C_2^{\frac{r}{q}} = C_1^{r(1-\frac{1}{p})} C_2^{\frac{r}{q}}$. After rearranging the various constants a bit, this gives the estimate in the statement of the theorem. \square

REMARK 4.5. — In the above proof, one could replace the Schur test (Lemma 4.1) by the weaker condition (15) based on Hölder's inequality. This would have the effect of replacing the suprema by integrals throughout the proof. This not only leads to suboptimal estimates, but more importantly, it would not allow handling of the critical case when (17) holds with equality.

Combining Theorems 3.1 and 4.2, we obtain the following boundedness result for the operators $\exp(-zL)$.

COROLLARY 4.6. — *Let $s \in \mathbb{C}$ with $\operatorname{Re} s > 0$ satisfy the conditions of the theorem and define $z \in \mathbb{C}$ by $s = \frac{1-e^{-z}}{1+e^{-z}}$. Then,*

$$\begin{aligned} & \|\exp(-zL)\|_{\mathcal{L}(L^p(\mathbb{R}^d, \gamma_\alpha), L^q(\mathbb{R}^d, \gamma_\beta))} \\ & \leq \frac{2^d C}{|1 - e^{-2z}|^{\frac{d}{2}}} \frac{1}{\left(1 - \frac{2}{\alpha p} + \operatorname{Re} \frac{1+e^{-2z}}{1-e^{-2z}}\right)^{\frac{d}{2}\left(1-\frac{1}{p}\right)} \left(\frac{2}{\beta q} - 1 + \operatorname{Re} \frac{1+e^{-2z}}{1-e^{-2z}}\right)^{\frac{d}{2}\frac{1}{q}}}, \end{aligned}$$

where C is the numerical constant in Theorem 4.2 (cf. Remark 4.3).

Proof. — Noting that $2/(1+e^{-z}) = 1+s$, we have

$$\begin{aligned} \|\exp(-zL)\| & \leq |1+s|^d \|\exp(-s(P^2+Q^2))\| \\ & \leq C|1+s|^d \frac{1}{|s|^{d/2}} \frac{1}{|1 - \frac{2}{\alpha p} + r_+(s)|^{\frac{d}{2}\left(1-\frac{1}{p}\right)} \left|\frac{2}{\beta q} - 1 + r_+(s)\right|^{\frac{d}{2}\frac{1}{q}}}. \end{aligned}$$

The result follows from this by substituting $r_+(s) = \frac{1}{2} \operatorname{Re}(\frac{1}{s} + s) = \operatorname{Re} \frac{1+e^{-2z}}{1-e^{-2z}}$. \square

5. Restricted L^p - L^2 boundedness and Sobolev embedding

As a first application of Theorem 4.2, we have the following “hyperboundedness” result for real times $t > 0$:

COROLLARY 5.1. — *For $p \in [1, 2]$ and $t > 0$ set $\alpha_{p,t} := (1+e^{-2t})/p$.*

- (1) *For all $t > 0$, the operator $\exp(-tL)$ is bounded from $L^1(\mathbb{R}^d, \gamma_{\alpha_{1,t}})$ to $L^2(\mathbb{R}^d, \gamma)$, with norm*

$$\|\exp(-tL)\|_{\mathcal{L}(L^1(\mathbb{R}^d, \gamma_{\alpha_{1,t}}), L^2(\mathbb{R}^d, \gamma))} \lesssim_d (1 - e^{-4t})^{-d/4}.$$

- (2) *For all $p \in [1, 2]$ and $t > 0$ the operator $\exp(-tL)$ is bounded from $L^p(\mathbb{R}^d, \gamma_{\alpha_{p,t}})$ to $L^2(\mathbb{R}^d, \gamma)$, with norm*

$$\|\exp(-tL)\|_{\mathcal{L}(L^p(\mathbb{R}^d, \gamma_{\alpha_{p,t}}), L^2(\mathbb{R}^d, \gamma))} \lesssim_{d,p} t^{-\frac{d}{2}\left(\frac{1}{p}-\frac{1}{2}\right)} \quad \text{as } t \downarrow 0.$$

Proof. — Elementary algebra shows that with $\alpha_{p,t} = \frac{2}{p} \left(1 + \frac{2s}{1+s^2}\right)$ and $s = \frac{1-e^{-t}}{1+e^{-t}}$, the criterion of Theorem 4.2 holds for all $t \geq 0$ (with equality in (17)). Both norm estimates follow from Corollary 4.6, the first by taking $p = 1$, the second by noting that $\operatorname{Re} \left(\frac{1+e^{-2t}}{1-e^{-2t}}\right) \sim \frac{1}{t}$ for small values of t . \square

A sharp version of this corollary is due to Bakry, Bolley and Gentil [1, Section 4.2, Eq. (28)], who showed (for $p = 1$) the hypercontractivity bound

$$\|\exp(-tL)\|_{\mathcal{L}(L^1(\mathbb{R}^d, \gamma_{\alpha_{1,t}}), L^2(\mathbb{R}^d, \gamma))} \leq (1 - e^{-4t})^{-d/4}.$$

Their proof relies on entirely different techniques, which seem not to generalize to complex time so easily.

The next corollary gives “ultraboundedness” of the operators $\exp(-zL)$ for arbitrary $\operatorname{Re} z > 0$ from $L^p(\mathbb{R}^d, \gamma_{2/p})$ into $L^2(\mathbb{R}^d, \gamma)$:

COROLLARY 5.2. — *Let $p \in [1, 2]$. For all $z \in \mathbb{C}$ with $\operatorname{Re} z > 0$ the operator $\exp(-zL)$ maps $L^p(\mathbb{R}^d, \gamma_{2/p})$ into $L^2(\mathbb{R}^d, \gamma)$. As a consequence, the semigroup generated by $-L$ extends to a strongly continuous holomorphic semigroup of angle $\frac{1}{2}\pi$ on $L^p(\mathbb{R}^d, \gamma_{2/p})$. For each $\theta \in (0, \frac{1}{2}\pi)$ this semigroup is uniformly bounded on the sector $\{z \in \mathbb{C} : z \neq 0, |\arg(z)| < \theta\}$.*

Proof. — This follows from Corollary 4.6 upon realizing that the assumptions of Theorem 4.2 are satisfied when $q = 2$, $\equiv 1$ and $\alpha = \frac{2}{p}$, or $q = p$ and $\alpha = \beta = \frac{2}{p}$. \square

A notable consequence of Corollary 4.6 is the following (restricted) Sobolev embedding result. It is interesting because $(I + L)^{-1}$ maps $L^p(\mathbb{R}^d, \gamma)$ into $L^2(\mathbb{R}^d, \gamma)$ only when $p = 2$ (i.e. no full Sobolev embedding theorem holds in the Ornstein-Uhlenbeck context).

COROLLARY 5.3 (Restricted Sobolev embedding). — *Let $p \in \left(\frac{2d}{d+2}, 2\right]$. The resolvent $(I + L)^{-1}$ maps $L^p(\mathbb{R}^d, \gamma_{2/p})$ into $L^2(\mathbb{R}^d, \gamma)$.*

Proof. — Let $p \in \left(\frac{2d}{d+2}, 2\right]$ and fix $u \in L^p(\mathbb{R}^d, \gamma_{2/p}) \cap L^2(\mathbb{R}^d, \gamma)$. Then

$$\|\exp(-t(I + L))u\|_{L^2(\mathbb{R}^d, \gamma)} \lesssim_{d,p} \|u\|_{L^p(\mathbb{R}^d, \gamma_{2/p})} \exp(-t)t^{-\frac{d}{2}(\frac{1}{p}-\frac{1}{2})} \quad \forall t \geq 0,$$

and thus

$$\begin{aligned} \|(I + L)^{-1}u\|_{L^2(\mathbb{R}^d, \gamma)} &\lesssim_{d,p} \|u\|_{L^p(\mathbb{R}^d, \gamma_{2/p})} \int_0^\infty \exp(-t)t^{-\frac{d}{2}(\frac{1}{p}-\frac{1}{2})} dt \\ &\lesssim \|u\|_{L^p(\mathbb{R}^d, \gamma_{2/p})}, \end{aligned}$$

since $p \in \left(\frac{2d}{d+2}, 2\right]$ implies $\frac{d}{2} \left(\frac{1}{p} - \frac{1}{2}\right) < 1$. \square

6. L^p - L^q Boundedness

We now turn to the classical setting of the spaces $L^p(\mathbb{R}^d, \gamma)$, where γ is the standard Gaussian measure. For $\alpha = \beta = 1$ and $s = x + iy$, the first positivity condition of Theorem 4.2 takes the form

$$(19) \quad 1 - \frac{2}{p} + r_+(s) > 0 \iff 1 - \frac{2}{p} + \frac{1}{2} \left(\frac{x}{x^2 + y^2} + x \right) > 0$$

whereas condition (17) is seen to be equivalent to the condition

$$(20) \quad (p - q) \left(x + \frac{x}{x^2 + y^2} \right) + pq \left(\frac{x^2}{x^2 + y^2} - 1 \right) + 2p + 2q - 4 \geq 0.$$

Let us also observe that if these two conditions hold, together they enforce the second positivity condition $\frac{2}{q} - 1 + r_+(s) > 0$; this is apparent from the representation in (17).

As a warm-up for the general case, let us first consider real times $t \in (0, 1)$ in the z -plane, which correspond to the values $s = x \in (0, 1)$ in the s -plane. The conditions (19) and (20) then reduce to

$$\left(1 - \frac{2}{p} \right) x + \frac{1}{2}(x^2 + 1) > 0$$

and

$$(p - q) \left(x + \frac{1}{x} \right) + 2p + 2q - 4 \geq 0,$$

respectively. The first condition is automatic. Substituting $x = \frac{1-e^{-t}}{1+e^{-t}}$ in the second and solving for e^{-t} , assuming $p \leq q$, we find that it is equivalent to the condition

$$e^{-2t} \leq \frac{p-1}{q-1}.$$

Thus we recover the boundedness part of Nelson's celebrated hypercontractivity result [6].

Turning to complex time, with some additional effort we also recover the following result due to Weissler [12] (see also Epperson [2] for further refinements), essentially as a Corollary of Theorem 4.2.

THEOREM 6.1 (L^p - L^q -boundedness of $\exp(-zL)$). — *Let $1 < p \leq q < \infty$. If $z \in \mathbb{C}$ satisfies $\operatorname{Re} z > 0$,*

$$(21) \quad |e^{-z}|^2 < p/q$$

and

$$(22) \quad (q-1)|e^{-z}|^4 + (2-p-q)(\operatorname{Re} e^{-z})^2 - (2-p-q+pq)(\operatorname{Im} e^{-z})^2 + p-1 > 0,$$

then the operator $\exp(-zL)$ maps $L^p(\mathbb{R}^d, \gamma)$ into $L^q(\mathbb{R}^d, \gamma)$.

Before turning to the proof, we make a couple of preliminary observations. By a simple argument involving quadratic forms (see [2, page 3]), the conditions (21) and (22) taken together are equivalent to the single condition

$$(23) \quad (\operatorname{Im}(we^{-z}))^2 + (q-1)(\operatorname{Re}(we^{-z}))^2 < (\operatorname{Im} w)^2 + (p-1)(\operatorname{Re} w)^2 \quad \forall w \in \mathbb{C}.$$

Let us denote the set of all $z \in \mathbb{C}$, $\operatorname{Re} z \geq 0$, for which (23) holds by $E_{p,q}$. The following two facts hold:

FACTS 6.2. —

- $E_{p,p} = E_p$.
- $E_{p,q} \subseteq E_p$ and $E_{p,q} \subseteq E_q$.

The first is implicit in [2, 3], can be proved by elementary means, and is taken for granted. The second is an immediate consequence of the assumption $p \leq q$.

Let us now start with the proof of Theorem 6.1. It is useful to dispose of the positivity condition (19) in the form of a lemma; see also Figure 6.1.




FIGURE 6.1. The region $R_p := \{s \in \mathbb{C} : 1 - \frac{2}{p} + r_+(s) > 0\}$ (red/orange) and the sector Σ_p (orange), both for $p = 4/3$. Lemma 6.3 implies that Σ_p is indeed contained in R_p .

Let $z \in \mathbb{C}$ satisfy $\operatorname{Re} z > 0$. By the remarks at the end of Section 3, z belongs to E_p if and only if $s = \frac{1-e^{-z}}{1+e^{-z}}$ belongs to $\Sigma_{\phi_p} \setminus \{1\}$.

LEMMA 6.3. — *Every $s \in \Sigma_{\phi_p}$ satisfies the positivity condition (19).*

Proof. — Writing $s = x + iy$, we then have

$$\frac{x^2}{x^2 + y^2} = \cos^2 \theta_p > \left(1 - \frac{2}{p}\right)^2,$$

where the angle θ_p is given by (3). To see that this implies (19), note that

$$(24) \quad 1 - \frac{2}{p} + \frac{1}{2} \left(\frac{x}{x^2 + y^2} + x \right) > 2 \left(1 - \frac{2}{p} \right) x + \left(1 - \frac{2}{p} \right)^2 + x^2 = \left(x + \left(1 - \frac{2}{p} \right) \right)^2$$

and the latter is trivially true. \square

Proof of Theorem 6.1. — Fix $\operatorname{Re} z > 0$ and set $s := \frac{1-e^{-z}}{1+e^{-z}}$. We show that the assumptions of the theorem imply the conditions of Theorem 4.2, so that $\exp(-s(P^2 + Q^2))$ maps $L^p(\mathbb{R}^d, \gamma)$ into $L^q(\mathbb{R}^d, \gamma)$. In combination with Theorem 3.1, this gives the result.

We begin by checking the condition (19). For this, the second fact tells us that there is no loss of generality in assuming that $q = p$. In that situation, the first fact tells us that z belongs to E_p . But then Lemma 6.3 gives us the desired result.

It remains to check (20). Multiplying both sides with $x^2 + y^2$, this can be rewritten as

$$(25) \quad (p - q)x(1 + x^2 + y^2) + pqx^2 - (pq - 2p - 2q + 4)(x^2 + y^2) \geq 0.$$

The proof of the theorem is completed by showing that (22) implies (25).

Towards this end, we rewrite (22) in a similar way. Setting $e^{-z} = \frac{1-s}{1+s}$ with $s = x + iy$, and using that

$$\operatorname{Re} \frac{1 - x - iy}{1 + x + iy} = \frac{1 - (x^2 + y^2)}{(1 + x)^2 + y^2}, \quad \operatorname{Im} \frac{1 - x - iy}{1 + x + iy} = -\frac{2y}{(1 + x)^2 + y^2},$$

(22) takes the form

$$(q - 1)((1 - (x^2 + y^2))^2 + 4y^2)^2 + (2 - p - q)(1 - (x^2 + y^2))^2((1 + x)^2 + y^2)^2 - (2 - p - q + pq)4y^2((1 + x)^2 + y^2)^2 + (p - 1)((1 + x)^2 + y^2)^4 > 0.$$

This factors as

$$(26) \quad \begin{aligned} & [4((1 + x)^2 + y^2)^2] [(p - q)x(1 + x^2 + y^2) \\ & + (2p + 2q - 4)x^2 - (pq - 2p - 2q + 4)y^2]. \end{aligned}$$

Quite miraculously, the second term in straight brackets precisely equals the term in (25). Since $4((1 + x)^2 + y^2)^2 > 0$, it follows that (26) (and hence (22)) implies (25) (and hence (17)). \square

It is shown in [2] (see also [5]) that the operator $\exp(-zL)$ is bounded from $L^p(\mathbb{R}^d, \gamma)$ to $L^q(\mathbb{R}^d, \gamma)$ if and only if $z \in \overline{E_p}$, and then the operators $\exp(-zL)$ are in fact contractions. Our proof does not recover the contractivity of $\exp(-zL)$. Nevertheless, it is remarkable that the boundedness part does follow from our method, which uses only (6), elementary calculus, the Schur test, and some algebraic manipulations.

For $p = q$, Theorem 6.1 combined with the fact that $E_{p,p} = E_p$ contains as a special case that, for a given $z \in \mathbb{C}$ with $\operatorname{Re} z > 0$, the operator $\exp(-zL)$ is bounded on $L^p(\mathbb{R}^d, \gamma)$ if z belongs to E_p . A more direct – and more transparent – proof of this fact may be obtained as a consequence of the following theorem.

THEOREM 6.4. — *For all $1 < p < \infty$ and $s \in \Sigma_{\theta_p}$, the operator $\exp(-s(P^2 + Q^2))$ is bounded on $L^p(\mathbb{R}^d, \gamma)$.*

As we explained in Section 3, this result translates into Epperson's result that the semigroup $\exp(-tL)$ on $L^p(\mathbb{R}^d, \gamma)$ can be analytically extended to E_p .

Proof. — Lemma 6.3 shows that (19) holds. Since $q = p$, (20) reduces to the condition

$$p^2 \left(\frac{x^2}{x^2 + y^2} - 1 \right) + 4p - 4 > 0,$$

which is equivalent to saying that $s \in \Sigma_{\theta_p}$. □

REMARK 6.5. — More generally, for an arbitrary pair $(\alpha, p) \in [1, \infty) \times [1, \infty)$ satisfying $\alpha p > 2$, by the same method we obtain that $\exp(-zL)$ is bounded on $L^p(\mathbb{R}^d, \gamma_\alpha)$ if $s = \frac{1-e^{-z}}{1+e^{-z}}$ satisfies

$$\frac{\operatorname{Re} s}{|s|} > 1 - \frac{2}{\alpha p}.$$

This corresponds to the sector of angle $\theta_{\alpha,p} = \arccos\left(1 - \frac{2}{\alpha p}\right)$ in the s -plane. In the z -plane, this corresponds to the Epperson region $E_{\alpha p}$.

Acknowledgment. — We thank Emiel Lorist for generating the figures.

BIBLIOGRAPHY

- [1] D. BAKRY, F. BOLLEY & I. GENTIL, “Dimension dependent hypercontractivity for Gaussian kernels”, *Probability Theory and Related Fields* **154** (2012), p. 845–874
- [2] J. B. EPPERSON, “The hypercontractive approach to exactly bounding an operator with complex Gaussian kernel”, *J. Funct. Anal.* **87** (1989), p. 1–30
- [3] J. GARCÍA-CUERVA, G. MAUCERI, S. MEDA, P. SJÖGREN & J. L. TORREA, “Functional calculus for the Ornstein-Uhlenbeck operator”, *J. Funct. Anal.* **183** (2001), p. 413–450
- [4] C. GERRY & P. KNIGHT, “Introductory quantum optics”, Cambridge University Press, 2005

- [5] S. JANSON, “On complex hypercontractivity”, *J. Funct. Anal.* **151** (1997), p. 270–280
- [6] E. NELSON, “A quartic interaction in two dimensions”, in *Mathematical theory of elementary particles*, 1966, p. 69–73
- [7] K. R. PARTHASARATHY, “An introduction to quantum stochastic calculus”, Modern Birkhäuser Classics, 1992
- [8] I. E. SEGAL, “Tensor algebras over Hilbert spaces I”, *Trans. Amer. Math. Soc.* **81** (1956), p. 106–134
- [9] W. SCHLAG, “Introduction to Ψ DOs and FIOs: the Weyl calculus”, math.uchicago.edu/~schlag/Notes3.pdf
- [10] C. D. SOGGE, “Fourier integral operators in classical analysis”, Cambridge University Press, 1993
- [11] E. M. STEIN, “Harmonic analysis: Real-variable methods, orthogonality, and oscillatory integrals”, Princeton University Press, 1993
- [12] F. B. WEISLER, “Two-point inequalities, the Hermite semigroup and the Gauss-Weierstrass semigroup”, *J. Funct. Anal.* **32** (1979), p. 102–121