

EDOARDO BALLICO

**A splitting theorem for the Kupka component of
a foliation of $\mathbb{CP}^n$, $n \geq 6$. Addendum to a paper
by O. Calvo-Andrade and N. Soares**

Annales de l'institut Fourier, tome 45, n° 4 (1995), p. 1119-1121

[<http://www.numdam.org/item?id=AIF_1995__45_4_1119_0>](http://www.numdam.org/item?id=AIF_1995__45_4_1119_0)

© Annales de l'institut Fourier, 1995, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

**A SPLITTING THEOREM FOR THE KUPKA
COMPONENT OF A FOLIATION OF $\mathbf{CP}^n$, $n \geq 6$.
ADDENDUM TO A PAPER BY
O. CALVO-ANDRADE AND N. SOARES**

by Edoardo BALLICO

A codimension one singular holomorphic foliation F of $\mathbf{CP}^n$ is given by $\omega \in H^0(\mathbf{CP}^n, \Omega(k))$ (for some k) with $\omega \neq 0$, ω not vanishing on a hypersurface. The Kupka subset $K(F) := \{P \in \mathbf{CP}^n : \omega(P) = 0, d\omega(P) \neq 0\}$ of the singular set $S(F) := \{P \in \mathbf{CP}^n : \omega(P) = 0\}$ of F has remarkable properties (e.g. if not empty it is a smooth submanifold of pure codimension 2 with strong stability properties with respect to deformations of F). For much more on this topic, see [GLM] and [CS]. Let $K \neq \emptyset$ be a Kupka component of F , i.e. ([CS]) a connected component of $K(F)$. It was proved in [CL] that if K is a complete intersection, then F has a meromorphic first integral. Motivated by this result in [CS] it was conjectured and proved in some cases that every Kupka component is a complete intersection. Here we prove the following result.

THEOREM. — *Let F be a codimension 1 singular holomorphic foliation of $\mathbf{CP}^n$, $n \geq 6$, induced by $\omega \in H^0(\mathbf{CP}^n, \Omega^1(k))$ and such that the codimension 2 component of the singular set of F consists of a single compact Kupka component K with $\deg(K) \neq k^2/4$. Then K is a complete intersection.*

Key words: Singular foliations – Codimension 1 foliations – Kupka component – Complete intersection – Unstable vector bundle – Rank 2 vector bundle – Splitting of a vector bundle – Meromorphic first integral.
Math. classification: 58F18 – 14F05 – 14M07.

The proof of this result uses in an essential way the results proven in [CS] and [GML]. We consider this paper as an addendum to [CS] and we invite the reader to turn to [GML] and [CS] for background, motivations, several results used here, and so on. For the results used on vector bundles and codimension 2 submanifolds of $\mathbf{CP}^n$, see [OSS], [FL] and [CS].

Assume that F is induced by $\omega \in H^0(\mathbf{CP}^n, \Omega^1(k))$. Let N_K be the normal bundle of K in $\mathbf{CP}^n$. By [CS], Corollary 3.5, N_K is the restriction $E|_K$ to K of a rank 2 vector bundle E on $\mathbf{CP}^n$. K is a complete intersection if and only if E is the direct sum of two line bundles ([OSS]). If $n \geq 6$ every line bundle on K is the restriction of a line bundle on $\mathbf{CP}^n$ (see [FL]). Hence, by a very nice result of Faltings ([F]) if $n \geq 6$ and N_K is the direct sum of two line bundles, K is a complete intersection. By [CS], Cor. 4.5 (2), we may assume $k > 0$. By [CS], Th. 3.4 (2) to prove our result we may distinguish two cases, according to the transversal type of K . First assume that the transversal type of K is given by $\eta = pxdy - qydy$ with p, q positive relatively prime integers. Look at [GML], Th. 2.3 and its proof at page 321 (in particular the two lines before eq. (2.6)) and use that K is simply connected if $n \geq 6$ ([FL], Cor. 6.3). The quoted result [GML], Th. 2.3, was the essential input for the proof of [CS], Th. 3.4; then [CS], Th. 3.4, and the calculations in [CS], §4, on the applications of the Baum-Bott formulas to K gave the proof of [CS], Cor. 4.5. By [GML], page 321, N_K is in this case the direct sum of two line bundles. Hence our theorem is proved in this case. Now assume that the transversal type of K is given by $\eta = pxdy - qydy$ with $p = q = 1$. By [CS], Th. 4.2, we have $\deg(K) = k^2/4$. Hence our theorem is proved even in this case.

The author was partially supported by MURST and GNSAGA of CNR (Italy).

BIBLIOGRAPHY

- [CS] O. CALVO-ANDRADE, M. SOARES, Chern numbers of a Kupka component, *Ann. Inst. Fourier*, 44-4 (1994), 1237–1242.
- [CL] D. CERVAU, A. LINS, Codimension one foliations in $\mathbf{CP}^n$, $n \geq 3$, with Kupka components, in: *Complex analytic methods in dynamical systems*, *Astérisque* (1994), 93–133.
- [F] G. FALTINGS, Ein Kriterium für vollständige Durchschnitte, *Invent. Math.*, 62 (1981), 393–401.
- [FL] W. FULTON, R. LAZARSFELD, Connectivity in algebraic geometry, in: *Algebraic Geometry, Proceedings Chicago 1980*, *Lect. Notes in Math.* 862, Springer-Verlag (1981), 26–92.

- [GML] X. GOMEZ-MONT, N. LINS, A structural stability of foliations with a meromorphic first integral, *Topology*, 30 (1990), 315–334.
- [OSS] Ch. OKONEK, M. SCHNEIDER, H. SPINDLER, Vector bundles on complex projective spaces, *Progress in Math.*, 3, Birkhäuser, Basel, 1978.

Manuscrit reçu le 6 mars 1995,
accepté le 7 avril 1995.

Edoardo BALLICO,
Dept. of Mathematics
University of Trento
38050 Povo (TN) (Italie).