ANNALES DE L'INSTITUT FOURIER

SATOSHI KOIKE

On condition (a_f) of a stratified mapping

Annales de l'institut Fourier, tome 33, n° 1 (1983), p. 177-184

<http://www.numdam.org/item?id=AIF_1983__33_1_177_0>

© Annales de l'institut Fourier, 1983, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (http://annalif.ujf-grenoble.fr/) implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/conditions). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
http://www.numdam.org/

Ann. Inst. Fourier, Grenoble
33, 1 (1983), 177-184

ON CONDITION (a_f) OF A STRATIFIED MAPPING

by Satoshi KOIKE

In [3], D.J.A. Trotman showed that Whitney's condition (a) on the pair of adjacent strata is equivalent to condition (a^s) which has more obvious geometric content. These conditions can be generalized to the conditions of the kernel of the mapping called a stratified mapping. The generalization of condition (a) is condition (a_f) which is well-known in the stratification theory. On the other hand, we shall call the generalization of condition (a^s) condition (a_f^s). Then, we have already known that (a_f) implies (a_f^s) from the proof of Lemma 11.4 in J.N. Mather [2] (or Lemma (2.4) of Chapter II in [1]). In this paper, we show that (a_f) is equivalent to (a_f^s). In § 2 we prove this result, and in § 3 we give the illustrative example of the fact.

1. Definitions and the result.

Let X, Y be disjoint C^1 submanifolds of $\mathbf{R}^n$, and let y_0 be a point in $Y \cap \overline{X}$. We say the pair (X, Y) satisfies *Whitney's condition* (a) at y_0 if for any sequence of points $\{x_i\}$ in X tending to y_0 such that the tangent space $T_{x_i}X$ tends to τ, we have $T_{y_0}Y \subset \tau$. As stated above, this condition is equivalent to the following condition; (a^s): For any local C^1 retraction at y_0, $\pi_Y : \mathbf{R}^n \longrightarrow Y$, there exists a neighborhood W of y_0 in $\mathbf{R}^n$ such that $\pi_{Y|W\cap X}$ is a submersion.

Let $f : A \longrightarrow \mathbf{R}^p$ be a smooth mapping defined in a neighborhood A of $X \cup Y$ in $\mathbf{R}^n$. Suppose that the restricted mappings

$f|_X : X \longrightarrow \mathbf{R}^p$ and $f|_Y : Y \longrightarrow \mathbf{R}^p$ are of constant ranks. Then we say the pair (X, Y) satisfies *condition* (a_f) at y_0 if for any sequence of points $\{x_i\}$ in X tending to y_0 such that the plane $\ker d(f|_X)_{x_i}$ tends to κ, we have $\ker d(f|_Y)_{y_0} \subset \kappa$, where $\ker d(f|_X)_x$ denotes the kernel of the differential of $f|_X$ at x.

Let U, V be C^1 submanifolds of $\mathbf{R}^p$ such that $U \cap V = \phi$ or $U = V$. Further, suppose that $f(X)$, $f(Y)$ are contained in U, V respectively, and that $f|_X : X \longrightarrow U$ and $f|_Y : Y \longrightarrow V$ are submersions. Then we call this mapping f a *stratified mapping.* From now, we shall think of a stratified mapping.

We say that a local C^1 retraction at y_0, $\pi_Y : \mathbf{R}^n \longrightarrow Y$, and a local C^1 retraction at $f(y_0)$, $\pi_V : \mathbf{R}^p \longrightarrow V$, satisfy the commutation relation (CRf) if it holds that $f \circ \pi_Y = \pi_V \circ f$ in a neighborhood of y_0.

Remark 1. – For a stratified mapping, the following facts hold.

1) For any local C^1 retraction at $f(y_0)$, $\pi_V : \mathbf{R}^p \longrightarrow V$, there exists a local C^1 retraction at y_0, $\pi_Y : \mathbf{R}^n \longrightarrow Y$ such that they satisfy (CRf). Consider the mapping $\pi_V \circ f$ in a neighborhood of y_0. Since $\pi_V \circ f|_Y : Y \longrightarrow V$ is a submersion, there exists a local C^1 retraction at y_0, $\pi_Y : \mathbf{R}^n \longrightarrow Y$, such that $\pi_V \circ f \circ \pi_Y = \pi_V \circ f$. Thus, we see that they satisfy (CRf) in a neighborhood of y_0.

2) On the other hand, it is not true that for any local C^1 retraction at y_0, there exists a local C^1 retraction at $f(y_0)$ such that they satisfy (CRf): See the example in 3.

Here we introduce the next condition;

(a_f^s): For any local C^1 retraction at y_0, $\pi_Y : \mathbf{R}^n \longrightarrow Y$, and local C^1 retraction at $f(y_0)$, $\pi_V : \mathbf{R}^p \longrightarrow V$, satisfying (CRf), there exists a neighborhood W of y_0 in $\mathbf{R}^n$ such that for any $x \in W \cap X$,

$$d(\pi_{YX})_x : \ker d(f|_X)_x \longrightarrow \ker d(f|_Y)_y$$

is onto, where $\pi_{YX} = \pi_Y|_X$ and $y = \pi_Y(x)$.

THEOREM. – *For a stratified mapping,* (a_f) *is equivalent to* (a_f^s).

Remark 2. – Theorem A in [3] is the case where $U = V = \{f(y_0)\}$ in the above theorem. Because, in that case, the kernel is the tangent

space, and (CRf) is satisfied for any local C^1 retraction at y_0, $\pi_Y : \mathbf{R}^n \longrightarrow Y$.

2. Proof of the theorem.

Let f be a stratified mapping i.e.

$$f|_X : X \longrightarrow U \quad \text{and} \quad f|_Y : Y \longrightarrow V$$

are submersions. We introduce the condition of "transverse foliation" defined locally in a neighborhood of y_0 in $\mathbf{R}^n$;

$(\mathcal{H}^1)$: For any local C^1 foliation $\mathcal{F}$ which is transversal to the fiber of $f|_Y$ at y_0, and whose leaves are unions of fibers of a local C^1 retraction π_Y satisfying the relation (CRf), there exists a neighborhood W of y_0 in $\mathbf{R}^n$ such that $\mathcal{F}$ is transversal to the fibers of $f|_X$ in W.

LEMMA. – *(a_f^s) is equivalent to $(\mathcal{H}^1)$.*

Proof. – As it is trivial that (a_f^s) implies $(\mathcal{H}^1)$, we shall show that $(\mathcal{H}^1)$ implies (a_f^s). Consider a local C^1 retraction at y_0, $\pi_Y : \mathbf{R}^n \longrightarrow Y$, and a local C^1 retraction at $f(y_0)$, $\pi_V : \mathbf{R}^p \longrightarrow V$, which satisfy (CRf) in a neighborhood of y_0. Let N_{y_0} denote the normal space of $\ker d(f|_Y)_{y_0}$ in $T_{y_0}Y$. Then, there exist a neighborhood W_1 of y_0 in Y, and a local C^1 foliation $\tilde{\mathcal{F}}$ of W_1 such that $N_{y_0} = T_{y_0}\tilde{F}_{y_0}$, where $\tilde{F}_{y_0}$ denotes the leaf of $\tilde{\mathcal{F}}$ which contains y_0. Shrinking the neighborhood W_1 if necessary, $\mathcal{F} \equiv \{\{v \in \mathbf{R}^n \mid \pi_Y(v) \in \tilde{F}\}\}_{\tilde{F} \in \tilde{\mathcal{F}}}$ is a local C^1 foliation of $\mathbf{R}^n$ in a neighborhood of y_0. From the construction, we have

$$T_{y_0}F_{y_0} \oplus \ker d(f|_Y)_{y_0} = T_{y_0}\mathbf{R}^n,$$

where F is a leaf of $\mathcal{F}$. Since $f|_Y : Y \longrightarrow V$ is a submersion, $\ker d(f|_Y)_y$ is continuous in the Grassman manifold of

$$\dim \ker (f|_Y)_{y_0}\text{-planes in } n\text{-space.}$$

Further, $(\mathcal{H}^1)$ holds from the assumption. Therefore, there exists a neighborhood W_2 of y_0 in $\mathbf{R}^n$ such that for any $y \in W_2$,

$$T_yF_y \oplus \ker d(f|_Y)_y = T_y\mathbf{R}^n \tag{2.1}$$

and for any $x \in W_2 \cap X$,

$$T_xF_x + \ker d(f|_X)_x = T_x\mathbf{R}^n. \tag{2.2}$$

From the relation (CRf), we have

$$d(\pi_{YX})_x : \ker d(f|_X)_x \longrightarrow \ker d(f|_Y)_y \qquad (2.3)$$

in a neighborhood of y_0. By (2.1), (2.2), and (2.3), we see that the differential mapping (2.3) is onto near y_0.

Remark 3. – From the proof of Theorem A in [3], we can see easily that (a_f) is equivalent to the following condition;

$(\mathcal{F}^1)$: For any C^1 foliation $\mathcal{F}$ which is transversal to the fiber of $f|_Y$ at y_0, there exists a neighborhood W of y_0 in $\mathbf{R}^n$ such that $\mathcal{F}$ is transversal to the fibers of $f|_X$ in W.

PROPERTY 1. – *Let* $\pi_V : \mathbf{R}^p \longrightarrow V$ *be a local* C^1 *retraction at* $f(y_0)$.

1) *There exists a neighborhood* W *of* $f(y_0)$ *in* V *such that* $\{\pi_V \circ f)^{-1}(w)\}_{w\in W}$ *is a* C^1 *foliation of codimension* V *in a sufficiently small neighborhood of* y_0 *in* $\mathbf{R}^n$.
Since $\pi_Y \circ f|_Y : Y \longrightarrow V$ is a submersion, $\pi_V \circ f$ has a maximal rank (of dimension V) at $y_0 \in \mathbf{R}^n$. Thus (1) follows.

2) *If a point* q *in* $\mathbf{R}^n$ *is contained in* $(\pi_V \circ f)^{-1}(w)$, *then we have* $\ker df \subset T_q(\pi_V \circ f)^{-1}(w)$.
It is clear from the fact that $T_q(\pi_V \circ f)^{-1}(w) = \ker d(\pi_V \circ f)_q$.

PROPERTY 2. – *Let* $\pi_Y : \mathbf{R}^n \longrightarrow Y$ *be a local* C^1 *retraction at* y_0, *and* $\pi_V : \mathbf{R}^p \longrightarrow V$ *be a local* C^1 *retraction at* $f(y_0)$. *If any fiber of* π_Y *is contained in some fiber of* $\pi_V \circ f$ *in a neighborhood of* y_0, *then* (CRf) *holds.*
It is trivial.

From Lemma, it is sufficient to show that $(\mathcal{H}^1)$ implies (a_f). We suppose that the pair (X, Y) does not satisfy condition (a_f) at y_0. Then, there exists a sequence of points $\{x_i\}$ in X tending to y_0 with $\lim_i \ker d(f|_X)_{x_i} = K$, such that $K \not\supset \ker d(f|_Y)_{y_0}$. Thus, there exists a vector $k \in \ker d(f|_Y)_{y_0}$ such that $k \notin K$. By the similar way as the proof of Theorem A in [3], we can construct a C^1 foliation $\mathcal{F}$ of codimension 1 such that $T_{y_0}F_{y_0} \not\ni k$ and $T_{x_i}F_{x_i} \supset \ker d(f|_X)_{x_i}$ i.e. $\mathcal{F}$ is transversal to the fiber of $f|_Y$ at y_0, and $\mathcal{F}$ is not transversal to the fiber of $f|_X$ at x_i.

We take a local C^1 retraction at $f(y_0)$, $\pi_V : \mathbf{R}^n \longrightarrow V$, arbitarily. From Property 1 (2), we have

$$k \in \ker d(f|_Y)_{y_0} \subset \ker df_{y_0} \subset T_{y_0}(\pi_V \circ f)^{-1}(w_0),$$

where $w_0 = f(y_0)$. Therefore, the local foliations $\{(\pi_V \circ f)^{-1}(w)\}_{w \in W}$ and $\mathcal{F}$ are transversal near y_0. Thus,

$$\{(\pi_V \circ f)^{-1}(w) \cap F\}_{\substack{w \in W \\ F \in \mathcal{F}}} \tag{2.4}$$

is a C^1 foliation in a neighborhood of y_0 in $\mathbf{R}^n$.

PROPERTY 3. – $(\pi_V \circ f)^{-1}(w_0) \cap F_{y_0}$ *is transversal to* Y *at* y_0. Since $\pi_V \circ f|_Y : Y \longrightarrow V$ is a submersion and

$$T_{y_0}(\pi_V \circ f)^{-1}(w_0) = \ker d(\pi_V \circ f)_{y_0},$$

we have

$$T_{y_0}Y + T_{y_0}(\pi_V \circ f)^{-1}(w_0) = T_{y_0}\mathbf{R}^n. \tag{2.5}$$

As $(\pi_V \circ f)^{-1}(w_0)$ is transversal to F_{y_0} *at* y_0, we have

$$T_{y_0}((\pi_V \circ f)^{-1}(w_0) \cap F_{y_0}) = T_{y_0}(\pi_V \circ f)^{-1}(w_0) \cap T_{y_0}F_{y_0}.$$

Further, the vector k is not an element of $T_{y_0}F_{y_0}$. Therefore, we have

$$T_{y_0}((\pi_V \circ f)^{-1}(w_0) \cap F_{y_0}) + \langle k \rangle = \ker d(\pi_V \circ f)_{y_0} \tag{2.6}$$

where $\langle k \rangle$ denotes the subvector space spanned by the vector k of $T_{y_0}\mathbf{R}^n$. From (2.5), (2.6), and the fact that $k \in \ker d(f|_Y)_{y_0} \subset T_{y_0}Y$, we see that $T_{y_0}((\pi_V \circ f)^{-1}(w_0) \cap F_{y_0}) + T_{y_0}Y = T_{y_0}\mathbf{R}^n$.

By using Property 3, we can construct a local C^1 retraction at y_0, $\pi_Y : \mathbf{R}^n \longrightarrow Y$, along leaves of the local foliation (2.4). Then, these local retractions π_Y and π_V satisfy (CRf) in a neighborhood of y_0 in $\mathbf{R}^n$, from Property 2. Further, from the construction it is clear that each leaf of $\mathcal{F}$ is a union of fibers of π_Y. Thus, $(\mathcal{HC}^1)$ does not hold. This completes the proof of the theorem.

3. An Example.

In this section, we give an example which illustrates the proof of the theorem, and demonstrates Remark 1 (2).

Let $f = (f_1, f_2) : \mathbf{R}^3 \longrightarrow \mathbf{R}^2$ be a mapping defined by

$$f(x, y, z) = (x, y^4 + 2y^2 z^2).$$

We take $X = \{y \neq 0\}$ and $Y = \{y = 0\}$ as disjoint submanifolds in $\mathbf{R}^3$, and take $U = \{y \neq 0\}$ and $V = \{y = 0\}$ as disjoint submanifolds in $\mathbf{R}^2$. Then, restricted mappings $f|_X : X \longrightarrow U$ and $f|_Y : Y \longrightarrow V$ are submersions i.e. f is a stratified mapping.

Put $S = \{p = (x, y, z) \in X \mid y = z\}$. For any point $p \in S$, we have

$$\text{grad}\,(f_{1|X})_p = (1, 0, 0) \text{ and } \text{grad}\,(f_{2|X})_p = (0, 8y^3, 4y^3).$$

Therefore, we have $\ker d(f|_X)_p = \langle (0, 1, -2) \rangle$. We take a sequence of points $\{p_i\}$ in S tending to $0 = (0, 0, 0) \in Y$. We have

$$\lim_i \ker d(f|_X)_{p_i} = \langle (0, 1, -2) \rangle.$$

On the other hand, $\ker d(f|_Y)_0 = \langle (0, 0, 1) \rangle$. Therefore, (X, Y) does not satisfy condition (a_f) at 0.

In this case, (X, Y) does not satisfy condition (a_f^s) at 0 as a matter of course. For example, we take the canonical projection over Y as a local retraction at $f(0)$, $\pi_V : \mathbf{R}^2 \longrightarrow V$. Then, the foliation whose leaves are fibers of $\pi_V \circ f$ is

$$\mathcal{F}_1 = \{\{(x, y, z) \in \mathbf{R}^3 \mid x = k_1\}\}_{k_1 \in \mathbf{R}}.$$

Further, we consider the foliation

$$\mathcal{F}_2 = \{\{(x, y, z) \in \mathbf{R}^3 \mid z + 2y = k_2\}\}_{k_2 \in \mathbf{R}},$$

which is transversal to the fiber of $f|_Y$ at 0, and is not transversal to the fibers of $f|_X$ in S. As $\mathcal{F}_1$ and $\mathcal{F}_2$ are transversal, the intersection of $\mathcal{F}_1$ and $\mathcal{F}_2$ becomes a foliation of $\mathbf{R}^3$,

$$\{\{(x, y, z) \in \mathbf{R}^3 \mid x = k_1, z + 2y = k_2\}\}_{\substack{k_1 \in \mathbf{R} \\ k_2 \in \mathbf{R}}}.$$

It is clear that the leaves of this foliation induce a retraction π_Y which does not admit condition (a_f^s) at $0 \in \mathbf{R}^3$.

Nextly, we show that this example demonstrates Remark 1 (2). We consider a retraction at $0 \in \mathbf{R}^3$, $\pi_Y(x, y, z) = (x + yz^2, 0, z)$. Then, we have $f \circ \pi_Y(x, y, z) = (x + yz^2, 0)$. Let (x_0, y_1, z_1), (x_0, y_2, z_2) be points in X such that $0 < y_1 < y_2$ and

$$y_1^4 + 2y_1^2 z_1^2 = y_2^4 + 2y_2^2 z_2^2 = C > 0.$$


z
X
Y
(x_0, y_1, z_1)
(x_0, y_2, z_2)
0
y
$y^4 + 2y^2 z^2 = C$
x
the level line of f
f
V
U
0
y
x

From the fact that $0 < y_1 < y_2$, we have

$$x_0 + y_1 z_1^2 = x_0 + \frac{C - y_1^4}{2y_1} \neq x_0 + \frac{C - y_2^4}{2y_2} = x_0 + y_2 z_2^2$$

i.e. $$f \circ \pi_Y(x_0, y_1, z_1) \neq f \circ \pi_Y(x_0, y_2, z_2).$$

On the other hand, $f(x_0, y_1, z_1) = f(x_0, y_2, z_2)$. Therefore, there does not exist a local C^1 retraction at $0 \in \mathbf{R}^2$, $\pi_V : \mathbf{R}^2 \longrightarrow V$, such that they satisfy (CRf).

BIBLIOGRAPHY

[1] C.G. GIBSON, K. WIRTHMULLER, A.A. du PLESSIS, E.J.N. LOOIJENGA, Topological stability of smooth mappings, *Springer Lect. Notes*, Berlin Vol. 552 (1976).

[2] J. MATHER, Notes on topological stability, *mimeographed*, Harvard Univ. (1970).

[3] D.J.A. TROTMAN, Geometric versions of Whitney regularity for smooth stratifications, *Ann. scient. Ec. Norm. Sup.*, Vol. 12 (1979), 461-471.

Manuscrit reçu le 4 janvier 1982.

Satoshi KOIKE,
Department of Mathematics
Faculty of Sciences
Kyoto University
Kyoto 606 (Japan).