Journal de l'École polytechnique
Mathématiques

Jérémy Blanc, Pierre-Marie Poloni, & Immanuel Van Santen

Complements of hypersurfaces in projective spaces

Tome 11 (2024), p. 733-768.

https://doi.org/10.5802/jep.264

© Les auteurs, 2024.

Cet article est mis à disposition selon les termes de la licence Licence internationale d'attribution Creative Commons BY 4.0.
https://creativecommons.org/licenses/by/4.0/

Publié avec le soutien
du Centre National de la Recherche Scientifique

Publication membre du
Centre Mersenne pour l'édition scientifique ouverte
www.centre-mersenne.org
e-ISSN : 2270-518X

Journal de l'École polytechnique
Mathématiques

Tome 11, 2024, p. 733–768 DOI: 10.5802/jep.264

COMPLEMENTS OF HYPERSURFACES IN PROJECTIVE SPACES

by Jérémy Blanc, Pierre-Marie Poloni & Immanuel Van Santen

Abstract. — We study the complement problem in projective spaces $\mathbb{P}^n$ over any algebraically closed field: If $H, H' \subseteq \mathbb{P}^n$ are irreducible hypersurfaces of degree d such that the complements $\mathbb{P}^n \smallsetminus H$, $\mathbb{P}^n \smallsetminus H'$ are isomorphic, are the hypersurfaces H, H' isomorphic?

For $n = 2$, the answer is positive if $d \leqslant 7$ and there are counterexamples when $d = 8$. In contrast, we provide counterexamples for all $n, d \geqslant 3$ with $(n, d) \neq (3, 3)$. Moreover, we show that the complement problem has an affirmative answer for $d = 2$ and give partial results in case $(n, d) = (3, 3)$. In the course of the exposition, we prove that rational normal projective surfaces admitting a desingularization by trees of smooth rational curves are piecewise isomorphic if and only if they coincide in the Grothendieck ring, answering affirmatively a question posed by Larsen and Lunts for such surfaces.

Résumé (Complémentaires d'hypersurfaces dans les espaces projectifs)

Nous étudions le problème du complémentaire dans les espaces projectifs $\mathbb{P}^n$ sur tout corps algébriquement clos : Si $H, H' \subseteq \mathbb{P}^n$ sont des hypersurfaces irréductibles de degré d telles que les complémentaires $\mathbb{P}^n \smallsetminus H$, $\mathbb{P}^n \smallsetminus H'$ sont isomorphes, les hypersurfaces H, H' sont-elles isomorphes ?

Pour $n = 2$, la réponse est positive si $d \leqslant 7$ et il y a des contre-exemples lorsque $d = 8$. En revanche, nous fournissons des contre-exemples pour tous les entiers $n, d \geqslant 3$ avec $(n, d) \neq (3, 3)$. De plus, nous montrons que le problème du complémentaire a une réponse affirmative pour $d = 2$ et donnons des résultats partiels dans le cas où $(n, d) = (3, 3)$. Au cours de l'exposition, nous prouvons que les surfaces projectives normales rationnelles admettant une désingularisation par des arbres de courbes rationnelles lisses sont isomorphes par morceaux si et seulement si elles coïncident dans l'anneau de Grothendieck, répondant ainsi positivement à une question posée par Larsen et Lunts pour de telles surfaces.

Contents

1. Introduction . . . 734
2. Lift to affine hypersurfaces . . . 736
3. Non-isomorphic hypersurfaces having isomorphic complements . . . 740
4. Topological Euler characteristic and piecewise isomorphisms . . . 747
5. Partial answers in low degree . . . 755
References . . . 767

Mathematical subject classification (2020). — 14J70, 14R05, 14E07, 19A99.
Keywords. — Complement problem, cylinder over Danielewski surfaces, piecewise isomorphisms.

The first author acknowledges support by the Swiss National Science Foundation Grant "Geometrically ruled surfaces" 200020–192217.

1. Introduction

Let $\mathbf{k}$ be a field, let $n \geqslant 2$ be an integer and let $f \in \mathbf{k}[x_0, \dots, x_n]$ be an irreducible homogeneous polynomial of degree $d \geqslant 1$. The variety

$$\mathbb{P}^n_f = \{[x_0 : \cdots : x_n] \in \mathbb{P}^n \mid f(x_0, \dots, x_n) \neq 0\}$$

is the complement in $\mathbb{P}^n$ of the hypersurface given by $f = 0$. It is an affine open subset of $\mathbb{P}^n$ with Picard group $\mathbb{Z}/d\mathbb{Z}$. Moreover, any open subset of $\mathbb{P}^n$ isomorphic to it is of the form $\mathbb{P}^n_g$ for some irreducible homogeneous polynomial $g \in \mathbf{k}[x_0, \dots, x_n]$ of degree d (see Lemma 2.6). One may then ask the following natural questions, for all irreducible homogeneous polynomials $f, g \in \mathbf{k}[x_0, \dots, x_n]$ of the same degree $d \geqslant 1$.

(1) Does any isomorphism $\mathbb{P}^n_f \xrightarrow{\simeq} \mathbb{P}^n_g$ extend to an element of $\mathrm{Aut}(\mathbb{P}^n)$?

(2) If $\mathbb{P}^n_f$ and $\mathbb{P}^n_g$ are isomorphic, is there an element of $\mathrm{Aut}(\mathbb{P}^n)$ that sends $\mathbb{P}^n_f$ onto $\mathbb{P}^n_g$?

(3) If $\mathbb{P}^n_f$ and $\mathbb{P}^n_g$ are isomorphic, are the zero loci $V_{\mathbb{P}^n}(f)$ and $V_{\mathbb{P}^n}(g)$ of f and g isomorphic?

Of course, any positive answer to one of the questions also gives a positive answer to the next one(s).

If $d = 1$, the first question has a negative answer, as there are many automorphisms of $\mathbb{P}^n_f \simeq \mathbb{A}^n$ that do not extend to automorphisms of $\mathbb{P}^n$. However, the two other questions have a positive answer.

For $d = 2$, we can obtain similar results, in any dimension, when $\mathbf{k}$ is algebraically closed. Indeed, we will answer the first question in the negative for any irreducible quadric, see Example 5.6, and provide positive answers to the two other questions, see Theorem B below.

The situation for $d = 3$ is already more complicated, at least in dimension $n = 3$. The first question has a negative answer for each singular cubic with Du Val singularities over an algebraically closed field of characteristic zero [CDP18, Th. C & Th. 4.3], and more generally for each singular cubic over any algebraically closed field (Proposition 5.22), but is wide open for any smooth cubic surface [GSY05, p. 6]. The two other questions are open as well.

In dimension $n = 2$, it was conjectured in [Yos84] that the answer to the second question should be always positive for algebraically closed fields of characteristic zero. This conjecture was proved for all algebraically closed fields if the curve $C \subseteq \mathbb{P}^2$ given by $f = 0$ is such that $C \smallsetminus L \simeq \mathbb{A}^1$ for some line (see [Yos84, Th.] for a proof in characteristic zero and [Hem19, Th. 1] for a proof in any characteristic). The conjecture was however proved to be false in general, with a first example of degree 39 given in [Bla09]. A second family of counterexamples of degree $9+4m$ for each $m \geqslant 0$ was given later in [Cos12]. Finally, an example of degree 8 and a proof that the conjecture is true for curves of degree $\leqslant 7$ appeared in [Hem19, Cor. 1.2, Th. 3] for algebraically closed fields of any characteristic. Moreover, the examples given in [Hem19] and [Bla09] also give a negative answer to the third question.

In dimension $n \geqslant 3$, there was no negative answer to Questions (2) and (3) until now. In this article, we show by constructing explicit examples, that the answer of the third question, and thus of the two others, is negative for large degree and large dimension. More precisely, in any dimension $n \geqslant 3$, the smallest degree for counterexamples is 3, except maybe for $n = 3$, where the smallest degree might be 4. This contrasts the case of dimension $n = 2$ where counterexamples for algebraically closed fields always have degree $\geqslant 8$.

Theorem A. — *Let* $\mathbf{k}$ *be a field and let* $d, n \geqslant 3$ *be integers such that* d *is not a multiple of* $\mathrm{char}(\mathbf{k})$ *and such that* $(d, n) \neq (3, 3)$.

There are two non-isomorphic irreducible hypersurfaces $H, H' \subseteq \mathbb{P}^n_{\mathbf{k}}$ *of degree* d*, that have isomorphic complements.*

We will prove Theorem A at the end of Section 3.2. If $d = 2$, then Question (2) has an affirmative answer, at least when $\mathbf{k}$ is algebraically closed:

Theorem B. — *Let* $n \geqslant 2$ *and let* $\mathbf{k}$ *be an algebraically closed field. If the complements of two irreducible quadric hypersurfaces* H, H' *in* $\mathbb{P}^n_{\mathbf{k}}$ *are isomorphic, then there exists an automorphism of* $\mathbb{P}^n_{\mathbf{k}}$ *that sends* H *onto* H'.

We will prove Theorem B at the end of Section 5.1. In the case where $n = 3$, we have the following partial positive result concerning Question (2):

Theorem C. — *Let* $\mathbf{k}$ *be an algebraically closed field and let* $H, H' \subseteq \mathbb{P}^3_{\mathbf{k}}$ *be irreducible hypersurfaces such that their complements in* $\mathbb{P}^3_{\mathbf{k}}$ *are isomorphic. Then we have:*

(1) *If both* H *and* H' *are normal, rational and each admits a desingularization by trees of smooth rational curves, then* H, H' *are piecewise isomorphic, see Definition* 4.6*;*

(2) *If* H *or* H' *is a cubic, then* H, H' *are piecewise isomorphic. Moreover,* H *is smooth if and only if* H' *is smooth;*

(3) *If* H *or* H' *is a non-rational cubic, then there exists* $\varphi \in \mathrm{Aut}(\mathbb{P}^3_{\mathbf{k}})$ *with* $\varphi(H) = H'$.

We will prove Theorem C at the end of Section 5.2. For $\mathbf{k} = \mathbb{C}$, it is proved in [LS12] that if $\mathbb{P}^3_f \simeq \mathbb{P}^3_g$ for irreducible polynomials f, g, then $V_{\mathbb{P}^2}(f)$ and $V_{\mathbb{P}^2}(g)$ are piecewise isomorphic, which gives essentially Theorem C over the field of complex numbers.

To prove Theorem C, we study piecewise isomorphisms between surfaces. We prove in particular that every normal rational projective surface admitting a desingularization by trees of smooth rational curves (e.g. Du Val singularities) is piecewise isomorphic to the disjoint union of $\mathbb{A}^2$, one point and n copies of $\mathbb{A}^1$ for some $n \geqslant 1$ (Proposition 4.19). A similar result does not hold for all rational normal projective surfaces (Example 4.20). Using the Grothendieck ring $K_0(\mathrm{Var}_{\mathbf{k}})$ and the topological Euler characteristic (see Section 4 for the definition), one sees that the integer n

determines the piecewise isomorphism class. As a consequence of this, we give an affirmative answer to a question posed by Larsen and Lunts (see [LL03, Quest. 1.2]) for rational projective normal surfaces admitting a desingularization by trees of smooth rational curves (see Proposition 5.14). This also partially generalizes [LS10, Th. 4].

Proposition D. — *Let* $\mathbf{k}$ *be algebraically closed and let* X, Y *be rational normal projective surfaces admitting a desingularization by trees of smooth rational curves. If the classes* $[X]$ *and* $[Y]$ *coincide in the Grothendieck ring* $K_0(\mathrm{Var}_{\mathbf{k}})$*, then* X *and* Y *are piecewise isomorphic.*

We remark that in general the question of Larsen and Lunts has a negative answer. The first example was given in [Bor18, Th. 2.14] (in dimension 13) and later in [Mar16, Th. 1.1] (in dimension 9) and in [KS18, Th. 1.9] (in dimension 3).

Acknowledgements. — The authors would like to thank the referee for the very helpful remarks, suggestions and questions that improved the exposition.

2. Lift to affine hypersurfaces

Let $n \geqslant 1$. For each homogeneous polynomial $f \in \mathbf{k}[x_0, \dots, x_n]$ of degree $d \geqslant 1$ and each $\mu \in \mathbf{k}^*$, we denote by $X_{f,\mu} \subseteq \mathbb{A}^{n+1}$ the hypersurface given by

$$X_{f,\mu} = \mathrm{Spec}(\mathbf{k}[x_0, \dots, x_n]/(f - \mu)) \subseteq \mathbb{A}^{n+1},$$

and obtain a canonical finite morphism

$$\begin{aligned} \pi_{f,\mu} \colon X_{f,\mu} &\longrightarrow \mathbb{P}^n_f \\ (x_0, \dots, x_n) &\longmapsto [x_0 : \cdots : x_n] \end{aligned}$$

of degree d, which is an étale covering if $\mathrm{char}(\mathbf{k})$ does not divide d.

If $\mathbf{k}$ is algebraically closed, all the varieties $X_{f,\mu}$ are isomorphic to $X_{f,1}$ by homotheties.

If $\mathbf{k}$ is equal to the field of complex numbers $\mathbb{C}$, then $\pi_{f,\mu}$ is the universal abelian covering, as the abelianization of the fundamental group of $\mathbb{P}^n_f$ with respect to the Euclidean topology is equal to $\mathbb{Z}/d\mathbb{Z}$ [Lib05, Prop. 2.3].

We now prove that for each field $\mathbf{k}$, the isomorphisms between complements of hypersurfaces lift to these coverings. This will be useful in the sequel, in order to study isomorphisms between varieties $\mathbb{P}^n_f$ and $\mathbb{P}^n_g$.

Proposition 2.1. — *Let* $\mathbf{k}$ *be a field, let* $n \geqslant 1$ *and let* $f, g \in \mathbf{k}[x_0, \dots, x_n]$ *be irreducible homogeneous polynomials of degree* $d \geqslant 1$*. Let* $\Phi_0, \dots, \Phi_n \in \mathbf{k}[x_0, \dots, x_n]$ *be homogeneous polynomials of degree* $\ell \geqslant 1$*. Then, the following statements are equivalent:*

(1) *The rational map*

$$\begin{aligned} \Phi \colon \mathbb{P}^n_f &\xrightarrow{\simeq} \mathbb{P}^n_g, \\ [x_0 : \cdots : x_n] &\longmapsto [\Phi_0(x_0, \dots, x_n) : \cdots : \Phi_n(x_0, \dots, x_n)] \end{aligned}$$

is an isomorphism and the gcd of $\Phi_0, \dots, \Phi_n$ *is a power of* f*.*

(2) *There exists $\mu \in \mathbf{k}^*$ such that the following map is an isomorphism*

$$\begin{aligned}\varphi\colon X_{f,1} &\xrightarrow{\simeq} X_{g,\mu},\\ (x_0,\dots,x_n) &\longmapsto (\Phi_0(x_0,\dots,x_n),\dots,\Phi_n(x_0,\dots,x_n)).\end{aligned}$$

Moreover, if these statements hold, then ℓ is invertible modulo d.

Proof. — "(1) $\Rightarrow$ (2)": There exists $r \geqslant 0$ and $\Phi'_0,\dots,\Phi'_n \in \mathbf{k}[x_0,\dots,x_n]$ without common factor such that $\Phi_i = f^r\Phi'_i$ for all i. The domain $\mathrm{dom}(\Phi)$ of the rational self-map $\Phi\colon \mathbb{P}^n \dashrightarrow \mathbb{P}^n$ is thus equal to $\mathbb{P}^n \smallsetminus V_{\mathbb{P}^n}(\Phi'_0,\dots,\Phi'_n)$ and contains $\mathbb{P}^n_f$. As g is nowhere zero on $\mathbb{P}^n_g$, the polynomial $g(\Phi_0,\dots,\Phi_n) = f^{dr}g(\Phi'_0,\dots,\Phi'_n)$ is nowhere zero on $\mathbb{P}^n_f$ and is thus equal to μf^t for some $\mu \in \mathbf{k}^*$ and some integer $t \geqslant 1$. We then obtain a morphism

$$\varphi\colon X_{f,1} \longrightarrow X_{g,\mu}, \quad (x_0,\dots,x_n) \longmapsto (\Phi_0(x_0,\dots,x_n),\dots,\Phi_n(x_0,\dots,x_n)).$$

It remains to see that φ is an isomorphism.

The inverse of Φ is given by

$$\Psi\colon \mathbb{P}^n_g \xrightarrow{\simeq} \mathbb{P}^n_f, \quad [x_0:\cdots:x_n] \longmapsto [\Psi_0(x_0,\dots,x_n):\cdots:\Psi_n(x_0,\dots,x_n)],$$

where $\Psi_0,\dots,\Psi_n \in \mathbf{k}[x_0,\dots,x_n]$ are homogeneous polynomials of the same degree $\ell' \geqslant 1$, without common factor.

As $\Psi\circ\Phi = \mathrm{id}_{\mathbb{P}^n_f}$ and $\Phi\circ\Psi = \mathrm{id}_{\mathbb{P}^n_g}$, there exist homogeneous polynomials $A, B \in \mathbf{k}[x_0,\dots,x_n]$ of degree $\ell\ell' - 1$ such that

$$\Psi_i(\Phi_0,\dots,\Phi_n) = Ax_i \quad \text{and} \quad \Phi_i(\Psi_0,\dots,\Psi_n) = Bx_i \qquad \text{for all } i \in \{0,\dots,n\}.$$

Hence, the zero locus $V_{\mathbb{P}^n_f}(A)$ of A in $\mathbb{P}^n_f$ satisfies the following: Ψ_i vanishes on $\Phi(V_{\mathbb{P}^n_f}(A))$ for all $i \in \{0,\dots,n\}$ and $\Phi(V_{\mathbb{P}^n_f}(A))$ lies in the domain of Ψ; i.e., $V_{\mathbb{P}^n_f}(A)$ is empty. Analogously, one can see that $V_{\mathbb{P}^n_g}(B)$ is empty. We get $A = \lambda f^s$ and $B = \lambda' g^s$ where $\lambda, \lambda' \in \mathbf{k}^*$ and where $s \geqslant 1$ is such that $ds = \ell\ell' - 1$. In particular, ℓ is invertible modulo d. We replace Ψ_i with Ψ_i/λ for each $i \in \{0,\dots,n\}$ and reduce to the case where $\lambda = 1$.

Consider now the morphism

$$\psi\colon X_{g,\mu} \longrightarrow \mathbb{A}^{n+1}, \quad (x_0,\dots,x_n) \longmapsto (\Psi_0(x_0,\dots,x_n),\dots,\Psi_n(x_0,\dots,x_n)).$$

Using that $\Psi_i(\Phi_0,\dots,\Phi_n) = f^s x_i$ for each $i \in \{0,\dots,n\}$, we obtain that $\psi\circ\varphi\colon X_{f,1} \to \mathbb{A}^{n+1}$ is the natural closed embedding. Hence, $(\psi\circ\varphi)^* = \varphi^*\circ\psi^*\colon \mathbf{k}[\mathbb{A}^{n+1}] \to \mathbf{k}[X_{f,1}]$ is surjective and φ is dominant, i.e., $\varphi^*\colon \mathbf{k}[X_{g,\mu}] \to \mathbf{k}[X_{f,1}]$ is injective. This implies that φ^* and thus φ is an isomorphism.

"(2) $\Rightarrow$ (1)": As $\Phi_0,\dots,\Phi_n$ are homogeneous of degree ℓ, we may define a rational self-map $\Phi\colon [x_0:\cdots:x_n] \mapsto [\Phi_0(x_0,\dots,x_n):\cdots:\Phi_n(x_0,\dots,x_n)]$ of $\mathbb{P}^n$. We now prove that Φ restricts to an isomorphism $\mathbb{P}^n_f \xrightarrow{\simeq} \mathbb{P}^n_g$ and that the gcd of $\Phi_0,\dots,\Phi_n$ is a power of f.

By assumption $g(\Phi_0,\dots,\Phi_n) = (f-1)p + \mu$ for some polynomial $p \in \mathbf{k}[x_0,\dots,x_n]$ of degree $r = d(\ell-1) \geqslant 0$. Write $p = \sum_{i=0}^r p_i$ where p_i is homogeneous of degree i.

Since $g(\Phi_0,\dots,\Phi_n)$ is homogeneous, it follows that $p_0=\mu$ and $p_{di}=fp_{d(i-1)}$ for all $1\leqslant i\leqslant \ell-1$. This implies that

$$g(\Phi_0,\dots,\Phi_n)=\mu f^\ell.$$

Since g is homogeneous and f is irreducible the gcd of $\Phi_0,\dots,\Phi_n$ is a power of f. Moreover, Φ restricts to a morphism $\Phi\colon \mathbb{P}^n_f\to\mathbb{P}^n_g$.

Using that $\pi_{g,\mu}\circ\varphi=\Phi\circ\pi_{f,1}$, we get the following commutative diagram

$$\begin{array}{ccc}
\mathbf{k}(X_{g,\mu}) & \xrightarrow{\varphi^*} & \mathbf{k}(X_{f,1}) \\
{\scriptstyle\pi^*_{g,\mu}}\uparrow & & \uparrow{\scriptstyle\pi^*_{f,1}} \\
\mathbf{k}(\mathbb{P}^n_g) & \xrightarrow{\Phi^*} & \mathbf{k}(\mathbb{P}^n_f)
\end{array}$$

and that $\Phi\colon \mathbb{P}^n_f\to\mathbb{P}^n_g$ is a quasi-finite surjection. Note that $\pi^*_{f,1}$ and $\pi^*_{g,\mu}$ are field extensions of degree d. Since φ^* is an isomorphism, Φ^* is an isomorphism as well, i.e., $\Phi\colon \mathbb{P}^n_f\to\mathbb{P}^n_g$ is birational. Zariski's Main Theorem [Gro61, Cor. 4.4.9] gives that $\Phi\colon \mathbb{P}^n_f\to\mathbb{P}^n_g$ is an isomorphism. □

Remark 2.2. — If there exists an isomorphism $\Phi\colon \mathbb{P}^n_f\to\mathbb{P}^n_g$, there are homogeneous polynomials $\Phi_0,\dots,\Phi_n\in\mathbf{k}[x_0,\dots,x_n]$ of degree $\ell\geqslant 1$ without common factor such that $\Phi([x_0:\cdots:x_n])=[\Phi_0(x_0,\dots,x_n):\cdots:\Phi_n(x_0,\dots,x_n)]$. Proposition 2.1 then gives the existence of $\mu\in\mathbf{k}^*$ such that

$$\begin{aligned}\varphi\colon X_{f,1}&\xrightarrow{\simeq} X_{g,\mu},\\ (x_0,\dots,x_n)&\longmapsto(\Phi_0(x_0,\dots,x_n),\dots,\Phi_n(x_0,\dots,x_n))\end{aligned}$$

is an isomorphism. If $\mathbf{k}$ is moreover algebraically closed, then we may multiply each Φ_i with a root of μ^{-1} and assume that $\mu=1$. Hence, $X_{f,1}$ and $X_{g,1}$ are isomorphic.

Remark 2.3. — Let $\mathbf{k}$ be a field, $n\geqslant 1$ be an integer and let $f,g\in\mathbf{k}[x_0,\dots,x_n]$ be irreducible homogeneous polynomials of degree $d\geqslant 1$. Let $e\geqslant 1$ be an integer and let $\Phi_0,\dots,\Phi_n\in\mathbf{k}[x_0,\dots,x_n]$ be polynomials such that the degree of each of their monomials is congruent to e modulo d. If

$$X_{f,1}\xrightarrow{\simeq} X_{g,1},\quad (x_0,\dots,x_n)\longmapsto(\Phi_0(x_0,\dots,x_n),\dots,\Phi_n(x_0,\dots,x_n))$$

is an isomorphism, then we may multiply the monomials of Φ_i with some powers of f and assume that $\Phi_0,\dots,\Phi_n$ are homogeneous of the same degree $\ell\geqslant 1$, and then apply Proposition 2.1 to obtain an isomorphism $\mathbb{P}^n_f\xrightarrow{\simeq}\mathbb{P}^n_g$.

Lemma 2.4. — *Let $\mathbf{k}$ be a field, $n\geqslant 1$ be an integer, $f,g\in\mathbf{k}[x_0,\dots,x_n]$ be irreducible homogeneous polynomials of degree $d\geqslant 1$, and let $\Phi\colon \mathbb{P}^n_f\xrightarrow{\simeq}\mathbb{P}^n_g$ be an isomorphism given by homogeneous polynomials of degree ℓ such that their gcd is a power of f. If ℓ is congruent to 1 modulo d, then $\mathbb{P}^m_f$ and $\mathbb{P}^m_g$ are isomorphic for each $m\geqslant n$.*

Proof. — We write Φ as $[x_0 : \cdots : x_n] \mapsto [\Phi_0(x_0, \ldots, x_n) : \cdots : \Phi_n(x_0, \ldots, x_n)]$, where $\Phi_0, \ldots, \Phi_n \in \mathbf{k}[x_0, \ldots, x_n]$ are homogeneous of degree ℓ and the gcd of $\Phi_0, \ldots, \Phi_n$ is a power of f. By Proposition 2.1, the morphism

$$\varphi\colon X_{f,1} \xrightarrow{\simeq} X_{g,\mu}, \quad (x_0, \ldots, x_n) \longmapsto (\Phi_0(x_0, \ldots, x_n), \ldots, \Phi_n(x_0, \ldots, x_n))$$

is an isomorphism, where $X_{f,1}, X_{g,\mu} \subseteq \mathbb{A}^{n+1}$ are defined as before. If ℓ is congruent to 1 modulo d, we can write $\ell = 1 + dr$ for some integer $r \geqslant 0$. For each $m \geqslant n$, we thus get an isomorphism $X_{f,1} \times \mathbb{A}^{m-n} \xrightarrow{\simeq} X_{g,\mu} \times \mathbb{A}^{m-n}$ given by

$$(x_0, \ldots, x_m) \longmapsto (\varphi(x_0, \ldots, x_m), x_{n+1} f(x_0, \ldots, x_n)^r, \ldots, x_m f(x_0, \ldots, x_n)^r)).$$

Applying again Proposition 2.1, now in the converse direction, we get that the homogeneous polynomials $\Phi_0, \ldots, \Phi_n, x_{n+1} f^r, \ldots, x_m f^r$ of degree ℓ also define an isomorphism $\mathbb{P}^m_f \xrightarrow{\simeq} \mathbb{P}^m_g$. □

The next examples shows that $\ell \neq 1$ in Proposition 2.1 is possible.

Example 2.5. — Let $\mathbf{k}$ be a field and let $f = g = xyz + x^3 + y^3 \in \mathbf{k}[x, y, z]$, which is the equation of a nodal cubic curve $\Gamma \subset \mathbb{P}^2$. By blowing-up the singular point, then blowing-up successively points on the exceptional divisor created lying on the curve, we obtain a birational morphism $X \to \mathbb{P}^2$ with a strict transform $\widetilde{\Gamma} \subseteq X$ of Γ being a (-1)-curve. We then contract this curve and contract the exceptional divisors created, except the last one, to get an automorphism of $\mathbb{P}^2 \smallsetminus \Gamma$ that does not extend to $\mathbb{P}^2$ and has degree 8. Explicitly, we define $\Phi_0, \Phi_1, \Phi_2 \in \mathbf{k}[x, y, z]$, homogeneous of degree 8, by

$$\begin{aligned}
\Phi_0 &= (-x^4 z + 2x^3 y^2 - 2x^2 y z^2 + 2xy^3 z + y^5 - y^2 z^3) \cdot f, \\
\Phi_1 &= (x^2 + yz) \cdot f^2, \\
\Phi_2 &= x^7 y - x^6 z^2 + 6x^5 y^2 z - x^4 y^4 - 3x^4 y z^3 + 9x^3 y^3 z^2 \\
&\qquad + x^2 y^5 z - 3x^2 y^2 z^4 - xy^7 + 4xy^4 z^3 + 2y^6 z^2 - y^3 z^5,
\end{aligned}$$

and calculate that these define an involution

$$\mathbb{P}^2_f \xrightarrow{\simeq} \mathbb{P}^2_f, \quad [x : y : z] \longmapsto [\Phi_0(x, y, z) : \Phi_1(x, y, z) : \Phi_2(x, y, z)].$$

Here, the common degree is 8, that is not congruent to 1 modulo 3, but invertible in $\mathbb{Z}/3\mathbb{Z}$, as Proposition 2.1 says.

Lemma 2.6. — *Let $\mathbf{k}$ be a field, $n \geqslant 1$, $f \in \mathbf{k}[x_0, \ldots, x_n]$ an irreducible homogeneous polynomial of degree $d \geqslant 1$ and let $U \subseteq \mathbb{P}^n$ be an open subvariety. If there exists an isomorphism $\mathbb{P}^n_f \xrightarrow{\simeq} U$, then there exists an irreducible homogeneous polynomial $g \in \mathbf{k}[x_0, \ldots, x_n]$ of degree d with $U = \mathbb{P}^n_g$.*

Before we start the proof we recall the following fact: If X is an affine irreducible normal variety over a field $\mathbf{k}$ and if $U \subseteq X$ is a big open subset, i.e., the codimension of $X \smallsetminus U$ in X is at least 2, then $\mathbf{k}[U] = \mathbf{k}[X]$ (see [Mat86, Th. 11.5]). In particular, if U is affine as well, then $U = X$.

Proof of Lemma 2.6. — If U is big open in $\mathbb{P}^n$, then for any open affine subvariety $V \subseteq \mathbb{P}^n$, the affine subvariety $U \cap V$ is big open in V and hence $U \cap V = V$. This would imply that $U = \mathbb{P}^n$ is affine, a contradiction since $n \geqslant 1$. Hence the union Z of the $(n-1)$-dimensional irreducible components of $\mathbb{P}^n \smallsetminus U$ is non-empty. As $\mathbb{P}^n \smallsetminus Z$ is an affine big open subset of U we get $\mathbb{P}^n \smallsetminus Z = U$. Moreover, $\mathbf{k}[\mathbb{P}^n_f]^* = \mathbf{k}^*$ yields $\mathbf{k}[U]^* = \mathbf{k}^*$, which implies that Z is irreducible. Hence, there exists an irreducible homogeneous polynomial $g \in \mathbf{k}[x_0, \dots, x_n]$ such that Z is the zero locus of g, i.e., $U = \mathbb{P}^n_g$. Finally, $\mathbb{Z}/\deg(g)\mathbb{Z} \simeq \mathrm{Pic}(\mathbb{P}^n_g) \simeq \mathrm{Pic}(\mathbb{P}^n_f) \simeq \mathbb{Z}/d\mathbb{Z}$ implies that $\deg(g) = d$. □

3. Non-isomorphic hypersurfaces having isomorphic complements

3.1. Explicit isomorphisms between complements of projective cones. — The next formulas are inspired by isomorphisms between cylinders over Danielewski surfaces given in [MJP21].

Lemma 3.1. — *Let* $\mathbf{k}$ *be a field, let* $s, m, n \geqslant 0$ *be integers and let* $S = \mathbf{k}[x_0, x_1, \dots, x_s]$ *be a polynomial ring in* $s+1$ *variables over* $\mathbf{k}$*. Let* $P, Q \in S[z]$ *and denote by* H_P *and* H_Q *the hypersurfaces of* $\mathbb{A}^{s+3}_{\mathbf{k}} = \mathrm{Spec}(S[y,z])$ *defined by the equations*

$$x_0^n y = P(z) \quad \text{and} \quad x_0^n y = Q(z),$$

respectively. Suppose that there exist $A, B \in S[z]$ *such that*

(1) $z - A(B(z))$ *and* $z - B(A(z))$ *both belong to the ideal* $x_0^m S[z]$*,*
(2) *We have*

$$Q\left(A(z) + x_0^m w\right) \in x_0^n S[z,w] + P(z)S[z,w]$$
$$\text{and} \qquad P\left(B(z) + x_0^m w\right) \in x_0^n S[z,w] + Q(z)S[z,w],$$

where w *is transcendental over* $S[y,z]$*.*

Then, the following are inverse isomorphisms.

$$\Phi\colon H_P \times \mathbb{A}^1_{\mathbf{k}} \xrightarrow{\simeq} H_Q \times \mathbb{A}^1_{\mathbf{k}}$$
$$(x_0, \dots, x_s, y, z, w) \longmapsto \Big(x_0, \dots, x_s, \frac{Q\left(A(z) + x_0^m w\right)}{x_0^n}, A(z) + x_0^m w, \frac{z - B\left(A(z) + x_0^m w\right)}{x_0^m}\Big),$$

$$\Psi\colon H_Q \times \mathbb{A}^1_{\mathbf{k}} \xrightarrow{\simeq} H_P \times \mathbb{A}^1_{\mathbf{k}}$$
$$(x_0, \dots, x_s, y, z, w) \longmapsto \Big(x_0, \dots, x_s, \frac{P\left(B(z) + x_0^m w\right)}{x_0^n}, B(z) + x_0^m w, \frac{z - A\left(B(z) + x_0^m w\right)}{x_0^m}\Big).$$

Proof. — The hypotheses on A and B imply that Φ and Ψ are morphisms. Moreover, the compositions are given by

$$(\Psi \circ \Phi)(x_0, \dots, x_s, y, z, w) = (x_0, \dots, x_s, P(z)/x_0^n, z, w),$$
$$(\Phi \circ \Psi)(x_0, \dots, x_s, y, z, w) = (x_0, \dots, x_s, Q(z)/x_0^n, z, w). \qquad \square$$

Proposition 3.2. — *Let* $\mathbf{k}$ *be a field,* $d \geqslant 4$ *be integers and let* $f, g \in \mathbf{k}[x, y, z]$ *be the homogeneous polynomials of degree* d *defined by*

$$f = x^{d-1}y + z^d \quad \textit{and} \quad g = x^{d-1}y + z^d + dx^{d-2}z^2.$$

Then, for each $r \geqslant 1$*, the open subvarieties* $\mathbb{P}^{r+2}_f, \mathbb{P}^{r+2}_g \subseteq \mathbb{P}^{r+2}$ *are isomorphic.*

Remark 3.3. — The open subvarieties $\mathbb{P}^2_f, \mathbb{P}^2_g \subseteq \mathbb{P}^2$ are not isomorphic when $\mathbf{k} = \mathbb{C}$ is the field of complex numbers. This follows from Proposition 2.1 and from the fact that, for every $\mu \in \mathbb{C}^*$, the (Danielewski) hypersurfaces $X_{f,1}, X_{g,\mu} \subseteq \mathbb{A}^3_{\mathbb{C}}$ defined by the equations $f = 1$ and $g = \mu$, respectively, are not isomorphic (see [Pol11, Th. 9]).

Proof. — We define $s = 0$, $x_0 = x$, $S = \mathbf{k}[x]$, $m = d$, $n = d - 1$,

$$P(z) = 1 - z^d, \qquad Q(z) = 1 - z^d - dx^{d-2}z^2,$$
$$A(z) = z - x^{d-2}z^3, \qquad B(z) = z + x^{d-2}z^3,$$

and prove that the hypotheses of Lemma 3.1 are satisfied for the above polynomials in $S[z]$.

Firstly, we have $A(z) \equiv z \equiv B(z) \pmod{x^{d-2}}$. Thus, since $d \leqslant 2(d-2)$, as $d \geqslant 4$, we get $x^{d-2}A(z)^3 \equiv x^{d-2}z^3 \equiv x^{d-2}B(z)^3 \pmod{x^d}$ and we obtain

$$z - A(B(z)) = z - B(z) + x^{d-2}B(z)^3 \equiv 0 \pmod{x^d},$$
$$z - B(A(z)) = z - A(z) - x^{d-2}A(z)^3 \equiv 0 \pmod{x^d}.$$

We then check that $P\left(B(z) + x^d w\right) \in x^{d-1}S[z, w] + Q(z)S[z, w]$:

$$\begin{aligned} P(B(z) + x^d w) &\equiv P(B(z)) \\ &\equiv 1 - (z + x^{d-2}z^3)^d \\ &\equiv 1 - z^d - dx^{d-2}z^{d+2} \\ &\equiv (1 - z^d - dx^{d-2}z^2) \cdot (1 + dx^{d-2}z^2) \\ &\equiv Q(z) \cdot (1 + dx^{d-2}z^2) \pmod{x^{d-1}}. \end{aligned}$$

Similarly, we have $Q(A(z) + x^d w) \in x^{d-1}S[z, w] + P(z)S[z, w]$:

$$\begin{aligned} Q(A(z) + x^d w) &\equiv Q\left(A(z)\right) \\ &\equiv 1 - (z - x^{d-2}z^3)^d - dx^{d-2}(z - x^{d-2}z^3)^2 \\ &\equiv 1 - z^d + dx^{d-2}z^{d+2} - dx^{d-2}z^2 \\ &\equiv (1 - z^d) \cdot (1 - dx^{d-2}z^2) \\ &\equiv P(z) \cdot (1 - dx^{d-2}z^2) \pmod{x^{d-1}}. \end{aligned}$$

Noting that $f-1 = x^{d-1}y - P(z)$ and $g-1 = x^{d-1}y - Q(z)$, we may now apply Lemma 3.1 to obtain an isomorphism

$$\begin{aligned}\varphi\colon \operatorname{Spec}(\mathbf{k}[x,y,z,w]/(f-1)) &\xrightarrow{\simeq} \operatorname{Spec}(\mathbf{k}[x,y,z,w]/(g-1))\\ (x,y,z,w) &\longmapsto \Big(x, \frac{Q(A(z)+x^dw)}{x^{d-1}}, A(z)+x^dw,\\ &\qquad\qquad \frac{z-B(A(z)+x^dw)}{x^d}\Big).\end{aligned}$$

Arguing as in Remark 2.3, we may then use this isomorphism to construct an isomorphism $\mathbb{P}^3_f \xrightarrow{\simeq} \mathbb{P}^3_g$. Indeed, the second component of φ can be replaced by a polynomial, since

$$Q\left(A(z)+x^dw\right) \equiv P\cdot(1-dx^{d-2}z^2) \pmod{x^{d-1}}$$

and $P \equiv x^{d-1}y \pmod{f-1}$. Moreover, the fourth component of φ is a polynomial, since $z - A(B(z)) \equiv 0 \pmod{x^d}$. Doing this, we obtain an expression for φ given by four polynomials. It remains to check that each monomial appearing in that polynomial expression of φ has degree congruent to 1 modulo d. Working modulo d with the degrees, we observe that A, B, P, Q are homogeneous of degree $1, 1, 0, 0$, respectively, hence that the polynomials $z - B(A(z)+x^dw)$, $Q(A(z)+x^dw)$ and $P - x^{d-1}y$ are homogeneous of degree 1, 0 and 0, respectively. Therefore, we can conclude that $\mathbb{P}^3_f \xrightarrow{\simeq} \mathbb{P}^3_g$ as in Remark 2.3 and furthermore that $\mathbb{P}^{r+2}_f \xrightarrow{\simeq} \mathbb{P}^{r+2}_g$ for all $r \geqslant 1$ by Lemma 2.4. □

Proposition 3.4. — *Let* $\mathbf{k}$ *be a field,* $d \geqslant 3$ *be an integer and let* $f, g \in \mathbf{k}[x_0, x_1, y, z]$ *be the homogeneous polynomials of degree* d *defined by*

$$f = x_0^{d-1}y + z^d \quad \textit{and} \quad g = x_0^{d-1}y + z^d + dx_0^{d-2}x_1^2.$$

Then, for each $r \geqslant 1$*, the open subvarieties* $\mathbb{P}^{r+3}_f, \mathbb{P}^{r+3}_g \subseteq \mathbb{P}^{r+3}$ *are isomorphic.*

Remark 3.5. — We don't know, whether the open subvarieties $\mathbb{P}^3_f, \mathbb{P}^3_g \subseteq \mathbb{P}^3$ are isomorphic, or even if the hypersurfaces $X_{f,1}, X_{g,1} \subseteq \mathbb{A}^3$ given by $f = 1$ and $g = 1$, respectively, are isomorphic or not.

Proof of Proposition 3.4. — We define $s = 1$, $S = \mathbf{k}[x_0, x_1]$, $m = d$, $n = d-1$, $\Delta = x_0^{d-2}x_1^2$,

$$\begin{aligned} P(z) &= 1 - z^d, & Q(z) &= 1 - z^d - d\Delta,\\ A(z) &= z(1 - \Delta + \Delta^2), & B(z) &= z(1+\Delta),\end{aligned}$$

and prove that the hypotheses of Lemma 3.1 are satisfied for the above polynomials in $S[z]$.

Firstly, we calculate $A(B(z)) = B(A(z)) = z(1-\Delta+\Delta^2)(1+\Delta) = z(1+\Delta^3)$. As $d \geqslant 3$ we have $3d - 6 \geqslant d$, so Δ^3 is divisible by x_0^d. Hence, $A(B(z)) = B(A(z))$ is congruent to z modulo x_0^d.

We then check that $P(B(z)+x_0^d w) \in x_0^{d-1}S[z,w] + Q(z)S[z,w]$. As $d \geqslant 3$, Δ^2 is divisible by x_0^{d-1}. Hence:

$$\begin{aligned} P(B(z)+x_0^d w) &\equiv P(B(z)) \\ &\equiv 1 - z^d(1+\Delta)^d \\ &\equiv 1 - z^d(1+d\Delta) \\ &\equiv (1 - z^d - d\Delta)\cdot(1+d\Delta) \\ &\equiv Q\cdot(1+d\Delta) \pmod{x_0^{d-1}}. \end{aligned}$$

Similarly, we have $Q(A(z)+x_0^d w) \in x_0^{d-1}S[z,w] + P(z)S[z,w]$:

$$\begin{aligned} Q(A(z)+x_0^d w) &\equiv Q(A(z)) \\ &\equiv 1 - z^d(1-\Delta)^d - d\Delta \\ &\equiv 1 - z^d(1-d\Delta) - d\Delta \\ &\equiv (1 - z^d)\cdot(1-d\Delta) \\ &\equiv P\cdot(1-d\Delta) \pmod{x_0^{d-1}}. \end{aligned}$$

Since $f-1 = x_0^{d-1}y - P(z)$ and $g-1 = x_0^{d-1}y - Q(z)$, we may now apply Lemma 3.1 to obtain an isomorphism

$$\begin{aligned} \varphi\colon \operatorname{Spec}(\mathbf{k}[x_0,x_1,y,z,w]/(f-1)) &\xrightarrow{\simeq} \operatorname{Spec}(\mathbf{k}[x_0,x_1,y,z,w]/(g-1)) \\ (x_0,x_1,y,z,w) &\longmapsto \Big(x_0, x_1, \frac{Q(A(z)+x_0^d w)}{x_0^{d-1}}, A(z)+x_0^d w, \\ &\qquad \frac{z - B(A(z)+x_0^d w)}{x_0^d}\Big). \end{aligned}$$

We again proceed as in Remark 2.3. The third component of φ can be expressed by a polynomial, since

$$Q(A(z)+x_0^d w) \equiv P\cdot(1 - dx_0^{d-2}x_1^2) \pmod{x_0^{d-1}}$$

and $P \equiv x_0^{d-1}y \pmod{f-1}$. The last component of φ is already a polynomial, as its numerator is a multiple of its denominator. It remains to check that each monomial appearing in the expression of φ has degree congruent to 1 modulo d. For this, we simply work with the degrees modulo d and observe that Δ is homogeneous of degree 0, so that A, B, P, Q are homogeneous of degree $1, 1, 0, 0$, respectively. Hence, $Q(A(z)+x_0^d w)$, $A(z)+x_0^d w$ and $z - B(A(z)+x_0^d w)$ are homogeneous of degree 0, 1 and 1, respectively. Therefore, we can conclude by Remark 2.3 and Lemma 2.4. □

3.2. Non-isomorphic hypersurfaces. — We now want to prove that the hypersurfaces of Propositions 3.2 and 3.4 are not isomorphic (except when the characteristic divides d, in which case both are in fact equal).

Recall that the multiplicity $\operatorname{mult}_p(X)$ of a variety X at a point p is the multiplicity of the maximal ideal $\mathfrak{m}_{X,p}$ in the local ring $\mathcal{O}_{X,p}$. If $X \subseteq \mathbb{A}^n$ is a hypersurface given by a reduced polynomial $f \in \mathbf{k}[x_1,\dots,x_n]$ of degree d, then X has multiplicity d at

the origin, if and only if f is homogeneous. In particular, if $X \subseteq \mathbb{P}^n$ is a hypersurface given by a reduced homogeneous polynomial $F \in \mathbf{k}[x_0, \dots, x_n]$ of degree d, then X has multiplicity d at $p = [0 : \cdots : 0 : 1 : 0 : \cdots : 0]$ if and only if F does not depend on x_i, where $i \in \{0, \dots, n\}$ is the index of the unique non-zero coordinate of p.

Lemma 3.6. — *Let* $\mathbf{k}$ *be a field, let* $n \geqslant 1$ *and let* $h \in \mathbf{k}[x_0, \dots, x_n]$ *be a reduced homogeneous polynomial of degree* $d \geqslant 1$. *Let* $q \in V_{\mathbb{P}^{n+1}}(h)(\mathbf{k})$ *be a* $\mathbf{k}$*-rational point of multiplicity* d. *Then, the exceptional divisor* E_q *of the blow-up of* $V_{\mathbb{P}^{n+1}}(h)$ *at* q *is isomorphic to* $V_{\mathbb{P}^n}(h)$.

Proof. — We first write $p = [0 : \cdots : 0 : 1]$ and observe that it is a point of multiplicity d of $V_{\mathbb{P}^{n+1}}(h)$. The blow-up of $V_{\mathbb{P}^{n+1}}(h)$ at p is then given by

$$\left\{([x_0 : \cdots : x_{n+1}], [y_0 : \cdots : y_n]) \in V_{\mathbb{P}^{n+1}}(h) \times V_{\mathbb{P}^n}(h) \;\middle|\; \begin{array}{c} x_i y_j = x_j y_i \\ \forall i, j \in \{0, \dots, n\} \end{array}\right\},$$

so the exceptional divisor E_p of p is given by $\{p\} \times V_{\mathbb{P}^n}(h)$ and is thus isomorphic to $V_{\mathbb{P}^n}(h)$. This gives the result in the case where $q = p$.

Suppose now that $q \neq p$. Applying an automorphism of $\mathbb{P}^{n+1}$ that fixes p, we may assume that $q = [0 : \cdots : 0 : 1 : 0]$. This implies that $h \in \mathbf{k}[x_0, \dots, x_{n-1}]$. Hence, the exchange of x_n and x_{n+1} is an automorphism of $\mathbb{P}^{n+1}$ that exchanges p and q and preserves $V_{\mathbb{P}^{n+1}}(h)$. The exceptional divisors of p and q are then isomorphic. □

Lemma 3.7. — *Let* $\mathbf{k}$ *be a field, let* $n \geqslant 1$ *and let* $f, g \in \mathbf{k}[x_0, \dots, x_n]$ *be reduced homogeneous polynomials of the same degree* $d \geqslant 1$. *If* $V_{\mathbb{P}^{n+1}}(f)$ *and* $V_{\mathbb{P}^{n+1}}(g)$ *are isomorphic, then* $V_{\mathbb{P}^n}(f)$ *and* $V_{\mathbb{P}^n}(g)$ *are isomorphic.*

Proof. — An isomorphism $V_{\mathbb{P}^{n+1}}(f) \xrightarrow{\simeq} V_{\mathbb{P}^{n+1}}(g)$ sends $p = [0 : \cdots : 0 : 1]$ onto a $\mathbf{k}$-rational point of multiplicity d. It then induces an isomorphism between the exceptional divisors, and thus induces an isomorphism $V_{\mathbb{P}^n}(f) \xrightarrow{\simeq} V_{\mathbb{P}^n}(g)$, by Lemma 3.6. □

Lemma 3.8. — *Let* $\mathbf{k}$ *be a field, let* $d \geqslant 4$ *be an integer that is not a multiple of* $\operatorname{char}(\mathbf{k})$ *and let*

$$f = x^{d-1}y + z^d \quad \text{and} \quad g = x^{d-1}y + z^d + \mu x^{d-2}z^2,$$

where $\mu \in \mathbf{k}$. *Suppose that* $\alpha \colon V_{\mathbb{P}^2}(f) \xrightarrow{\simeq} V_{\mathbb{P}^2}(g)$ *is an isomorphism. Then,* $\mu = 0$ *and* α *is equal to* $[x : y : z] \mapsto [a^d x : y : a^{d-1}z]$ *for some* $a \in \mathbf{k}^*$.

Remark 3.9. — If $d \in \{2, 3\}$ is not a multiple of $\operatorname{char}(\mathbf{k})$, then there is an automorphism of $\mathbb{P}^2$ that sends $V_{\mathbb{P}^2}(f)$ onto $V_{\mathbb{P}^2}(g)$.

Proof. — Expressing y in terms of x, z, we obtain two birational morphisms

$$\begin{array}{l} \tau \colon \mathbb{P}^1 \longrightarrow V_{\mathbb{P}^2}(f) \\ [u : v] \longmapsto [u^d : -v^d : u^{d-1}v] \end{array} \quad \text{and} \quad \begin{array}{l} \tau' \colon \mathbb{P}^1 \longrightarrow V_{\mathbb{P}^2}(g) \\ [u : v] \longmapsto [u^d : -v^d - \mu u^{d-2}v^2 : u^{d-1}v] \end{array}$$

whose inverses are given by $[x : y : z] \mapsto [x : z]$. They induce isomorphisms

$$\mathbb{P}^1 \smallsetminus \{[0 : 1]\} \xrightarrow{\simeq} V_{\mathbb{P}^2}(f) \smallsetminus \{[0 : 1 : 0]\} \quad \text{and} \quad \mathbb{P}^1 \smallsetminus \{[0 : 1]\} \xrightarrow{\simeq} V_{\mathbb{P}^2}(g) \smallsetminus \{[0 : 1 : 0]\}.$$

Both τ and τ' are bijective and they send $[0:1]$ onto $[0:1:0]$, which is the unique singular point of $V_{\mathbb{P}^2}(f)$ and $V_{\mathbb{P}^2}(g)$, respectively. Since the isomorphism $\alpha\colon V_{\mathbb{P}^2}(f) \xrightarrow{\simeq} V_{\mathbb{P}^2}(g)$ must fix the point $[0:1:0]$, the birational map $\widehat{\alpha} = (\tau')^{-1}\alpha\tau$ is an automorphism of $\mathbb{P}^1$ that fixes the point $[0:1]$. Thus, $\widehat{\alpha}$ is of the form $[u:v] \mapsto [u : av + bu]$ for some $a \in \mathbf{k}^*$, $b \in \mathbf{k}$.

Note that $s_a\colon [x:y:z] \mapsto [a^d x : y : a^{d-1}z]$ is an automorphism of $V_{\mathbb{P}^2}(f)$, that lifts to $\widehat{s}_a = (\tau)^{-1}s_a\tau = [u:v] \mapsto [au:v]$. Replacing α with αs_a, we replace $\widehat{\alpha}$ with $\widehat{\alpha}\widehat{s}_a$, and we may assume that $a = 1$. It remains to see that $b = \mu = 0$. We calculate

$$\begin{aligned}\alpha([x:y:z]) &= \tau'\widehat{\alpha}\tau^{-1}([x:y:z])\\ &= \tau'\widehat{\alpha}([x:z])\\ &= \tau'([x:z+bx])\\ &= [x^d : -(z+bx)^d - \mu x^{d-2}(z+bx)^2 : x^{d-1}(z+bx)].\end{aligned}$$

As α fixes $[0:1:0]$, the pull-back by α of x/y and z/y are rational functions

$$\frac{x^d}{-(z+bx)^d - \mu x^{d-2}(z+bx)^2} \qquad \text{and} \qquad \frac{x^{d-1}}{-(z+bx)^{d-1} - \mu x^{d-2}(z+bx)},$$

whose restrictions to $V_{\mathbb{P}^2}(f)$ are regular at $[0:1:0]$. For each $j \in \{d-1, d\}$, there are thus two homogeneous polynomials $P_j, Q_j \in \mathbf{k}[x,y,z]$, of the same degree, such that $Q_j(0,1,0) \neq 0$ and such that

$$\frac{P_j(x,y,z)}{Q_j(x,y,z)} = \frac{x^j}{-(z+bx)^j - \mu x^{d-2}(z+bx)^{j+2-d}}$$

on the curve $V_{\mathbb{P}^2}(f)$. Using the morphism

$$\mathbb{A}^1 \longrightarrow V_{\mathbb{P}^2}(f), \quad t \longmapsto \tau([t:1]) = [t^d : -1 : t^{d-1}] = [t : -t^{1-d} : 1],$$

we find

$$\frac{P_j(t^d, -1, t^{d-1})}{Q_j(t^d, -1, t^{d-1})} = \frac{t^j}{B_j},$$

where $B_j = -(1+bt)^j - \mu t^{d-2}(1+bt)^{j+2-d} \in \mathbf{k}[t]$. As $Q_j(0,1,0) \neq 0$, the polynomial $Q_j(t^d, -1, t^{d-1}) \in \mathbf{k}[t]$ is not divisible by t. As $B_j(0) \neq 0$ (we use here $d \geqslant 4$), t^j divides $P_j(t^d, -1, t^{d-1})$. There is thus $A_j \in \mathbf{k}[t]$ such that

$$P_j(t^d, -1, t^{d-1}) = A_j \cdot t^j \quad \text{and} \quad Q_j(t^d, -1, t^{d-1}) = A_j \cdot B_j.$$

We consider now the case $j = d$ and prove that $b = 0$. We shall afterward prove $\mu = 0$ by considering the case $j = d-1$.

As $d \geqslant 4$, the equality $P_d(t^d, -1, t^{d-1}) = A_d \cdot t^d$ gives the existence of $\varepsilon \in \mathbf{k}$ such that $A_d \equiv \varepsilon \pmod{t^2}$. We moreover have $B_d \equiv -1 - bdt \pmod{t^2}$. This gives

$$Q_d(t^d, -1, t^{d-1}) = A_d \cdot B_d \equiv -\varepsilon - \varepsilon bdt \pmod{t^2}.$$

The polynomial $Q(t^d, -1, t^{d-1})$ is not divisible by t, so $\varepsilon \neq 0$, and its coefficient of t is zero, as $d \geqslant 4$. Since $\mathrm{char}(\mathbf{k})$ does not divide d, this gives $b = 0$.

We now use $j = d-1$. As $d \geqslant 4$, the equality $P_{d-1}(t^d, -1, t^{d-1}) = A_{d-1} \cdot t^{d-1}$ gives the existence of $\xi, \xi' \in \mathbf{k}$ such that $A_{d-1} \equiv \xi + t\xi' \pmod{t^{d-1}}$. As $b = 0$, we find $B_{d-1} = B_d = -1 - \mu t^{d-2}$. This gives

$$Q_{d-1}(t^d, -1, t^{d-1}) = A_{d-1} \cdot B_{d-1} \equiv -\xi - t\xi' - \mu\xi t^{d-2} \pmod{t^{d-1}}.$$

As t does not divide this polynomial, $\xi \neq 0$. Moreover, we obtain $\xi' = 0$ and $\mu = 0$, as $d \geqslant 4$. □

Proposition 3.10. — *Let* $\mathbf{k}$ *be a field, let* $d \geqslant 4$ *be an integer that is not a multiple of* $\mathrm{char}(\mathbf{k})$ *and let* $f, g \in \mathbf{k}[x, y, z]$ *be defined by*

$$f = x^{d-1}y + z^d \text{ and } g = x^{d-1}y + z^d + \mu x^{d-2}z^2,$$

where $\mu \in \mathbf{k}^*$. *Then, for each* $r \geqslant 0$, *the two hypersurfaces* $V_{\mathbb{P}^{r+2}}(f)$, $V_{\mathbb{P}^{r+2}}(g)$ *in* $\mathbb{P}^{r+2}$ *are not isomorphic.*

Proof. — By Lemma 3.8, the result is true when $r = 0$. Using Lemma 3.7, we can then argue by induction and obtain the result for every integer r. □

Proposition 3.11. — *Let* $\mathbf{k}$ *be a field,* $d \geqslant 3$ *be an integer and let* $f, g \in \mathbf{k}[x_0, x_1, y, z]$ *be defined by*

$$f = x_0^{d-1}y + z^d, \quad g = x_0^{d-1}y + z^d + \mu x_0^{d-2}x_1^2,$$

where $\mu \in \mathbf{k}^*$. *Then, for each* $r \geqslant 0$, *the two hypersurfaces* $V_{\mathbb{P}^{r+3}}(f)$, $V_{\mathbb{P}^{r+3}}(g)$ *in* $\mathbb{P}^{r+3}$ *are not isomorphic.*

Proof. — By Lemma 3.7, it suffices to consider the case $r = 0$ and to prove that $V_{\mathbb{P}^3}(f)$ and $V_{\mathbb{P}^3}(g)$ are not isomorphic. Looking at the derivative with respect to y, the singular locus of both hypersurfaces is contained in the line $\ell \subseteq \mathbb{P}^3$ given by $x_0 = z = 0$. The surface $V_{\mathbb{P}^3}(f)$ has multiplicity d at the point where $x_0 = y = z = 0$.

It remains to see that $V_{\mathbb{P}^3}(g)$ has multiplicity $< d$ at every point and we have to check this only for points in ℓ. For this, write $g = z^d + x_0^{d-2}(x_0y + \mu x_1^2)$ and observe that $V_{\mathbb{P}^3}(x_0y + \mu x_1^2)$ is smooth outside $x_0 = x_1 = y = 0$ and thus on ℓ. Hence, $x_0^{d-2}(x_0y + \mu x_1^2)$ has multiplicity $d-2$ or $d-1$ at any point of ℓ, which implies that g has multiplicity $d-2$ or $d-1$ at any point of ℓ. □

We may now prove Theorem A, which is a direct consequence of Propositions 3.2, 3.4, 3.10 and 3.11.

Proof of Theorem A. — As in the statement, we take a field $\mathbf{k}$, and integers $d, n \geqslant 3$ such that d is not a multiple of $\mathrm{char}(\mathbf{k})$ and such that $(d, n) \neq (3, 3)$. We are looking for hypersurfaces $H = V_{\mathbb{P}^n}(f), H' = V_{\mathbb{P}^n}(g) \subseteq \mathbb{P}^n_{\mathbf{k}}$ that are not isomorphic but have isomorphic complements.

If $d \geqslant 4$, we may choose $f = x^{d-1}y + z^d$ and $g = x^{d-1}y + z^d + dx^{d-2}z^2$, that we consider in $\mathbf{k}[x, y, z, w_1, \ldots, w_r]$ with $r = n - 2 \geqslant 1$. By Propositions 3.2, the complements of H and H' are isomorphic, and by Proposition 3.10, the hypersurfaces H and H' are not isomorphic.

If $n \geqslant 4$, we may choose $f = x_0^{d-1}y + z^d$ and $g = x_0^{d-1}y + z^d + dx_0^{d-2}x_1^2$, that we consider in $\mathbf{k}[x_0, x_1, y, z, w_1, \ldots, w_r]$ with $r = n-3 \geqslant 1$. By Propositions 3.4, the complements of H and H' are isomorphic, and by Proposition 3.11, the hypersurfaces H and H' are not isomorphic. □

4. Topological Euler characteristic and piecewise isomorphisms

Throughout this section we assume that $\mathbf{k}$ is algebraically closed. In the sequel we recall the definition and some basic facts of the topological Euler characteristic. In order to do this, we also recall some facts from étale ℓ-adic cohomology with compact support. As a reference we take [Mil13] and [Mil80].

Let ℓ be a prime number that is different from the characteristic of the ground field $\mathbf{k}$. For a variety X, the group $H_c^i(X_{\text{ét}}, \mathbb{Q}_\ell)$ denotes the i-th étale ℓ-adic cohomology with compact support, i.e.,

$$H_c^i(X_{\text{ét}}, \mathbb{Q}_\ell) \coloneqq \left(\varprojlim H_c^i(X_{\text{ét}}, \mathbb{Z}/\ell^n\mathbb{Z})\right) \otimes_{\mathbb{Z}_\ell} \mathbb{Q}_\ell,$$

where $\mathbb{Z}_\ell \coloneqq \varprojlim \mathbb{Z}/\ell^n\mathbb{Z}$ and $\mathbb{Q}_\ell$ is the quotient field of $\mathbb{Z}_\ell$, see e.g. [Mil13, §18, §19]. We have:

Lemma 4.1. — *Let X be a variety. Then the $\mathbb{Q}_\ell$-vector space $H_c^i(X_{\text{ét}}, \mathbb{Q}_\ell)$ has finite dimension and vanishes for $i > 2\dim X$.*

Proof. — Let $k\colon X \to \overline{X}$ be a completion. By definition (see e.g. [Mil13, Def. 18.1]) we have

$$H_c^i(X_{\text{ét}}, \mathbb{Z}/\ell^n\mathbb{Z}) = H^i(\overline{X}_{\text{ét}}, k_!(\mathbb{Z}/\ell^n\mathbb{Z})).$$

By [Mil13, Th. 19.2] the limit $\varprojlim H^i(\overline{X}_{\text{ét}}, k_!(\mathbb{Z}/\ell^n\mathbb{Z}))$ is a finitely generated $\mathbb{Z}_\ell$-module; this implies the first statement. By [Mil13, Th. 15.1], we have that the $\mathbb{Z}_\ell$-module $H^i(\overline{X}_{\text{ét}}, k_!(\mathbb{Z}/\ell^n\mathbb{Z}))$ vanishes for all $i > 2\dim X$ and hence the second statement follows. □

The topological Euler-characteristic of a variety X is defined by

$$\chi(X) \coloneqq \sum_{i=0}^{2\dim X} (-1)^i \dim_{\mathbb{Q}_\ell} H_c^i(X_{\text{ét}}, \mathbb{Q}_\ell).$$

The following properties are very useful in order to compute the topological Euler characteristic. For lack of a reference with proof, we give an argument here:

Lemma 4.2

(1) *Let X be a variety and let $Z \subseteq X$ be a closed subvariety. Then $\chi(X) = \chi(X \smallsetminus Z) + \chi(Z)$.*

(2) *Let X, Y be varieties. Then $\chi(X \times Y) = \chi(X) \cdot \chi(Y)$.*

Proof

(1) Let $U \coloneqq X \smallsetminus Z$. By [AGV73, Ch. XVII, §5.1.16] we have a long exact sequence

$$\cdots \longrightarrow H^i_c(U_{\text{ét}}, \mathbb{Z}/\ell^n\mathbb{Z}) \longrightarrow H^i_c(X_{\text{ét}}, \mathbb{Z}/\ell^n\mathbb{Z}) \longrightarrow H^i_c(Z_{\text{ét}}, \mathbb{Z}/\ell^n\mathbb{Z}) \longrightarrow H^{i+1}_c(U_{\text{ét}}, \mathbb{Z}/\ell^n\mathbb{Z}) \longrightarrow \cdots .$$

Taking the limit over n and tensoring with $\mathbb{Q}_\ell$ over $\mathbb{Z}_\ell$ gives a long exact sequence

$$\cdots \longrightarrow H^i_c(U_{\text{ét}}, \mathbb{Q}_\ell) \longrightarrow H^i_c(X_{\text{ét}}, \mathbb{Q}_\ell) \longrightarrow H^i_c(Z_{\text{ét}}, \mathbb{Q}_\ell) \longrightarrow H^{i+1}_c(U_{\text{ét}}, \mathbb{Q}_\ell) \longrightarrow \cdots .$$

The statement follows now by using Lemma 4.1.

(2) By the Künneth-formula (see [Mil80, Ch. VI, Cor. 8.23]) we get for $0 \leqslant k \leqslant 2(\dim X + \dim Y)$

$$\dim_{\mathbb{Q}_\ell} H^k_c((X \times Y)_{\text{ét}}, \mathbb{Q}_\ell) = \sum_{i=0}^{k} \dim_{\mathbb{Q}_\ell} H^i_c(X_{\text{ét}}, \mathbb{Q}_\ell) \otimes_{\mathbb{Q}_\ell} H^{k-i}_c(Y_{\text{ét}}, \mathbb{Q}_\ell),$$

which gives $\chi(X \times Y) = \chi(X) \cdot \chi(Y)$. □

If X is a complete variety, then $H^k_c(X_{\text{ét}}, \mathbb{Q}_\ell) = H^k(X_{\text{ét}}, \mathbb{Q}_\ell)$ (see e.g. [Mil13, Def. 18.1]). This will be used in the following examples.

Example 4.3. — Let C be a smooth irreducible projective curve of genus g. Then

$$H^0(C_{\text{ét}}, \mathbb{Q}_\ell) = H^2(C_{\text{ét}}, \mathbb{Q}_\ell) = \mathbb{Q}_\ell \quad \text{and} \quad H^1(C_{\text{ét}}, \mathbb{Q}_\ell) = \mathbb{Q}_\ell^{2g},$$

see [Mil13, Prop. 14.2]. Using that $H^i(C_{\text{ét}}, \mathbb{Q}_\ell) = 0$ for all $i > 2$ (see Lemma 4.1), it follows that $\chi(C) = 2 - 2g$.

Example 4.4. — Let $m \geqslant 0$. Then

$$H^i(\mathbb{P}^m_{\text{ét}}, \mathbb{Q}_\ell) = \begin{cases} \mathbb{Q}_\ell, & 0 \leqslant i \leqslant 2m \text{ and } i \text{ is even}, \\ 0, & \text{otherwise}, \end{cases}$$

see [Mil13, Ex. 16.3]. Hence, $\chi(\mathbb{P}^m) = m + 1$. Using Lemma 4.2 and that $\mathbb{A}^m \simeq \mathbb{P}^m \smallsetminus \mathbb{P}^{m-1}$ for all $m \geqslant 0$ (where $\mathbb{P}^{m-1}$ is linearly embedded in $\mathbb{P}^m$ and $\mathbb{P}^{-1} = \varnothing$), it follows that $\chi(\mathbb{A}^m) = 1$ for all $m \geqslant 0$.

Example 4.5. — Let $n \geqslant 0$ and let $\pi\colon X \to B$ be a locally trivial $\mathbb{A}^n$-bundle with respect to the Zariski topology, where B is any variety. Then $\chi(X) = \chi(B)$. Indeed, we proceed by induction on $\dim B$ and note that the case $\dim B = 0$ is clear. By assumption there exists a closed subvariety $B' \subseteq B$ such that π is trivial over $B \smallsetminus B'$ and $\dim B' < \dim B$. Hence,

$$\chi(X) = \chi(\mathbb{A}^n \times (B \smallsetminus B')) + \chi(\pi^{-1}(B')) = \chi(\mathbb{A}^n \times B) = \chi(\mathbb{A}^n)\chi(B) = \chi(B),$$

where for the second equality we used the induction hypothesis $\chi(\pi^{-1}(B')) = \chi(B') = \chi(\mathbb{A}^n \times B')$.

Let us denote by $\mathrm{Var}_{\mathbf{k}}$ the set of isomorphism classes of varieties (over $\mathbf{k}$) and denote by $\mathbb{Z}\,\mathrm{Var}_{\mathbf{k}}$ the free abelian group over $\mathrm{Var}_{\mathbf{k}}$. Moreover, let $I \subseteq \mathbb{Z}\,\mathrm{Var}_{\mathbf{k}}$ be the subgroup that is generated by

$$[X] - [Z] - [X \smallsetminus Z]$$

for all varieties X and closed subvarieties Z (here $[W]$ denotes the isomorphism class of a variety W). By bilinear extension of the operation $[X] \cdot [Y] \coloneqq [X \times Y]$ we get a ring structure on $\mathbb{Z}\,\mathrm{Var}_{\mathbf{k}}$ and I is an ideal of it. The *Grothendieck ring* is then the quotient of $\mathbb{Z}\,\mathrm{Var}_{\mathbf{k}}$ by I:

$$K_0(\mathrm{Var}_{\mathbf{k}}) \coloneqq \mathbb{Z}\,\mathrm{Var}_{\mathbf{k}}\,/I,$$

see e.g. [LS10, §2.2.1]. By abuse of notation we denote by $[X]$ the class of a variety X in $K_0(\mathrm{Var}_{\mathbf{k}})$. Using Lemma 4.2, the topological Euler characteristic gives a ring homomorphism

$$\chi\colon K_0(\mathrm{Var}_{\mathbf{k}}) \longrightarrow \mathbb{Z}, \qquad \sum_{i=0}^{n} a_i[X_i] \longmapsto \sum_{i=0}^{n} a_i\chi(X_i).$$

Definition 4.6. — Let X, Y be varieties. Then X, Y are called *piecewise isomorphic* if X and Y can be decomposed as disjoint unions of $n \geqslant 1$ locally closed subsets $X_1, \ldots, X_n$ and $Y_1, \ldots, Y_n$, respectively, such that $X_i \simeq Y_i$ for all $1 \leqslant i \leqslant n$. Note that we take on X_i the reduced scheme structure.

Note that "piecewise isomorphic" defines an equivalence relation among the varieties. It seems that these ideas of "cutting and pasting" appeared the first time in a letter of Grothendieck to Serre, see [CS01, Letter from 16.8.1964].

Remark 4.7. — In the above definition, if some X_i is not irreducible, we replace X_i in the decomposition by one of its irreducible components and its complement, and do the same for their images in Y_i. After finitely many such steps we may assume that all X_i are irreducible, and thus all Y_i are irreducible.

Remark 4.8. — If X, Y are piecewise isomorphic, then $\dim X = \dim Y$.

Example 4.9. — Let $\Gamma \subseteq \mathbb{P}^2$ be an irreducible cuspidal cubic curve. Then Γ is piecewise isomorphic to $\mathbb{P}^1$. Indeed, we choose $X_1, X_2 \subseteq \Gamma$ as X_1 to be the singular locus (one point) and $X_2 = \Gamma \smallsetminus X_1$. Then, X_1 is isomorphic to $Y_1 = [0:1] \in \mathbb{P}^1$ and X_2 is isomorphic to $Y_2 = \mathbb{P}^2 \smallsetminus Y_1 \simeq \mathbb{A}^1$.

Lemma 4.10. — *Let X, Y be varieties and consider the following statements:*

(1) *X and Y are piecewise isomorphic;*
(2) *There are open subsets $U \subseteq X$, $V \subseteq Y$ such that*

$$U \simeq V, \quad \dim X \smallsetminus U = \dim Y \smallsetminus V < \dim X = \dim Y$$

and $X \smallsetminus U$, $Y \smallsetminus V$ are piecewise isomorphic. Moreover, if $X' \subseteq X$, $Y' \subseteq Y$ are locally closed subsets such that

$$\dim X \smallsetminus X', \dim Y \smallsetminus Y' < \dim X = \dim Y,$$

we may assume that $U \subseteq X'$ and $V \subseteq Y'$.

(3) $[X] = [Y]$ *inside* $K_0(\mathrm{Var}_{\mathbf{k}})$*;*

(4) $\chi(X) = \chi(Y)$.

Then we have (1) $\Leftrightarrow$ (2) $\Rightarrow$ (3) $\Rightarrow$ (4).

Proof. — "(1) $\Rightarrow$ (2)": Let $X_1, \dots X_n$ and $Y_1, \dots Y_n$ be disjoint locally closed subsets of X and Y, respectively, such that their union is equal to X and Y, respectively, and such that there are isomorphisms $\psi_i \colon X_i \xrightarrow{\simeq} Y_i$ for $1 \leqslant i \leqslant n$. By replacing X_i with $X_i \cap X'$ and $X_i \smallsetminus X'$ and analogously for Y_i we may assume that each X_i is either contained in X' or $X \smallsetminus X'$ and Y_i is either contained in Y' or $Y \smallsetminus Y'$. Using Remark 4.7, we may moreover assume that all X_i and Y_i are irreducible.

Let $d \coloneqq \dim X = \dim Y$. After reordering, we may assume that the closures $\overline{X_1}, \dots, \overline{X_m}$ in X are the irreducible components of X of dimension d. Then $\overline{Y_1}, \dots, \overline{Y_m}$ are irreducible closed subsets of Y of dimension d and since $Y_1, \dots, Y_m$ are disjoint and locally closed in Y, $\overline{Y_1}, \dots, \overline{Y_m}$ are mutually different. This implies that $\overline{Y}_1, \dots, \overline{Y}_m$ are the irreducible components of Y of dimension d. For $1 \leqslant i \leqslant m$, note that X_i and Y_i is contained in X' and Y', respectively, since $X \smallsetminus X'$ and $Y \smallsetminus Y'$ have strictly lower dimension that X and Y.

For each $i \in \{1, \dots, m\}$, the locally closed subset $X_i \subseteq X$ is equal to $X_i = A \cap \overline{X_i}$, where $A \subseteq X_i$ is open. Choosing $U_i' \subseteq X$ to be the open subset $A \smallsetminus I$, where I is the union of all irreducible components of X different from $\overline{X_i}$, we find a subset $U_i' \subseteq X_i$ that is open in X with $\overline{U_i'} = \overline{X}_i$. By construction, $U_1', \dots, U_m'$ are disjoint. Similarly, there exist open disjoint subsets $V_1', \dots, V_m'$ in Y such that for all $1 \leqslant i \leqslant m$ we get $V_i' \subseteq Y_i'$ and $\overline{V_i'} = \overline{Y_i}$. For $1 \leqslant i \leqslant m$, let

$$U_i \coloneqq U_i' \cap \psi_i^{-1}(V_i') \subseteq X \quad \text{and} \quad V_i \coloneqq \psi_i(U_i') \cap V_i' \subseteq Y.$$

Then the restriction $\psi_i|_{U_i} \colon U_i \xrightarrow{\simeq} V_i$ is an isomorphism. Let $U \coloneqq \bigcup_{i=1}^m U_i$ and $V \coloneqq \bigcup_{i=1}^m V_i$. By construction, $U \subseteq X'$ and $V \subseteq Y'$. Since $U_1, \dots, U_m$ are disjoint open subsets of X and $V_1, \dots, V_m$ are disjoint open subsets of Y, we have $U \simeq V$. Moreover, $X \smallsetminus U$ and $Y \smallsetminus V$ are piecewise isomorphic (take the locally closed subsets $X_1 \smallsetminus U_1, \dots, X_m \smallsetminus U_m, X_{m+1}, \dots, X_n$ and $Y_1 \smallsetminus V_1, \dots, Y_m \smallsetminus V_m, Y_{m+1}, \dots, Y_n$ of X and Y, respectively). By construction, $\dim X \smallsetminus U < \dim X$.

"(2) $\Rightarrow$ (1)": The decomposition of $X \smallsetminus U$, together with U, gives the decomposition of X, and we do similarly for Y.

"(2) $\Rightarrow$ (3)": By assumption we have $[U] = [V]$ inside $K_0(\mathrm{Var}_{\mathbf{k}})$. As $X \smallsetminus U$, $Y \smallsetminus V$ are piecewise isomorphic and since $\dim X \smallsetminus U = \dim Y \smallsetminus V < \dim X = \dim Y$, we may proceed by induction on d in order to get $[X \smallsetminus U] = [Y \smallsetminus V]$ inside $K_0(\mathrm{Var}_{\mathbf{k}})$. This implies (3).

"(3) $\Rightarrow$ (4)": Follows from Lemma 4.2. □

Corollary 4.11. — *If X and Y are two varieties that are both equidimensional and piecewise isomorphic, then X and Y are birational to each other.*

Proof. — Follows from the implication (1) $\Rightarrow$ (2) in Lemma 4.10. □

Example 4.12. — Corollary 4.11 really needs both X and Y to be equidimensional. Indeed, $\mathbb{A}^1 \amalg \{\mathrm{pt}\}$ and $\mathbb{P}^1$ are piecewise isomorphic, but not birational.

Lemma 4.13. — *For $i \in \{1,2\}$, let Γ_i be a variety of dimension 1, and let $\Gamma_i' \subseteq \Gamma_i$ be the union of the irreducible components of Γ_i of dimension 1 (we remove isolated points). Then, the following are equivalent:*

(1) *Γ_1 and Γ_2 are piecewise isomorphic;*
(2) *Γ_1' and Γ_2' are birational, and $\chi(\Gamma_1) = \chi(\Gamma_2)$.*

Proof. — "(1) $\Rightarrow$ (2)": Suppose that Γ_1 and Γ_2 are piecewise isomorphic. The implication (1) $\Rightarrow$ (2) in Lemma 4.10 implies that Γ_1' and Γ_2' are birational and the implication (1) $\Rightarrow$ (4) yields $\chi(\Gamma_1) = \chi(\Gamma_2)$.

"(2) $\Rightarrow$ (1)": As Γ_1' and Γ_2' are birational, there are open dense subsets $U_1 \subseteq \Gamma_1'$ and $U_2 \subseteq \Gamma_2'$ such that $U_1 \simeq U_2$. For each $i \in \{1,2\}$, the set U_i is open in Γ_i, and $F_i = \Gamma_i \smallsetminus U_i$ is finite. As $\chi(\Gamma_1) = \chi(\Gamma_2)$ and $\chi(U_1) = \chi(U_2)$, Lemma 4.2 implies that $\chi(F_1) = \chi(F_2)$. The two finite sets F_1 and F_2 are thus isomorphic, which proves that Γ_1 and Γ_2 are piecewise isomorphic. □

Lemma 4.14

(1) *Let X be a disjoint union of trees of smooth projective rational curves, with r irreducible components and s connected components. Then, X is piecewise isomorphic to the disjoint union of r copies of $\mathbb{A}^1$ and s points.*

(2) *Let X, Y be two disjoint unions of trees of smooth projective rational curves. Then, X and Y are piecewise isomorphic if and only if they have the same number of irreducible components and the same number of connected components.*

Proof

(1) If T is a tree of smooth projective rational curves, with r irreducible components, it is piecewise isomorphic to the disjoint union of one point and r copies of $\mathbb{A}^1$. Hence, if X has r irreducible components and s connected components, then X is piecewise isomorphic to the disjoint union of r copies of $\mathbb{A}^1$ and s points, and thus satisfies $\chi(X) = r + s$ (Lemma 4.2 and Example 4.4)

(2) By Lemma 4.13, X and Y are piecewise isomorphic if and only if they are birational and $\chi(X) = \chi(Y)$. The first assertion is equivalent to ask that X and Y have the same number of irreducible components. The above calculation shows that the two assertions are equivalent to ask that X and Y have the same number of irreducible components and the same number of connected components. □

Lemma 4.15. — *Let U be a smooth surface. For $i = 1, 2$, let*

$$X_i \coloneqq U \amalg S_i \amalg R_i,$$

where S_i is a disjoint union of $s_i \geqslant 0$ points and R_i is the disjoint union of $r_i \geqslant 0$ irreducible curves. If X_1 and X_2 are piecewise isomorphic, then:

(1) *R_1 and R_2 are birational; in particular $r_1 = r_2$.*
(2) *If moreover all irreducible components of R_1, R_2 are isomorphic, then $s_1 = s_2$.*

Proof. — By Lemma 4.10 there exist open dense subsets $U_1, U_2 \subseteq U$ that are isomorphic and such that $E_1 \coloneqq X_1 \smallsetminus U_1$ and $E_2 \coloneqq X_2 \smallsetminus U_2$ are piecewise isomorphic.

For any one-dimensional variety Z denote by Z' the union of irreducible components of dimension one of Z. The isomorphism $U_1 \xrightarrow{\simeq} U_2$ gives a birational map $\overline{U} \dashrightarrow \overline{U}$ of a smooth completion $\overline{U}$ of U that decomposes into n blow-ups and n blow-downs. Hence, $(\overline{U} \smallsetminus U_1)'$ and $(\overline{U} \smallsetminus U_2)'$ are birational and thus $(U \smallsetminus U_1)'$ and $(U \smallsetminus U_2)'$ are birational as well (by using that $(U \smallsetminus U_i)' \cup (\overline{U} \smallsetminus U)'$ is dense in $(\overline{U} \smallsetminus U_i)'$). By Lemma 4.13, the unions of one-dimensional irreducible components of E_1 and E_2 are birational and since $E_i = (U \smallsetminus U_i) \amalg S_i \amalg R_i$ it follows that R_1 and R_2 are birational. If the irreducible components of R_1 and R_2 are all isomorphic to C, we get $\chi(E_i) = \chi(U) - \chi(U_i) + s_i + r_i\chi(C)$ and since $\chi(E_1) = \chi(E_2)$, $\chi(U_1) = \chi(U_2)$ we get $s_1 = s_2$. □

Example 4.16. — We show that Lemma 4.13 does not generalize to higher dimension.

The irreducible surfaces $X = \mathbb{A}^2$ and $Y = \mathbb{P}^2 \smallsetminus \{[1:0:0], [0:1:0]\}$ are birational and satisfy $\chi(X) = \chi(Y) = 1$ (Lemma 4.2 and Example 4.4), but X and Y are not piecewise isomorphic. Indeed, this last assertion follows from Lemma 4.15 and the fact that X is piecewise isomorphic to $\mathbb{A}^2 \smallsetminus \{(0,0)\} \amalg \{pt\}$ and Y is piecewise isomorphic to $\mathbb{A}^2 \smallsetminus \{(0,0)\} \amalg \mathbb{A}^1$.

Proposition 4.17. — *Let X, Y be two varieties, that are piecewise isomorphic and let $E \subseteq X$, $F \subseteq Y$ be locally closed subvarieties of dimension at most 1. If E and F are piecewise isomorphic, then $X \smallsetminus E$ and $Y \smallsetminus F$ are piecewise isomorphic.*

Proof. — By definition, X and Y can be decomposed as disjoint unions of $n \geqslant 1$ locally closed subsets $X_1, \dots, X_n$ and $Y_1, \dots, Y_n$, respectively, such that there is an isomorphism $\varphi_i \colon X_i \xrightarrow{\simeq} Y_i$ for each $i \in \{1, \dots, n\}$.

(A) We first consider the case where E is one point (and thus F is also one point). In this case, there is exactly one $i \in \{1, \dots, n\}$ and one $j \in \{1, \dots, n\}$ such that $E \subseteq X_i$ and $F \subseteq Y_j$. If $i = j$, the isomorphism φ_i induces an isomorphism from $X_i \smallsetminus (E \cup \varphi_i^{-1}(F))$ to $Y_i \smallsetminus (F \cup \varphi_i(E))$, and we are done if $\varphi_i(E) = F$ or otherwise use an isomorphism from the point $\varphi_i^{-1}(F)$ to the point $\varphi_i(E)$. If $i \neq j$, then φ_i gives an isomorphisms $X_i \smallsetminus E \xrightarrow{\simeq} Y_i \smallsetminus \varphi_i(E)$ and φ_j gives an isomorphism $X_j \smallsetminus \varphi_j^{-1}(F) \xrightarrow{\simeq} Y_j \smallsetminus F$. It then remains to use an isomorphism from the point $\varphi_j^{-1}(F)$ to the point $\varphi_i(E)$.

(B) If E (and thus F) is finite, we proceed by induction on the number of points, applying (A), and obtain the result.

(C) In the general case, we proceed by induction on the number of irreducible components of E of dimension 1. When no such component exists, we are done by (B). As E and F are piecewise isomorphic, there are open subsets $U' \subseteq E$ and $V' \subseteq F$ such that $E \smallsetminus U'$ and $F \smallsetminus V'$ are finite, with the same number of points, and there is an isomorphism $\psi \colon U' \xrightarrow{\simeq} V'$ (Lemma 4.10). We take an open subset $U \subseteq U'$, that is irreducible and infinite, take $V = \psi(U) \subseteq V'$, which is also open, irreducible

and infinite, and obtain that $E \smallsetminus U$ and $F \smallsetminus V$ are piecewise isomorphic. As U, V are irreducible curves, there is exactly one $i \in \{1, \dots, n\}$ and one $j \in \{1, \dots, n\}$ such that $U \cap X_i$ and $V \cap Y_j$ are infinite. By removing finitely many points of U and V respectively, we may assume that $U \subseteq X_i$, $V \subseteq Y_j$ and still assume that U, V are isomorphic (we remove points outside of X_i or Y_j and their images under ψ or ψ^{-1}), and that $E \smallsetminus U$, $F \smallsetminus V$ are piecewise isomorphic. We now prove that $X \smallsetminus U$ and $Y \smallsetminus V$ are piecewise isomorphic, which will achieve the proof, as $E \smallsetminus U$ and $F \smallsetminus V$ are piecewise isomorphic, with one irreducible component of dimension 1 less.

If $i = j$, the isomorphism φ_i restricts to an isomorphism

$$X_i \smallsetminus (U \cup \varphi_i^{-1}(V)) \xrightarrow{\simeq} Y_i \smallsetminus (V \cup \varphi_i(U)),$$

so we only need to see that $\varphi_i^{-1}(V) \smallsetminus (U \cap \varphi_i^{-1}(V))$ and $\varphi_i(U) \smallsetminus (V \cap \varphi_i(U))$ are piecewise isomorphic. Applying φ_i to the first set, we need to show that $V \smallsetminus (\varphi_i(U) \cap V)$ and $\varphi_i(U) \smallsetminus (V \cap \varphi_i(U))$ are piecewise isomorphic. If $\varphi_i(U) \cap V$ is finite, this follows from (B). If $\varphi_i(U) \cap V$ is infinite, then $V \smallsetminus (\varphi_i(U) \cap V)$ and $\varphi_i(U) \smallsetminus (V \cap \varphi_i(U))$ are both finite, with the same number of points by using the topological Euler characteristic (Lemma 4.2).

If $i \neq j$, then φ_i gives an isomorphism $X_i \smallsetminus U \xrightarrow{\simeq} Y_i \smallsetminus \varphi_i(U)$ and φ_j gives an isomorphism $X_j \smallsetminus \varphi_j^{-1}(V) \xrightarrow{\simeq} Y_j \smallsetminus V$. It then remains to use an isomorphism from the curve $\varphi_j^{-1}(V)$ to the curve $\varphi_i(U)$. □

Lemma 4.18. — *Let X be a smooth projective rational surface with Picard group $\mathrm{Pic}(X) \simeq \mathbb{Z}^n$. Then, there is an open subset $U \subseteq X$ isomorphic to $\mathbb{A}^2$, a point $p \in X$ and locally closed curves $C_1, \dots, C_n \subseteq X$ such that each C_i is isomorphic to $\mathbb{A}^1$ and X is the disjoint union*

$$X = U \amalg C_1 \amalg \cdots \amalg C_n \amalg \{p\}.$$

Moreover, for each finite set $\Delta \subseteq X$ and each closed curve $E \subseteq X$, we may choose the above decomposition such that $\Delta \subseteq U$, $\{p\} \cap (\Delta \cup E) = \varnothing$ and $C_i \not\subseteq E$ for each $i \in \{1, \dots, n\}$.

Proof. — If $X = \mathbb{P}^2$, we choose a general line $\ell \subseteq \mathbb{P}^2$ and a general point $p \in \ell$, and let $U = \mathbb{P}^2 \smallsetminus \ell$, $C_1 = \ell \smallsetminus \{p\}$. This gives the statement in case $X = \mathbb{P}^2$.

We may thus assume that X is not isomorphic to $\mathbb{P}^2$, so there is a birational morphism $X \to \mathbb{F}_d$ for some Hirzebruch surface $\mathbb{F}_d$ where $d \geqslant 0$. We proceed by induction on the number of blow-ups of $X \to \mathbb{F}_d$. If $X = \mathbb{F}_d$, we choose a general section ℓ_1 of $\mathbb{F}_d \to \mathbb{P}^1$ of self-intersection d and a general fiber ℓ_2 of $\mathbb{F}_d \to \mathbb{P}^1$ and let $p \coloneqq \ell_1 \cap \ell_2$, $U \coloneqq \mathbb{F}_d \smallsetminus (\ell_1 \cup \ell_2)$. Then, $C_i = \ell_i \smallsetminus \{p\} \simeq \mathbb{A}^1$ for each $i \in \{1, 2\}$.

We then do successive blow-ups, where we may always assume that the points blown-up are in the subset $U \simeq \mathbb{A}^2$. Then, taking a general line of $\mathbb{A}^2$ passing through the point, the strict transform of the line is a closed curve $C_i \simeq \mathbb{A}^1$ of the blow-up of $\mathbb{A}^2$, with complement isomorphic to $\mathbb{A}^2$. □

Proposition 4.19. — *Let Y be a normal projective rational surface, that admits a desingularization $X \to Y$ with $\mathrm{Pic}(X) \simeq \mathbb{Z}^n$ and exceptional divisor being a finite disjoint union of trees of smooth rational curves with $r \geqslant 0$ irreducible components (this holds for instance when Y only has Du Val singularities). Then, Y is piecewise isomorphic to the disjoint union of $\mathbb{A}^2$, one point, and $n-r$ copies of $\mathbb{A}^1$, with $n-r \geqslant 1$.*

Proof. — If Y is smooth, then the result, with $r = 0$, directly follows from Lemma 4.18. We then assume that Y is singular. Let $E \subseteq X$ be the exceptional divisor of the desingularization $X \to Y$, let r be the number of irreducible components of E and let s be the number of connected components of E. Then s is equal to the number of points of the image of E under $X \to Y$, since all fibres of $X \to Y$ are connected by the normality of Y, see Zariski's main theorem, [Har77, Ch. III, Cor. 11.4]. Moreover, $r, s \geqslant 1$, since otherwise every fiber of the proper birational morphism $X \to Y$ onto the normal variety Y would be a single point, i.e., $X \to Y$ would be an isomorphism, see [Gro61, Cor. 4.4.9].

We observe that $n - r \geqslant 1$. Indeed, the r irreducible components $E_1, \dots, E_r$ of E, together with the pull-back P of an irreducible curve of Y not passing through the image of E under $Y \to X$ are $r+1$ effective divisors on X, that are linearly independent: Assume $D \coloneqq aP + \sum_{i=1}^{r} a_i E_i = 0$ in $\mathrm{Pic}(X)$ for some $a, a_1, \dots, a_r \in \mathbb{Z}$. Then $0 = L \cdot D = aL^2$, which shows that $a = 0$. Since the intersection matrix $(E_i \cdot E_j)$ is negative definite (see [Art62, §2, (2.1)]), we get $a_1 = \cdots = a_r = 0$.

By Lemma 4.141, E is piecewise isomorphic to the disjoint union of r copies of $\mathbb{A}^1$ and s points. By Lemma 4.18, X is piecewise isomorphic to the disjoint union of $\mathbb{A}^2$ with one point and n copies of $\mathbb{A}^1$.

Hence, Proposition 4.17 shows that $X \smallsetminus E$ is piecewise isomorphic to the disjoint union of $\mathbb{A}^2 \smallsetminus \Delta$ with $n - r$ copies of $\mathbb{A}^1$, where Δ is a finite set of $s - 1$ points.

Let S be the image of E under $X \to Y$. As $Y \smallsetminus S$ is isomorphic to $X \smallsetminus E$ and S contains s points, this proves that Y is piecewise isomorphic to the disjoint union of $\mathbb{A}^2$ with one point and $n - r$ copies of $\mathbb{A}^1$. □

Example 4.20. — Let $f, g \in \mathbf{k}[x, y, z] \smallsetminus \{0\}$ be homogeneous polynomials of degree 3 and 4, respectively, without common factor. Then, $V_{\mathbb{P}^2}(f, g)$ consists of r points, with $1 \leqslant r \leqslant 12$. We define $X \subseteq \mathbb{P}^3$ to be the quartic rational surface given by

$$X = \{[w : x : y : z] \in \mathbb{P}^3 \mid wf(x, y, z) = g(x, y, z)\},$$

and consider the open subset $X_f \subseteq X$ where f is non-zero and its complement $F \coloneqq X \smallsetminus X_f = V_{\mathbb{P}^3}(f, g)$. The hypersurface X is normal, since the singular locus is finite, see e.g. [Har77, Ch. II, Prop. 8.23]. We have an isomorphism

$$X_f \xrightarrow{\simeq} \mathbb{P}^2_f, \quad [w : x : y : z] \longmapsto [x : y : z],$$

and F is the union of r lines through $q = [1 : 0 : 0 : 0]$. Hence, X is piecewise isomorphic to the disjoint union of $\mathbb{P}^2_f$ with one point and r copies of $\mathbb{A}^1$. We now distinguish different cases for the zero locus $\Gamma \coloneqq V_{\mathbb{P}^2}(f) \subseteq \mathbb{P}^2$.

If Γ consists of one, two or three lines meeting in one point, then X is piecewise isomorphic to the disjoint union of $\mathbb{A}^2$ with one point and s copies of $\mathbb{A}^1$, for some $s \leqslant r$, similarly as in Proposition 4.19. The same holds if Γ is the union of a conic and a line, meeting at only one point.

If Γ consists of three general lines, then X is piecewise isomorphic to the disjoint union of $\mathbb{A}^2$ with *two* points and $r-2$ copies of $\mathbb{A}^1$. It is then *not* piecewise isomorphic to a smooth projective rational surface. Indeed, this follows from Lemma 4.15, since a smooth projective rational surface is piecewise isomorphic to a disjoint union of $\mathbb{A}^2$, *one* point and some copies of $\mathbb{A}^1$, see Lemma 4.18. A similar result holds if Γ is the union of a conic and a line meeting at two points.

If Γ is a cuspidal cubic curve, it is piecewise isomorphic to a line, so $\mathbb{P}^2_f$ is piecewise isomorphic to $\mathbb{A}^2$ (Proposition 4.17). Hence, this case is similar to the one of a line. If Γ is a nodal cubic, it is piecewise isomorphic to $\mathbb{A}^1 \smallsetminus \{0\}$ and thus X is piecewise isomorphic to the disjoint union of $\mathbb{A}^2$ with *two* points and r copies of $\mathbb{A}^1$, and thus X is again not piecewise isomorphic to a smooth rational projective surface.

If Γ is a smooth irreducible cubic, this one is not rational. Let $\mathbb{P}^1 \subseteq \mathbb{P}^2$ be a line that intersects Γ in exactly one point $p \in \mathbb{P}^2$ and let $\Gamma_0 \coloneqq \Gamma \smallsetminus p$. As X is piecewise isomorphic to $\mathbb{A}^2 \smallsetminus \Gamma_0 \amalg \{pt\} \amalg \coprod_{i=1}^{r+1} \mathbb{A}^1$, it follows that X is not piecewise isomorphic to a smooth rational projective surface Y, since such a Y is piecewise isomorphic to $\mathbb{A}^2 \smallsetminus \Gamma_0 \amalg \{pt\} \amalg \Gamma_0 \amalg \coprod_{i=1}^{s} \mathbb{A}^1$ for some $s \geqslant 1$, see Lemma 4.15.

Example 4.21. — In Example 4.20, we may choose f to obtain a singular normal rational quartic $X \subseteq \mathbb{P}^3$ which is not piecewise isomorphic to a smooth rational projective surface, and thus which does not admit a desingularization by trees of smooth rational curves (follows from Proposition 4.19).

We now choose $f = xy(x+y)$ and $g \in \mathbf{k}[x,y,z]_4$ general, and check that this gives an example of a normal rational quartic

$$X = \{[w : x : y : z] \in \mathbb{P}^3 \mid wf(x,y,z) = g(x,y,z)\},$$

which admits a desingularization by trees of smooth rational curves, but that does not have Du Val singularities. For this, we consider the birational map

$$\psi \colon \mathbb{P}^2 \dashrightarrow X, \qquad [x : y : z] \longmapsto [g(x,y,z) : xf(x,y,z) : yf(x,y,z) : zf(x,y,z)],$$

and observe that it has exactly twelve base-points, being $V_{\mathbb{P}^2}(f,g)$. Denoting by $\eta \colon Y \to \mathbb{P}^2$ the blow-up of the twelve points, we obtain a birational morphism $\psi \circ \eta \colon Y \to X$, that contracts the strict transforms of the three lines defined by $V_{\mathbb{P}^2}(f)$. These curves are smooth rational curves on Y with self-intersection -3, intersecting into a common point. This tree of smooth rational curves is contracted to the singular point of multiplicity three of X.

5. Partial answers in low degree

In this chapter, we give for certain cases affirmative answers to Questions (2) and (3). Moreover, over an algebraically closed field, we show for any irreducible

hypersurface $H \subseteq \mathbb{P}^n$ of degree d that there exists an element of $\mathrm{Aut}(\mathbb{P}^3 \smallsetminus H)$ that does not extend to an automorphism of $\mathbb{P}^3$ in case $d = 2$ or $n = d = 3$ and H is singular. This gives thus a negative answer to Question (1) in the above mentioned cases.

5.1. Degree two. — The goal of this subsection is to prove Theorem B, which gives an affirmative answer to Question (2) (and thus (3)) for irreducible quadric hypersurfaces in $\mathbb{P}^n_{\mathbf{k}}$ over an algebraically closed field $\mathbf{k}$. For doing this, we count the $\mathbb{F}_q$-rational points, where $\mathbb{F}_q$ denotes the finite field with q elements.

We start with a lemma that is certainly well-known to the specialists. For lack of a reference, we insert a proof.

Lemma 5.1. — *Let $\mathbb{Z} \subseteq A$, $\mathbb{Z} \subseteq B$ be finitely generated ring extensions such that A, B are integral domains. If there exists a field K and a K-isomorphism between $K \otimes_{\mathbb{Z}} A$ and $K \otimes_{\mathbb{Z}} B$, then $\mathbb{F}_q \otimes_{\mathbb{Z}} A$ and $\mathbb{F}_q \otimes_{\mathbb{Z}} B$ are $\mathbb{F}_q$-isomorphic for some prime power q.*

Proof. — As $\mathbb{Z}$ is a principal ideal domain, it is a regular ring. Using that A, B are integral domains, we get that A, B are flat $\mathbb{Z}$-modules; see [Har77, Ch. III, Prop. 9.7]. Hence, every ring extension $R_1 \subseteq R_2$ induces injections

$$R_1 \otimes_{\mathbb{Z}} A \subseteq R_2 \otimes_{\mathbb{Z}} A \quad \text{and} \quad R_1 \otimes_{\mathbb{Z}} B \subseteq R_2 \otimes_{\mathbb{Z}} B.$$

Let $a_1, \dots, a_n \in A$ with $A = \mathbb{Z}[a_1, \dots, a_n]$ and $b_1, \dots, b_m \in B$ with $B = \mathbb{Z}[b_1, \dots, b_m]$. Denote by $\varphi\colon K \otimes_{\mathbb{Z}} A \xrightarrow{\sim} K \otimes_{\mathbb{Z}} B$ a K-isomorphism. There exist finitely many $c_1, \dots, c_r \in K$ such that the $\mathbb{Z}$-algebra R spanned by $c_1, \dots, c_r$ inside K satisfies $\varphi(a_i) \in R \otimes_{\mathbb{Z}} B$ for all i with $1 \leqslant i \leqslant n$ and $\varphi^{-1}(b_j) \in R \otimes_{\mathbb{Z}} A$ for all j with $1 \leqslant j \leqslant m$. In particular, φ restricts to an R-isomorphism $R \otimes_{\mathbb{Z}} A \simeq R \otimes_{\mathbb{Z}} B$. We may assume that R is non-zero and thus contains a maximal ideal $\mathfrak{m}$. Then we get an $R/\mathfrak{m}$-isomorphism

$$(R/\mathfrak{m}) \otimes_{\mathbb{Z}} A \xrightarrow{\sim} (R/\mathfrak{m}) \otimes_{\mathbb{Z}} B.$$

Denote by F the prime field of $R/\mathfrak{m}$, i.e., the quotient field of the image of the unique ring homomorphism $\mathbb{Z} \to R/\mathfrak{m}$. As R is finitely generated as a $\mathbb{Z}$-algebra, $R/\mathfrak{m}$ is finitely generated as an F-algebra and by Noether's normalization theorem, the field extension $F \subseteq R/\mathfrak{m}$ is finite. By the Artin-Tate lemma (see [AM69, Prop. 7.8]) it follows that F is a finitely generated $\mathbb{Z}$-algebra. Hence, F cannot be isomorphic to $\mathbb{Q}$ and thus $F = \mathbb{F}_p$ where $p = \mathrm{char}(R/\mathfrak{m})$. As the field extension $F \subseteq R/\mathfrak{m}$ is finite, $R/\mathfrak{m} = \mathbb{F}_{p^r}$ for some $r \geqslant 1$. □

Corollary 5.2. — *Let X, Y be affine integral schemes of finite type over $\mathbb{Z}$ such that $X \to \mathrm{Spec}(\mathbb{Z})$, $Y \to \mathrm{Spec}(\mathbb{Z})$ are dominant. If the pull-backs X_K, Y_K are K-isomorphic for some field K, then there exists a prime power q and a bijection between the $\mathbb{F}_q$-rational points $X(\mathbb{F}_q) \to Y(\mathbb{F}_q)$ that maps the regular $\mathbb{F}_q$-rational points of X bijectively onto the regular $\mathbb{F}_q$-rational points of Y.*

Proof. — This is a direct consequence of Lemma 5.1. □

In the following lemmas, we denote by $Z(\mathbb{F}_q)$ ($Z(\mathbb{F}_q)_{\mathrm{reg}}, Z(\mathbb{F}_q)_{\mathrm{sing}}$) the set of (regular, singular) $\mathbb{F}_q$-rational points of a scheme Z that is defined over $\mathbb{Z}$. Note that $Z(\mathbb{F}_q)_{\mathrm{reg}} = Z(\mathbb{F}_q) \smallsetminus Z(\mathbb{F}_q)_{\mathrm{sing}}$.

In case Z is a hypersurface in $\mathbb{A}^{n+1}_{\mathbb{Z}}$ given by a polynomial $f \in \mathbb{Z}[x_0, \dots, x_n]$, the set $Z(\mathbb{F}_q)$ consists of those points in $(a_0, \dots, a_n) \in \mathbb{F}_q^{n+1}$ that satisfy $f(a_0, \dots, a_n) = 0$ and $(a_0, \dots, a_n) \in Z(\mathbb{F}_q)$ is regular, if not all the partial derivatives $\partial f/\partial x_0, \dots, \partial f/\partial x_n \in \mathbb{Z}[x_0, \dots, x_n]$ vanish at $(a_0, \dots, a_n)$.

Lemma 5.3. — *For all integers m, n with $1 \leqslant 2m-1 \leqslant n$, let $X_{m,n}$ be the hypersurface in $\mathbb{A}^{n+1}_{\mathbb{Z}}$ given by*

$$\sum_{i=0}^{m-1} x_{2i}x_{2i+1} = 1.$$

Then for any prime power q we get

$$\#X_{m,n}(\mathbb{F}_q) = \#X_{m,n}(\mathbb{F}_q)_{\mathrm{reg}} = q^{n-m}(q^m - 1).$$

Proof. — As $X_{m,n}$ is smooth over $\mathrm{Spec}(\mathbb{Z})$, $X_{m,n}(\mathbb{F}_q)_{\mathrm{reg}} = X_{m,n}(\mathbb{F}_q)$. We proceed now by induction on $m \geqslant 1$. For any $n \geqslant 1$, we get $\#X_{1,n}(\mathbb{F}_q) = q^{n-1}(q-1)$, as $X_{1,n}$ and $(\mathbb{A}^1_{\mathbb{Z}} \smallsetminus \{0\}) \times \mathbb{A}^{n-1}_{\mathbb{Z}}$ are isomorphic over $\mathbb{Z}$. Moreover, for $m > 1$:

$$X_{m,n}(\mathbb{F}_q) = V_{X_{m,n}}(x_{2m-1})(\mathbb{F}_q) \coprod (X_{m,n})_{x_{2m-1}}(\mathbb{F}_q).$$

As $(X_{m,n})_{x_{2m-1}}$ and $(\mathbb{A}^1_{\mathbb{Z}} \smallsetminus \{0\}) \times \mathbb{A}^{n-1}_{\mathbb{Z}}$ are isomorphic over $\mathbb{Z}$, it follows that

$$\#(X_{m,n})_{x_{2m-1}}(\mathbb{F}_q) = q^{n-1}(q-1).$$

As $V_{X_{m,n}}(x_{2m-1})$ and $X_{m-1,n-1}$ are isomorphic over $\mathbb{Z}$, we get by induction

$$\begin{aligned}\#X_{m,n}(\mathbb{F}_q) &= \#X_{m-1,n-1}(\mathbb{F}_q) + q^{n-1}(q-1)\\ &= q^{n-m}(q^{m-1} - 1) + q^{n-1}(q-1) = q^{n-m}(q^m - 1).\end{aligned}$$ □

Lemma 5.4. — *For all integers m, n with $0 \leqslant 2m \leqslant n$, let $Y_{m,n}$ be the hypersurface in $\mathbb{A}^{n+1}_{\mathbb{Z}}$ that is given by*

$$\left(\sum_{i=0}^{m-1} x_{2i}x_{2i+1}\right) + x_{2m}^2 = 1.$$

Then for every odd *prime power q we get*

$$\#Y_{m,n}(\mathbb{F}_q) = \#Y_{m,n}(\mathbb{F}_q)_{\mathrm{reg}} = q^{n-m}(q^m + 1),$$

and for every even *prime power q, we have*

$$\#(Y_{m,n})(\mathbb{F}_q) = q^n \quad \textit{and} \quad \#(Y_{m,n})(\mathbb{F}_q)_{\mathrm{reg}} = q^{n-2m}(q^{2m} - 1).$$

Proof. — Suppose first that q is odd. Here, $Y_{m,n}$ is smooth over $\mathrm{Spec}(\mathbb{Z}) \smallsetminus \{(2)\}$, so $Y_{m,n}(\mathbb{F}_q)_{\mathrm{reg}} = Y_{m,n}(\mathbb{F}_q)$. We proceed by induction on $m \geqslant 0$. For any $n \geqslant 1$, we get $\#Y_{0,n} = 2q^n$, as $Y_{0,n}$ is given by $x_0^2 - 1$ in $\mathbb{A}^{n+1}_{\mathbb{Z}}$ over $\mathbb{Z}$. Assume $m > 0$. As in

Lemma 5.3, we count the points where $x_{2m-1} = 0$ and the points where $x_{2m-1} \neq 0$, and obtain:

$$\begin{aligned}\#Y_{m,n}(\mathbb{F}_q) &= \#Y_{m-1,n-1}(\mathbb{F}_q) + q^{n-1}(q-1)\\ &= q^{n-m}(q^{m-1}+1) + q^{n-1}(q-1) = q^{n-m}(q^m+1).\end{aligned}$$

Suppose now that q is a power of 2. As $\mathbb{F}_q \to \mathbb{F}_q, t \to t^2$ is bijective, the projection $Y_{m,n} \to \mathbb{A}^n_{\mathbb{Z}}$, $(x_0,\dots,x_n) \mapsto (x_0,\dots,x_{2m-1},x_{2m+1},\dots,x_n)$ induces a bijection $Y_{m,n}(\mathbb{F}_q) \xrightarrow{\simeq} \mathbb{A}^n_{\mathbb{Z}}(\mathbb{F}_q)$, which yields $\#(Y_{m,n})(\mathbb{F}_q) = q^n$. The set of singular $\mathbb{F}_q$-rational points of $Y_{m,n}$ is equal to

$$(Y_{m,n})(\mathbb{F}_q)_{\mathrm{sing}} = \{(0,\dots,0,1,x_{2m+1},\dots,x_n) \mid x_{2m+1},\dots,x_n \in \mathbb{F}_q\}$$

and hence $(Y_{m,n})(\mathbb{F}_q)_{\mathrm{sing}} = q^{n-2m}$. This gives the result. □

Proposition 5.5. — *If* $\mathbf{k}$ *is algebraically closed, every irreducible quadric hypersurface in* $\mathbb{P}^n$ *is given, under a suitable change of coordinates, by*

$$\sum_{i=0}^{m-1} x_{2i}x_{2i+1} \quad \textit{or} \quad \left(\sum_{i=0}^{m-1} x_{2i}x_{2i+1}\right) + x_{2m}^2,$$

where $1 < 2m-1 \leqslant n$ *in the first case and* $0 < 2m \leqslant n$ *in the second case.*

Proof. — This follows from [Pfi95, Th. 1.8] if $\mathrm{char}(\mathbf{k}) \neq 2$ and from [Pfi95, Th. 4.3] if $\mathrm{char}(\mathbf{k}) = 2$. □

Proof of Theorem B. — Let

$$\mathcal{M} = \mathcal{M}_X \coprod \mathcal{M}_Y,$$

where

$$\mathcal{M}_X \coloneqq \{X_{m,n} \mid 1 < 2m-1 \leqslant n\} \quad \text{and} \quad \mathcal{M}_Y \coloneqq \{Y_{m,n} \mid 0 < 2m \leqslant n\}$$

and $X_{m,n}, Y_{m,n}$ are defined in Lemmas 5.3 and 5.4, respectively. Using Proposition 2.1, Remark 2.2 and Proposition 5.5, it is enough to show that distinct elements of $\mathcal{M}$ are non-isomorphic over $\mathbf{k}$. By Corollary 5.2, an isomorphism over $\mathbf{k}$ between two elements from $\mathcal{M}$ would give a bijection between the $\mathbb{F}_q$-rational points and between the regular $\mathbb{F}_q$-rational points, for some prime power q.

We consider for every prime power q the map

$$\Phi_q \colon \mathcal{M} \longrightarrow \mathbb{N}_0, \quad Z \longmapsto \#Z(\mathbb{F}_q)_{\mathrm{reg}}.$$

By Lemmas 5.3–5.4, we get

$$\Phi_q(X_{m,n}) = q^{n-m}(q^m-1) \quad \text{for all prime powers } q$$

and

$$\Phi_q(Y_{m,n}) = \begin{cases} q^{n-m}(q^m+1) & \text{if } q \text{ is an odd prime power,}\\ q^{n-2m}(q^{2m}-1) & \text{if } q \text{ is a power of 2.}\end{cases}$$

Hence, for odd prime powers q, the map Φ_q is injective. If $q = 2^r$ for some $r \geqslant 1$, then the restrictions $\Phi_{2^r}|_{\mathcal{M}_X}$ and $\Phi_{2^r}|_{\mathcal{M}_Y}$ are still injective. By the Lemmas 5.3–5.4, all

$\mathbb{F}_{2^r}$-rational points of $X_{m,n}$ are regular, whereas $Y_{m,n}$ has singular $\mathbb{F}_{2^r}$-rational points. This achieves to prove that distinct elements from $\mathcal{M}$ are non-isomorphic over $\mathbf{k}$. □

The following example shows that Question (1) has a negative answer for degree 2 and any $n \geqslant 2$:

Example 5.6. — Let $n \geqslant 2$. For each $f \in \mathbf{k}[x_3, \dots, x_n]$, homogeneous of degree 2 (we choose $f = 0$ if $n = 2$), the following morphism defines a $\mathbb{G}_a$-action

$$\begin{aligned} \rho\colon \mathbb{G}_a \times \mathbb{P}^n_{x_0x_1+x_2^2+f} &\longrightarrow \mathbb{P}^n_{x_0x_1+x_2^2+f} \\ (t, [x_0 : x_1 : \cdots : x_n]) &\longmapsto [x_0 : x_1 - 2tx_2 - t^2x_0 : x_2 + tx_0 : x_3 : \cdots : x_n]. \end{aligned}$$

Since $x_0^2/(x_0x_1 + x_2^2 + f)$ is a $\mathbb{G}_a$-invariant function on $\mathbb{P}^n_{x_0x_1+x_2^2+f}$, it follows that for every non-constant univariate polynomial q the following map

$$[x_0 : x_1 : \cdots : x_n] \longmapsto \rho\Big(q\Big(\frac{x_0^2}{x_0x_1 + x_2^2 + f}\Big), [x_0 : x_1 : \cdots : x_n]\Big)$$

gives an element of $\mathrm{Aut}(\mathbb{P}^n_{x_0x_1+x_2^2+f})$ that doesn't extend to an element of $\mathrm{Aut}(\mathbb{P}^n)$.

Similarly, for each $n \geqslant 3$ and each polynomial $g \in \mathbf{k}[x_4, \dots, x_n]$ (again $g = 0$ if $n = 3$), the following morphism defines a $\mathbb{G}_a$-action

$$\begin{aligned} \rho\colon \mathbb{G}_a \times \mathbb{P}^n_{x_0x_1+x_2x_3+g} &\longrightarrow \mathbb{P}^n_{x_0x_1+x_2x_3+g} \\ (t, [x_0 : x_1 : \cdots : x_n]) &\longmapsto [x_0 : x_1 - tx_3 : x_2 + tx_0 : x_3 : \cdots : x_n]. \end{aligned}$$

Since $x_0^2/(x_0x_1 + x_2x_3 + g)$ is a $\mathbb{G}_a$-invariant function on $\mathbb{P}^n_{x_0x_1+x_2x_3+g}$, it follows that for every non-constant univariate polynomial q the following map

$$[x_0 : x_1 : \cdots : x_n] \longmapsto \rho\Big(q\Big(\frac{x_0^2}{x_0x_1 + x_2x_3 + g}\Big), [x_0 : x_1 : \cdots : x_n]\Big)$$

defines an automorphism of $\mathbb{P}^n_{x_0x_1+x_2x_3+g}$ that doesn't extend to an element of $\mathrm{Aut}(\mathbb{P}^n)$.

If $\mathbf{k}$ is algebraically closed, it follows from Proposition 5.5 that every irreducible quadric is of one of the above form, up to change of coordinates.

5.2. Degree three and beyond in dimension three. — Throughout this subsection we assume always that $\mathbf{k}$ is algebraically closed.

Recall that an irreducible hypersurface in $\mathbb{P}^3$ is normal if and only if its singular locus is finite [Har77, Ch. II, Prop. 8.23].

Remark 5.7. — Let $X \subseteq \mathbb{P}^2$ be a normal rational irreducible cubic hypersurface. Then there exist six (possibly infinitely near) points $p_1, \dots, p_6$ in $\mathbb{P}^2$ such that the blow-up $\pi\colon \widetilde{X} \to \mathbb{P}^2$ of them satisfies: There exists a birational morphism $\eta\colon \widetilde{X} \to X$ and it is the minimal desingularization of X. In case X is singular this follows from the proof of Lemma 5.9 and in case X is smooth this follows from [Har77, Ch. V, Cor. 4.7, Rem. 4.7.1]. Moreover in the smooth case $p_1, \dots, p_6$ belong to $\mathbb{P}^2$ (and are in general position).

Lemma 5.8. — *An irreducible cubic hypersurface in $\mathbb{P}^3$ is non-rational if and only if it is the cone over a smooth cubic curve in $\mathbb{P}^2$.*

Proof. — Let $X \subseteq \mathbb{P}^3$ be a non-rational irreducible cubic hypersurface. Then X has a singularity (as otherwise it would be the blow-up of $\mathbb{P}^2$ at six points, see Remark 5.7), say at $P = [1 : 0 : 0 : 0] \in \mathbb{P}^3$ and it is given by a homogeneous polynomial of the form $f_2(x, y, z)w + f_3(x, y, z)$, where $[w : x : y : z]$ denote the homogeneous coordinates of $\mathbb{P}^3$ and $f_i \in \mathbf{k}[x, y, z]$ is homogeneous of degree i. As X is non-rational, we get $f_2 = 0$ (otherwise the projection to x, y, z gives a birational map to $\mathbb{P}^2$) and X is the cone in $\mathbb{P}^3$ over a cubic curve $\Gamma \subset \mathbb{P}^2$ given by $f_3 = 0$. As X is irreducible, so is Γ. Then, X is birational to $\Gamma \times \mathbb{P}^1$, and thus Γ is non-rational as X is. This implies that Γ is smooth, and that X is the cone in $\mathbb{P}^3$ over a smooth cubic curve in $\mathbb{P}^2$.

Assume now that X is a cone over a smooth cubic curve C. Blowing up the vertex yields a projective smooth ruled surface S over C. In particular $H^1(S, \mathcal{O}_S)$ doesn't vanish (see [Har77, Ch. V, Cor. 2.5]) and hence X is not rational (see [KSC04, Th. 3.2]). □

The next lemma is certainly known to the specialists. For lack of a proof in any characteristic we provide one:

Lemma 5.9. — *Let $X \subseteq \mathbb{P}^3$ be a normal rational cubic hypersurface. Then X has only Du Val singularities and is piecewise isomorphic to the disjoint union of $\mathbb{A}^2$, one point, and n copies of $\mathbb{A}^1$, with $1 \leqslant n \leqslant 7$. Moreover, $3 \leqslant \chi(X) = 2 + n \leqslant 9$ and $\chi(X) = 9$ if and only if X is smooth.*

Proof. — If X is smooth, then X is the blow-up of 6 points of $\mathbb{P}^2$ (Remark 5.7) and is thus piecewise isomorphic to the disjoint union of $\mathbb{A}^2$, one point and seven copies of $\mathbb{A}^1$ by Lemma 4.18. In particular, $\chi(X) = 9$.

Thus we may assume that X is singular and after some coordinate change we may assume that X is given by $f_2(x, y, z)w + f_3(x, y, z)$, where $f_2, f_3 \in \mathbf{k}[x, y, z]$ are homogeneous polynomials of degree $2, 3$, respectively, and $[w : x : y : z]$ denote the homogeneous coordinates in $\mathbb{P}^3$. If $f_2 = 0$, then, X is a cone over $V_{\mathbb{P}^2}(f_3)$. As X is a rational cubic, f_3 is irreducible and $V_{\mathbb{P}^2}(f_3)$ has to be singular (Lemma 5.8), contradicting the fact that X is normal. Hence, $f_2 \neq 0$.

The projection $X \smallsetminus \{[1 : 0 : 0 : 0]\} \to \mathbb{P}^2$, $[w : x : y : z] \mapsto [x : y : z]$ gives a birational morphism whose inverse $\Phi \colon \mathbb{P}^2 \dashrightarrow X$ is given by

$$[x : y : z] \longmapsto [f_3(x, y, z) : xf_2(x, y, z) : yf_2(x, y, z) : zf_2(x, y, z)].$$

By [Sak10, Prop. 1.5], there exists a smooth projective surface $\widetilde{X}$ which is the blow-up $\pi \colon \widetilde{X} \to \mathbb{P}^2$ of six (possibly infinitely near) points in the zero locus $S = V(f_2, f_3)$ such that $\eta \coloneqq \Phi \circ \pi \colon \widetilde{X} \to X$ is a morphism and η is given by the complete linear system $|-K_{\widetilde{X}}|$, where $K_{\widetilde{X}}$ denotes the canonical divisor in $\widetilde{X}$. Let X_{reg} be the open subvariety of regular points in X and let X_{sing} be its complement in X.

By applying the adjunction formula to the smooth closed hypersurface X_{reg} in $\mathbb{P}^3 \smallsetminus X_{\mathrm{sing}}$, we get that the canonical divisor K_X of X is equal to $-H$, where H is a

hyperplane section of $X \subseteq \mathbb{P}^3$. This gives $\eta^* K_X = K_{\widetilde{X}}$. By [Art62, Th. 2.7] it follows that the exception locus of η is a finite union of trees and that in fact X has only Du Val singularities.

By Proposition 4.19, X is piecewise isomorphic to a disjoint union of $\mathbb{A}^2$, one point and $7-r$ copies of $\mathbb{A}^1$, with $1 \leqslant r \leqslant 6$. In particular, $3 \leqslant \chi(X) \leqslant 8$. □

We will use the following classification of non-normal irreducible cubic surfaces in $\mathbb{P}^3$:

Proposition 5.10 (see [LPS11, Th. 3.1]). — *Let $X \subseteq \mathbb{P}^3$ be a non-normal irreducible cubic surface. Denote by $[w : x : y : z]$ the homogeneous coordinates in $\mathbb{P}^3$.*

– If X is a cone over a curve in $\mathbb{P}^2$, then up to a coordinate change of $\mathbb{P}^3$, the hypersurface X is given by

$$(*) \qquad x^2w + y^3 \quad or \quad x^2w + y^3 + xy^2 \quad or \quad xyw + x^3 + y^3.$$

Moreover, if $\operatorname{char}(\mathbf{k}) \neq 3$, then $x^2w + y^3$ and $x^2w + y^3 + xy^2$ are the same up to a linear coordinate change in w, x, y.

– If X is not a cone over a curve in $\mathbb{P}^2$, then up to a coordinate change of $\mathbb{P}^3$, the hypersurface X is given by

$$(**) \qquad x^2w + y^2z \quad or \quad xyw + y^2z + x^3 \quad or \quad xyw + (x^2+y^2)z.$$

Moreover, if $\operatorname{char}(\mathbf{k}) \neq 2$, then $x^2w + y^2z$ and $xyw + (x^2+y^2)z$ are the same up to a linear coordinate change in w, x, y, z.

In particular, X always contains a line with multiplicity 2 (the line given by $x = y = 0$ in the above equations).

In the following lemma, we give a similar decompositions into locally closed subsets as in Lemma 5.9 for irreducible non-normal cubic hypersurfaces in $\mathbb{P}^3$:

Lemma 5.11. — *Let*

$$f_1 = xyw + x^3 + y^3, \quad f_2 = x^2w + y^3, \quad f_3 = x^2w + y^3 + xy^2,$$
$$f_4 = xyw + y^2z + x^3, \quad f_5 = x^2w + y^2z, \quad f_6 = xyw + (x^2+y^2)z.$$

Then, $X_i = V_{\mathbb{P}^3}(f_i) \subseteq \mathbb{P}^3$ is piecewise isomorphic to the disjoint union of $\mathbb{A}^2$, one point, and n_i copies of $\mathbb{A}^1$, with $n_1 = 0$, $n_2 = n_3 = n_4 = 1$ and $n_5 = n_6 = 2$. In particular, $\chi(X_i) = 2 + n_i \in \{2, 3, 4\}$.

Proof. — We can decompose X_i into $X_i \cap \mathbb{P}^3_x$ and $X_i \cap V_{\mathbb{P}^3}(x)$. Then we get Table 1, where $\mathbb{P}^1 \vee \mathbb{P}^1$ denotes two copies of $\mathbb{P}^1$ in $\mathbb{P}^2$ (that intersect transversally in exactly one point). This implies the statement. □

Corollary 5.12. — *Let X be a rational cubic hypersurface of $\mathbb{P}^3$. Then, X is piecewise isomorphic to the disjoint union of $\mathbb{A}^2$, one point, and n copies of $\mathbb{A}^1$, with $0 \leqslant n \leqslant 7$. Moreover, $2 \leqslant \chi(X) = 2 + n \leqslant 9$.*

Proof. — If X is normal, this follows from Lemma 5.9. Otherwise, this follows from Lemma 5.11 in combination with Proposition 5.10. □

Table 1.

Equation	$X_i \cap \mathbb{P}^3_x$	$X_i \cap V_{\mathbb{P}^3}(x)$
$xyw + x^3 + y^3$	$\mathbb{A}^1 \times (\mathbb{A}^1 \smallsetminus \{0\})$	$\mathbb{P}^1$
$x^2w + y^3$	$\mathbb{A}^2$	$\mathbb{P}^1$
$x^2w + y^3 + xy^2$	$\mathbb{A}^2$	$\mathbb{P}^1$
$xyw + y^2z + x^3$	$\mathbb{A}^1 \times (\mathbb{A}^1 \smallsetminus \{0\})$	$\mathbb{P}^1 \vee \mathbb{P}^1$
$x^2w + y^2z$	$\mathbb{A}^2$	$\mathbb{P}^1 \vee \mathbb{P}^1$
$xyw + (x^2 + y^2)z$	$\mathbb{A}^2$	$\mathbb{P}^1 \vee \mathbb{P}^1$

Lemma 5.13. — *Let $X \subseteq \mathbb{P}^3$ be an irreducible cubic hypersurface. If X is rational, then $\chi(X) \geqslant 2$. If X is not rational, then $\chi(X) = 1$.*

Proof. — If X is rational, then Corollary 5.12 gives $\chi(X) \geqslant 2$.

Assume now that X is non-rational. By Lemma 5.8, X is the cone over a smooth cubic curve C in $\mathbb{P}^2$. Let $p \in X$ be the unique singularity. Then $X \smallsetminus \{p\}$ is a locally trivial $\mathbb{A}^1$-bundle over C (with respect to the Zariski topology). Hence $\chi(X) = 1 + \chi([X \smallsetminus p]) = 1 + \chi(C) = 1$, where the second and third equality follow from Example 4.5 and Example 4.3, respectively. □

Proposition 5.14. — *Let X_1, X_2 be normal projective rational surfaces having only singularities that can be resolved by trees of smooth rational curves (e.g. Du Val singularities). Then the following statements are equivalent*

(1) *X_1 and X_2 are piecewise isomorphic;*
(2) *$[X_1] = [X_2]$ inside $K_0(\mathrm{Var}_{\mathbf{k}})$;*
(3) *$\chi(X_1) = \chi(X_2)$.*

Proof. — The implications "(1) ⇒ (2)" and "(2) ⇒ (3)" both follow from Lemma 4.10. Proposition 4.19 says that X_i is piecewise isomorphic to the disjoint union of $\mathbb{A}^2$, one point and $s_i \geqslant 0$ copies of $\mathbb{A}^1$. If $\chi(X_1) = \chi(X_2)$, then we get $s_1 = s_2$, i.e., X_1, X_2 are piecewise isomorphic. This gives the implication "(3) ⇒ (1)". □

Corollary 5.15. — *Let $f, g \in \mathbf{k}[x, y, z, w]$ be irreducible homogeneous polynomials such that $\mathbb{P}^3_f \simeq \mathbb{P}^3_g$. Assume moreover that the zero loci $X \coloneqq V_{\mathbb{P}^3}(f)$, $Y \coloneqq V_{\mathbb{P}^3}(g)$*

(a) *are both normal, rational and each admits a desingularization by trees of smooth rational curves or*
(b) *are both of degree 3 and rational.*

Then X, Y are piecewise isomorphic.

Proof. — Since $\mathbb{P}^3_f \simeq \mathbb{P}^3_g$ we get $[X] = [Y]$ inside $K_0(\mathrm{Var}_{\mathbf{k}})$ and thus $\chi(X) = \chi(Y)$, see Lemma 4.10. In case (a), the statement follows from Proposition 5.14. So assume we are in case (b). Then the statement follows from Proposition 4.19 in combination

with Lemma 5.9 if X and Y are normal and from Proposition 5.10 in combination with Lemma 5.11 in the other case. □

We now focus on cones over smooth cubic curves, i.e., on non-rational cubic hypersurfaces of $\mathbb{P}^3$. This constitutes the last unproved part of Theorem C. To prove that for such hypersurfaces, Question (2) has an affirmative answer, we will need the following result:

Lemma 5.16. — *Let $f \in \mathbf{k}[x,y,z]$ be an irreducible homogeneous polynomial of degree 3 such that $V_{\mathbb{P}^2}(f)$ is a smooth cubic curve. Then, the surface*

$$X = V_{\mathbb{A}^3}(f-1)$$

admits no non-trivial $\mathbb{G}_a$-action.

Proof. — Suppose for contradiction that X admits a non-trivial $\mathbb{G}_a$-action. Applying [BFT23, Prop. 2.5.1], we obtain a $\mathbb{G}_a$-invariant affine dense open subset $X' \subseteq X$, that is a $\mathbb{G}_a$-cylinder, i.e., X' is $\mathbb{G}_a$-isomorphic to $\mathbb{G}_a \times U$, where U is a smooth affine curve and where the $\mathbb{G}_a$-action on $\mathbb{G}_a \times U$ is given by $s \cdot (t,u) \coloneqq (s+t,u)$ for $s,t \in \mathbb{G}_a$, $u \in U$.

Using the canonical embedding $\mathbb{A}^3 \hookrightarrow \mathbb{P}^3, (x,y,z) \mapsto [1:x:y:z]$ we can view X as an open subset of the irreducible surface $Y = V_{\mathbb{P}^3}(f-w^3)$, where $[w:x:y:z]$ denote the homogeneous coordinates on $\mathbb{P}^3$. Here, $X = Y_w$ is the complement in Y of a smooth curve Γ and $\Gamma \simeq V_{\mathbb{P}^2}(f)$.

Suppose first that Y is a cone over a smooth cubic curve C. Writing $p \in Y$ the singular point, the projection away from p gives to $Y \smallsetminus \{p\}$ an $\mathbb{A}^1$-bundle structure over C. Hence, every closed rational curve on Y is one of the lines through p. As Γ is a hyperplane section of Y and as Γ is not rational, it intersects a general such line into at least one point outside of p. As the open subset $X' \subseteq Y \smallsetminus \Gamma$ is contained in the smooth locus of Y, it contains only finitely many curves isomorphic to $\mathbb{A}^1$. This contradicts the fact that X' is a $\mathbb{G}_a$-cylinder.

In the remaining case, Y is a rational cubic surface (Lemma 5.8). Hence, U is a smooth affine rational curve, and is thus isomorphic to $\mathbb{A}^1 \smallsetminus \Delta$ for some finite set Δ of $r \geqslant 0$ points. Hence, X' is isomorphic to the open subset $V' = \mathbb{A}^1 \times (\mathbb{A}^1 \smallsetminus \Delta) \subseteq \mathbb{A}^2$. By Corollary 5.12, Y is piecewise isomorphic to the disjoint union of $\mathbb{A}^2$, one point, and n copies of $\mathbb{A}^1$, with $0 \leqslant n \leqslant 7$, and thus to the disjoint union of $X' \simeq V'$ with one point and $n+r$ copies of $\mathbb{A}^1$. As the one-dimensional variety $Y \smallsetminus X'$ contains Γ, that is not rational, we get a contradiction, by applying Lemma 4.15, where $U \coloneqq X'$, S_1 is a point, R_1 is the disjoint union of $n+r$ copies of $\mathbb{A}^1$, and S_2 is a finite set, R_2 is a disjoint union of irreducible curves such that $Y \smallsetminus X'$ is piecewise isomorphic to $S_2 \amalg R_2$. □

We are now able to prove part 3 of Theorem C, that is Proposition 5.17. If $\mathbf{k} = \mathbb{C}$, then Proposition 5.17 follows essentially from [LS12].

Proposition 5.17. — *Let $f, g \in \mathbf{k}[x,y,z,w]$ be irreducible homogeneous polynomials of degree three such that $V_{\mathbb{P}^3}(f)$, $V_{\mathbb{P}^3}(g)$ are non-rational. If $\mathbb{P}^3_f \simeq \mathbb{P}^3_g$, then there exists $\varphi \in \mathrm{Aut}(\mathbb{P}^3)$ with $\varphi(V_{\mathbb{P}^3}(f)) = V_{\mathbb{P}^3}(g)$.*

Proof. — By Lemma 5.8 we may assume after applying automorphisms of $\mathbb{P}^3$ to f and g that $f, g \in \mathbf{k}[x,y,z]$ and that the zero loci $V_{\mathbb{P}^2}(f)$, $V_{\mathbb{P}^2}(g)$ are smooth cubic curves.

By Lemma 5.16, every $\mathbb{G}_a$-action on the affine surfaces $X_0 \coloneqq V_{\mathbb{A}^3}(f-1)$ and $Y_0 \coloneqq V_{\mathbb{A}^3}(g-1)$ is trivial.

Let $\theta\colon \mathbb{P}^3_f \to \mathbb{P}^3_g$ be an isomorphism. By Proposition 2.1 and Remark 2.2, we get an isomorphism

$$\varphi\colon \mathbb{A}^1 \times X_0 = V_{\mathbb{A}^4}(f-1) \longrightarrow V_{\mathbb{A}^4}(g-1) = \mathbb{A}^1 \times Y_0$$

such that $\pi_{g,1} \circ \varphi = \theta \circ \pi_{f,1}$, where the morphisms $\pi_{f,1}\colon V_{\mathbb{A}^4}(f-1) \to \mathbb{P}^3_f$ and $\pi_{g,1}\colon V_{\mathbb{A}^4}(g-1) \to \mathbb{P}^3_g$ denote the canonical projections.

Using that X_0 admit no non-trivial $\mathbb{G}_a$-action, by [Cra04, Prop. 4.7] it follows that the intersection of the invariant subalgebras $\mathbf{k}[\mathbb{A}^1 \times X_0]^{\mathbb{G}_a}$ inside $\mathbf{k}[\mathbb{A}^1 \times X_0]$ over all $\mathbb{G}_a$-actions on $\mathbb{A}^1 \times X_0$ is equal to $\mathbf{k}[X_0]$. Using that a similar statement holds for $Y_0 \times \mathbb{A}^1$, we get that φ maps the fibres of $\mathbb{A}^1 \times X_0 \to X_0$ onto the fibres of $\mathbb{A}^1 \times Y_0 \to Y_0$. Using the commutative diagrams

$$\begin{array}{ccc} \mathbb{A}^1 \times X_0 & \xrightarrow{\pi_{f,1}} & \mathbb{P}^3_f \\ {\scriptstyle (t,x_0)\mapsto x_0}\downarrow & & \downarrow{\scriptstyle \text{projection from } [1:0:0:0]} \\ X_0 & \longrightarrow & \mathbb{P}^2_f \end{array} \qquad \begin{array}{ccc} \mathbb{A}^1 \times Y_0 & \xrightarrow{\pi_{g,1}} & \mathbb{P}^3_g \\ {\scriptstyle (t,y_0)\mapsto y_0}\downarrow & & \downarrow{\scriptstyle \text{projection from } [1:0:0:0]} \\ Y_0 & \longrightarrow & \mathbb{P}^2_g \end{array}$$

we get that $\theta\colon \mathbb{P}^3_f \to \mathbb{P}^3_g$ maps the fibres of the locally trivial $\mathbb{A}^1$-bundle $\mathbb{P}^3_f \to \mathbb{P}^2_f$ (with respect to the Zariski topology) onto those of $\mathbb{P}^3_g \to \mathbb{P}^2_g$. Hence, we get an isomorphism $\mathbb{P}^2_f \simeq \mathbb{P}^2_g$. As $V_{\mathbb{P}^2}(f)$ and $V_{\mathbb{P}^2}(g)$ are non-rational this isomorphism extends to an automorphism of $\mathbb{P}^2$. □

Now we are able to give the proof of Theorem C:

Proof of Theorem C

(1) This follows from Corollary 5.15.

(3) If H is a non-rational cubic surface, then H' is a cubic as well (by using the order of the Picard groups of the complements, see also Lemma 2.6). Since $\chi(H) = \chi(H')$, Lemma 5.13 implies that H' is non-rational. Thus 3 follows from Proposition 5.17.

(2) If H or H' is a cubic surface, then as before, both are cubics. If either H or H' is non-rational, then 3 implies that H and H' are isomorphic, and thus also piecewise isomorphic. We may thus assume that both H and H' are rational. By Corollary 5.12 it follows that H (respectively H') is piecewise isomorphic to a disjoint union of $\mathbb{A}^2$, one point and n (respectively n') copies of $\mathbb{A}^1$. As $\mathbb{P}^3 \smallsetminus H$ and $\mathbb{P}^3 \smallsetminus H'$ are isomorphic, we get $2 + n = \chi(H) = \chi(H') = 2 + n'$ and hence H, H' are piecewise isomorphic.

If moreover H is smooth, then $\chi(H) = 9$ by Lemma 5.9. If H' would be non-normal, then Lemma 5.11 and Proposition 5.10 would imply that $\chi(H') \leqslant 4$, which contradicts $\chi(H) = \chi(H')$. Thus Lemma 5.9 implies that H' is smooth. This finishes the proof of 2. □

We finish this subsection with the following concrete question motivated by Proposition 3.4:

Question 5.18. — *Let $f, g \in \mathbf{k}[x, y, z, w]$ be given by*

$$f = x^2y + z^3 \quad \text{and} \quad g = x^2y + z^3 + xw^2.$$

Are the varieties $\mathbb{P}^3_f$ and $\mathbb{P}^3_g$ isomorphic?

An affirmative answer would give a negative answer to Question (3) in degree three and dimension three, since the zero locus of f in $\mathbb{P}^3$ is non-normal, whereas the zero locus of g in $\mathbb{P}^3$ is normal. Moreover, $\chi(V_{\mathbb{P}^3}(f)) = \chi(V_{\mathbb{P}^3}(g))$, since $\mathbb{P}^4_f \simeq \mathbb{P}^4_g$ by Proposition 3.4 and since $\mathbb{P}^4_f \to \mathbb{P}^3_f$, $\mathbb{P}^4_g \to \mathbb{P}^3_g$ are locally trivial $\mathbb{A}^1$-bundles with respect to the Zariski topology, see Example 4.5.

5.3. Automorphisms of complements of singular cubic surfaces. — Throughout this subsection we always assume that $\mathbf{k}$ is algebraically closed.

We finish this text by giving some examples, that give a negative answer to Question (1), for singular irreducible cubic hypersurfaces of $\mathbb{P}^3$. As explained in the introduction, this question is wide open for smooth cubics and known to have a negative answer for singular cubic surfaces with Du Val singularities in characteristic zero [CDP18, Th. C & Th. 4.3]. We now extend this to other singular cubics. Each irreducible cubic hypersurface of $\mathbb{P}^3$ that does not have Du Val singularities is either the cone over a smooth cubic curve in $\mathbb{P}^2$ or is rational and non-normal (this follows from Lemmas 5.8 and 5.9). In this latter case, it always contains a line with multiplicity 2 (Proposition 5.10). These two cases are done in the next two simple lemmas, that work in any dimension.

Lemma 5.19. — *Let $d \geqslant 1$, $n \geqslant 3$ and let $X \subseteq \mathbb{P}^n$ be an irreducible hypersurface of degree d, having a point of multiplicity d (i.e., being a cone). Then, there exists an element $\varphi \in \mathrm{Aut}(\mathbb{P}^n \smallsetminus X)$ that does not extend to an element of $\mathrm{Aut}(\mathbb{P}^n)$.*

Proof. — Changing coordinates, we may assume that $[0 : \cdots : 0 : 1]$ is a point of X of multiplicity d. Hence, $X = V_{\mathbb{P}^n}(f)$, where $f \in \mathbf{k}[x_0, \ldots, x_{n-1}]$ is irreducible and homogeneous of degree d. We define $\varphi \in \mathrm{Aut}(\mathbb{P}^n_f)$ to be the involution given by

$$[x_0 : \cdots : x_n] \longmapsto \Big[x_0 : \cdots : x_{n-1} : -x_n + \frac{x_0^{d+1}}{f(x_0, \ldots, x_{n-1})}\Big]. \qquad \square$$

Lemma 5.20. — *Let $d \geqslant 1$, $n \geqslant 3$ and let $X \subseteq \mathbb{P}^n$ be an irreducible hypersurface of degree d, being of multiplicity $d-1$ along a linear subspace $L \subseteq \mathbb{P}^n$ of dimension $r \in \{1, \ldots, n-2\}$. Then, there exists an element $\varphi \in \mathrm{Aut}(\mathbb{P}^n \smallsetminus X)$ that does not extend to an element of $\mathrm{Aut}(\mathbb{P}^n)$.*

Proof. — Changing coordinates, we may assume that $L = V_{\mathbb{P}^n}(x_{r+1}, \dots, x_n)$. Hence, $X = V_{\mathbb{P}^n}(f)$, where $f \in \mathbf{k}[x_0, \dots, x_n]$ is irreducible, homogeneous of degree d and of the form

$$f = x_0 a_0 + \cdots + x_r a_r + b,$$

where $a_0, \dots, a_r, b \in \mathbf{k}[x_{r+1}, \dots, x_n]$. We define a $\mathbb{G}_a$-action on $\mathbb{P}^n_f$ by

$$\begin{aligned} \mathbb{G}_a \times \mathbb{P}^n_f &\longrightarrow \mathbb{P}^n_f \\ (t, [x_0 : \cdots : x_n]) &\longmapsto \Big[x_0 + t\frac{a_1 x_n^2}{f(x_0, \dots, x_n)} : x_1 - t\frac{a_0 x_n^2}{f(x_0, \dots, x_n)} : x_2 : \cdots : x_n\Big] \end{aligned}$$

and choose $\varphi \in \mathrm{Aut}(\mathbb{P}^n_f)$ to be any non-trivial element of $\mathbb{G}_a \subseteq \mathrm{Aut}(\mathbb{P}^n_f)$. □

The following lemma follows from [CDP18, Th. C & Th. 4.3] in characteristic zero. We insert a proof (in any characteristic), as the argument is simple.

Lemma 5.21. — *Let $X \subseteq \mathbb{P}^3$ be a normal rational singular cubic hypersurface. Then $\mathbb{P}^3 \smallsetminus X$ contains an open affine subset U that is an $\mathbb{A}^1$-cylinder, i.e., $U \simeq \mathbb{A}^1 \times U'$ for some affine variety U'.*

Proof. — Since X is singular, we may assume (after a coordinate change) that X is given by $f \coloneqq f_2(x, y, z)w + f_3(x, y, z)$ where $f_i \in \mathbf{k}[x, y, z]$ is homogeneous of degree i and $[w : x : y : z]$ denote the homogeneous coordinates of $\mathbb{P}^3$. As X is normal and rational, $f_2 \neq 0$, see Lemma 5.8. Let $C \coloneqq V_{\mathbb{P}^2}(f_2)$. We may choose the coordinates (x, y, z) in such a way, that $V_{\mathbb{P}^2}(z)$ is contained in C if f_2 is reducible, and that $V_{\mathbb{P}^2}(z)$ is tangent to C if f_2 is irreducible. Then $\mathbb{P}^2_{zf_2} \simeq \mathbb{A}^2_{f_2(x,y,1)} \simeq D \times \mathbb{A}^1$, where $D \in \{\mathbb{A}^1, \mathbb{A}^1 \smallsetminus \{0\}\}$, and we get an isomorphism

$$\begin{aligned} \mathbb{P}^3_{zff_2} \simeq \mathbb{A}^3_{f(w,x,y,1)f_2(x,y,1)} &\xrightarrow{\simeq} (\mathbb{A}^1 \smallsetminus \{0\}) \times \mathbb{A}^2_{f_2(x,y,1)} \\ (w, x, y) &\longmapsto (f_2(x, y, 1)w + f_3(x, y, 1), x, y) \\ \Big(\frac{u - f_3(x, y, 1)}{f_2(x, y, 1)}, x, y\Big) &\longmapsfrom (u, x, y). \end{aligned}$$

□

Proposition 5.22. — *For each singular irreducible cubic hypersurface $X \subseteq \mathbb{P}^3$, there exists an element $\varphi \in \mathrm{Aut}(\mathbb{P}^3 \smallsetminus X)$ that does not extend to an element of* $\mathrm{Aut}(\mathbb{P}^3)$.

Proof. — If X is a cone, this follows from Lemma 5.19. If X is not normal, it contains a line with multiplicity 2 (Proposition 5.10), and the result follows from Lemma 5.20. In the remaining case, X is rational (Lemma 5.8) and normal. As $\mathrm{Pic}(\mathbb{P}^3 \smallsetminus X) \simeq \mathbb{Z}/3\mathbb{Z}$ is finite, Lemma 5.21 implies that $\mathbb{P}^3 \smallsetminus X$ admits a non-trivial $\mathbb{G}_a$-action, see e.g. [DK15, Prop. 2].

Since $\mathbb{P}^3 \smallsetminus X$ is affine and of dimension $\geqslant 2$, the Krull-dimension of the $\mathbb{G}_a$-invariant functions is $\geqslant 1$. As one may multiply the $\mathbb{G}_a$-action by any $\mathbb{G}_a$-invariant function on $\mathbb{P}^3 \smallsetminus X$, we get a faithful $\mathbb{G}_a^r$-action on $\mathbb{P}^3 \smallsetminus X$ for any $r \geqslant 1$. On the other hand, the elements of $\mathrm{Aut}(\mathbb{P}^3 \smallsetminus X)$ that extend to automorphisms of $\mathbb{P}^3$ form an algebraic group. This implies the result. □

References

[Art62] M. Artin – "Some numerical criteria for contractability of curves on algebraic surfaces", *Amer. J. Math.* **84** (1962), p. 485–496.

[AGV73] M. Artin, A. Grothendieck & J.-L. Verdier – *Théorie des topos et cohomologie étale des schémas. Tome 3*, Lect. Notes in Math., vol. 305, Springer-Verlag, Berlin-New York, 1972–1973, Séminaire de Géométrie Algébrique du Bois–Marie 1963–64 (SGA 4).

[AM69] M. F. Atiyah & I. G. Macdonald – *Introduction to commutative algebra*, Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1969.

[Bla09] J. Blanc – "The correspondence between a plane curve and its complement", *J. reine angew. Math.* **633** (2009), p. 1–10.

[BFT23] J. Blanc, A. Fanelli & R. Terpereau – "Connected algebraic groups acting on three-dimensional Mori fibrations", *Internat. Math. Res. Notices* (2023), no. 2, p. 1572–1689.

[Bor18] L. A. Borisov – "The class of the affine line is a zero divisor in the Grothendieck ring", *J. Algebraic Geom.* **27** (2018), no. 2, p. 203–209.

[CDP18] I. Cheltsov, A. Dubouloz & J. Park – "Super-rigid affine Fano varieties", *Compositio Math.* **154** (2018), no. 11, p. 2462–2484.

[CS01] P. Colmez & J.-P. Serre (eds.) – *Correspondance Grothendieck-Serre*, Documents Mathématiques, vol. 2, Société Mathématique de France, Paris, 2001.

[Cos12] P. Costa – "New distinct curves having the same complement in the projective plane", *Math. Z.* **271** (2012), no. 3-4, p. 1185–1191.

[Cra04] A. J. Crachiola – "On the AK invariant of certain domains", PhD Thesis, Wayne State University, Ann Arbor, MI, 2004, ProQuest LLC, p. 84.

[DK15] A. Dubouloz & T. Kishimoto – "Log-uniruled affine varieties without cylinder-like open subsets", *Bull. Soc. math. France* **143** (2015), no. 2, p. 383–401.

[Gro61] A. Grothendieck – "Éléments de géométrie algébrique. III. Étude cohomologique des faisceaux cohérents. I", *Publ. Math. Inst. Hautes Études Sci.* (1961), no. 11, p. 167.

[GSY05] J. Gutierrez, V. Shpilrain & J.-T. Yu (eds.) – *Affine algebraic geometry*, Contemp. Math., vol. 369, American Mathematical Society, Providence, RI, 2005, Papers from the Special Session at the 1st Joint AMS-RSME Meeting held in Seville, June 18–21, 2003.

[Har77] R. Hartshorne – *Algebraic geometry*, Graduate Texts in Math, vol. 52, Springer-Verlag, New York-Heidelberg, 1977.

[Hem19] M. Hemmig – "Isomorphisms between complements of projective plane curves", *Épijournal de Géom. Alg.* **3** (2019), article no. 16 (44 pages).

[KSC04] J. Kollár, K. E. Smith & A. Corti – *Rational and nearly rational varieties*, Cambridge Studies in Advanced Math., vol. 92, Cambridge University Press, Cambridge, 2004.

[KS18] A. Kuznetsov & E. Shinder – "Grothendieck ring of varieties, D- and L-equivalence, and families of quadrics", *Selecta Math. (N.S.)* **24** (2018), no. 4, p. 3475–3500.

[LS12] S. Lamy & J. Sebag – "Birational self-maps and piecewise algebraic geometry", *J. Math. Sci. Univ. Tokyo* **19** (2012), no. 3, p. 325–357.

[LL03] M. Larsen & V. A. Lunts – "Motivic measures and stable birational geometry", *Moscow Math. J.* **3** (2003), no. 1, p. 85–95.

[LPS11] W. Lee, E. Park & P. Schenzel – "On the classification of non-normal cubic hypersurfaces", *J. Pure Appl. Algebra* **215** (2011), no. 8, p. 2034–2042.

[Lib05] A. Libgober – "Lectures on topology of complements and fundamental groups", 2005, `arXiv:math/0510049`.

[LS10] Q. Liu & J. Sebag – "The Grothendieck ring of varieties and piecewise isomorphisms", *Math. Z.* **265** (2010), no. 2, p. 321–342.

[Mar16] N. Martin – "The class of the affine line is a zero divisor in the Grothendieck ring: an improvement", *Comptes Rendus Mathématique* **354** (2016), no. 9, p. 936–939.

[Mat86] H. Matsumura – *Commutative ring theory*, Cambridge Studies in Advanced Math., vol. 8, Cambridge University Press, Cambridge, 1986.

[Mil80] J. S. Milne – *Étale cohomology*, Princeton Math. Series, vol. 33, Princeton University Press, Princeton, NJ, 1980.

[Mil13] ———, "Lectures on etale cohomology (v2.21)", 2013, Available at `www.jmilne.org/math/`, p. 202.

[MJP21] L. Moser-Jauslin & P.-M. Poloni – "Isomorphisms between cylinders over Danielewski surfaces", *Beitr. Algebra Geom.* **62** (2021), no. 4, p. 755–771.

[Pfi95] A. Pfister – *Quadratic forms with applications to algebraic geometry and topology*, London Math. Soc. Lect. Note Series, vol. 217, Cambridge University Press, Cambridge, 1995.

[Pol11] P.-M. Poloni – "Classification(s) of Danielewski hypersurfaces", *Transform. Groups* **16** (2011), no. 2, p. 579–597.

[Sak10] Y. Sakamaki – "Automorphism groups on normal singular cubic surfaces with no parameters", *Trans. Amer. Math. Soc.* **362** (2010), no. 5, p. 2641–2666.

[Yos84] H. Yoshihara – "On open algebraic surfaces $\mathbf{P}^2 - C$", *Math. Ann.* **268** (1984), no. 1, p. 43–57.

Manuscript received 28th July 2023
accepted 17th June 2024

Jérémy Blanc, Universität Basel, Departement Mathematik und Informatik,
Spiegelgasse 1, CH-4051 Basel, Switzerland
E-mail : jeremy.blanc@unibas.ch
Url : http://algebra.dmi.unibas.ch/blanc

Pierre-Marie Poloni, Universität Basel, Departement Mathematik und Informatik,
Spiegelgasse 1, CH-4051 Basel, Switzerland
E-mail : poloni.pierremarie@gmail.com
Url : http://algebra.dmi.unibas.ch/poloni

Immanuel Van Santen, Universität Basel, Departement Mathematik und Informatik,
Spiegelgasse 1, CH-4051 Basel, Switzerland
E-mail : immanuel.van.santen@unibas.ch
Url : http://algebra.dmi.unibas.ch/vansanten