

JOURNAL

de Théorie des Nombres

de BORDEAUX

anciennement Séminaire de Théorie des Nombres de Bordeaux

Federico PELLARIN

On a variant of Schanuel conjecture for the Carlitz exponential

Tome 29, n° 3 (2017), p. 845-873.

<http://jtnb.cedram.org/item?id=JTNB_2017__29_3_845_0>

© Société Arithmétique de Bordeaux, 2017, tous droits réservés.

L'accès aux articles de la revue « Journal de Théorie des Nombres de Bordeaux » (<http://jtnb.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://jtnb.cedram.org/legal/>). Toute reproduction en tout ou partie de cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>

On a variant of Schanuel conjecture for the Carlitz exponential

par FEDERICO PELLARIN

RÉSUMÉ. Nous introduisons et décrivons une variante de la conjecture de Schanuel dans le cadre de l'exponentielle de Carlitz sur des algèbres de Tate et de fonctions similaires. Un autre objectif de ce travail est de stimuler des possibles investigations en transcendance et indépendance algébrique en caractéristique non nulle.

ABSTRACT. We introduce and discuss a variant of Schanuel conjecture in the framework of the Carlitz exponential function over Tate algebras and allied functions. Another purpose of the present paper is to widen the horizons of possible investigations in transcendence and algebraic independence in positive characteristic.

1. Introduction

Schanuel's conjecture is an unproven statement predicting the behavior of the intersections with algebraic sub-varieties defined over \mathbb{Q} of a certain analytic subvariety of $\mathbb{G}_a^n(\mathbb{C}) \times \mathbb{G}_m^n(\mathbb{C})$ of dimension n , built on the graph of the classical exponential function. Somehow, the conjecture expects that these intersections behave in the simplest possible way:

Conjecture 1.1 (Schanuel). *Let u_1, \dots, u_n be complex numbers which are linearly independent over \mathbb{Q} . Then the transcendence degree over \mathbb{Q} of the subfield of \mathbb{C} :*

$$\mathbb{Q}(u_1, \dots, u_n, e^{u_1}, \dots, e^{u_n})$$

is $\geq n$.

This conjecture first appeared in print in Lang's book [15]. It is surprising to see such a syntactically simple statement governing an intricate constellation of results of independence of classical mathematical constants. We mention that the *Lindemann–Weierstrass Theorem* confirms Schanuel's

Manuscrit reçu le 13 septembre 2016, accepté le 3 mars 2017.

2010 *Mathematics Subject Classification.* 11M38.

Mots-clés. Multiple zeta values, Carlitz module, Schanuel's conjecture.

This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme under the Grant Agreement No 648132.

conjecture in the case of u_1, \dots, u_n algebraic numbers, and can be seen as a particular case of Siegel–Shidlovski Theorem for the values of Siegel E -functions at algebraic numbers (see Beukers’ [7]). Baker’s Theorem on linear forms of logarithms of algebraic numbers, on the other hand, partially clarifies the case in which e^{u_1}, \dots, e^{u_n} are algebraic numbers. Outside these two cases, u_1, \dots, u_n algebraic or e^{u_1}, \dots, e^{u_n} algebraic, fragmentary information is available. Among those, a corollary of Nesterenko’s Theorem asserts that the numbers π and e^π are algebraically independent over \mathbb{Q} . We do not give precise references for these statements. Instead, we refer to the survey of Waldschmidt [25] and its detailed bibliography. The paper of Scanlon [21] gives a nice introduction of related topics.

As a final, maybe less known and more recent consequence of Schanuel’s conjecture, we also mention that in [17, Theorem 1.6], Marker deduces from the Schanuel conjecture that every non-zero exponential polynomial $P(x, e^x) \in \mathbb{Q}[x, e^x]$ such that $P(X, Y)$ depends on both X and Y , has infinitely many algebraically independent roots in \mathbb{C} . In particular, one deduces, from Schanuel’s conjecture, that any finite set of distinct fixed points of the exponential are algebraically independent over \mathbb{Q} .

In the present paper, we shall discuss variants of another conjecture, very similar to Schanuel’s conjecture in aspect, but in the framework of function fields of positive characteristic. Although at first sight, our interest in this framework may seem rather artificial (why looking at an analogue conjecture while we already have so many interesting open problems in the framework of the classical Schanuel conjecture?), we hope that the reader, after having looked into the present paper, will be finally convinced that some new structures emerge in this framework which would have remained hidden in the classical setting, more related to the theory of difference algebra and fields, perhaps giving a new view on the original Schanuel conjecture itself.

This text grew up from the tentative of the author to understand, in the viewpoint of the theory of transcendence and algebraic independence, the meaning of the fact that certain L -values introduced in [20] also behave as “Stark-Anderson units” and are thus sent to a polynomial by the Carlitz exponential function, extended over Tate algebras (see [2, 5, 4]). Our main purpose is to propose a statement which could play the role of a Schanuel conjecture in this setting; on the other hand, we warn the reader that the present paper does not contain substantial mathematical proofs in this direction.

It is a pleasure to dedicate the present work to David Goss, that helped in a fundamental way to shape the intuition in these investigations and constantly encouraged the progression of these works with determinant energy, and to Michel Waldschmidt, that inspired the “transcendental thinking” of

the author, encouraged and accompanied him constantly and generously, from his first steps, as a number theorist.

1.1. Some background. We set $A = \mathbb{F}_q[\theta]$ and $K = \mathbb{F}_q(\theta)$, where \mathbb{F}_q is the finite field with q elements and θ is an indeterminate. In all the following, we also denote by p the characteristic of \mathbb{F}_q so that $q = p^e$ for an integer $e > 0$. We denote by K_∞ the local field completion of K at the infinity place of K ; then K_∞ can be identified with the field of formal Laurent series $\mathbb{F}_q((1/\theta))$, with valuation v_∞ normalized by $v_\infty(\theta) = -1$. Let K_∞^{ac} be an algebraic closure of K_∞ . Then we denote by \mathbb{C}_∞ the completion of K_∞^{ac} for the unique extension of v_∞ . Note that \mathbb{C}_∞ is a K_∞^{ac} -vector space of infinite dimension, and that K_∞^{ac} is of infinite dimension over K_∞ . The field \mathbb{C}_∞ carries a unique extension

$$\mathbb{C}_\infty \xrightarrow{v_\infty} \mathbb{Q} \cup \{\infty\}$$

of the valuation v_∞ . Sometimes, we will also use the associated norm $|\cdot| = q^{-v_\infty(\cdot)}$. Just as the ring \mathbb{Z} is discrete and co-compact in \mathbb{R} , we have that the \mathbb{F}_q -algebra A is discrete and co-compact in K_∞ ; note also that the infinity place is the only place of K with this property.

1.1.1. The Carlitz module. Let $\iota : A \rightarrow B$ be a commutative A -algebra. Then B is equipped with the \mathbb{F}_q -algebra endomorphism $\tau : B \rightarrow B$ which sends $b \in B$ to $b^q \in B$ (the raising to the power q is relative to the algebra structure of B and is \mathbb{F}_q -linear). Let $B\{\tau\}$ be the skew ring of finite sums $\sum_{i \geq 0} b_i \tau^i$ with $b_i \in B$ for all i , and product defined by the rule $\tau b = \tau(b)\tau$ for $b \in B$; the ring $B\{\tau\}$ acts on B , as well as on any B -algebra, by *evaluation*. If $f = \sum_i \tau^i \in B\{\tau\}$, then for all $b \in B$, we define the evaluation of f at b as:

$$f(b) := \sum_i f_i \tau^i(b) = \sum_i f_i b^{q^i}.$$

The *Carlitz module* $C(B)$ over B is by definition the A -module whose underlying \mathbb{F}_q -vector space is B , in which the multiplication by $\theta \in A$, sufficient to define the entire A -module structure, is given by the evaluation of the skew polynomial $\iota(\theta) + \tau$. We shall write $C_a(b)$ for the multiplication of an element b of B by $a \in A$. For example, we have

$$C_\theta(b) = \iota(\theta)b + \tau(b), \quad b \in B.$$

In particular, \mathbb{C}_∞ is an A -algebra with ι the identity map, and the above construction gives rise to the A -module $C(\mathbb{C}_\infty)$.

1.1.2. The Carlitz exponential. The Carlitz exponential

$$\exp_C : \mathbb{C}_\infty \rightarrow \mathbb{C}_\infty$$

allows to *analytically uniformize* $C(\mathbb{C}_\infty)$. In what concerns the foundation

theory of this function, we are going to follow Goss' treatise [14] which can be consulted by any interested reader. We first remark, to define this function, that there is, available in the theory over A , an analogue of the sequence of numbers $n!$, defined as follows:

$$d_n = \prod_a a,$$

where the product runs over the monic polynomials a of A of degree n (this is, more properly, an analogue of the number $q^n!$). We set, for $z \in \mathbb{C}_\infty$,

$$\exp_C(z) := \sum_{n \geq 0} d_n^{-1} \tau^n(z) \in \mathbb{C}_\infty.$$

One verifies that $v_\infty(d_n) = nq^n$, from which we deduce that \exp_C is an entire (hence necessarily surjective by simple considerations of Newton polygons), \mathbb{F}_q -linear map $\mathbb{C}_\infty \rightarrow \mathbb{C}_\infty$. Since it is entire, its kernel Λ , an \mathbb{F}_q -vector space, determines \exp_C uniquely, and we have the convergent Weierstrass product expansion

$$(1.1) \quad \exp_C(z) = z \prod_{\lambda \in \Lambda \setminus \{0\}} \left(1 - \frac{z}{\lambda}\right), \quad z \in \mathbb{C}_\infty.$$

This function \exp_C in fact is the unique entire \mathbb{F}_q -linear map $\mathbb{C}_\infty \xrightarrow{F} \mathbb{C}_\infty$ which satisfies $F' = 1$, and which induces an exact sequence of A -modules:

$$0 \rightarrow \Lambda \rightarrow \mathbb{C}_\infty \rightarrow C(\mathbb{C}_\infty) \rightarrow 0,$$

where $\Lambda \subset \mathbb{C}_\infty$ is an A -sub-module of \mathbb{C}_∞ . The study of the Newton polygon of \exp_C implies that Λ is of rank one. In particular, there exists an element $\tilde{\pi} \in \mathbb{C}_\infty^\times$ such that $\Lambda = \tilde{\pi}A$; it is defined up to multiplication by an element of $\mathbb{F}_q^\times = \mathbb{F}_q \setminus \{0\}$. Note that for all $a \in A$ and $z \in \mathbb{C}_\infty$,

$$C_a(\exp_C(z)) = \exp_C(az),$$

hence providing an analytic isomorphism of A -modules

$$C(\mathbb{C}_\infty) \cong \frac{\mathbb{C}_\infty}{\tilde{\pi}A}.$$

The element $\tilde{\pi} \in \mathbb{C}_\infty$ can be constructed explicitly by limit processes in several ways. For instance, we recall from [14] that $\tilde{\pi}$ is the value in \mathbb{C}_∞ of a convergent infinite product

$$(1.2) \quad \tilde{\pi} := -(-\theta)^{\frac{q}{q-1}} \prod_{i=1}^\infty (1 - \theta^{1-q^i})^{-1} \in (-\theta)^{\frac{1}{q-1}} K_\infty,$$

uniquely defined up to the choice of a root $(-\theta)^{\frac{1}{q-1}}$. It has been proved in

a variety of ways (see [19] for a survey) that $\tilde{\pi}$ is moreover transcendental over K ; the first proof of which was obtained by Wade in [24].

2. A Carlitzian analogue of Schanuel’s conjecture

The following conjecture is due to Laurent Denis (see [12]).

Conjecture 2.1 (Denis). *Let u_1, \dots, u_n be elements of \mathbb{C}_∞ which are A -linearly independent. Then the transcendence degree over K of the sub-field of \mathbb{C}_∞*

$$K(u_1, \dots, u_n, \exp_C(u_1), \dots, \exp_C(u_n))$$

is $\geq n$.

The case in which u_1, \dots, u_n are algebraic over K has been solved by Thiery in [22] and can be viewed as the analogue of Lindemann–Weierstrass theorem for the Carlitz exponential. Note that the analogue for the Carlitz exponential of the Theorem of Hermite–Lindemann already appears in Wade [24] (see also [19, Theorem 2.5]). In the arithmetic theory of function fields in positive characteristic, what it is usually called the “analogue of Hermite–Lindemann Theorem” is a corollary of a very general transcendence result by Yu in [26] (see also [19, Theorem 2.2]). Years later, Papanikolas solved, in [18], the case in which $\exp_C(u_1), \dots, \exp_C(u_n)$ are algebraic over K by using the “criterion of linear independence of Anderson–Brownawell–Papanikolas”. Denis also obtained [13], at the same time as Papanikolas, a partial result in this direction, by using a characteristic p variant of “Mahler’s method”. Additionally, in [12, Corollaire 2(a)], Denis proved that $\tilde{\pi}$ and $\tilde{\varepsilon} := \exp_C(1)$ are algebraically independent over K if $q \geq 3$ (see [20] for an overview of these results).

2.1. Digression: strengthening. We mention, briefly, a natural way to reinforce Conjecture 2.1, but this has to be considered as a digression, since the nature of the statements we are primarily interested in, is different. Denis, in [12], proposes a strengthening of Conjecture 2.1 in order to give an interpretation of a multitude of results of algebraic independence he was obtaining, not only for special constants related to the Carlitz exponential, but also for their derivatives in the variable θ . Although not central for our further investigations, these strengthenings tell us that in the present settings (over \mathbb{C}_∞), there are more apparent structures that there seems to be over \mathbb{C} , and this allows us to propose more elaborate statements.

Denote by $\exp_C^{(i)} \in K[[z]]$ the i -th higher derivative of $\exp_C \in K[[z]]$ with respect to θ (note indeed that the formal series \exp_C can also be viewed as a double formal series in $\mathbb{F}_q[[\theta^{-1}, z]]$); it is an \mathbb{F}_q -linear entire function. We recall that p is the prime dividing q , that is, the characteristic of \mathbb{F}_q .

Conjecture 2.2 (Denis). *Let $u_1, \dots, u_n \in \mathbb{C}_\infty$ be elements which are A -linearly independent. Then the transcendence degree over K of the sub-field of \mathbb{C}_∞*

$$K(u_1, \dots, u_n, \exp_C(u_1), \dots, \exp_C(u_n), \dots, \exp_C^{(p-1)}(u_1), \dots, \exp_C^{(p-1)}(u_n))$$

is $\geq pn$.

In fact, the original statement in [12] was for $u_1, \dots, u_n \in K_\infty^{ac}$. However, we do not see any reason for which Conjecture 2.2 should fail for $u_i \in \mathbb{C}_\infty \setminus K_\infty^{ac}$ for some i . This conjecture obviously implies Conjecture 2.1.

Remark 2.3. Note that $\tilde{\pi}$ belongs to the separable closure K_∞^{sep} of K_∞ in \mathbb{C}_∞ . It is easy to see that the usual \mathbb{F}_q -linear higher derivatives in θ , defined by $D_n(\theta^m) = \binom{m}{n}\theta^{m-n}$, extend in an unique way to \mathbb{F}_q^{ac} -linear higher derivatives $K_\infty^{\text{sep}} \rightarrow K_\infty^{\text{sep}}$ (with \mathbb{F}_q^{ac} denoting the algebraic closure of \mathbb{F}_q in \mathbb{C}_∞). P. Voutier informed the author [23] of a work of D. Brownawell and A. van der Poorten, in which they proved that $\tilde{\pi}, D_1(\tilde{\pi}), \dots$ are algebraically independent over K , hence suggesting yet another way in which one could generalize Conjecture 2.1.

3. The basic settings for our operator-theoretic conjecture

In the following, we will keep focusing on statements similar to Conjecture 2.1. We want to formulate (this is the aim of the present paper) an *operator-theoretic conjecture* (see Conjecture 3.14 below) implying Conjecture 2.1. In order to do so, we first need to introduce certain difference Banach algebras. In §3.2, we will extend the Carlitz exponential to these algebras. This will give rise to several notions of independence, generalizing algebraic independence over K essential for our statement, and studied in §3.4. The statement of our conjecture will appear in §3.5 and we will give examples of application in §4.

We shall start with the algebra $\mathbb{C}_\infty[t_1, \dots, t_s]$ for variables t_1, \dots, t_s . If $q = p^e$ for an integer $e > 0$, then we set $\tau = \mu^e$ which is $\mathbb{F}_q[t_1, \dots, t_s]$ -linear. It is customary at this point, to take, when it is well defined, the completion of the above \mathbb{C}_∞ -algebra for the unique extension of the valuation v_∞ which is trivial over $\mathbb{F}_q[t_1, \dots, t_s]$, giving rise to a *Tate algebra*. Observe that if $s = 0$, then we just have the field \mathbb{C}_∞ . Let

$$\mathbb{C}_\infty[t_1, \dots, t_s] \xrightarrow{v_\infty} \mathbb{Q} \cup \{\infty\}$$

be the unique extension of the valuation v_∞ over \mathbb{C}_∞ which is trivial over $\mathbb{F}_q[t_1, \dots, t_s]$. The completion of $\mathbb{C}_\infty[t_1, \dots, t_s]$ with respect to this valuation is the standard *s-dimensional Tate algebra* denoted by \mathbb{T}_s in all the following. It is also called, in several papers, the *free s-dimensional affinoid*

algebra over \mathbb{C}_∞ . It is well known that \mathbb{T}_s is a ring which is Noetherian, factorial, of Krull dimension s (see [8] for the general theory of these algebras). It is isomorphic to the \mathbb{C}_∞ -algebra of the formal series

$$f = \sum_{i_1, \dots, i_s \geq 0} f_{i_1, \dots, i_s} t_1^{i_1} \dots t_s^{i_s} \in \mathbb{C}_\infty[[t_1, \dots, t_s]]$$

which satisfy

$$\lim_{\min\{i_1, \dots, i_s\} \rightarrow \infty} f_{i_1, \dots, i_s} = 0.$$

Thus, we have, for f a formal series of \mathbb{T}_s expanded as above, and non-zero, that

$$v_\infty(f) = \inf_{i_1, \dots, i_s} v_\infty(f_{i_1, \dots, i_s}) = \min_{i_1, \dots, i_s} v_\infty(f_{i_1, \dots, i_s}).$$

We also set, for convenience,

$$\|\cdot\| := q^{-v_\infty(\cdot)}$$

and $0 = \|0\| = q^{-\infty}$. In all the following, we are going to view the algebras \mathbb{T}_s as one embedded in the other, so that

$$\mathbb{C}_\infty = \mathbb{T}_0 \subset \mathbb{T}_1 \subset \dots \subset \mathbb{T}_s \subset \dots$$

It is easy to see that, for all $i > 0$,

$$\mathbb{T}_i = \left\{ f = \sum_{j \geq 0} f_j t_i^j; f_j \in \mathbb{T}_{i-1}, v_\infty(f_j) \rightarrow \infty \right\},$$

where v_∞ denotes here the Gauss valuation extending the valuation of \mathbb{C}_∞ over \mathbb{T}_{i-1} .

Then the map $\mathbb{C}_\infty \xrightarrow{\mu} \mathbb{C}_\infty, x \mapsto \mu(x) = x^p$ extends, uniquely, to a continuous, open $\mathbb{F}_p[t_1, \dots, t_s]$ -linear automorphism of \mathbb{T}_s (for all s) such that, for all $f \in \mathbb{T}_s$, we have that

$$v_\infty(\mu(f)) = p v_\infty(f).$$

It is also easy to prove that the subring $\mathbb{T}_s^{\mu=1} = \{f \in \mathbb{T}_s; \mu(f) = f\}$ is equal to $\mathbb{F}_p[t_1, \dots, t_s]$ (all these properties are proved in detail in [4]).

3.1. Some complete difference fields. It is suitable to also have, at hand, complete difference fields containing \mathbb{T}_s , not just difference algebras. Let L be any commutative field. We denote by $L\langle\langle\mathbb{Q}\rangle\rangle_\infty$ the set of formal series

$$\sum_{i \in \mathcal{I}} c_i \theta^{-i}, \quad c_i \in L,$$

where $\mathcal{I} \subset \mathbb{Q}$ is a *well-ordered* subset, that is, any non-empty subset has a minimum element. Then with the natural valuation v_∞ trivial over L and such that $v_\infty(\theta) = -1$, it is well known that $L\langle\langle\mathbb{Q}\rangle\rangle_\infty$ is a valued field which is complete and has residual field L (it is in fact henselian). Moreover, it has no proper immediate extensions and, if L is algebraically closed,

then $L\langle\langle\mathbb{Q}\rangle\rangle_\infty$ is algebraically closed. Since the residue field of $\mathbb{C}_\infty, \mathbb{F}_q^{ac}$, is algebraically closed, there is an isometric map of \mathbb{C}_∞ in $\mathbb{F}_q^{ac}\langle\langle\mathbb{Q}\rangle\rangle_\infty$, and therefore, if $L = \mathbb{F}_q^{ac}(t_1, \dots, t_s)$, then $\mathbb{C}_\infty(t_1, \dots, t_s)$ maps isometrically in

$$\mathbb{K}_s := \mathbb{F}_q^{ac}(t_1, \dots, t_s)\langle\langle\mathbb{Q}\rangle\rangle_\infty.$$

The complete field \mathbb{K}_s has thus residue field $\mathbb{F}_q^{ac}(t_1, \dots, t_s)$ and valuation group \mathbb{Q} . In particular, there is an isometric map from the completion of the fraction field of \mathbb{T}_s to \mathbb{K}_s .

The map $\mu : \mathbb{K}_s \rightarrow \mathbb{K}_s$ which sends an element $x = \sum_{i \in \mathbb{I}} c_i \theta^{-i} \in \mathbb{K}_s$ to the element $\mu(x) = \sum_{i \in \mathbb{I}} \mu(c_i) \theta^{-pi}$ extends the previously introduced $\mathbb{F}_p[t_1, \dots, t_s]$ -linear automorphism of \mathbb{T}_s (seen as embedded in \mathbb{K}_s as indicated above) is a continuous, open $\mathbb{F}_q(t_1, \dots, t_s)$ -linear automorphism. Further, we have the identity

$$\mathbb{K}_s^{\mu=1} = \{f \in \mathbb{K}_s : \mu(f) = f\} = \mathbb{F}_p(t_1, \dots, t_s)$$

which follows directly from the definition of the field \mathbb{K}_s .

3.2. Extensions of the Carlitz exponential function. We briefly recall some basic facts about the Carlitz exponential over Tate algebras, from [4]. We can construct a continuous, open $\mathbb{F}_q[t_1, \dots, t_s]$ -linear endomorphism

$$\mathbb{T}_s \xrightarrow{\text{exp}_C} \mathbb{T}_s,$$

by setting

$$\text{exp}_C(f) = \sum_{i \geq 0} d_i^{-1} \tau^i(f) = \sum_{i \geq 0} d_i^{-1} \mu^{ei}(f), \quad f \in \mathbb{T}_s.$$

Note that the restriction of exp_C to the subring $\mathbb{T}_0 = \mathbb{C}_\infty \cong \mathbb{C}_\infty.1 \subset \mathbb{T}_s$ returns the Carlitz exponential function defined in §1.1.2. Following the arguments of §1.1.1, we endow \mathbb{T}_s with the structure of an $A[t_1, \dots, t_s]$ -module $C(\mathbb{T}_s)$ in the following way. The underlying $\mathbb{F}_q[t_1, \dots, t_s]$ -module is just that of \mathbb{T}_s . Moreover, the multiplication by $\theta \in A$ is given by $C_\theta = \theta + \tau$ and all this produces a structure of $A[t_1, \dots, t_s]$ -module in an unique way.

Example. To give a concrete example of how this $A[t_1, \dots, t_s]$ -module structure works, let us suppose that $s = 1$. In this case, we more simply write $t = t_1$ and $\mathbb{T} = \mathbb{T}_1$. Let us consider $f = t_1 - \theta = t - \theta$, which belongs to $A[t] \subset \mathbb{T}$. Then $\tau(f) = t - \theta^q$ and

$$C_\theta(f) = t(\theta + 1) - (\theta^2 + \theta^q).$$

The following result is proved in [3].

Proposition 3.1. *The map exp_C induces an exact sequence of $A[t_1, \dots, t_s]$ -modules:*

$$(3.1) \quad 0 \rightarrow \tilde{\pi}A[t_1, \dots, t_s] \rightarrow \mathbb{T}_s \xrightarrow{\text{exp}_C} C(\mathbb{T}_s) \rightarrow 0.$$

Remark 3.2. For $s = 0$, we have seen that \exp_C defines an entire function $\mathbb{C}_\infty \rightarrow \mathbb{C}_\infty$. In particular, \exp_C , as an entire function, is uniquely defined by the divisor of its zeroes and has the Weierstrass product expansion (1.1) with $\Lambda = \tilde{\pi}A$. However, for $s > 0$, the extension of \exp_C to the Tate algebra \mathbb{T}_s that we have defined above, is no longer entire. For example, it has no Weierstrass product expansion over \mathbb{T}_s in contrast with the case $s = 0$.

More generally, we have the following.

Proposition 3.3. *The Carlitz exponential \exp_C gives rise to an exact sequence of $\mathbb{F}_q(t_1, \dots, t_s)[\theta]$ -modules:*

$$0 \rightarrow \tilde{\pi}\mathbb{F}_q(t_1, \dots, t_s)[\theta] \rightarrow \mathbb{K}_s \xrightarrow{\exp_C} C(\mathbb{K}_s) \rightarrow 0.$$

Observe that the $\mathbb{F}_q(t_1, \dots, t_s)[\theta]$ -module $C(\mathbb{K}_s)$ is well defined; the multiplication of $f \in \mathbb{K}_s$ by θ for this module structure is $C_\theta(f) = \theta f + \tau(f) = \theta f + \mu^e(f)$, and all the operators are extended $\mathbb{F}_q(t_1, \dots, t_s)$ -linearly. Restricted over the image of \mathbb{T}_s in \mathbb{K}_s , this exact sequence gives back the exact sequence (3.1).

Proof of Proposition 3.3. Since \mathbb{K}_s is complete, \exp_C is well defined. It is easy to see, writing down explicit generalized series of \mathbb{K}_s , that if $g \in \mathbb{K}_s$, then there exists a solution $f \in \mathbb{K}_s$ of the equation $C_\theta(f) = \theta f + \tau(f) = g$. This means that $C(\mathbb{K}_s)$ is θ -divisible and we can construct a continuous section of \exp_C ; this implies that \exp_C is surjective. Now, it is immediate that $\mathbb{F}_q(t_1, \dots, t_s)[\theta]\tilde{\pi} \subset \text{Ker}(\exp_C)$. To show the opposite inclusion, observe that, if $\mu \in \mathbb{K}_s^\times$ is such that $\exp_C(\mu) = 0$, then, there exists $N \geq 0$ such that $v_\infty(\frac{\mu}{\theta^N}) > -\frac{q}{q-1}$ and $\lambda = \exp_C(\frac{\mu}{\theta^N}) \neq 0$ is in the kernel of C_{θ^N} . It is easy to show that this kernel is equal to

$$\mathbb{F}_q(t_1, \dots, t_s) \exp_C\left(\frac{\tilde{\pi}}{\theta}\right) \oplus \dots \oplus \mathbb{F}_q(t_1, \dots, t_s) \exp_C\left(\frac{\tilde{\pi}}{\theta^N}\right)$$

(recall that $\mathbb{K}_s^{\tau=1} = \mathbb{F}_q(t_1, \dots, t_s)$). Since \exp_C induces an $\mathbb{F}_q(t_1, \dots, t_s)$ -linear isometry of the disk $\{f \in \mathbb{K}_s : v_\infty(f) > -\frac{q}{q-1}\}$, we can conclude. \square

3.3. A first conjectural statement. We set

$$K_s = \mathbb{F}_p(\theta, t_1, \dots, t_s).$$

This field will play the role of the field $K_0 = \mathbb{F}_p(\theta)$ (note that $K = K_0$ if and only if $q = p$; we hope that the fact that the notations K and K_s are so similar will not confuse the reader). In the settings of §3.2, it is natural to state the following conjecture, which generalizes Denis' conjecture 2.1:

Conjecture 3.4. *Let u_1, \dots, u_n be elements of \mathbb{K}_s which are $A[t_1, \dots, t_s]$ -linearly independent. Then the transcendence degree over K_s of the sub-field of \mathbb{K}_s*

$$K_s(u_1, \dots, u_n, \exp_C(u_1), \dots, \exp_C(u_n))$$

$is \geq n$.

We now explain why the above statement does not look so interesting; we set $s = 1$ for commodity (we write thus $\mathbb{T} = \mathbb{T}_1, t = t_1$ etc.). The *Anderson–Thakur function* can be defined as the element

$$\omega = \exp_C \left(\frac{\tilde{\pi}}{\theta - t} \right) \in \mathbb{T}^\times.$$

It is the generator of the $\mathbb{F}_q[t]$ -module $\text{Ker}(C_{\theta-t}) \cap \mathbb{T}$, free of rank one (see [3]). Since

$$(3.2) \quad \tau(\omega) = (t - \theta)\omega$$

(this is equivalent to saying that $\omega \in \text{Ker}(C_{\theta-t})$), we also deduce the next Proposition in quite an elementary way (see [3] for more details).

Proposition 3.5. *The following properties hold:*

- (1) *We have the product expansion*

$$\omega = (-\theta)^{\frac{1}{q-1}} \prod_{i \geq 0} \left(1 - \frac{t}{\theta^{q^i}} \right)^{-1},$$

convergent in \mathbb{T}^\times .

- (2) ω , *identified with a function of the variable $t \in \mathbb{C}_\infty$ with $\|t\| \leq 1$, extends to a meromorphic function over \mathbb{C}_∞ and has, as unique singularities, simple poles at the points $t = \theta, \theta^q, \theta^{q^2}, \dots$. The residues can be explicitly computed. In particular, we have $\text{Res}_{t=\theta}(\omega) = -\tilde{\pi}$.*
- (3) *The function $1/\omega$ extends to an entire function $\mathbb{C}_\infty \rightarrow \mathbb{C}_\infty$ with unique zeros located at the poles of ω .*

Since ω has infinitely many poles, it is transcendental over $K(t)$ and it is easy to see that $\tilde{\pi}, \omega$ are algebraically independent over $K(t)$, but the Conjecture 3.4 in the case $n = 1$ and $s = 1$ only implies that one among the two elements $\tilde{\pi}$ and ω is transcendental.

In fact, ω is transcendental over $K(t)$ but is also “ τ -algebraic” in the following sense: it satisfies the difference equation (3.2) which is a kind of algebraic relation involving τ , even though τ is not an algebraic morphism. What is missing to Conjecture 3.4 is the sensitiveness to such difference equations, and this is what we want to explore now.

3.4. A suitable notion of independence. Our operator-theoretic generalization of Denis–Schanuel’s conjecture 2.2 and of Conjecture 3.4 that will take place in the valued field \mathbb{K}_s . Before presenting it, we need to describe the relevant relations.

We recall Conjectures 1.1, 2.2 and 3.4. In the simplest portrait, we have a complete *environment field* (\mathbb{C} in the first above mentioned conjecture, \mathbb{C}_∞ in the second conjecture and \mathbb{K}_s in the third) a *countable base subfield*

of coefficients (\mathbb{Q} in the first conjecture, K in the second conjecture and K_s in the third). Moreover, we have notions of *linear relation* (over \mathbb{Z} , A and $A[t_1, \dots, t_s]$) and of *independence* (over the fraction fields of these rings). Our statement will take place in the environment field \mathbb{K}_s , concerns the extension of the Carlitz exponential discussed in the previous section, and will vaguely sound as follows, with a (momentarily) non-specified countable subfield of coefficients $L_s \subset \mathbb{K}_s$ and a non-specified notion of dependence over L_s :

Conjecture (Prototype). *Let f_1, \dots, f_n be elements of \mathbb{K}_s which are linearly independent over the ring $\mathbb{F}_q(t_1, \dots, t_s)[\theta]$. Then, among the $2n$ elements f_1, \dots, f_n and $\exp_C(f_1), \dots, \exp_C(f_n)$, n are “ μ -independent” over L_s .*

Of course, since our conjecture is tailored to imply Conjectures 2.2 and 3.4, we want that our relations over L_s to extend algebraic dependence over $K = \mathbb{F}_q(\theta)$, which is equivalent to algebraic dependence over $\mathbb{F}_p(\theta)$. Now, we are going to suppose that $s \geq 1$. In our μ -difference settings, the difference field extension (\mathbb{K}_s, μ) of the difference field (K_s, μ) , there are several noticeable classical notions of independence over K_s but we will see soon that they must be refined for our purposes.

3.4.1. Transformal independence. The first notion we want to discuss is that of *transformal independence* over K_s (read Levin, [16, §2.2]); as announced, it will turn out to be too coarse.

Let L be a μ -difference subfield of \mathbb{K}_s containing K_s .

Definition 3.6. Elements $f_1, \dots, f_n \in \mathbb{K}_s$ are *transformally μ -independent* over L if the elements

$$f_1, \mu(f_1), \dots, f_2, \mu(f_2), \dots, \dots, f_n, \mu(f_n), \dots$$

are algebraically independent over L ; the notions of *transformal μ -dependence*, *transformal μ -algebraicity* and *transformal μ -transcendence* can be defined accordingly (see [16]). This also leads to a notion of *transformal μ -independence degree*. Let f_1, \dots, f_n be elements of \mathbb{K}_s and let us denote by:

$$\mathcal{L} = L(f_1, \dots, f_n)_\mu$$

the smallest μ -difference subfield of \mathbb{K}_s containing L and f_1, \dots, f_n . Then the transformal μ -independence degree of \mathcal{L} over L

$$\text{transf deg}_L(\mathcal{L})$$

is by definition the minimal cardinality of a transformally μ -independent subset of \mathcal{L} over L . If f_1, \dots, f_n are transformally μ -independent over L , then

$$\text{transf deg}_L(\mathcal{L}) = n.$$

The notions above are not tailored for the statement we want to produce. To illustrate this, observe that, since $c^p = \mu(c)$ for all $c \in \mathbb{C}_\infty$, every element of \mathbb{C}_∞ is transformally μ -algebraic over K , independently of the fact that it is algebraic over K or not. Worse, for all $c \in \mathbb{C}_\infty$ and for all $f \in K_s$, the element $g = cf \in \mathbb{C}_\infty(t_1, \dots, t_s)$ is transformally μ -algebraic. Indeed, $\mu(f)g^p = f^p\mu(g)$. We deduce from [16, Theorem 4.1.2] that every element of $\mathbb{C}_\infty(t_1, \dots, t_s)$ is transformally μ -algebraic, no matter if the coefficients of the rational fractions are algebraic or not.

Note also that in particular, the subfield generated by the elements $x \in \mathbb{K}_s$ which are transformally μ -algebraic over K_s is not countable. The generalization of Conjecture 2.1 (see Conjecture 3.14) that we have in mind is of arithmetic nature, and is similar to [6, Hypothesis (Σ) p. 252], which is equivalent to the Schanuel conjecture itself (see [6, Theorem 2]). There, a countable base subfield is required (the field \mathbb{Q}). It seems to us that for statements of arithmetic nature such as Schanuel’s Conjecture¹, it is natural to have a countable base field; this means that the notion of transformal μ -independence is not suitable for our framework. We now make an attempt to give a more refined notion of relation.

3.4.2. Analytically critical and regular μ -polynomials. We choose a finite set of symbols

$$\underline{X} = (X_1, \dots, X_n)$$

and we introduce infinitely many indeterminates

$$\mu^0(X_i), \mu^1(X_i), \mu^2(X_i), \dots \quad i = 1, \dots, n.$$

To simplify the notation and to make it more natural, we shall write $X_i = \mu^0(X_i)$ and $\mu(X_i) = \mu^1(X_i)$. Let L be a μ -difference field. We consider the ring in infinitely many indeterminates

$$L[\underline{X}]_\mu = L[X_i, \mu(X_i), \mu^2(X_i), \dots : i = 1, \dots, n],$$

that we turn into a μ -difference ring by setting

$$\mu(\mu^j(X_i)) = \mu^{j+1}(X_i).$$

The elements of $L[\underline{X}]_\mu$ are called the μ -polynomials in the symbols \underline{X} with coefficients in L . Any element $P \in L[\underline{X}]_\mu$ can be written in a unique way as

$$(3.3) \quad P(\underline{X}) = \sum_{\underline{i} \in \mathbb{N}^{(k+1)n}} c_{\underline{i}} X_1^{i_{1,0}} \mu(X_1)^{i_{1,1}} \dots \mu^k(X_1)^{i_{1,k}} \dots \\ \dots X_n^{i_{n,0}} \mu(X_n)^{i_{n,1}} \dots \mu^k(X_n)^{i_{n,k}}, \quad c_{\underline{i}} \in L,$$

and the smallest $k \in \mathbb{N}$ for which the expansion (3.3) holds is called the *depth* of P .

¹Observe that we are not considering here, statements such as [6, (SP)].

Let P be in $L[\underline{X}]_\mu$ as in (3.3) and let $\underline{f} = (f_1, \dots, f_n)$ be an element of \mathbb{K}_s^n . Then the *evaluation of P at \underline{f}* is the element of \mathbb{K}_s :

$$P(\underline{f}) := \sum_{\underline{i} \in \mathbb{N}^{(k+1)n}} c_{\underline{i}} f_1^{i_{1,0}} \mu(f_1)^{i_{1,1}} \dots \mu^k(f_1)^{i_{1,k}} \dots f_n^{i_{n,0}} \mu(f_n)^{i_{n,1}} \dots \mu^k(f_n)^{i_{n,k}}.$$

If $P(\underline{f}) = 0$, we say that f is a *zero* or a *root* of P . We choose a μ -difference subfield L' of \mathbb{K}_s containing the coefficients of a μ -polynomial P . The *μ -variety of zeroes of P in L'* is the subset of $\underline{f} \in L'^n$ such that $P(\underline{f}) = 0$. It is denoted by $Z_{L'}(P)$ or more simply, by $Z(\overline{P})$ when $L' = \mathbb{K}_s$. We can also consider varieties of simultaneous zeroes of a collection of μ -polynomials. We observe that $Z(P) = Z(\mu(P))$ for any μ -polynomial. Let P be a μ -polynomial. We denote by P^μ the μ -polynomial obtained by applying μ to all the coefficients of P . Then, it is easy to see that $Z(P^\mu) = \mu(Z(P))$.

We set $D := D(0, 1) = \{z \in \mathbb{C}_\infty : v_\infty(z) \geq 0\}$. Let $n \geq 1$ be an integer. We set $\underline{X} = (X_1, \dots, X_n)$ and $\underline{z} = (z_1, \dots, z_n)$. We denote by $\widehat{\mathbb{T}}_{\underline{z}}$ the Tate algebra $\widehat{\mathbb{T}}_{\underline{z}} = \widehat{\mathbb{C}_\infty[z_1, \dots, z_n]}$, completion of the polynomial ring $\mathbb{C}_\infty[z_1, \dots, z_n]$ for the Gauss norm.

Definition 3.7. Let P be a non-zero μ -polynomial of $\mathbb{K}_s[\underline{X}]_\mu$. We say that it is *analytically critical* if there exists a map

$$D^n \xrightarrow{F} Z(P) \subset \mathbb{K}_s^n,$$

$\underline{z} \in D^n \mapsto F(\underline{z}) = (F_1(\underline{z}), \dots, F_n(\underline{z}))$ which is a *rigid immersion* in the following sense. The coordinate functions $\underline{F} = (F_1, \dots, F_n)$ are series

$$F_i = F_i(z_1, \dots, z_n) = \sum_{\substack{\underline{k}=(k_1, \dots, k_n) \\ k_i \geq 0, \forall i=1, \dots, n}} F_{i, \underline{k}} z_1^{k_1} \dots z_n^{k_n} \in \mathbb{K}_s \widehat{\otimes}_{\mathbb{C}_\infty} \widehat{\mathbb{T}}_n,$$

(such that, for all i, \underline{k} , $F_{i, \underline{k}} \in \mathbb{K}_s$ and $\|F_{i, \underline{k}}\| \rightarrow 0$ as $|\underline{k}| = \sum_j k_j \rightarrow \infty$), and moreover, the Jacobian

$$J_{\underline{z}}(\underline{F}) = \begin{pmatrix} \frac{\partial F_1}{\partial z_1} & \dots & \frac{\partial F_n}{\partial z_1} \\ \vdots & & \vdots \\ \frac{\partial F_1}{\partial z_n} & \dots & \frac{\partial F_n}{\partial z_n} \end{pmatrix}$$

has everywhere maximal rank over D^n . A μ -polynomial which is not analytically critical is said *analytically regular*.

Equivalently, P is analytically critical if a point \underline{F} with elements of $\mathbb{K}_s \widehat{\otimes}_{\mathbb{C}_\infty} \widehat{\mathbb{T}}_{\underline{z}}$ as entries, satisfying the above conditions, belongs to the set $Z_{\mathbb{K}_s \widehat{\otimes}_{\mathbb{C}_\infty} \widehat{\mathbb{T}}_{\underline{z}}}(P)$. From this, we deduce that the set of analytically critical μ -polynomials of $\mathbb{K}_s[\underline{X}]_\mu$ is a non-zero prime ideal.

3.4.3. Examples. Any polynomial $P \in \mathbb{K}_s[\underline{X}]_\mu$ which is multi-homogeneous for the unique multi-graduation which assigns to $\mu^i(X_j)$ the degree p^i , is analytically critical as soon as it has a root $\underline{f} \in (\mathbb{K}_s \setminus \{0\})^n$. Indeed, if $\underline{f} = (f_1, \dots, f_n) \in \mathbb{K}_s^n$ is a root, then also $(c_1 f_1, \dots, c_n f_n)$ for all $c_1, \dots, c_n \in \mathbb{C}_\infty$ is a root, due to the fact that $c^p = \mu(c)$ if $c \in \mathbb{C}_\infty$, so that there is a rigid immersion map $D^n \rightarrow Z(P)$. In particular, the μ -polynomial in one variable $\mu(X) - X^p$ is analytically critical. Note that there also are analytically critical μ -polynomials which are not homogeneous for the above graduations.

3.4.4. Examples of analytically regular μ -polynomials.

Lemma 3.8. *Any non-zero polynomial $P \in \mathbb{K}_s[\underline{X}]$ is analytically regular. Moreover, a μ -polynomial $P \in \mathbb{K}_s[\underline{X}]_\mu$ is analytically critical if and only if $\mu(P)$ is analytically critical. Similarly, P is analytically critical if and only if P^μ is analytically critical.*

Proof. It is easy to see, by the Jacobian criterion, that the components of any rigid immersion are algebraically independent over \mathbb{K}_s . This proves the first property. Now, recall that $Z(\mu(P)) = Z(P)$ for any μ -polynomial P ; this suffices to show that P is analytically critical if and only if $\mu(P)$ is analytically critical. To prove the last property, recall that $Z(P^\mu) = \mu(Z(P))$. Assume first that P is analytically critical. Then, there is a rigid immersion $D^n \xrightarrow{F} Z(P)$. We write $\underline{F} = (F_1, \dots, F_n)$ with $F_i = \sum_{\underline{k}} F_{i,\underline{k}} z_1^{k_1} \dots z_n^{k_n}$ with $F_{i,\underline{k}} \in \mathbb{K}_s$, for all i . We observe that μ induces a bijection of D^n and, if we set

$$\tilde{F}_i := \sum_{\underline{k}} \mu(F_{i,\underline{k}}) z_1^{k_1} \dots z_n^{k_n}, \quad i = 1, \dots, n,$$

then $\tilde{\underline{F}} = (\tilde{F}_1, \dots, \tilde{F}_n)$ defines a rigid immersion $D^n \rightarrow Z(P^\mu)$. Similarly, let us suppose that P^μ is analytically critical, and let $\underline{F} = (F_1, \dots, F_n)$ be a rigid immersion $D^n \rightarrow Z(\mu(P))$. Then, setting now

$$\tilde{F}_i := \sum_{\underline{k}} \mu^{-1}(F_{i,\underline{k}}) z_1^{k_1} \dots z_n^{k_n}, \quad i = 1, \dots, n$$

defines a rigid immersion $D^n \rightarrow Z(P)$ and P is analytically critical. □

Hence, a μ -polynomial of depth zero is analytically regular, but these are not the only examples.

The next definition provides a larger class of analytically regular μ -polynomials.

Definition 3.9. Let $P \in \mathbb{K}_s[\underline{X}]_\mu$ be as in (3.3), and non-zero. We say that it is *tame* if $c_{\underline{i}} \neq 0$ implies that, writing $\underline{i} = (i_{l,j})_{\substack{0 \leq l \leq n \\ 0 \leq j \leq k}}$, we have that

$$0 \leq i_{l,j} < p \text{ for all } l, j.$$

In other, more loosely words, a tame μ -polynomial is one such that no exponent exceeds, or is equal to p .

Lemma 3.10. *Every tame μ -polynomial is analytically regular.*

Proof. We suppose again that $\underline{X} = (X_1, \dots, X_n)$. We shall proceed by induction over the depth k ; the statement is clear for depth $k = 0$ (see Lemma 3.8). Let $P \in \mathbb{K}_s[\underline{X}]_\mu$ be a tame μ -polynomial with μ -expansion as in (3.3) and depth $k > 0$. By Lemma 3.8, we can suppose that P depends of the indeterminates X_1, \dots, X_n (more precisely, of the indeterminates $\mu^0(X_1), \dots, \mu^0(X_n)$). Let us suppose by contradiction that it is analytically critical. Then there is a rigid immersion

$$D^n \xrightarrow{F=(F_1, \dots, F_n)} Z(P).$$

Since the Tate algebra \mathbb{T}_z in the variables z_1, \dots, z_n is stable for the partial derivatives $\frac{\partial}{\partial z_i}$ for all i , the map $z \mapsto P(\underline{F}(z))$ is differentiable in each variable z_i (in the polydisk D^n). We note that

$$\frac{\partial}{\partial z_j} (\mu^l(X_k)^i(\underline{F}(z))) = 0, \quad j = 1, \dots, n, \quad i, l > 0$$

(here, μ acts as the p -power on \mathbb{T}_z). By the usual chain rule and the invertibility of the Jacobian of \underline{F} we deduce that

$$\frac{\partial P}{\partial X_1}(\underline{F}(z)) = \dots = \frac{\partial P}{\partial X_n}(\underline{F}(z)) = 0.$$

Observe that $0 < \deg_{X_i}(P) < p$ for all i . Then we have proved that there exists $i \in \{1, \dots, n\}$ such that $\frac{\partial P}{\partial X_i}$ is tame and analytically critical. Iterating this process, we can successively construct a sequence of analytically critical tame polynomials such that the sequence of the total degrees in the indeterminates X_1, \dots, X_n is strictly decreasing and we ultimately find a tame analytically critical μ -polynomial of depth k which does not depend of the indeterminates X_1, \dots, X_n . Then, by Lemma 3.8, $P = \mu^r(Q)$ for some $r > 0$ and Q tame, analytically critical of depth $< k$ which is impossible. □

Remark 3.11. Note that a polynomial $P \in \mathbb{K}_s[\underline{X}]$, that is, a μ -polynomial of depth zero, needs not to be tame. Also, the product of two tame μ -polynomials needs not to be tame; however, there exists a natural \mathbb{K}_s -algebra structure over the \mathbb{K}_s -vector space of tame μ -polynomials in $\mathbb{K}_s[\underline{X}]_\mu$ which makes it isomorphic to $\mathbb{K}_s[\underline{X}]$. See the appendix §A with the details of the construction (it will not be used in this paper). Note also that the maps $\mathbb{C}_\infty^n \rightarrow \mathbb{K}_s$ that tame polynomials define, are in natural \mathbb{K}_s -linear isomorphism with the polynomial maps $\mathbb{C}_\infty^n \rightarrow \mathbb{K}_s$ induced by polynomials of $\mathbb{K}_s[\underline{X}]$.

Definition 3.12. Let L be a μ -difference subfield of \mathbb{K}_s containing K_s . Let f_1, \dots, f_n be elements of \mathbb{K}_s . We say that they are *regularly μ -independent over L* if for all $P \in L[\underline{X}]_\mu$ analytically regular (hence non-zero), $P(\underline{f}) \neq 0$, where $\underline{f} = (f_1, \dots, f_n)$. We say that f_1, \dots, f_n are *tamely μ -independent over L* if $P(\underline{f}) \neq 0$ for all tame μ -polynomial $P \in L[\underline{X}]_\mu$.

The definition also produces notions of μ -dependence, tame μ -dependence, μ -transcendence, μ -algebraicity, etc.

If f_1, \dots, f_m are tamely μ -dependent, then they are regularly μ -dependent. Also, if f_1, \dots, f_m are regularly μ -dependent, then they are transformally μ -dependent in the sense of Levin [16]. But the reverse implications are all false. We collect a few more properties in the following proposition.

Proposition 3.13. *Let f_1, \dots, f_n be elements of \mathbb{K}_s . We have:*

- (1) *If f_1, \dots, f_n are regularly μ -independent over L , then they are algebraically independent over L .*
- (2) *If f_1, \dots, f_n are in \mathbb{C}_∞ and are algebraically independent over K , then they are regularly μ -independent over K_s .*

There exist algebraic elements $f \in \mathbb{K}_s$ which are transcendental over K_s but tamely μ -algebraic over K_s ; examples are provided by certain torsion points of the Carlitz module, see §4.1.

Proof of Proposition 3.13.

(1). It follows from Lemma 3.8.

(2). We suppose to have elements $f_1, \dots, f_n \in \mathbb{C}_\infty$ and an analytically regular μ -polynomial $P \in K_s[\underline{X}]_\mu$ with $P(\underline{f}) = 0$ and $\underline{f} = (f_1, \dots, f_n)$. This implies, in particular, that the map $\mathbb{C}_\infty^n \rightarrow \mathbb{K}_s$ induced by P is not identically zero. This map is further equal, by the fact that μ and $x \mapsto x^p$ agree on \mathbb{C}_∞ , to a polynomial map induced by a polynomial $Q \in K_s[\underline{X}]$; since the map is not identically zero by hypothesis, Q is non-zero. Since K^s is dense in D^s , there exists an element $\underline{x} = (x_1, \dots, x_s) \in D^s \cap K^s$ such that the evaluation of the coefficients of Q , which are rational functions of K_s , is well defined and does not map to $\{0\}$. This yields a non-trivial algebraic dependence relation of f_1, \dots, f_n over K . □

3.5. A refinement of Denis–Schanuel’s conjecture. We set $q = p^e$ with $e > 0$ and we fix an integer $s \geq 0$. We recall that

$$K_s = \mathbb{F}_p(\theta, t_1, \dots, t_s) \subset \mathbb{K}_s.$$

We denote by L_s the smallest μ -difference subfield of \mathbb{K}_s containing K_s and all the $f \in \mathbb{K}_s$ which are regularly μ -algebraic over K_s ; by construction, it is countable. The exponential function $\exp_C : \mathbb{K}_s \rightarrow C(\mathbb{K}_s)$ is defined by

$$\exp_C = \sum_{i \geq 0} d_i^{-1} \tau^i = \sum_{i \geq 0} d_i^{-1} \mu^{ei}$$

as in §3.2. We can now state our conjecture.

Conjecture 3.14 (Operator-theoretic generalization of Denis–Schanuel’s Conjecture). *Let f_1, \dots, f_n be elements of \mathbb{K}_s and let us write*

$$g_i := \exp_C(f_i) \in \mathbb{K}_s$$

for all $i = 1, \dots, n$. If f_1, \dots, f_n are $\mathbb{F}_q(t_1, \dots, t_s)[\theta]$ -linearly independent, then n among the elements

$$f_1, \dots, f_n, g_1, \dots, g_n$$

are regularly μ -independent over L_s .

We see that in the case $s = 0$ or if f_1, \dots, f_n all belong to \mathbb{C}_∞ , then this conjecture reduces to Conjecture 2.1 by Proposition 3.13(2). Also, for any s , Conjecture 3.14 implies Conjecture 2.1 and Conjecture 3.4. We do not know if, reciprocally, Conjecture 2.1 implies Conjecture 3.14 in analogy with the hypothesis (Σ) of [6], which is equivalent to the Schanuel conjecture; this looks unlikely.

4. Some examples

We are going to give some examples of consequences of our Conjecture 3.14. We first review basic properties of certain special functions that are used in our examples.

4.1. Torsion. Our Carlitz exponential $\exp_C : \mathbb{K}_s \rightarrow C(\mathbb{K}_s)$ gives rise to quite a rich torsion structure that we review here (see [4] for more properties). Let $a \in \mathbb{F}_q(t_1, \dots, t_s)[\theta]$ be of degree $d \geq 0$ in θ . Then for all $j = 0, \dots, \deg_\theta(a) - 1$, $\frac{\tilde{\pi}\theta^j}{a} \in \mathbb{K}_s$. In particular $\exp_C\left(\frac{\tilde{\pi}\theta^j}{a}\right) \in \mathbb{K}_s$ and one sees that

$$C_a\left(\exp_C\left(\frac{\tilde{\pi}\theta^j}{a}\right)\right) = 0, \quad j = 0, \dots, \deg_\theta(a) - 1.$$

In fact, the elements $\exp_C\left(\frac{\tilde{\pi}\theta^j}{a}\right) \in \mathbb{K}_s$ constitute an $\mathbb{F}_q(t_1, \dots, t_s)$ -basis of the vector space underlying the $\mathbb{F}_q(t_1, \dots, t_s)[\theta]$ -submodule $\text{Ker}(C_a) \subset C(\mathbb{K}_s)$. Every such a torsion point satisfies a non-trivial linear τ -difference equation with coefficients in $\mathbb{F}_q(t_1, \dots, t_s)[\theta]$, hence it satisfies a non-trivial linear μ -difference equation with coefficients in K_s . If moreover a is a polynomial of $\mathbb{F}_q[t_1, \dots, t_s, \theta]$ which is monic in θ , then $a^{-1} \in \mathbb{T}_s$ and we have $\exp_C\left(\frac{\tilde{\pi}\theta^j}{a}\right) \in \mathbb{T}_s$ for all j . We see that, for $j = 0, \dots, d - 1$, the above elements span the rank d free $\mathbb{F}_q[t_1, \dots, t_s]$ -module $\text{Ker}(C_a) \cap \mathbb{T}_s$.

The simplest case (with $s = 1$, case in which we write t for t_1) is given by the Anderson–Thakur function and comes with the choice of $a = \theta - t$. Note that it is also tamely μ -algebraic; indeed, from (3.2), we deduce that

the μ -polynomial $\mu^e(X) - (t - \theta)X = C_{\theta-t}(X) \in K_s[X]_\mu$ vanishes at ω and is tame.

4.2. Zeta values. The so-called *Carlitz zeta values* are defined as follows, for $n \geq 1$ an integer, where A^+ denotes the multiplicative monoid of monic polynomials of A , and the product runs over the irreducible polynomials $P \in A^+$:

$$\zeta_A(n) = \sum_{a \in A^+} a^{-n} = \prod_P \left(1 - \frac{1}{P^n}\right)^{-1} \in 1 + \theta^{-1}\mathbb{F}_q[[\theta^{-1}]].$$

Carlitz essentially proved in [9] that

$$(4.1) \quad \exp_C(\zeta_A(1)) = 1.$$

Conjectures 2.1 and 3.14 immediately imply that $\zeta_A(1)$ is transcendental over K and indeed, this follows from the transcendence theory of the Carlitz module in a variety of ways. First of all, this can be viewed as a consequence of the analogue of Hermite–Lindemann Theorem that can be found in [24]. Also, this follows from the Theorem of Papanikolas [18], see also Denis [13] and the survey [19] by the author, not to mention other proofs, with more diophantine, or automatic flavor. More generally, Chieh-Yu Chang and Jing Yu have proved in [11] the following result.

Theorem 4.1 (Chieh-Yu Chang & Jing Yu). *The element $\tilde{\pi}$ and all the Carlitz zeta-values $\zeta_A(n)$ for all $n \geq 1$ not divisible by p and $q - 1$, are algebraically independent over K .*

Note that the elements of the above theorem generate the K -algebra

$$K(\tilde{\pi}, \zeta_A(1), \zeta_A(2), \dots).$$

Indeed, on one side, we have the “trivial sum-shuffle relations”

$$(4.2) \quad \zeta_A(pn) = \zeta_A(n)^p = \mu(\zeta_A(n)),$$

and on the other hand, we have the result of Carlitz

$$(4.3) \quad \zeta_A(k(q - 1)) \in K^\times \tilde{\pi}^{k(q-1)}$$

for all $k \geq 1$. In other words, The so-called Bernoulli–Carlitz relations $\zeta_A(n) \in K^\times$ for $n > 0$ such that $q - 1 \mid n$ and the relations (4.2) exhaust all the algebraic dependence relations between $\tilde{\pi}$ and $\zeta_A(n)$ for all $n \geq 1$.

4.2.1. Zeta values in Tate algebras. We recall from [4] the construction of *Carlitz zeta values in \mathbb{T}_s* . They are defined as follows, for $n \geq 1$ and an integer $s \geq 0$:

$$\zeta_A(n; s) = \sum_{a \in A^+} a^{-n} a(t_1) \dots a(t_s) = \prod_P \left(1 - \frac{P(t_1) \dots P(t_s)}{P^n}\right)^{-1}.$$

The above product converges in the complete topological subgroup

$$1 + \theta^{-1} \mathbb{F}_q(t_1, \dots, t_s) \llbracket \theta^{-1} \rrbracket \subset \mathbb{T}_s^\times.$$

The classical Carlitz zeta values are a special case of our construction with $s = 0$; in our notations, $\zeta_A(n) = \zeta_A(n; 0)$. In terms of the variables t_1, \dots, t_s , these series define entire functions $\mathbb{C}_\infty^s \rightarrow \mathbb{C}_\infty$ (see [2]). Therefore, evaluation at $t_i = \theta^{q^{k_i}}$, $i = 1, \dots, s$ and $k_i \in \mathbb{Z}$ makes sense and, for $n > 0$, we have the identities

$$\zeta_A(n) = \zeta_A(n; 0) = \zeta_A(n + q^{k_1} + \dots + q^{k_s}; s) \Big|_{t_i = \theta^{q^{k_i}}}.$$

In this respect, we can view these functions as entire *interpolations of Carlitz zeta values*. Moreover, note that, for all $m > 0$, there exists $N \geq 0$ and s such that

$$\zeta_A(m) = \tau^N(\zeta_A(1; s))_{t_i = \theta, \forall i = 1, \dots, s}.$$

4.2.2. Consequences of Conjecture 3.14. We begin by reviewing an important property of the Carlitz zeta values $\zeta_A(1; s)$. In [4], the following generalization of Carlitz’s identity (4.1) is proved:

Theorem 4.2. *For $s \geq 0$, we have*

$$\exp_C(\zeta_A(1; s) \omega(t_1) \dots \omega(t_s)) = P_s(t_1, \dots, t_s) \omega(t_1) \dots \omega(t_s),$$

where $P_s \in A[t_1, \dots, t_s]$. Moreover, for $s > 1$, we have $P_s = 0$ if and only if $s \equiv 1 \pmod{q - 1}$. In this case, we have

$$\zeta_A(1; s) = \frac{\tilde{\pi} B_s}{\omega(t_1) \dots \omega(t_s)},$$

with $B_s \in A[t_1, \dots, t_s] \setminus \{0\}$.

We shall now use this result to prove:

Theorem 4.3. *Conjecture 3.14 implies the truth of the following statement: The element $\tilde{\pi}$ and all the elements $\zeta_A(1; s)$ for $s \not\equiv 1 \pmod{q - 1}$ are regularly μ -independent over K_s .*

In particular, we deduce, conditionally upon verification of Conjecture 3.14, that the values considered are algebraically independent and also tamely μ -independent over K_s .

Proof of Theorem 4.3. We choose $s_0 > 0$ and we denote by \mathfrak{S} the set of integers s with $s_0 \geq s \geq 0$ such that $s \not\equiv 1 \pmod{q - 1}$; we choose s_0 so that \mathfrak{S} is non-empty. We also denote by ω_s , for $s \in \mathfrak{S}$, the product $\omega(t_1) \dots \omega(t_s)$, which belongs to \mathbb{T}_{s_0} . Observe that, by the difference equation (3.2), for all $s \in \mathfrak{S}$,

$$\mu^e(\omega_s) = (t_1 - \theta) \dots (t_s - \theta) \omega_s,$$

which implies that ω_s is tamely, thus regularly μ -algebraic over K_s .

For fixed $s \in \mathfrak{S}$, the function ω_s also defines a meromorphic function $\mathbb{C}_\infty^{s_0} \rightarrow \mathbb{C}_\infty$ whose polar divisor D_s is, by Proposition 3.5, given by the free sum (with multiplicity one) of the affine subsets of co-dimension one

$$D_{i,k,s} = \{(x_1, \dots, x_{i-1}, \theta^{q^k}, x_{i+1}, \dots, x_s, x_{s+1}, \dots, x_{s_0}) : x_j \in \mathbb{C}_\infty \forall j\} \subset \mathbb{C}_\infty^{s_0},$$

for $k \geq 0$ and $1 \leq i \leq s$, $s \in \mathfrak{S}$. We note that if $s < s'$ and $s, s' \in \mathfrak{S}$, then $D_s \subset D_{s'}$ and $D_{s'} \setminus D_s$ has infinitely many irreducible components.

We claim that the elements $\tilde{\pi}$ and $\omega_s \zeta_A(1; s)$ with $s \in \mathfrak{S}$ are linearly independent $\mathbb{F}_q(t_1, \dots, t_{s_0})[\theta]$. We write $\mathfrak{S} = \{s_1, \dots, s_n\}$. Let us assume by contradiction that there exists a non-trivial linear dependence relation

$$a_0 \tilde{\pi} + a_1 \omega_{s_1} \zeta_A(1; s_1) + \dots + a_n \omega_{s_n} \zeta_A(1; s_n) = 0, \\ a_i \in A[t_1, \dots, t_{s_0}], s_i \in \mathfrak{S}.$$

For all $\underline{k} := (k_1, \dots, k_{s_0}) \in \mathbb{N}^{s_0}$, the congruence condition on $s \in \mathfrak{S}$ implies that the evaluation $\text{ev}_{\underline{k}}(\zeta_A(1; s))$ of $\zeta_A(1; s)$ at the point $(t_1, \dots, t_s) = (\theta^{q^{k_1}}, \dots, \theta^{q^{k_s}})$ is non-zero (this follows from a well known result of Goss, in [14]). This means that for all j , the function $\omega_{s_j} \zeta_A(1, s_j)$ has polar divisor given by D_{s_j} . Since the locus of the zeros of a_j has only finitely many irreducible components and since the polar divisors of the functions $\omega_{s_j} \zeta_A(1, s_j)$ are embedded one in the other along the total order of \mathfrak{S} induced by $<$ on \mathbb{N} , we deduce a contradiction. This shows that $a_0 = a_1 = \dots = a_n = 0$ and the elements $\tilde{\pi}$ and $\omega_{s_j} \zeta_A(1, s_j)$ are linearly independent as expected.

We set $f_j = \omega_{s_j} \zeta_A(1, s_j)$, $f_0 = \tilde{\pi}$ and $g_j = \exp_C(f_j)$ for $0 \leq j \leq n$ (note that $g_0 = 0$). By Theorem 4.2,

$$\exp_C[\omega_s \zeta_A(1; s)] \in \omega_s A[t_1, \dots, t_s], \quad s \geq 0.$$

In particular, g_1, \dots, g_n are tamely μ -algebraic over K_{s_0} (they satisfy linear μ -difference equations with coefficients in K_{s_0}). Conjecture 3.14 then implies that f_1, \dots, f_n are regularly μ -independent over L_{s_0} . Now, the theorem follows from the fact that $\omega_s \in L_{s_0}$ for all $s \leq s_0$ and that $K_{s_0} \subset L_{s_0}$. \square

The next Proposition goes in the direction of Theorem 4.3. The statement is so much weaker, but holds unconditionally and is deduced from an entirely different statement: Chang and Yu’s Theorem 4.1.

Proposition 4.4. *Assuming that $q = p$, the elements $\tilde{\pi}$ and $\zeta_A(1; s)$ for $0 \leq s \leq p - 1, s \neq 1$ are tamely μ -independent over K_s .*

It is easy to see that, if $s = 1$, $\zeta_A(1; 1)$ and $\tilde{\pi}$ are regularly μ -dependent. Indeed, by the formula

$$\zeta_A(1; 1) = \frac{\tilde{\pi}}{(\theta - t)\omega}$$

that can be found in [20], we see that the regular μ -polynomial

$$P(X_1, X_2) = (\theta^q - t)\mu^e(X_2) - X_1^{q-1}X_2 \in K_1[X_1, X_2]_\mu$$

satisfies $(\tilde{\pi}, \zeta(1; 1)) \in Z(P)$.

Proof of Proposition 4.4. Assume by contradiction that the statement is false. Then there exists a non-trivial relation $P(f_0, f_1, f_3, \dots, f_p) = 0$ with $P \in K(t_1, \dots, t_s)[\underline{X}]_\mu$ tame and non-zero, with $f_0 = \tilde{\pi}, f_1 = \zeta_A(1), f_3 = \zeta_A(1; 2), \dots, f_p = \zeta_A(1; p - 1)$ (this time, $\underline{X} = (X_0, X_1, X_3, \dots, X_{p-1})$ and note that X_2 is missing because it corresponds to $\zeta_A(1; 1)$ that we have discarded). Since the subset

$$\{(\theta^{p-k_1}, \dots, \theta^{p-k_{p-1}}); \underline{k} = (k_1, \dots, k_{p-1}) \in \mathbb{N}^{p-1}\} \subset \mathbb{C}_\infty^{p-1}$$

is Zariski-dense, there exists a subset \mathcal{I} of \mathbb{N}^{q-1} (of positive density) such that the evaluation $\text{ev}_{\underline{k}}(c_i)$ of a coefficient c_i of P at the point

$$(t_1, \dots, t_{p-1}) = (\theta^{p-k_1}, \dots, \theta^{p-k_{p-1}})$$

for all $\underline{k} = (k_1, \dots, k_{p-1}) \in \mathcal{I}$ is non-zero. We choose such a p -tuple. Further we have, for $m \geq 0, n > 0, 0 \leq s' \leq p - 1$ and $k := \max\{k_1, \dots, k_{p-1}\}$,

$$(4.4) \quad \begin{aligned} \text{ev}_{\underline{k}}(\mu^m(\zeta_A(n; s'))) &= \zeta_A(n - p^{-m-k_1} - \dots - p^{-m-k_{s'}})p^m \\ &= \zeta_A(p^{m+k}n - p^{k-k_1} - \dots - p^{k-k_{s'}})p^{-k}. \end{aligned}$$

Now, it is easy to see, due to our simple choice of parameters (especially the fact that $s' \leq p - 1$ and $s' \neq 1$), that our assumption yields a non-trivial algebraic dependence relation among $\tilde{\pi}$ and the values

$$\zeta_A \left(p^{m+k} - \sum_{i=1}^{s'} p^{k-k_i} \right), \quad m \geq 0, \quad 0 \leq s' \leq p - 1, s' \neq 1,$$

where $k := \max\{k_1, \dots, k_{p-1}\}$. First of all, we can choose $\underline{k} = (k_1, \dots, k_{p-1})$ so that, for all $i, k_i > 0$. This implies that for all m and s' , the integer

$$n_{m,s'} := p^{m+k} - \sum_{i=1}^{s'} p^{k-k_i}$$

is positive. Further, for all such m, s' , we have that $n_{m,s'} \not\equiv 0 \pmod{p-1}$; this already avoids the existence of a relation of the Euler–Carlitz-type (4.3). In order to avoid the degenerate sum shuffle relations (4.2) we need to show that $n_{m,s'} = n_{m',s''}p^j$ for $1 \leq s', s'' \leq p - 1$ and $j, m, m' \geq 0$ implies $m = m', j = 0$ and $s' = s''$. But assuming that an identity

$$p^{m+k} - \sum_{i=1}^{s'} p^{k-k_i} = p^j \left(p^{m'+k} - \sum_{i=1}^{s''} p^{k-k_i} \right)$$

holds with $j \geq 0$ implies $j = 0$. Indeed, there exists some i such that $k_i = k$, and the set of such indices i , non-empty, has cardinality $< p$, so that the left-hand side is not divisible by p . Note that the condition $s', s'' \leq p - 1$ is crucial. The conclusion follows from Chang and Yu’s Theorem 4.1. \square

In a similar vein, we have the next Lemma.

Lemma 4.5. *Supposing that $p = q > 2$, we have that*

$$f = \sum_{a \in A^+} a(t)^2 a^{-p} \in \mathbb{T}$$

is transformally μ -transcendental, hence regularly μ -transcendental over K_1 .

Proof. We assume by contradiction that there is a non-trivial relation of algebraic dependence

$$\sum_{\underline{i} \in \mathbb{N}^k} c_{\underline{i}} f^{i_0} \mu(f)^{i_1} \dots \mu^{k-1}(f)^{i_{k-1}}, \quad c_{\underline{i}} \in K_1$$

(the sum is finite). We can suppose, without loss of generality, that all the coefficients $c_{\underline{i}}$ are in $A[t]$ and that they are not all divisible by $t - \theta$. This means that the evaluation at $t = \theta$ of the coefficients $c_{\underline{i}}$ yields a non-zero vector with entries in A . Then observing that the evaluation at $t = \theta$ of $\mu^j(f)$ is equal to the Carlitz zeta value $\zeta_A(p^{j+1} - 2)$, we have thus a non-trivial relation of algebraic dependence with coefficients in A of the Carlitz zeta values $\zeta_A(p^j - 2)$ for $j = 1, \dots, k$. Since $p - 1, p \nmid p^j - 2$ for all $j > 0$, this is again in contradiction with Chang and Yu’s Theorem 4.1. \square

4.2.3. Two more conjectures. Additionally, we propose the following, for general q :

Conjecture 4.6. *Any finite subset of the set whose elements are $\tilde{\pi}, \zeta_A(n; s)$ with $p \nmid n$ and $n \not\equiv s \pmod{q - 1}$ is regularly μ -independent over K_s .*

This conjecture does not seem to be a consequence of Conjecture 3.14 and is probably also quite a difficult one. In order to present something which is perhaps provable in the near future, we mention that Conjecture 3.14 implies, for $n = 1$, the following conjecture:

Conjecture 4.7. *If $f \in \mathbb{K}_s \setminus \{0\}$, then either f or $\exp_C(f)$ is regularly μ -transcendental over K_s .*

The author presently does not know if this result can be directly deduced from the analogue of the theorem of Hermite–Lindemann for the Carlitz exponential over \mathbb{C}_∞ of [22].

5. Transcendence degree of difference subfields

Let $\underline{f} = (f_1, \dots, f_n)$ be an element of \mathbb{K}_s^n . We suppose that for all i , $g_i = \exp_C(f_i) \in \mathbb{K}_s$ are regularly μ -algebraic. We address the following loosely question.

Question 5.1. Assume that $s > 0$. Compute the transcendence degree over K_s of the μ -field

$$K_s(g_1, \dots, g_n)_\mu$$

from the knowledge of f_1, \dots, f_n .

Answering is likely to be difficult. The transcendence degree behaves wildly if $s > 0$ and does not seem to be in transparent (conjectural) relation with the linear forms over $\mathbb{F}_q(t)[\theta]$ satisfied by the various f , as we are going to see. From now on, we are going to assume that $s = 1$ for commodity. Therefore, we are going to write $t = t_1$, $\mathbb{T} = \mathbb{T}_1$ and $\mathbb{K} = \mathbb{K}_1$.

We recall that the \mathbb{C}_∞ -algebra \mathbb{T} is stable under the action of the divided derivatives \mathcal{D}_n , which are, for all $n \geq 0$, the \mathbb{C}_∞ -linear endomorphisms uniquely defined by the rule

$$\mathcal{D}_n t^m = \binom{n}{m} t^{m-n}.$$

Proposition 5.2 (Analogue of Hölder’s Theorem for the function ω). *We set $f_i = \frac{\tilde{\pi}}{(\theta-t)^{i+1}}$ for $i \geq 0$. The following properties hold:*

- (1) For all $i \geq 0$, $\exp_C(f_i) = \mathcal{D}_i(\omega)$.
- (2) The functions $\mathcal{D}_i(\omega)$ define the entries of a solution of the τ -difference system

$$\tau(X_i) = (t - \theta)X_i + X_{i-1}, \quad i \geq 0, \quad X_{-1} := 0.$$

- (3) The functions $\mathcal{D}_i(\omega)$ are algebraically independent over $K(t)$. More precisely we have, for all $n \geq 0$,

$$\begin{aligned} \text{transf deg}_{K(t)} K(t)(\mathcal{D}_i(\omega) : 0 \leq i \leq n)_\tau &= 0, \\ \text{tr deg}_{K(t)} K(t)(\mathcal{D}_i(\omega) : 0 \leq i \leq n)_\tau &= n + 1. \end{aligned}$$

Proof. The first two properties follow easily from the fact that, over \mathbb{T} ,

$$\exp_C \frac{d}{dt} = \frac{d}{dt} \exp_C,$$

and from the difference equation (3.2). For (3), since ω is transcendental over $K(t)$ we can argue by induction and suppose that the elements $\omega, \mathcal{D}_1(\omega), \dots, \mathcal{D}_{n-1}(\omega)$ are algebraically independent over $K(t)$. Let us suppose by contradiction that $\mathcal{D}_n(\omega)$ is algebraically dependent of the previous functions; then there exists a non-zero irreducible polynomial P of $K(t)[X_0, \dots, X_n]$ which vanishes at the point determined by the functions

$\omega, \dots, \mathcal{D}_n(\omega)$. The property (2) of the proposition implies that, if we denote by P^τ the polynomial of $K(t)[X_0, \dots, X_n]$ obtained by letting τ act on the coefficients, Then $P \mid P^\tau$ and in fact, $P^\tau = (t - \theta)^d P$ with d the total degree of P in its indeterminates, looking at the homogeneous part of highest degree of P . But looking at the monomials of smallest total degree of P , we see that this is impossible. The property concerning the transformal dependence is easy and left to the reader. \square

Hence, the linear relations over $\mathbb{F}_q(t)[\theta]$ of $f_1, \dots, f_n \in \mathbb{K}$ do not have much to say about the transcendence degree over $K(t)$ of the τ -difference field generated by $g_1 = \exp_C(f_1), \dots, g_n = \exp_C(f_n)$.

5.1. Solving linear difference equations in \mathbb{K} . We are going to discuss how the Carlitz exponential \exp_C can be used to solve the equations

$$\tau^{-1}(X) = X + g, \quad g \in \mathbb{K}.$$

Lemma 5.3. *Let g be an element of \mathbb{K} . Let $v \in \mathbb{K}$ be such that*

$$\exp_C(v) = -\tau(g)(t - \theta)\omega.$$

Then all the solutions $x \in \mathbb{K}$ of the equation

$$\tau^{-1}(X) = X + g$$

are the elements of the set

$$\mathcal{T}(g) := \omega^{-1} \exp_C\left(\frac{v}{\theta - t}\right) + \mathbb{F}_q(t).$$

Proof. All we need to show is that $f_v := \omega^{-1} \exp_C\left(\frac{v}{\theta - t}\right)$ is a solution of our equation. But:

$$\begin{aligned} \tau(f_v) &= \tau\left(\exp_C\left(\frac{v}{\theta - t}\right)\right) \tau(\omega)^{-1} \\ &= \left(C_{\theta - t}\left(\exp_C\left(\frac{v}{\theta - t}\right)\right) - (\theta - t) \exp_C\left(\frac{v}{\theta - t}\right)\right) \tau(\omega)^{-1} \\ &= \left(\exp_C(v) - (\theta - t) \exp_C\left(\frac{v}{\theta - t}\right)\right) ((t - \theta)\omega)^{-1} \\ &= f_v + \frac{\exp_C(v)}{(t - \theta)\omega} \\ &= f_v - \tau(g), \end{aligned}$$

and the equation $\tau(X) = X - \tau(g)$ has the same set of solutions as our equation. \square

Remark 5.4. We have that

$$\mathcal{T}(g) = \omega^{-1} \exp_C \frac{1}{\theta - t} \exp_C^{-1}((\theta - t)\omega\tau(g)).$$

We define two towers of field extensions of $K^{ac}(t)$, inductively. The first one is defined by $L_0 = K^{ac}(t)$ and, for all i ,

$$L_i = L_{i-1}(\omega)(f \in \mathbb{K} : \tau^{-1}(f) = f + \omega^l g : l \in \mathbb{Z}, g \in L_{i-1}).$$

For the second one, we set again, $M_0 = K^{ac}(t)$ and then inductively,

$$M_i = M_{i-1}(\cup_{l \in \mathbb{Z}} \exp_C(M_{i-1}\omega^l) \cup \cup_{l' \in \mathbb{Z}} \exp_C^{-1}(M_{i-1}\omega^{l'})).$$

Then writing $L_\infty = \cup_i L_i$ and $M_\infty = \cup_i M_i$, we obtain $L_\infty \subset M_\infty$ as a consequence of Lemma 5.3. In the next subsection, we are going to observe that the entries of the rigid analytic trivializations associated to *Chang's multiple polylogarithms* (as in [10]) are all in the field L_∞ .

5.2. Polylogarithm t -motives. An important feature of the entire function

$$\Omega := \tau(\omega)^{-1}$$

is highlighted in the paper of Anderson, Brownawell and Papanikolas [1] and in the paper of Papanikolas [18]: Ω is a *rigid analytic trivialization* of the *Carlitz dual t -motive C* . The reader can consult [18, §1.1.2 and §3.4] for the definition and the basic properties of *Anderson (dual) t -motives* (note that the operator τ^{-1} is used therein). The basic properties of the function Ω and of the Carlitz t -motive are discussed in [1, §3.1.2] and [18, §3.4.3, §3.3.4], while the notion of rigid analytic trivializations of t -motives is discussed in [18, §1.1.3 and §3.3].

In essence, the rigid analytic triviality of an object M of the Tannakian category \mathcal{T} over $\mathbb{F}_q(t)$ of *t -motives* (containing the category of Anderson's t -motives and introduced by Papanikolas, see [18, §1.1.6 and §3]) allows one to realize M as the solutions $\Psi \in \text{GL}_r(\mathbb{K})$ (for some r) of a linear τ^{-1} -difference system

$$(5.1) \quad \tau^{-1}(\Psi) = \Phi\Psi,$$

with $\Phi \in \text{GL}_r(\mathbb{C}_\infty(t))$.

Let Ψ be a rigid analytic trivialization associated to a multiple polylogarithm as in [10], that is, a matrix $\Psi \in \text{GL}_{d+1}(\mathbb{K})$ satisfying the τ^{-1} -difference system (5.1) with $\Phi \in \text{GL}_{d+1}(K^{ac}(t))$ as in [10, (5.3.3)] with $s_1, s_2, \dots, s_d \in \mathbb{Z}_{>0}$ and $Q_1, \dots, Q_d \in K^{ac}(t)$. We shall prove:

Proposition 5.5. *The entries of Ψ belong to L_∞ .*

Proof. If we write

$$\Psi = \begin{pmatrix} \Omega^{s_1+\dots+s_d} & 0 & \dots & 0 & 0 \\ x_{1,0} & \Omega^{s_2+\dots+s_d} & \dots & 0 & 0 \\ x_{2,0} & x_{2,1} & \dots & 0 & 0 \\ \dots & \vdots & & \vdots & \vdots \\ x_{d,0} & x_{d,1} & \dots & x_{d,d-1} & 1 \end{pmatrix},$$

solving the system (5.1) for this choice of Φ amounts to solve the iterated system of τ^{-1} -difference equations

$$(5.2) \quad \tau^{-1}(x_{i,j}) = \tau^{-1}(Q_i)(t - \theta)^{s_i + \dots + s_d} x_{i-1,j} + (t - \theta)^{s_{i+1} + \dots + s_d} x_{i,j},$$

$$0 \leq j < i \leq d,$$

setting also $x_{i,i} = \Omega^{s_{i+1} + \dots + s_d}$ for $i = 1, \dots, d$. This system is equivalent, by setting

$$y_{i,j} = \frac{x_{i,j}}{\Omega^{s_{i+1} + \dots + s_d}},$$

to the system of equations:

$$\tau^{-1}(y_{i,j}) = \tau^{-1}(Q_i)((t - \theta)\Omega)^{s_i} y_{i-1,j} + y_{i,j}, \quad 0 \leq j < i \leq d,$$

and the result follows. □

Before Schanuel’s conjecture, there was a conjectural statement by Gelfond (see appendix of Waldschmidt’s paper [25]) which looked as a very intricate statement involving iteration of exponentials and logarithms. This lets us appreciate the simplicity and the strength of Schanuel’s Conjecture. In what concerns, for example, a control of the transcendence degree of the field generated by the entries of a rigid analytic trivialization of a multiple polylogarithm motive, we are led to understand the algebraic relations between elements of the field M_∞ . It seems thus that we are somewhat back to Gelfond’s starting point.

Acknowledgements. The author is thankful to Laurent Denis and Paul Voutier for interesting and useful discussions, and Amador Martin-Pizarro for having suggested the references [17] and [21] and for several remarks that have helped to improve the paper. This research started while the author was visiting the MPIM (Bonn) in February 2016 and the IHES (Bures-sur-Yvette) in April 2016. The author is thankful to both institutions for the very good environment that positively influenced the quality of this work.

Appendix A. Some ring structures related to base- p digits

We begin with a commutative, unitary ring R . For a symbol Y , we introduce, in the R -module $R[Y_0, Y_1, \dots]$ of polynomials in infinitely many indeterminates Y_0, Y_1, \dots with coefficients in R , the monomials

$$\langle Y \rangle^i := \prod_{j=0}^{\infty} Y_j^{i_j}, \quad i \in \mathbb{N}, \quad i = i_0 + i_1 p + \dots + i_r p^r,$$

where the last expression is the expansion of i in base p , so that $0 \leq i_j \leq p - 1$ for all j (note that the product is finite for any i). For $i, j \in \mathbb{N}$, we set $k := i + j$ and:

$$\langle Y \rangle^i \cdot \langle Y \rangle^j := \langle Y \rangle^k.$$

the R -module

$$R\langle Y \rangle := \left\{ \sum_{i=0}^N c_i \langle Y \rangle^i : N \in \mathbb{N}, c_i \in R \right\}$$

is now equipped with the structure of an R -algebra, by using the product of monomials defined above. Explicitly, we have, for two polynomials $f = \sum_i f_i \langle Y \rangle^i$ and $g = \sum_j g_j \langle Y \rangle^j$,

$$f \cdot g = \sum_k \langle Y \rangle^k \sum_{i+j=k} f_i g_j$$

(note that $R\langle Y \rangle$ is a submodule but not a subring of $R[Y_0, Y_1, \dots]$). This means that $R\langle Y \rangle$ is isomorphic to the polynomial algebra $R[Z]$ in one indeterminate Z with coefficients in R . In particular, if R is a field, $R\langle Y \rangle$ is an R -algebra of dimension one.

Let R be an \mathbb{F}_p -algebra, together with an \mathbb{F}_p -linear injective endomorphism $R \xrightarrow{\mu} R$. We define a μ -difference structure over $R\langle Y \rangle$ by setting

$$\mu \left(\sum_i f_i \langle Y \rangle^i \right) := \sum_i \mu(f_i) \langle Y \rangle^{pi}.$$

It is easy to see that, in this way, the isomorphism $R\langle Y \rangle \cong R[Z]$ becomes a μ -difference algebra isomorphism.

We assume, from now on, that R is a commutative field extension of \mathbb{F}_q . Let \mathcal{P} be the ideal of $R[X]_\mu$ generated by the polynomials $X^{p^k} - \mu^k(X)$, for all $k \geq 0$. Then \mathcal{P} is a μ -invariant ideal.

Lemma A.1. *The ideal \mathcal{P} is a prime ideal, and the quotient $R[X]_\mu/\mathcal{P}$ is isomorphic, as an R -algebra which also is a μ -difference algebra, to the R -algebra $R\langle Y \rangle$.*

Proof. All we need to show, is that there is an R -algebra isomorphism $R[Z] \xrightarrow{\phi} R[X]_\mu/\mathcal{P}$. Indeed, it follows in this way that $R[X]_\mu/\mathcal{P}$ is integral.

We construct ϕ in the following way. Let $i \in \mathbb{N}$ be with base- p expansion $i = i_0 + i_1p + \dots + i_r p^r$, with $0 \leq i_j < p$ for all j . We define the map ϕ to be R -linear, with

$$\phi(Z^i) = X^{i_0} \mu(X)^{i_1} \dots \mu^r(X)^{i_r} \pmod{\mathcal{P}_X}.$$

We need to show that ϕ is multiplicative. Let $i, j, k \in \mathbb{N}$ be such that $i + j = k$. We further expand i, j, k in base p ; let r be an integer such that $i = \sum_{n=0}^r i_n p^n, j = \sum_{n=0}^r j_n p^n, k = \sum_{n=0}^r k_n p^n$, with $0 \leq i_n, j_n, k_n < p$. We define the sequence $(b_n)_{n \geq 0}$ of integers to be that characterizing the carry over of the base- p expansion of the sum $i + j = k$, namely, $b_n \in \mathbb{N}$ is defined by $b_{-1} := 0$ and, inductively for $n \geq 0$, by:

$$i_n + j_n + b_{n-1} = k_n + p b_n, \quad n \geq 0.$$

We have:

$$\begin{aligned}
& \phi(Z^i)\phi(Z^j) \\
&= X^{i_0+j_0}\mu(X)^{i_1+j_1}\dots\mu^r(X)^{i_r+j_r} \\
&\equiv X^{k_0}\mu(X)^{i_1+j_1+b_0}\mu^2(X)^{i_2+j_2}\dots\mu^r(X)^{i_r+j_r} \pmod{(X^p-\mu(X))} \\
&\equiv X^{k_0}\mu(X)^{k_1}\mu^2(X)^{i_2+j_2+b_1}\dots\mu^r(X)^{i_r+j_r} \pmod{(X^p-\mu(X),\mu(X)^p-\mu^2(X))} \\
&\equiv X^{k_0}\mu(X)^{k_1}\mu^2(X)^{i_2+j_2+b_1}\dots\mu^r(X)^{i_r+j_r} \pmod{(X^p-\mu(X),X^{p^2}-\mu^2(X))} \\
&\vdots \\
&\equiv X^{k_0}\mu(X)^{k_1}\dots\mu^r(X)^{k_r} \pmod{\mathcal{P}_X} \\
&= \phi(Z^k)
\end{aligned}$$

so that ϕ is multiplicative, and it is obviously surjective and injective (this is also easy to verify on the structure of the ring $R\langle Y \rangle$). The verification of the remaining properties is left to the reader. \square

References

- [1] G. W. ANDERSON, W. D. BROWNAWELL & M. A. PAPANIKOLAS, “Determination of the algebraic relations among special Γ -values in positive characteristic”, *Ann. Math.* **160** (2004), no. 1, p. 237-313.
- [2] B. ANGLÈS & F. PELLARIN, “Functional identities for L -series values in positive characteristic”, *J. Number Theory* **142** (2014), p. 223-251.
- [3] ———, “Universal Gauss-Thakur sums and L -series”, *Invent. Math.* **200** (2015), no. 2, p. 653-669.
- [4] B. ANGLÈS, F. PELLARIN & F. TAVARES RIBEIRO, “Arithmetic of positive characteristic L -series values in Tate algebras”, *Compos. Math.* **152** (2016), no. 1, p. 1-61, with an appendix of Florent Demeslay.
- [5] ———, “Anderson-Stark units for $\mathbb{F}_q[\theta]$ ”, *Trans. Am. Math. Soc.* (2017), <https://doi.org/10.1090/tran/6994> (to appear in print).
- [6] J. AX, “On Schanuel’s Conjectures”, *Ann. Math.* **93** (1971), p. 252-268.
- [7] F. BEUKERS, “A refined version of the Siegel-Shidlovskii theorem”, *Ann. Math.* **163** (2006), no. 1, p. 369-379.
- [8] S. BOSCH, U. GÜNTZER & R. REMMERT, *Non-Archimedean analysis. A systematic approach to rigid analytic geometry*, Grundlehren der Mathematischen Wissenschaften, vol. 261, Springer, 1984, xii+436 pages.
- [9] L. CARLITZ, “On certain functions connected with polynomials in a Galois field”, *Duke Math. J.* **1** (1935), p. 137-168.
- [10] C.-Y. CHANG, “Linear independence of monomials of multizeta values in positive characteristic”, *Compos. Math.* **150** (2014), no. 11, p. 1789-1808.
- [11] C.-Y. CHANG & J. YU, “Determination of algebraic relations among special zeta values in positive characteristic”, *Adv. Math.* **216** (2007), no. 1, p. 321-345.
- [12] L. DENIS, “Indépendance algébrique et exponentielle de Carlitz”, *Acta Arith.* **69** (1995), no. 1, p. 75-89.
- [13] ———, “Indépendance algébrique de logarithmes en caractéristique p ”, *Bull. Aust. Math. Soc.* **74** (2006), no. 3, p. 461-470.
- [14] D. GOSS, *Basic Structures of Function Field Arithmetic*, *Ergebnisse der Mathematik und ihrer Grenzgebiete.* 3, vol. 35, Springer, 1996, xiii+422 pages.
- [15] S. LANG, *Introduction to transcendental numbers*, Addison-Wesley Series in Mathematics, Addison-Wesley Publishing Company, 1966, vi+105 pages.

- [16] A. B. LEVIN, *Difference algebra*, Algebra and Applications, vol. 8, Springer, 2008, xi+519 pages.
- [17] D. MARKER, “A remark on Zilber’s pseudoexponentiation”, *J. Symb. Log.* **71** (2006), no. 3, p. 791-798.
- [18] M. A. PAPANIKOLAS, “Tannakian duality for Anderson-Drinfeld motives and algebraic independence of Carlitz logarithms”, *Invent. Math.* **171** (2008), no. 1, p. 123-174.
- [19] F. PELLARIN, “Aspects de l’indépendance algébrique en caractéristique non nulle”, in *Séminaire Bourbaki. Volume 2006/2007. Exposés 967–981*, Astérisque, vol. 317, Société Mathématique de France, 2008, p. 205-242, Exp no. 973.
- [20] ———, “On the generalized Carlitz module”, *J. Number Theory* **133** (2013), no. 5, p. 1663-1692.
- [21] T. SCANLON, “ o -minimality as an approach to the André-Oort conjecture”, To appear in *Panor. Synth.*
- [22] A. THIERY, “Indépendance algébrique des périodes et quasi-périodes d’un module de Drinfeld”, in *The arithmetic of function fields. Proceedings of the workshop at the Ohio State University*, Ohio State University Mathematical Research Institute Publications, vol. 2, Walter de Gruyter, 1992, p. 265-284.
- [23] P. M. VOUTIER, Letter to the author, June, 20, 2016.
- [24] L. I. WADE, “Certain quantities transcendental over $\text{GF}(p^n, x)$ ”, *Duke Math. J.* **8** (1941), p. 701-720.
- [25] M. WALDSCHMIDT, “Schanuel’s Conjecture: algebraic independence of transcendental numbers”, in *Colloquium De Giorgi 2013 and 2014*, Colloquia, vol. 5, Edizioni della Normale, xv+137, p. 129-137.
- [26] J. YU, “Analytic homomorphisms into Drinfeld modules”, *Ann. Math.* **145** (1997), no. 2, p. 215-233.

Federico PELLARIN
Institut Camille Jordan
UMR 5208 Site de Saint-Etienne
23 rue du Dr. P. Michelon
42023 Saint-Etienne, France
E-mail: federico.pellarin@univ-st-etienne.fr