

The Bramson delay in the non-local Fisher-KPP equation

Emeric Bouin ^a, Christopher Henderson ^{b,*}, Lenya Ryzhik ^c

^a CEREMADE - Université Paris-Dauphine, UMR CNRS 7534, Place du Maréchal de Lattre de Tassigny, 75775 Paris Cedex 16, France

^b Department of Mathematics, University of Arizona, Tucson, AZ 85721, United States of America

^c Department of Mathematics, Stanford University, Stanford, CA 94305, United States of America

Received 9 October 2017; received in revised form 6 March 2019; accepted 13 July 2019

Available online 30 July 2019

Abstract

We consider the non-local Fisher-KPP equation modeling a population with individuals competing with each other for resources with a strength related to their distance, and obtain the asymptotics for the position of the invasion front starting from a localized population. Depending on the behavior of the competition kernel at infinity, the location of the front is either $2t - (3/2) \log t + O(1)$, as in the local case, or $2t - O(t^\beta)$ for some explicit $\beta \in (0, 1)$. Our main tools here are a local-in-time Harnack inequality and an analysis of the linearized problem with a suitable moving Dirichlet boundary condition. Our analysis also yields, for any $\beta \in (0, 1)$, examples of Fisher-KPP type non-linearities f_β such that the front for the local Fisher-KPP equation with reaction term f_β is at $2t - O(t^\beta)$.

© 2019 L'Association Publications de l'Institut Henri Poincaré. Published by Elsevier B.V. All rights reserved.

Résumé

Dans cet article, nous considérons l'équation de Fisher-KPP non locale, qui modélise la dynamique d'une population où la force de compétition pour les ressources dépend de la distance entre les individus. Nous obtenons une asymptotique précise en temps long de la position d'une population qui est initialement localisée en espace. Selon la décroissance à l'infini du noyau de compétition, la position du front est soit $2t - (3/2) \log t + O(1)$, comme dans le cas de l'équation locale, soit $2t - O(t^\beta)$, pour un $\beta \in (0, 1)$ calculé explicitement. Les outils les plus importants utilisés dans cet article sont une version locale en temps d'une inégalité de Harnack parabolique ainsi qu'une analyse fine du problème linéarisé avec une condition de bord de Dirichlet dynamique. Notre analyse donne aussi, pour tout $\beta \in (0, 1)$, des exemples de non-linéarités de type Fisher-KPP pour lesquelles le front se trouve en $2t - O(t^\beta)$.

© 2019 L'Association Publications de l'Institut Henri Poincaré. Published by Elsevier B.V. All rights reserved.

MSC: 35K57; 35Q92; 45K05; 35C07

Keywords: Reaction-diffusion equations; Logarithmic delay; Parabolic Harnack inequality

* Corresponding author.

E-mail addresses: bouin@ceremade.dauphine.fr (E. Bouin), ckhenderson@math.arizona.edu (C. Henderson), ryzhik@math.stanford.edu (L. Ryzhik).

1. Introduction

The Fisher-KPP equation

$$u_t = u_{xx} + u(1 - u) \quad (1.1)$$

is one of the simplest models for population spreading, accounting for a competition for resources. However, (1.1) only accounts for a local competition between individuals. When this competition is non-local, one is led to the non-local Fisher-KPP equation

$$\begin{aligned} u_t - u_{xx} &= u(1 - \phi \star u), \quad t > 0, \ x \in \mathbb{R}, \\ u(0, \cdot) &= u_0. \end{aligned} \quad (1.2)$$

Here, ϕ is a probability density that represents the strength of the competition between individuals a given distance apart. Equation (1.2) has garnered much interest recently, mostly for two reasons. First, it does not admit a comparison principle, leading to inherent technical difficulties – even proving a uniform upper bound on u is non-trivial [19]. Second, unusual behavior may occur, such as the existence of oscillating wave trains behind the front [12–14,21].

Our interest is in the spreading of the solutions of (1.2) when the initial density u_0 is localized. To motivate our work, we recall the known results for the local Fisher-KPP equation (1.1). Going back to the work of Bramson, it is known that if u_0 is compactly supported, the front of u is located at

$$X(t) = 2t - \frac{3}{2} \log t + s_0, \quad (1.3)$$

where s_0 is a shift depending only on u_0 [5,6], with less precise asymptotics obtained earlier by Uchiyama [27]. These proofs have been simplified in recent years [17,26], with some refinements in [23,24], and also extended to the spatially periodic case [18]. One may think of $\bar{X}(t) = 2t$ as the position of a traveling wave, and $d(t) = (3/2) \log t$ as the delay due to the fact that the initial condition u_0 is compactly supported, so that the solution lags behind the traveling wave.

In the non-local case considered in the present paper, we show that the front position depends on the rate of decay of the kernel ϕ at infinity. When ϕ decays fast enough, solutions of (1.2) spread as those of the local equation: the front is at a position as in (1.3), up to a constant order error. However, when ϕ decays slowly, and the competition at large distances is relatively strong, the delay behind the traveling wave position $2t$ is not logarithmic but algebraic, of the order $O(t^\beta)$, with β depending only on the rate of decay of ϕ .

We now make our assumptions more precise. First, we assume that ϕ is an even, continuous, and bounded probability density:

$$\int_{\mathbb{R}} \phi(x) dx = 1, \quad \text{and} \quad \phi(x) = \phi(-x) \text{ for all } x \in \mathbb{R}. \quad (1.4)$$

In addition, ϕ has some “mass” near the origin, that is, there exists $\sigma_\phi > 0$ such that

$$\phi(x) \geq \sigma_\phi \mathbb{1}_{[-\sigma_\phi, \sigma_\phi]}. \quad (1.5)$$

The behavior of u depends strongly on the tail behavior of ϕ . Here we make two different assumptions, that are helpful for the upper and lower bounds, respectively. The first assumption, an upper bound on the tail of ϕ , is that there exists $A_\phi > 0$ and $r > 1$ such that, for all $R \geq 1$,

$$\int_R^\infty \phi(x) dx \leq A_\phi R^{-r+1}. \quad (1.6)$$

Sometimes we will need to complement this with a lower bound on the tail: for all $R \geq 1$, we have

$$\int_R^\infty \phi(x) dx \geq A_\phi^{-1} R^{-r+1}. \quad (1.7)$$

Roughly, (1.6) and (1.7) mean that $\phi \sim x^{-r}$ for $x \gg 1$.

For the initial condition, we assume that u_0 is localized to the left of some point x_0 :

$$0 \leq u_0 \leq 1, \quad \exists x_0 \text{ such that } u_0(x) = 0 \text{ for all } x \geq x_0, \quad \text{and} \quad \liminf_{x \rightarrow -\infty} u_0(x) > 0. \quad (1.8)$$

We expect our results to hold when u_0 has “fast” exponential decay, that is, $u_0(x)e^{(1+\epsilon)x} \rightarrow 0$ as $x \rightarrow 0$ for some $\epsilon > 0$, rather than compactly supported on the right. However, we recall that the front position asymptotics for solutions of (1.1) with u_0 that has a sufficiently slow exponential tail on the right is different from (1.3), see [5,6].

The main result of this paper is the following.

Theorem 1.1. *Suppose that u satisfies (1.2) and (1.8) with ϕ satisfying (1.4), (1.5), and (1.6). If $r > 3$, then the solution u propagates with a logarithmic delay:*

$$\liminf_{t \rightarrow \infty} \inf_{x \leq 0} u\left(t, 2t - \frac{3}{2} \log t + x\right) > 0, \quad (1.9)$$

and

$$\lim_{L \rightarrow \infty} \limsup_{t \rightarrow \infty} \sup_{x \geq L} u\left(t, 2t - \frac{3}{2} \log t + x\right) = 0. \quad (1.10)$$

If $r = 3$, then the solution u propagates with a larger logarithmic delay: there exists $S_\phi > s_\phi > 3/2$ such that

$$\liminf_{t \rightarrow \infty} \inf_{x \leq 0} u\left(t, 2t - S_\phi \log t + x\right) > 0, \quad (1.11)$$

and

$$\lim_{t \rightarrow \infty} \sup_{x \geq 0} u\left(t, 2t - s_\phi \log t + x\right) = 0. \quad (1.12)$$

If $r \in (1, 3)$, then the delay is algebraic: there exists $C_\phi > 0$, depending only on r , σ_ϕ , and A_ϕ , such that

$$\liminf_{t \rightarrow \infty} \inf_{x \leq 0} u\left(t, 2t - C_\phi t^{\frac{3-r}{1+r}} + x\right) > 0, \quad (1.13)$$

and, if additionally (1.7) holds, then there exists $c_\phi \in (0, C_\phi)$, depending only on σ_ϕ , r , and A_ϕ , such that

$$\lim_{t \rightarrow \infty} \sup_{x \geq 0} u\left(t, 2t - c_\phi t^{\frac{3-r}{1+r}} + x\right) = 0. \quad (1.14)$$

As we discuss later in greater detail, heuristically, the competition term $\phi \star u$ acts on the scale t^γ , with $\gamma = 2/(1+r)$. Note that

$$\frac{3-r}{1+r} = 2\gamma - 1, \quad (1.15)$$

and that, when $r > 3$, $\gamma < 1/2$, which, in turn, suggests that the competition scale is smaller than the diffusive scale \sqrt{t} . This is one way to see that there is a phase transition at $r = 3$.

As a by-product of our analysis, we also obtain results for the local Fisher-KPP equation

$$u_t = u_{xx} + f(u). \quad (1.16)$$

Let us assume that f is of the KPP class: $f(u)/u$ is decreasing in u near 0, $f \in C^1$, and $f'(0) = 1$. A natural question is whether these assumptions are sufficient to ensure that the front location is given by the logarithmic Bramson correction in (1.3). We show, roughly, the following: if

$$1 - \frac{f(u)}{u} \sim \left(\log \left(\frac{1}{u} \right) \right)^{1-r} \quad \text{with } r > 1,$$

then the conclusion of Theorem 1.1 holds, with the logarithmic delay for $r \geq 3$ and an algebraic delay of the order $O(t^{(3-r)/(1+r)})$ for $1 < r < 3$. These non-linearities are not purely mathematical curiosities: they are regularly used in

biology and are known as Gompertz models, see [8] and the vast body of literature around it. The statement and proof of this result are contained in Section 6.

Let us mention a few related works. The model (1.2) considered here was first introduced by Britton [7] and has a quite involved history, see the introduction of [1] for a brief overview. The non-local term $\phi \star u$ has different effects depending on whether one is studying the behavior of u behind the front or at the front. Behind the front, there is a possible Turing instability of the steady state of the local Fisher-KPP equation $u \equiv 1$, which complicates the behavior. For example, wave trains have been constructed by Faye and Holzer [12] and, in a related setting, in [21]. Such wave trains have also been observed numerically by Genieys, Volpert, and Auger in [13]. As a result, without finer assumptions on ϕ , one cannot hope for a stronger result than the lower bounds in Theorem 1.1. As far as the behavior at the front is concerned, the main result in this direction is that traveling waves of speed $c = 2$ exist [10, 14] and solutions to the Cauchy problem with compact initial data or which satisfy (1.8) propagate with speed $c(t) = 2 + o(1)$ as $t \rightarrow +\infty$ [19].

As far as algebraic delays are concerned, we point to the work of Fang and Zeitouni [11] and Maillard and Zeitouni [20], as well as [22] where a Fisher-KPP model with a diffusivity that changes slowly in time was studied, and a delay, roughly, of order $t^{1/3}$ was obtained. However, both the set-up and the mechanism for the large delay are quite different in these papers than in the present work. Finally, we also mention the recent paper of Ducrot [9] in which he constructs a class of non-linearities $f(x, u)$, which tend to $u(1 - u)$ as $|x| \rightarrow \infty$, such that if the nonlinearity $u(1 - u)$ in (1.1) is replaced by $f(x, u)$, then the front is at $2t - \lambda \log(t)$ for any $\lambda \geq 3/2$.

While in the final stages of preparing this paper, we learned of a very recent probabilistic study of the delay term by Penington [25]. In our notation, she obtains the log delay up to an error term $O(\log \log(t))$, when $r > 3$, and an algebraic delay $t^{(3-r)/(1+r) \pm \epsilon}$ for any $\epsilon > 0$ for $r \in (1, 3)$. Penington's assumptions on ϕ are the same as ours when $r \geq 3$. However, her assumptions are weaker when $r \in (1, 3)$: the $R^{-(r-1)}$ term in (1.6) is replaced by $R^{-(r-1)/2}$, at the expense of a slightly less precise power in the correction. The proofs in [25] are probabilistic, involving the Feynman-Kac formula and an in-depth study of the trajectories of Brownian motion. Overall, our work and [25] are quite different and reveal different features of the equation.

1.1. Heuristics and methods of proof

The upper bound (1.10) is obtained by a rather direct adaptation of the arguments in [17]. Let us outline a heuristic argument leading to the upper bound (1.14) for $r \in (1, 3)$. It also explains how the exponent $(3 - r)/(1 + r)$ comes about. Let the front have a delay $d(t)$ behind $2t$, so that

$$\inf_{x \leq 2t - d(t)} u(t, x) \geq \delta_0, \quad (1.17)$$

with some $\delta_0 > 0$. We expect that the solution looks like an exponential to the right of $x = 2t - d(t)$ and until the “front edge” at $x = 2t + e(t)$:

$$u(t, x) \sim \exp\{-(x - 2t + d(t))\}, \quad \text{for } x \in (2t - d(t), 2t + e(t)). \quad (1.18)$$

The diffusive Gaussian decay dominates the exponential “traveling wave” decay for $x > 2t + e(t)$. Using (1.17) and then (1.7), one may estimate $\phi \star u(t, x)$ when $x \in (2t - d(t), 2t + e(t))$ as

$$\phi \star u(t, x) \geq \delta_0 \int_{-\infty}^{2t - d(t)} \phi(x - y) dy \gtrsim (x - (2t - d(t)))^{1-r} \gtrsim (e(t) + d(t))^{1-r}.$$

Thus, in order for the exponential in (1.18) to be a super-solution to (1.2) inside $(2t - d(t), 2t + e(t))$, we need

$$(e(t) + d(t))^{1-r} \gtrsim d'(t). \quad (1.19)$$

We also need the exponential to be above $u(t, x)$ at the front edge. To control u there, we use that, letting $h = e^{-t}u$, h is a sub-solution to the heat equation. In other words,

$$h_t \leq h_{xx}$$

and, hence, for all $x \gg 1$,

$$e^{-t}u(t, x) = h(t, x) \leq \int e^{-\frac{|x-y|^2}{4t}} u_0(y) dy \lesssim \frac{\sqrt{t}}{x} e^{-\frac{x^2}{4t}}. \quad (1.20)$$

Thus, for u to sit below the exponential super-solution at $x = 2t + e(t)$, we require

$$\exp\left\{t - \frac{(2t + e(t))^2}{4t}\right\} \lesssim \exp\{-(e(t) + d(t))\},$$

that is,

$$e(t)^2 \geq 4td(t). \quad (1.21)$$

Since $e(t)$ should be $o(t)$, we get

$$\lim_{t \rightarrow +\infty} \frac{d(t)}{e(t)} = 0. \quad (1.22)$$

Combining (1.19), (1.21) and (1.22) gives, for t large,

$$d'(t) \lesssim e(t)^{1-r} \lesssim t^{\frac{1-r}{2}} d(t)^{\frac{1-r}{2}},$$

and thus necessarily

$$d(t) \lesssim t^{(3-r)/(1+r)}.$$

We deduce also $e(t) \gtrsim t^\gamma$, with γ as in (1.15).

A way to estimate the solution from below, to get the lower bounds, is to study the linearized Fisher-KPP equation with a Dirichlet boundary condition at $2t + e(t)$, as in [17]. The problem that comes up after removing the exponential factor is

$$\begin{aligned} z_t &= z_{xx} + e'(t)(z_x - z), \quad t > 0, x > 0, \\ z(t, 0) &= 0. \end{aligned} \quad (1.23)$$

Once again, the case $r > 3$ is treated similarly to [17]. In particular, while the term $e'(t)z$ is important and is responsible for the $3/2$ pre-factor in the logarithmic correction, the drift $e'(t)z_x$ is negligible. Roughly, we estimate $z(t, x)$ at $x \sim \sqrt{t}$, and use a “tracing back to a shifted traveling wave” argument, to construct a sub-solution for u .

When $r < 3$, we choose $e(t) = t^\gamma$. Since now $\gamma > 1/2$, the drift $e'(t)z_x$ can no longer be neglected, and the choice of the exact exponent γ is necessary to get matching asymptotics. We explicitly construct a sub-solution of u to estimate the solution at the far edge, and then perform a “tracing back” argument with a traveling wave.

Lastly, in the case when $r = 3$, the diffusive scale and the induced drift have the same order. Here, the balance of these two scales causes a somewhat larger delay.

1.2. The local in time Harnack inequality

The main tool that allows us to get “reasonably sharp” asymptotics for the front position is a local-in-time Harnack inequality that is of an independent interest.

Proposition 1.2. *Suppose that $u \in L^\infty([0, T] \times \mathbb{R})$ is a non-negative function that solves*

$$u_t = u_{xx} + c(t, x)u,$$

on $[0, T] \times \mathbb{R}$ with $c \in L^\infty([0, T] \times \mathbb{R})$ and $T > 0$. Then, for any $p \in (1, \infty)$, there exist positive constants α , β , and C , that depend only on $\|c\|_{L^\infty([0, T] \times \mathbb{R})}$ and p , such that, for all $x, y \in \mathbb{R}$ and $t \in (0, T]$, we have

$$u(T, x + y) \leq C \|u\|_{L^\infty([t, T] \times \mathbb{R})}^{1 - \frac{1}{p}} u(T, x)^{\frac{1}{p}} e^{\alpha t + \frac{\beta y^2}{t}}. \quad (1.24)$$

This inequality is an indispensable tool to obtain “reasonably sharp” results for non-local problems. We have used a less precise form of it to obtain the logarithmic delay for solutions of the cane toads equation in [4], and it has also been used to establish a precise lower bound on the propagation speed of solutions of a Keller-Segel-Fisher system [16]. As

far as we know, [4] is the only other non-local context where a delay asymptotics has been established. It allows us to bound solutions of the non-local Fisher-KPP equation (1.2) in terms of the solutions of a local Fisher-KPP equation with a local time-dependent nonlinearity $g(t, u)$, that is logarithmic in u (Gompertz type). This equation has inherent difficulties coming from the time dependence and the logarithmic behavior near zero, but it is much more tractable because it admits a comparison principle.

The rest of the paper is organized as follows. In Section 2, we present the proofs of the upper bounds (1.10) and (1.14). Section 3 is where the proofs of the lower bounds (1.9), (1.11) and (1.13) are given. In order to complete the proof of the lower bounds, some estimates on linearized problems with moving Dirichlet boundary conditions are obtained in Section 4 and Section 5. In Section 6, we state and prove the result concerning the local KPP equation with logarithmic nonlinearity. The Harnack inequality is proved in Section 7.

Acknowledgments The authors thank Nicolas Champagnat for the reference [8]. EB was supported by “INRIA Programme Explorateur”. LR was supported by NSF grants DMS-1311903 and DMS-1613603. Part of this work was performed within the framework of the LABEX MILYON (ANR-10-LABX-0070) of Université de Lyon, within the program “Investissements d’Avenir” (ANR-11-IDEX-0007) operated by the French National Research Agency (ANR). In addition, CH has received funding from the European Research Council (ERC) under the European Unions Horizon 2020 research and innovation program (grant agreement No 639638) and was partially supported by the NSF RTG grant DMS-1246999 and DMS-1907853.

2. Upper bounds on the location of the front

In this section, we prove the upper bounds (1.10) and (1.14) in Theorem 1.1.

2.1. The upper bound when $r > 3$

The case $r > 3$ is very close to the local Fisher-KPP equation. The $(3/2) \log t$ delay is the best case scenario – in fact, the delay has to be at least that large for any r , so the bound is a quite straightforward application of bounds obtained in [17].

Proof of (1.10). Take $t_0 > 0$ to be determined later. Working in the moving frame with the logarithmic correction, the function

$$u_{\text{mov}}(t, x) = u\left(t, 2t - \frac{3}{2} \log\left(1 + \frac{t}{t_0}\right) + x\right),$$

satisfies

$$\begin{aligned} (u_{\text{mov}})_t &\leq \left(2 - \frac{3}{2} \frac{1}{t + t_0}\right) (u_{\text{mov}})_x + (u_{\text{mov}})_{xx} + u_{\text{mov}}, & \text{for all } t > 0, x \in \mathbb{R}, \\ u_{\text{mov}}(0, x) &= u_0(x), & \text{for all } x \in \mathbb{R}. \end{aligned}$$

We construct a super-solution \bar{u} as in [17]. Let \bar{v} be the solution to the boundary value problem

$$\begin{aligned} \bar{v}_t &= \left(2 - \frac{3}{2} \frac{1}{t + t_0}\right) \bar{v}_x + \bar{v}_{xx} + \bar{v}, & \text{for all } t > 0 \text{ and } x > 0, \\ \bar{v}(t, 0) &= 0, & \text{for all } t > 0, \\ \bar{v}(0, x) &= \mathbb{1}_{(0,2)}(x) & \text{for all } x > 0. \end{aligned}$$

Then [17, Lemma 2.1] implies that, provided that t_0 is sufficiently large, there exists $A_0 \geq 1$ such that for all $t \geq 0$, we have

$$\bar{v}(t, 1) \geq A_0^{-1}.$$

We also have the following uniform bound on the solutions to (1.2).

Lemma 2.1. [19, Theorem 1.2] Suppose that u satisfies (1.2) with initial data u_0 satisfying (1.8). Then there exists $M > 0$ such that, $u(t, x) \leq M$ for all $t > 0$ and $x \in \mathbb{R}$.

Let us now define $\bar{u}(t, x)$ as

$$\bar{u}(t, x) = M \left(\mathbf{1}_{x \leq x_0} + \min \left(1, A_0 \bar{v}(t, x - x_0 + 1) \right) \mathbf{1}_{x \geq x_0} \right),$$

where M is as in Lemma 2.1. By construction, \bar{u} is a super-solution to u_{mov} , and by our assumptions on u_0 (1.8), we also have $\bar{u}(0, x) \geq u_{\text{mov}}(0, x)$ for all $x \in \mathbb{R}$. In addition, [17, Lemma 2.1] implies that there exists T_0 such that, for all z and all $t \geq T_0$,

$$\bar{v}(t, z) \leq A_0 z e^{-z}. \quad (2.1)$$

We are now in a position to conclude the proof. Indeed, as $u \leq \bar{u}$, the upper bound in (2.1) implies

$$\begin{aligned} \limsup_{L \rightarrow \infty} \limsup_{t \rightarrow \infty} \sup_{x \geq L} u \left(t, 2t - \frac{3}{2} \log t + x \right) &= \limsup_{L \rightarrow \infty} \limsup_{t \rightarrow \infty} \sup_{x \geq L} u_{\text{mov}}(t, x) \\ &\leq \limsup_{L \rightarrow \infty} \limsup_{t \rightarrow \infty} \sup_{x \geq L} \bar{u}(t, x) \leq \lim_{L \rightarrow \infty} M A_0 L e^{-L} = 0, \end{aligned} \quad (2.2)$$

which concludes the proof. \square

2.2. The upper bound when $r = 3$

In this section, we show how to derive the upper bound on the location of the front assuming the lower bound on the location of the front. In other words, we prove (1.12) assuming (1.11), which we prove in the next section.

Proof of (1.12) assuming (1.11). Our proof proceeds similarly as in the previous subsection. Set $\tilde{s}_\phi < S_\phi$ to be determined. Using (1.11) and (1.7), we find $L > 0$ such that, for all $x \geq 0$ and $t \geq L$,

$$\phi \star u(t, x + 2t - \tilde{s}_\phi \log(t)) \geq \frac{1}{L A_\phi} (x + (S_\phi - \tilde{s}_\phi) \log(t) + L)^{-2}. \quad (2.3)$$

Next, we use the following result that is proved in Appendix A.

Lemma 2.2. *There exists v , $\tilde{s}_\phi > 3/2$, and L such that*

$$\begin{aligned} v_t &\geq v_{xx} + v(1 - v(t, x - (2t - \tilde{s}_\phi \log(t + t_0))))), & t > L, x > 2t - s_\phi \log(t + t_0) + L, \\ v(L, x) &\geq u(L, x), & x > 2L - \tilde{s}_\phi \log(L + t_0) + L, \end{aligned} \quad (2.4)$$

$v(t, L + 2t - \tilde{s}_\phi \log(t)) \geq M + 1$ for all $t \geq L$, and $v(t, x + 2t - \tilde{s}_\phi \log(t)) \rightarrow 0$ as $x \rightarrow \infty$ uniformly in $t \geq L$.

With Lemma 2.2 in hand, we now conclude. Notice that, (2.3) implies that v is a super-solution of u in $\{(t, x) \in [L, \infty) \times \mathbb{R} : x \geq 2t - \tilde{s}_\phi \log(t) + L\}$. Let

$$\bar{u}(t, x) = \begin{cases} M + 1, & \text{if } x \leq 2t - s_\phi \log(t) + L, \\ \min\{M + 1, v(t, x)\} & \text{if } x \geq 2t - s_\phi \log(t) + L. \end{cases}$$

As in the previous case $r > 3$, the comparison principle implies that $\bar{u} \geq u$ on $[L, \infty) \times \mathbb{R}$. The result then follows taking $s_\phi \in (\tilde{s}_\phi, 3/2)$. \square

2.3. The upper bound when $r \in (1, 3)$

In this section, we show how to derive the upper bound on the location of the front from the lower bound on the location of the front. In other words, we prove (1.14) assuming (1.13), which we prove in the next section.

Proof of the upper bound (1.14) assuming the lower bound (1.13). Note that, by (2.2), we have

$$\lim_{t \rightarrow \infty} \sup_{x \geq 2t + t^\gamma} u(t, x) = 0.$$

As a consequence, taking into account (1.15), it suffices to show that

$$\lim_{t \rightarrow \infty} \sup_{x \in (2t - c_\phi t^{2\gamma-1}, 2t + t^\gamma)} u(t, x) = 0.$$

We do this by creating a relevant super-solution to u on the interval $(2t - c_\phi t^{2\gamma-1}, 2t + t^\gamma)$. Note that the constant c_ϕ is still to be determined at this stage. Define, for any $T > 0$ and C_ϕ as in (1.13), the space-time domain (recall that $\gamma > 1/2$ for $1 < r < 3$):

$$\mathcal{P}_T := \left\{ (t, x) : t \in (T, \infty), x \in (2t - C_\phi t^{2\gamma-1}, 2t + t^\gamma) \right\},$$

and, for $(t, x) \in \mathcal{P}_T$, the function

$$\bar{v}(t, x) := B \exp \left\{ - \left(x - 2t + 2c_\phi t^{2\gamma-1} \right) \right\}.$$

On \mathcal{P}_T , the function \bar{v} satisfies

$$\bar{v}_t = \bar{v}_{xx} + \bar{v} \left(1 - 2c_\phi (2\gamma - 1) t^{\gamma(1-r)} \right). \quad (2.5)$$

The rest of the proof is devoted to showing that u is, indeed, a subsolution to (2.5) when the various constants above are suitably chosen: specifically, we show that

$$u_t - u_{xx} - u(1 - 2c_\phi (2\gamma - 1) t^{\gamma(1-r)}) \leq 0 \text{ in } \mathcal{P}_T, \quad (2.6)$$

and

$$u(t, x) \leq \bar{v}(t, x), \quad \text{on } \partial \mathcal{P}_T. \quad (2.7)$$

First, we show that (2.6) holds. It follows from (1.13) that there exist C_ϕ and δ_ϕ , depending only on ϕ , and T_0 such that, for all $t \geq T_0$,

$$\inf_{x \leq 2t - C_\phi t^{2\gamma-1}} u(t, x) \geq \delta_\phi. \quad (2.8)$$

Using (2.8), we can estimate $\phi \star u$ from below, for $t \geq T_0$ and $x > 2t - C_\phi t^{2\gamma-1}$:

$$\begin{aligned} \phi \star u(t, x) &= \int_{\mathbb{R}} \phi(x - y) u(t, y) dy \geq \int_{-\infty}^{2t - C_\phi t^{2\gamma-1}} \phi(x - y) u(t, y) dy \\ &\geq \delta_\phi \int_{-\infty}^{2t - C_\phi t^{2\gamma-1}} \phi(x - y) dy = \delta_\phi \int_{x - 2t + C_\phi t^{2\gamma-1}}^{+\infty} \phi(z) dz \\ &\geq \delta_\phi A_\phi^{-1} \int_{x - 2t + C_\phi t^{2\gamma-1}}^{+\infty} z^{-r} dz = \frac{\delta_\phi}{A_\phi(r-1)} \left(x - 2t + C_\phi t^{2\gamma-1} \right)^{1-r}. \end{aligned} \quad (2.9)$$

Note that, as $r > 1$, we have

$$2\gamma - 1 = \frac{3-r}{1+r} = \gamma + \frac{1-r}{1+r} < \gamma.$$

Further increasing T , if necessary, the right-hand side in (2.9) can be estimated, for $t \geq T$, as

$$\begin{aligned} \frac{\delta_\phi}{A_\phi(r-1)} \left(x - 2t + C_\phi t^{2\gamma-1} \right)^{1-r} &\geq \frac{\delta_\phi}{A_\phi(r-1)} \left(t^\gamma + C_\phi t^{2\gamma-1} \right)^{1-r} \\ &= \frac{\delta_\phi}{A_\phi(r-1)} \left(1 + C_\phi t^{\frac{1-r}{1+r}} \right)^{1-r} t^{(1-r)\gamma} \geq \frac{\delta_\phi}{A_\phi(r-1)} \left(1 + C_\phi T^{\frac{1-r}{1+r}} \right)^{1-r} t^{(1-r)\gamma} \\ &\geq \frac{2^{1-r} \delta_\phi}{A_\phi(r-1)} t^{(1-r)\gamma} \geq 2c_\phi (2\gamma - 1) t^{\gamma(1-r)}, \end{aligned} \quad (2.10)$$

as long as c_ϕ is sufficiently small. Now, (2.6) follows from (1.2), (2.9) and (2.10).

To show (2.7), first, we consider the right spatial boundary $x = 2t + t^\gamma$, $t \geq T$. As this point is at the far edge of the front, it is natural to use the linearized problem

$$\begin{aligned}\bar{u}_t &= \bar{u}_{xx} + \bar{u}, \quad t > 0, x \in \mathbb{R}, \\ \bar{u}(t=0, \cdot) &= u_0.\end{aligned}$$

Then, with x_0 as in (1.8), we can write for $t \geq T$:

$$\begin{aligned}u(t, 2t + t^\gamma) &\leq \bar{u}(t, 2t + t^\gamma) = \frac{e^t}{\sqrt{4\pi t}} \int_{\mathbb{R}} e^{-\frac{(2t+t^\gamma-y)^2}{4t}} u_0(y) dy \leq \frac{e^t}{\sqrt{4\pi t}} \int_{-\infty}^{x_0} e^{-\frac{(2t+t^\gamma-y)^2}{4t}} dy \\ &= \frac{e^t}{\sqrt{\pi}} \int_{\frac{2t+t^\gamma-x_0}{2\sqrt{t}}}^{+\infty} e^{-y^2} dy \leq \frac{Ce^t \sqrt{t}}{2t + t^\gamma - x_0} e^{-\frac{(2t+t^\gamma-x_0)^2}{4t}} \\ &\leq C_0 \exp \left\{ -t^\gamma - \frac{1}{4} t^{2\gamma-1} \right\} \leq B \exp \left\{ -t^\gamma - 2c_\phi t^{2\gamma-1} \right\} = \bar{v}(t, 2t + t^\gamma),\end{aligned}\tag{2.11}$$

so long as $B \geq C_0$. Above, we have increased T and decreased c_ϕ if necessary. The constant C_0 depends only on γ and x_0 . Thus, (2.7) holds at $x = 2t + t^\gamma$ for all $t \geq T$ as long as $B \geq C_0$.

At the left boundary $x = 2t - C_\phi t^{2\gamma-1}$, we have

$$\bar{v}(t, 2t - C_\phi t^{2\gamma-1}) = B \exp \left\{ (C_\phi - 2c_\phi) t^{2\gamma-1} \right\} \geq M \geq u(t, 2t - C_\phi t^{2\gamma-1}),\tag{2.12}$$

as long as $2c_\phi \leq C_\phi$ and $B \geq M$. Here, M is the upper bound in Lemma 2.1.

Lastly, we check that (2.7) holds at $t = T$, for $2T - C_\phi T^{2\gamma-1} \leq x \leq 2T + T^\gamma$:

$$\bar{v}(T, x) = B \exp \left\{ -\left(x - 2T + 2c_\phi T^{2\gamma-1}\right) \right\} \geq B \exp \left\{ -T^\gamma - 2c_\phi T^{2\gamma-1} \right\}.$$

As long as $B \geq M \exp \left\{ T^\gamma + 2c_\phi T^{2\gamma-1} \right\}$, we have that, for all $x \in [2T - C_\phi T^{2\gamma-1}, 2T + T^\gamma]$

$$\bar{v}(T, x) \geq M \geq u(T, x),\tag{2.13}$$

and (2.7) holds on all of $\partial\mathcal{P}_T$.

It follows from (2.6) and (2.7) that, with T and B sufficiently large, and c_ϕ sufficiently small, we have

$$\lim_{t \rightarrow \infty} \sup_{x \geq 2t - c_\phi t^{2\gamma-1}} u(t, x) \leq \lim_{t \rightarrow \infty} \sup_{x \geq 2t - c_\phi t^{2\gamma-1}} \bar{v}(t, x) \leq \lim_{t \rightarrow \infty} B \exp \left\{ -\left(2c_\phi - c_\phi\right) t^{2\gamma-1} \right\} = 0,$$

which finishes the proof of the upper bound. \square

3. Lower bounds on the location of the front

The proofs of the lower bounds in Theorem 1.1 are much more involved. They hinge on estimating $\phi \star u$ in terms of u in a local way, and then deriving precise heat kernel type estimates on the resulting local equation.

3.1. Estimating the non-local term by a local counterpart

To begin, we estimate the convolution term $\phi \star u$ in terms of u under the assumptions of Theorem 1.1. The assumptions of these two theorems differ only in the range of r . In this section, we assume only that $r > 1$ so our computations apply to all cases.

Lemma 3.1. *There exists $C_{\text{conv}} > 0$, depending only on ϕ , such that, for all $t \geq 1$ and all $x \in \mathbb{R}$,*

$$\phi \star u(t, x) \leq C_{\text{conv}} \max \left\{ 1, \left[\frac{1}{t} \log \left(\frac{M}{u(t, x)} \right) \right]^{\frac{r-1}{2}} \log \left(\frac{M}{u(t, x)} \right)^{1-r} \right\}.\tag{3.1}$$

Proof. It is here that the local-in-time Harnack inequality is used crucially. Fix any time $t \geq 1$ and $x, y \in \mathbb{R}$. Proposition 1.2 with $p = 2$ implies that there exists $\alpha > 0$ so that

$$u(t, x + y) \leq C\sqrt{u(t, x)} \exp \left\{ \alpha t' + \frac{\alpha y^2}{t'} \right\}, \quad \text{for all } t' \in (0, t], \quad (3.2)$$

Above, we absorbed the uniform bound M of $\|u\|_\infty$ given by Lemma 2.1 into the constant C . By increasing M if necessary, we may assume that $M \geq \|u\|_\infty + 1$, which allows us to simplify notation in the sequel. Using (1.6) and (3.2), we obtain, for $R > 0$ and $t' \in (0, t]$ to be determined,

$$\begin{aligned} \phi \star u(t, x) &\leq \int_{\mathbb{R}} \phi(y) u(t, x - y) dy \leq C \int_{B_R} \phi(y) \sqrt{u(t, x)} e^{\alpha t' + \frac{\alpha R^2}{t'}} dy + M \int_{B_R^c} \phi(y) dy \\ &\leq C\sqrt{u(t, x)} R \exp \left\{ \alpha t' + \frac{\alpha R^2}{t'} \right\} + CM R^{-r+1}. \end{aligned} \quad (3.3)$$

The constant C changes line-by-line for the remainder of the proof and depends only on ϕ and α .

We now optimize the right-hand side in (3.3) with respect to $t' \in (0, t]$ and $R > 0$. If $t' = R$, then

$$\phi \star u(t, x) \leq C\sqrt{u(t, x)} R e^{2\alpha R} + CM R^{-r+1}. \quad (3.4)$$

To roughly balance the two terms in the right side of (3.4), we choose

$$R = \frac{1}{8\alpha} \log \left(\frac{M}{u(t, x)} \right), \quad (3.5)$$

the most important point being that R should be of order $\log u$. As we have set $t' = R$ in (3.4), and we need to have $0 \leq t' \leq t$, the choice (3.5) is possible only if

$$t \geq \frac{1}{8\alpha} \log \left(\frac{M}{u(t, x)} \right). \quad (3.6)$$

With this, we find, from (3.4):

$$\begin{aligned} \phi \star u(t, x) &\leq C\sqrt{u(t, x)} \log \left(\frac{M}{u(t, x)} \right) \exp \left\{ -\frac{1}{4} \log \left(\frac{u(t, x)}{M} \right) \right\} + C \left(\log \left(\frac{M}{u(t, x)} \right) \right)^{1-r} \\ &\leq C \left(u(t, x)^{1/4} \left(\log \left(\frac{M}{u(t, x)} \right) \right)^r + 1 \right) \left(\log \left(\frac{M}{u(t, x)} \right) \right)^{1-r} \leq C \left(\log \left(\frac{M}{u(t, x)} \right) \right)^{1-r}. \end{aligned} \quad (3.7)$$

When (3.6) does not hold, so that

$$t \leq \frac{1}{8\alpha} \log \left(\frac{M}{u(t, x)} \right), \quad (3.8)$$

we choose $t' = t$ and set

$$R = \left(\frac{t}{8\alpha} \log \left(\frac{M}{u(t, x)} \right) \right)^{1/2},$$

in (3.3), leading to

$$\begin{aligned} \phi \star u(t, x) &\leq C\sqrt{u(t, x)} \sqrt{t} \log^{1/2} \left(\frac{M}{u(t, x)} \right) \exp \left\{ \alpha t + \frac{1}{8} \log \left(\frac{M}{u(t, x)} \right) \right\} + C \left(t \log \left(\frac{M}{u(t, x)} \right) \right)^{-\frac{(r-1)}{2}} \\ &\leq C\sqrt{u(t, x)} \log \left(\frac{M}{u(t, x)} \right) \exp \left\{ \frac{1}{4} \log \left(\frac{M}{u(t, x)} \right) \right\} + C \left(t \log \left(\frac{M}{u(t, x)} \right) \right)^{-\frac{(r-1)}{2}} \\ &\leq C u(t, x)^{1/4} \log \left(\frac{M}{u(t, x)} \right) + C \left(t \log \left(\frac{M}{u(t, x)} \right) \right)^{-\frac{(r-1)}{2}} \\ &\leq C \left(1 + u(t, x)^{1/4} \log \left(\frac{M}{u(t, x)} \right) \left(t \log \left(\frac{M}{u(t, x)} \right) \right)^{\frac{r-1}{2}} \right) \left(t \log \left(\frac{M}{u(t, x)} \right) \right)^{-\frac{(r-1)}{2}} \\ &\leq C \left(u(t, x)^{1/4} \left(\log \left(\frac{M}{u(t, x)} \right) \right)^r + 1 \right) \left(t \log \left(\frac{M}{u(t, x)} \right) \right)^{-\frac{(r-1)}{2}} \leq C \left(t \log \left(\frac{M}{u(t, x)} \right) \right)^{\frac{1-r}{2}}. \end{aligned} \quad (3.9)$$

We used (3.8) several times above, as well as the upper bound $u(t, x) \leq M$ in the last inequality. The combination of (3.7) and (3.9) concludes the proof of the lemma. \square

3.2. A local equation and related bounds

In view of Lemma 3.1, it is natural to introduce the following nonlinearity. Fix $r > 1$, and for any positive constants θ_g and A_g , set $\Theta_g := \theta_g \exp\{-A_g^{1/(r-1)}\}$ and define $g \in C^{0,1}$ on $(0, \Theta_g)$ as

$$g(t, u) := A_g \max \left\{ 1, \left[\left(t + A_g^{\frac{1}{r-1}} \right)^{-1} \log \left(\frac{\theta_g}{u} \right) \right]^{\frac{r-1}{2}} \log \left(\frac{\theta_g}{u} \right)^{1-r}, \text{ if } u \in (0, \Theta_g). \right. \quad (3.10)$$

Outside $[0, \Theta_g]$ we set $g(t, u) = 0$ for $u < 0$ and $g(t, u) = 1$ for $u > \Theta_g$. By construction, $g(t, \cdot)$ is continuous. The “ $A_g^{1/(r-1)}$ ” term in the second part of the maximum in the definition of g does not affect the analysis in any way. In fact, any other choice of g that preserves the asymptotics as u and t tend to zero would have the desired properties that we prove in the sequel.

We will make use of the local equation with a moving boundary at the front edge:

$$\begin{aligned} w_t &= w_{xx} + w(1 - g(t, w)), & \text{in } \mathcal{P}_{g,\gamma} &:= \left\{ (t, x) : t > 0, x > 2t + (t + t_0)^\gamma - t_0^\gamma \right\}, \\ w(t, 2t + (t + t_0)^\gamma - t_0^\gamma) &= 0, & \text{for all } t > 0, \\ w(0, x) &= w_0(x) & \text{for all } x > 0. \end{aligned} \quad (3.11)$$

The following proposition contains the crucial lower bounds for the solutions of (3.11) we will need.

Proposition 3.2. *Assume that there exists $\delta_w > 0$ and $x_w \in \mathbb{R}^+$ such that the initial condition $w_0(x)$ for (3.11) satisfies $w_0(x) \geq \delta_w 1_{(0, x_w)}(x)$.*

1. *If $r > 3$, then there exists X_w and T_0 such that if $x_w \geq X_w$ and $t_0 \geq T_0$ then there exists a positive constant B_1 , depending only on $x_w, \delta_w, t_0, \gamma$, and g , such that, for all t sufficiently large, we have*

$$w(t, 2t + t^\gamma + \sqrt{t}) \geq B_1^{-1} t^{-1} e^{-\sqrt{t} - t^\gamma}.$$

2. *If $r = 3$, then set $t_0 = 1$. There exists $\tilde{N} > 0$ such that if $x_w \geq 1$ then there exists a positive constant B_2 , depending only on $\tilde{N}, \delta_w, \gamma$, and g , such that, for all t sufficiently large, we have*

$$w(t, 2t + 2\sqrt{t}) \geq B_2^{-1} t^{-1-\tilde{N}} e^{-\sqrt{t}}.$$

3. *If $r \in (1, 3)$, then set $t_0 = 1$. There exists $B_3 > 0$, depending only on δ_w and g , such that if $x_w \geq 1$ then, for all $t \geq 1$, we have*

$$w(t, 2t + t^\gamma + \sqrt{t}) \geq B_3^{-1} e^{-\sqrt{t} - t^\gamma - B_3 t^{2\gamma-1}}.$$

We delay the proof of this proposition until Section 4 and now continue the proof of the lower bounds of Theorem 1.1. Having reduced the problem to estimating a delay for a local equation, we now transfer known bounds of Theorem 1.1 on w to bounds on u .

3.3. From a bound on w to a bound on u

Let us take $\theta_g = M$ and $A_g = C_{\text{conv}}$ in the definition (3.10) of $g(t, u)$ and let the initial condition in (3.11) be $w_0(x) = e^{-M} u_0(x)$. A combination of Lemma 3.1 and Proposition 3.2 implies that u is a super-solution for w for $t \geq 1$. Further, it follows from considerations as in [4, Section 3], that $w(1, x) \leq u(1, x)$ for all $x \in \mathbb{R}$ due to the e^{-M} pre-factor in the definition of w_0 . The maximum principle then implies that $w(t, x) \leq u(t, x)$ for all $t \geq 1$ and all $x \in \mathbb{R}$.

Using the assumptions on the initial data (1.8), we can, up to translating u_0 , and thus w_0 as well, assume that w_0 satisfies the hypothesis $x_w = x_0 \geq X_w$ in Proposition 3.2. Translating further and using parabolic regularity we may remove the dependence on t_0 . As a direct consequence, we have established:

Corollary 3.3. Suppose that u satisfies (1.2) and (1.8) with ϕ satisfying (1.4), (1.5), and (1.6). Then there exists S_0 , depending only on u_0 and ϕ , such that:

1. If $r > 3$, then there exists a positive constant B_1 , depending only on u_0 and ϕ such that, for all t sufficiently large, we have

$$u(t, 2t + t^\gamma + \sqrt{t} - S_0) \geq B_1^{-1} t^{-1} e^{-\sqrt{t} - t^\gamma}.$$

2. If $r = 3$, then there exist positive constants \bar{N} , and B_2 such that, for all t sufficiently large, we have

$$u(t, 2t + 2\sqrt{t} - S_0) \geq B_2^{-1} t^{-(1+\bar{N})} e^{-2\sqrt{t}}.$$

3. If $r \in (1, 3)$, then there exists a positive constant B_3 , depending only on u_0 and ϕ , such that, for all $t \geq 1$, we have

$$u(t, 2t + t^\gamma + \sqrt{t} - S_0) \geq B_3^{-1} e^{-t^\gamma - B_3 t^{2\gamma-1}}.$$

3.4. From a bound on u on the right to the location of the front

We are now in a position to obtain the lower bounds (1.9), (1.11), and (1.13). Thanks to Corollary 3.3, we fit a suitable translate of a traveling wave solution for (3.11) underneath u , for $x \leq 2t + t^\gamma + \sqrt{t}$.

Fix any $A_V > C_{\text{conv}}$ and let V be a traveling wave solution of

$$-2V' = V'' + V \left(1 - A_V \log \left(\frac{M}{V} \right)^{1-r} \right), \quad V(-\infty) = M \exp \left\{ -A_V^{1/(r-1)} \right\} \text{ and } V(+\infty) = 0.$$

The existence, uniqueness up to translation, and monotonicity of V is given by, for example, [2]. We also recall (see [15,3]) the fact that there exists $\beta_{r,A_V} > 0$, depending only on r and A_V , and $\kappa > 0$ such that, as $\xi \rightarrow \infty$

$$V(\xi) \sim \begin{cases} \kappa \xi e^{-\xi}, & \text{if } r > 3, \\ \kappa \xi^{1+\beta_{r=3,A_V}} e^{-\xi}, & \text{if } r = 3, \end{cases} \quad (3.12)$$

and, if $r \in (1, 3)$,

$$\frac{\log(V(\xi)) + \xi}{\xi^{(3-r)/2}} \sim \beta_{r,A_V}. \quad (3.13)$$

Define v as

$$\begin{aligned} v(t, x) &= V \left(x - 2t + \frac{3}{2} \log t + s_0 \right), & \text{if } r > 3, \\ v(t, x) &= V \left(x - 2t + \left(\frac{3}{2} + \bar{N} + \frac{\beta_{r=3,A_V}}{2} \right) \log t + s_0 \right), & \text{if } r = 3, \\ v(t, x) &= V \left(x - 2t + (2B_3 + \beta_{r,A_V}) t^{2\gamma-1} + s_0 \right), & \text{if } r \in (1, 3), \end{aligned}$$

where the shift s_0 is to be determined below and λ is as in Proposition 3.2 and Corollary 3.3.

Lemma 3.4. There exists $T_1 > 0$ and \underline{s}_0 such that if $s_0 \geq \underline{s}_0$, then $v(t, x) \leq u(t, x)$ if $r \neq 3$, $t \geq T_1$, and $x \leq 2t + t^\gamma + \sqrt{t} - S_0$ where S_0 is the shift given in Corollary 3.3.

Proof. We prove the lemma for $r > 3$, so that $\gamma < 1/2$, the proof being the same in the other cases up to situational modifications. We use the parabolic maximum principle. First, we note that, up to increasing s_0 and A_V , we may ensure that

$$v(T_1, x) \leq u(T_1, x) \quad \text{for all } x \leq 2T_1 + \sqrt{T_1} + T_1^\gamma - S_0.$$

Second, we claim that, up to increasing s_0 , we have

$$v(t, 2t + \sqrt{t} + t^\gamma - S_0) \leq u(t, 2t + \sqrt{t} + t^\gamma - S_0) \quad \text{for all } t \geq T_1.$$

Indeed, for t sufficiently large, (3.12) implies, as $\gamma < 1/2$:

$$\begin{aligned} v(t, 2t + \sqrt{t} + t^\gamma - S_0) &= V\left(\sqrt{t} + t^\gamma + \frac{3}{2} \log t + s_0 - S_0\right) \\ &\leq 2\kappa\left(\sqrt{t} + t^\gamma + \frac{3}{2} \log t + s_0 - S_0\right) \exp\left\{-\left(\sqrt{t} + t^\gamma + \frac{3}{2} \log t + s_0 - S_0\right)\right\} \\ &\leq 4\kappa\sqrt{t}t^{-3/2} \exp\left\{-\sqrt{t} - t^\gamma - s_0 + S_0\right\} \leq 4\kappa t^{-1} \exp\left\{-\sqrt{t} - t^\gamma - s_0 + S_0\right\}. \end{aligned} \quad (3.14)$$

It follows that

$$v(t, 2t + \sqrt{t} + t^\gamma - S_0) \leq 4\kappa e^{S_0 - s_0} B_1 u(t, 2t + \sqrt{t} + t^\gamma - S_0) \leq u(t, 2t + \sqrt{t} + t^\gamma - S_0),$$

for T_1 sufficiently large and all $s_0 \geq S_0 + \log(4\kappa B_1)$.

Third, up to increasing A_V , the ordering holds true near $-\infty$. Indeed, using Lemma 3.1 and the assumptions (1.8) on u_0 , it is easy to see that there exists $\delta > 0$, depending only on u_0 and ϕ such that, for any $\bar{x} < 0$ with $|\bar{x}|$ is sufficiently large, the function

$$\underline{u}(x) = \delta \cos((x - \bar{x})/100)$$

is a sub-solution for u for all $t \geq 1$, so that $\delta = \underline{u}(\bar{x}) \leq u(t, \bar{x})$ for all $t \geq 1$. Thus, increasing A_V , if necessary, we have that, for all $t > 0$,

$$\lim_{x \rightarrow -\infty} v(t, x) < M e^{-A_V^{1/(r-1)}} < \delta \leq \inf_{t \geq 1} \liminf_{x \rightarrow -\infty} u(t, x).$$

Now, assume for the sake of a contradiction that there exists a first touching time $(t_{\text{ft}}, x_{\text{ft}})$ such that

$$t_{\text{ft}} \geq T_1, x_{\text{ft}} \leq 2t_{\text{ft}} + \sqrt{t_{\text{ft}}} + t_{\text{ft}}^\gamma - S_0,$$

and

$$u(t_{\text{ft}}, x_{\text{ft}}) = v(t_{\text{ft}}, x_{\text{ft}}),$$

and $u(t, x) > v(t, x)$ for all $t \in [T_1, t_{\text{ft}})$ and $x < 2t + \sqrt{t} + t^\gamma - S_0$. Our goal is to obtain a contradiction by estimating $\phi \star u$ and looking at the equation satisfied by $u - v$.

First, we estimate $\phi \star u(t_{\text{ft}}, x_{\text{ft}})$ using Lemma 3.1. By increasing s_0 if necessary, we obtain

$$\begin{aligned} v(t, 2t + \sqrt{t} + t^\gamma - S_0) &= V\left(\sqrt{t} + t^\gamma + \frac{3}{2} \log(t) + s_0 - S_0\right) \\ &\geq \frac{\kappa}{2}\left(\sqrt{t} + t^\gamma + \frac{3}{2} \log t + s_0 - S_0\right) \exp\left\{-\left(\sqrt{t} + t^\gamma + \frac{3}{2} \log t + s_0 - S_0\right)\right\} \\ &\geq \frac{\kappa}{2t^{3/2}}\left(\sqrt{t} + t^\gamma + \frac{3}{2} \log t + s_0 - S_0\right) \exp\left\{-\sqrt{t} - t^\gamma - s_0 + S_0\right\}. \end{aligned} \quad (3.15)$$

Since V is monotonic, $\gamma < 1$, and $x_{\text{ft}} \leq 2t_{\text{ft}} + t_{\text{ft}}^{1/2} + t_{\text{ft}}^\gamma - S_0$, it follows that up to increasing T_1 , we have that

$$u(t_{\text{ft}}, x_{\text{ft}}) = v(t_{\text{ft}}, x_{\text{ft}}) \geq M e^{-t_{\text{ft}}},$$

which, in turn, implies that

$$\log\left(\frac{M}{u(t_{\text{ft}}, x_{\text{ft}})}\right)^{-(r-1)} \geq t_{\text{ft}}^{-\frac{r-1}{2}} \log\left(\frac{M}{u(t_{\text{ft}}, x_{\text{ft}})}\right)^{-\frac{r-1}{2}}.$$

In view of the bound on $\phi \star u$ obtained in Lemma 3.1, we have that, at $(t_{\text{ft}}, x_{\text{ft}})$,

$$u_t - u_{xx} - u\left(1 - A_V \log\left(\frac{M}{u}\right)^{-(r-1)}\right) \geq (A_V - C_{\text{conv}})u\left(\log\left(\frac{M}{u}\right)\right)^{1-r} > 0, \quad (3.16)$$

where we used the fact that $A_V > C_{\text{conv}}$ in the last inequality. In addition, we note that

$$\begin{aligned}
v_t - v_{xx} - v \left(1 - A_V \log \left(\frac{M}{v} \right)^{-(r-1)} \right) \\
= \left(\frac{3}{2(t+1)} - 2 \right) V' - V'' - V \left(1 - A_V \log \left(\frac{M}{V} \right)^{-(r-1)} \right) = \frac{3}{2(t+1)} V' \leq 0.
\end{aligned} \tag{3.17}$$

Hence, setting $\psi = u - v$, (3.16) and (3.17), imply that

$$\psi_t - \psi_{xx} - c\psi > 0,$$

where we define

$$c := \frac{u \left(1 - A_V \log \left(\frac{M}{u} \right)^{-(r-1)} \right) - v \left(1 - A_V \log \left(\frac{M}{v} \right)^{-(r-1)} \right)}{u - v}.$$

Notice that, due to the Lipschitz continuity of $w \mapsto w \left(1 - A_V \log \left(\frac{M}{w} \right)^{-(r-1)} \right)$ on compact subsets of $[0, M)$, $c \in L^\infty$. On the other hand, using that t_{ft} is the first time that ψ touches zero and x_{ft} is the location of a minimum of ψ , we have that

$$\psi_t - \psi_{xx} - c\psi \leq 0.$$

This yields a contradiction, finishing the proof. \square

The lower bounds now follow easily.

Proof of (1.9), (1.11), and (1.13). We conclude the proof by noticing that, for all $t \geq T_1$,

$$\begin{aligned}
\inf_{x \leq 2t - (3/2) \log t} u(t, x) &= \inf_{x \leq 0} u \left(t, x + 2t - \frac{3}{2} \log t \right) \geq \inf_{x \leq 0} v \left(t, x + 2t - \frac{3}{2} \log t \right) \\
&= \inf_{x \leq 0} V(x + s_0) = V(s_0),
\end{aligned} \tag{3.18}$$

which means that (1.9) holds. The proofs of (1.13) and (1.11) are similar and, thus, omitted. \square

4. Proof of Proposition 3.2

To obtain estimates on the solution of (3.11), we consider the corresponding linearized problem with the Dirichlet boundary condition:

$$\begin{aligned}
\tilde{v}_t &= \tilde{v}_{xx} + \tilde{v}, & \text{on } \{(t, x) : t > 0, x > 2t + (t + t_0)^\gamma - t_0^\gamma\}, \\
\tilde{v}(t, 2t + (t + t_0)^\gamma - t_0^\gamma) &= 0, & \text{for all } t > 0, \\
\tilde{v}(0, x) &= w_0(x), & \text{for all } x > 0,
\end{aligned} \tag{4.1}$$

where w_0 is as in Proposition 3.2.

4.1. The case $r > 3$

The following key lemma about solutions to (4.1) allows us to prove Proposition 3.2 when $r > 3$. We prove this lemma in Section 5.1.

Lemma 4.1. Assume $r > 3$. If t_0 and x_w are sufficiently large, depending only on γ , there exist positive constants T and B , depending only on w_0 and t_0 , such that, for all $t \geq T$, we have $\|\tilde{v}(t, \cdot)\|_\infty \leq B e^{-t^\gamma}$ and

$$\tilde{v}(t, 2t + t^\gamma + \sqrt{t}) \geq B^{-1} t^{-1} \exp \left\{ -\sqrt{t} - t^\gamma \right\}.$$

We now finish the proof of Proposition 3.2. Let \tilde{v} be as in Lemma 4.1. We may assume, without loss of generality, that $T \geq 1$, and set

$$\delta = \min \left\{ B^{-1}, B^{-1} \theta_g e^{-A_g^{1/(r-1)}}, e^{-T} \right\}.$$

We also take a continuous function $a(t) \leq 1$ for all $t \geq 0$, to be determined, and set

$$\underline{v}(t, x) = \delta a(t) \tilde{v}(t, x). \quad (4.2)$$

Using (4.1), we obtain

$$\underline{v}_t - \underline{v}_{xx} - \underline{v}(1 - g(t, \underline{v})) = \delta a' \tilde{v} + \delta a \tilde{v}_t - \delta a \tilde{v}_{xx} - \delta a \tilde{v} + \delta a \tilde{v} g(t, \delta a \tilde{v}) = \delta \tilde{v} (a' + a g(t, \delta a \tilde{v})). \quad (4.3)$$

Thus, \underline{v} is a sub-solution of w for $t \geq T$ as long as

$$a' + a g(t, \delta a \tilde{v}) \leq 0.$$

Using the upper bound on \tilde{v} along with the definition of δ , we see that this inequality would hold if

$$a' + a A_g \max \left\{ 1, \left[\left(t + A_g^{\frac{1}{r-1}} \right)^{-1} \log \left(\frac{1}{a e^{-t^\gamma}} \right) \right]^{\frac{r-1}{2}} \right\} \log \left(\frac{1}{a e^{-t^\gamma}} \right)^{1-r} \leq 0. \quad (4.4)$$

A lengthy but straightforward computation using, in particular, that $A_g \geq 1$, shows that (4.4) is satisfied if we take

$$a(t) = \exp \left\{ \beta \left[(t+1)^{2\gamma-1} - 1 \right] \right\},$$

with a suitable $\beta > 0$.

Hence \underline{v} is a sub-solution of w . Further, arguing as in [4, Section 3] and using the choice of δ and a , we have that $\underline{v}(T, x) \leq w(T, x)$ for all $x \geq 2T + (T + t_0)^\gamma - t_0^\gamma$. The maximum principle then implies that $\underline{v}(t, x) \leq w(t, x)$ for all $t > T$ and $x > 2t + (t + t_0)^\gamma - t_0^\gamma$. The conclusion of the proposition follows immediately from Lemma 4.1 since $2t + t^\gamma \geq 2t + (t + t_0)^\gamma - t_0^\gamma$.

4.2. The case $r = 3$

We follow here the same strategy as for $r > 3$, but the estimates on \tilde{v} are obtained differently.

Lemma 4.2. *For $r = 3$ and t sufficiently large, there exist λ and $B > 0$ such that*

$$\|\tilde{v}(t, \cdot)\|_\infty \leq B e^{-\sqrt{t}},$$

and

$$\tilde{v}(t, 2t + 2\sqrt{t}) \geq B^{-1} t^{-1-\lambda} \exp \left\{ -2\sqrt{t} \right\}.$$

With this lemma, proved in Section 5.2, one may repeat the argument for $r > 3$, building a sub-solution $\underline{v}(t, x)$ as in (4.2), with $\delta > 0$ sufficiently small, and $a(t)$ such that

$$a' + a A_g \max \left\{ 1, \left(t + \sqrt{A_g} \right)^{-1} \log \left(\frac{e^{\sqrt{t}}}{C \delta a} \right) \right\} \log \left(\frac{e^{\sqrt{t}}}{C \delta a} \right)^{-2} \leq 0.$$

The above inequality is satisfied with $a(t) = (t + \sqrt{A_g})^{-N}$ for all $t \geq 1$ so long as δ is chosen small enough and N is chosen large enough, depending only on A_g and C .

4.3. The estimate when $r \in (1, 3)$

Here we directly construct a sub-solution of w . We seek a sub-solution \tilde{v} solving

$$\begin{aligned} \tilde{v}_t &\leq \tilde{v}_{xx} + \tilde{v}, & \text{for } t > 0, x > 2t + (t+1)^\gamma - 1, \\ \tilde{v}(t, 2t + (t+1)^\gamma - 1) &= 0, & \text{for } t > 0. \end{aligned} \quad (4.5)$$

Recall that $t_0 = 1$ in parts 2 and 3 of Proposition 3.2. Given $a > 0$, set

$$v(t, x) = \frac{x}{(1+t)^3} \exp \left\{ -x - \frac{\gamma}{2} x (1+t)^{\gamma-1} - (1+t)^\gamma - \left[\frac{\gamma^2}{4(2\gamma-1)} + a \right] (1+t)^{2\gamma-1} - \frac{x^2}{2(1+t)} \right\}. \quad (4.6)$$

Here, the key computation is the following:

Lemma 4.3. *There exists $a_0 > 0$ such that if $a \geq a_0$ then $\tilde{v}(t, x) = v(t, x - (2t + (t+1)^\gamma - 1))$ solves (4.5).*

We delay the proof of Lemma 4.3 until Section 5.3 and proceed with the proof of Proposition 3.2.

A bound for small times Unfortunately, v is not compactly supported at $t = 0$, so we need to “fit it under” w at a later time. To do this, we first obtain a preliminary lower bound on w at time 1 by using the infinite speed of propagation of the heat equation. Recall that $w_0 \geq \delta_w \mathbb{1}_{(-\infty, x_w)}$ and $1 - g(t, w) \geq 0$. Hence, we have

$$w_t - w_{xx} \geq 0,$$

so that w is a super-solution to the heat equation with a Dirichlet boundary condition fixed at

$$\bar{x}_0 := 2 \cdot 2 + (2 + t_0)^\gamma - t_0^\gamma = 3^\gamma + 3,$$

on the time interval $[0, 2]$. It follows that

$$\begin{aligned} w(2, x + \bar{x}_0) &\geq \frac{1}{\sqrt{8\pi}} \int_0^\infty w_0(y + \bar{x}_0) \left[e^{-|x-y|^2/8} - e^{-|x+y|^2/8} \right] dy \\ &\geq \frac{\delta_w e^{-x^2/8}}{\sqrt{8\pi}} \int_0^{x_w - \bar{x}_0} e^{-y^2/8} \left[e^{xy/4} - e^{-xy/4} \right] dy \\ &\geq \frac{\delta_w e^{-x^2/8 - (x_w - \bar{x}_0)^2/8}}{\sqrt{8\pi}} \frac{2}{x} \left(\cosh \left(\frac{x(x_w - \bar{x}_0)}{4} \right) - 1 \right) \geq \frac{x}{C} \delta_w e^{-x^2/8}, \end{aligned} \quad (4.7)$$

for some C independent of all parameters, as long as $x_w \geq \bar{x}_0 + 1$. We used here that $\cosh(x) - 1 \geq x^2/C$ for some universal $C > 0$. On the other hand, from the explicit expression (4.6) for v , we get

$$v(2, x - \bar{x}_0) \leq C(x - \bar{x}_0) \exp \left\{ -x - x \frac{\gamma}{2} 3^{\gamma-1} - \frac{x^2}{6} + \frac{x\bar{x}_0}{3} \right\}.$$

Thus, there exists $\epsilon > 0$ such that

$$\epsilon \tilde{v}(2, x) = \epsilon v(2, x - \bar{x}_0) \leq w(2, x) \text{ for } x \geq \bar{x}_0 = 3^\gamma + 3.$$

The subsolution We now follow the same strategy as before, constructing a sub-solution of the form $\underline{v}(t, x) = \delta a(t) \tilde{v}(t, x)$ on

$$\mathcal{P} := \{(t, x) : t \geq 1, x > 2t + (1+t)^\gamma - 1\}.$$

Another lengthy but straightforward computation shows that $\underline{v}(t, x)$ is a sub-solution for w on \mathcal{P} if we choose $a(t) = \exp \left\{ -\beta t^{2\gamma-1} \right\}$ for a suitable β and δ sufficiently small.

Note also that \tilde{v} and w satisfy the same boundary conditions at $x = 2t + (1+t)^\gamma - 1$. Finally, choosing $\delta \leq \epsilon$ and using the computation (4.7) and the discussion following it, we see that

$$\underline{v}(2, x) \leq w(2, x) \text{ for all } x > 3 + 3^\gamma.$$

The conclusion of the proposition when $r \in (1, 3)$ follows by simply using the explicit form of $\underline{v}(t, x)$.

5. Estimates on the linearized KPP equation

In this section, we adopt the convention that any constant denoted C may change line-by-line but depends only on ϕ and u_0 .

5.1. The case $r > 3$: the proof of Lemma 4.1

The key observation is that $\gamma < 1/2$ when $r > 3$. Thus, the t^γ term is of a lower order than the diffusive scale \sqrt{t} . This allows us to use the strategy in [17], obtaining energy estimates in self-similar variables. Since the present proof is similar to that in [17], we provide a rather brief treatment.

Proof of Lemma 4.1. We begin by removing an exponential factor from \tilde{v} and changing to the moving frame: let

$$z(t, x) := e^x \tilde{v}(t, 2t + (t + t_0)^\gamma - t_0^\gamma + x), \quad x > 0.$$

This function satisfies

$$\begin{aligned} z_t &= z_{xx} + \gamma(t + t_0)^{\gamma-1}(z_x - z), & \text{for } t > 0, x > 0, \\ z(t, 0) &= 0, & \text{for } t > 0, \\ z(0, x) &= e^x w_0(x), & \text{for } x > 0. \end{aligned} \tag{5.1}$$

We now turn to self-similar variables, which are natural for the diffusive process. Let

$$\tau = \log\left(1 + \frac{t}{t_0}\right), \quad y = (t + t_0)^{-1/2}x,$$

and $\zeta(\tau, y) = z(t_0(e^\tau - 1), t_0^{1/2}e^{\tau/2}y)$. Then ζ satisfies the equation

$$\zeta_\tau = \zeta_{yy} + \frac{y}{2}\zeta_y + 1 + \gamma(t_0 e^\tau)^{\gamma-1/2}\zeta_y - \left(1 + \gamma(t_0 e^\tau)^\gamma\right)\zeta.$$

We remove the integrating factor above, setting

$$\zeta(\tau, y) = e^{-(\tau + t_0^\gamma(e^{\gamma\tau} - 1))} \bar{\zeta}(\tau, y),$$

so that $\bar{\zeta}$ satisfies

$$\bar{\zeta}_\tau = L\bar{\zeta} + \gamma t_0^{\gamma-1/2} e^{(\gamma-1/2)\tau} \bar{\zeta}_y, \tag{5.2}$$

with

$$L := \partial_y^2 + \frac{y}{2}\partial_y + 1. \tag{5.3}$$

It is now heuristically clear that the last term in (5.2) should be not important due to the $e^{(\gamma-1/2)\tau}$ term and the fact that $\gamma < 1/2$. The following lemma is proved in Appendix A.

Lemma 5.1. *Let $\bar{\zeta}$ solve*

$$\bar{\zeta}_\tau = L\bar{\zeta} + \varepsilon e^{(\gamma-1/2)\tau} \bar{\zeta}_y,$$

with initial data $\bar{\zeta}(\tau = 0, \cdot) = \bar{\zeta}_0$. There exists $\varepsilon_0 > 0$ such that for all compact subsets $K \subset \mathbb{R}_+$ there exists $C_K > 0$ such that for all $\varepsilon < \varepsilon_0$,

$$\bar{\zeta}(\tau, y) = y \left(\frac{e^{-y^2/4}}{2\sqrt{\pi}} \left(\int_0^\infty \xi \bar{\zeta}_0(\xi) d\xi + O(\varepsilon) \right) + e^{(\gamma-1/2)\tau} \bar{h}(\tau, y) \right),$$

for all $y > 0$, $\tau > 0$, and such that $|\bar{h}(\tau, y)| \leq C_K$ for all $\tau > 0$ and $y \in K$.

Undoing the various changes of variable, we get

$$\begin{aligned} \tilde{v}(t, 2t + (t + t_0)^\gamma - t_0^\gamma + x) &= e^{-x} z(t, x) = e^{-x} \zeta \left(\log \left(1 + \frac{t}{t_0} \right), \frac{x}{(t + t_0)^{1/2}} \right) \\ &= \frac{x e^{-x} t_0 e^{-((t+t_0)^\gamma - t_0^\gamma)}}{(t + t_0)^{3/2}} \left(\frac{e^{-\frac{x^2}{4(t+t_0)}}}{2\sqrt{\pi}} \left(\int_0^\infty \xi e^\xi w_0(\sqrt{t_0}\xi) d\xi + O(t_0^{\gamma-\frac{1}{2}}) \right) + \left(1 + \frac{t}{t_0} \right)^{\gamma-\frac{1}{2}} h(t, x) \right), \end{aligned} \quad (5.4)$$

where $h(t, x) = \bar{h} \left(\log \left(1 + \frac{t}{t_0} \right), (t + t_0)^{-\frac{1}{2}} x \right)$.

First, notice that the L^∞ bound on \tilde{v} in Lemma 4.1 follows immediately from the expression above on sets of the form $[2t + t^\gamma, 2t + t^\gamma + \sigma\sqrt{t}]$. To obtain bounds on sets of the form $[2t + t^\gamma + \sigma\sqrt{t}, \infty)$, we simply use that $e^{-t}\tilde{v}$ is a sub-solution to the heat equation on \mathbb{R} (see, e.g., (1.20)). Hence, we obtain that, for $x \geq 0$,

$$\tilde{v}(t, 2t + t^\gamma + \sigma\sqrt{t} + x) \leq \frac{C e^t}{\sqrt{t}} \exp \left\{ -\frac{(2t + t^\gamma + \sigma\sqrt{t} + x)^2}{4t} \right\} \leq C e^{-\sqrt{t}-t^\gamma}, \quad (5.5)$$

where C is some constant depending only on the initial data and γ .

Second, we have

$$\int_0^\infty \xi e^\xi w_0(t_0^{1/2}\xi) d\xi + O(t_0^{\gamma-\frac{1}{2}}) \geq \delta_w \int_0^{x_w/t_0^{1/2}} \xi e^\xi d\xi + O(t_0^{\gamma-\frac{1}{2}}).$$

Choosing first $x_w \geq \sqrt{t_0}$ and $t_0 \gg 1$ so that the first two terms in the parentheses in (5.4) are positive and then choosing T_0 large depending on t_0 and α , we have that, for all $0 \leq x \leq \sqrt{t+t_0}$ and $t \geq T_0$,

$$\tilde{v}(t, 2t + (t + t_0)^\gamma - t_0^\gamma + x) \geq \frac{x}{C} \frac{e^{-x-((t+t_0)^\gamma - t_0^\gamma)}}{(t + t_0)^{3/2}}.$$

The lower bound on $\tilde{v}(t, 2t + t^\gamma + \sqrt{t})$ is immediate after evaluating at $x = t^\gamma - ((t + t_0)^\gamma - t_0^\gamma) + \sqrt{t}$. This concludes the proof. \square

5.2. The case $r = 3$: the proof of Lemma 4.2

Note that in this case $\gamma = 1/2$. As a consequence, the drift induced by the moving boundary has the same order as the diffusion. It is thus useful to modify the t^γ term in the moving boundary by a small multiplicative factor.

Proof of Lemma 4.2. To begin, fix $\varepsilon > 0$. Work in the moving frame $2t + [(t + 1)^{1/2} - 1]$ and remove an exponential factor, as previously:

$$z(t, x) := e^x \tilde{v}(t, x + 2t + [(1 + t)^{1/2} - 1]).$$

Passing then to self-similar coordinates

$$\tau = \log(t + 1) \quad \text{and} \quad y = (t + 1)^{-1/2} x,$$

so that

$$\zeta(\tau, y) := z(e^\tau - 1, e^{\tau/2} y),$$

we see that ζ satisfies

$$\zeta_\tau = L\zeta + \frac{1}{2}\zeta_y - \left(1 + \frac{1}{2}e^{\tau/2}\right)\zeta,$$

with L as in (5.3). Finally, pulling out the zeroth order factor

$$\zeta(\tau, y) = e^{-\tau - (e^{\tau/2} - 1)} \bar{\zeta}(\tau, y),$$

we see that $\bar{\zeta}$ solves

$$\bar{\zeta}_\tau = L\bar{\zeta} + \frac{1}{2}\bar{\zeta}_y. \quad (5.6)$$

We finish using the following lemma, proved in Appendix A. This result falls outside of [17] and Lemma 5.1 because the $\bar{\zeta}_y$ term in (5.6) is no longer a remainder term.

Lemma 5.2. *Let $\bar{\zeta}$ solve (5.6), then it can be represented as*

$$\bar{\zeta}(\tau, y) = \exp\left\{-\frac{y^2}{8} - \frac{y}{4}\right\} \left(\left(\int_{\mathbb{R}^+} \psi(y) e^{\frac{y^2}{8} + \frac{y}{4}} \bar{\zeta}(0, y) dy \right) \psi(y) e^{-\lambda\tau} + y \bar{h}(\tau, y) e^{-\mu\tau} \right). \quad (5.7)$$

Here, $\mu > \lambda > 0$, $\|\psi\|_2 = 1$, and ψ and $\bar{h}(\tau, y)$ are bounded on all compact subsets of $[0, \infty)$. Also, for every compact set $K \subset [0, \infty)$, there exists $C_K > 0$ such that $C_K^{-1}y \leq \psi(y) \leq C_K y$, for all $y \in K$.

We are now able to establish the upper bound on \tilde{v} for all $x \in [2t + ((1+t)^{1/2} - 1), 2t + 2\sqrt{t}]$. Indeed, returning to the original variables in (5.7), we find

$$\begin{aligned} |\tilde{v}(t, x)| &= e^{-(1+t)^{1/2}y} \zeta(\tau, y) = e^{-(1+t)^{1/2}y - \tau - (e^{\tau/2} - 1)} \bar{\zeta}(\tau, y) \\ &\leq C e^{-(1+t)^{1/2}y - \tau - (e^{\tau/2} - 1) - \frac{y^2}{8} - \frac{y}{4}} (\psi(y) e^{-\lambda\tau} + h(\tau, y) e^{-\mu\tau}) \leq C e^{-\sqrt{t}}. \end{aligned} \quad (5.8)$$

In fact, the estimate (5.8) holds for all $x \geq 2t + [(1+t)^{1/2} - 1]$ since, as above, \tilde{v} may be estimated for $x \in [2t + 2\sqrt{t}, \infty)$ using the same approach as in (5.5) (see also (1.20)). Indeed,

$$\tilde{v}(t, x) \leq \frac{C\sqrt{t}}{x} e^{t - \frac{x^2}{2t}} \leq \frac{C}{\sqrt{t}} e^{t - \frac{(2t+2\sqrt{t})^2}{4t}} = \frac{C}{\sqrt{t}} e^{-2\sqrt{t}-1} \leq C e^{-\sqrt{t}}.$$

We now establish the lower bound from (5.7). Taking t sufficiently large and evaluating at $x = 2t + 2\sqrt{t}$, we see that

$$\tilde{v}(t, 2t + 2\sqrt{t}) \geq \frac{\alpha}{t^{1+\lambda}} e^{-2\sqrt{t}},$$

for some α depending only on u_0 . This concludes the proof. \square

5.3. The case $r \in (1, 3)$: the proof of Lemma 4.3

To motivate some of the steps in the following proof, we briefly discuss a heuristic. In the stationary frame, as we did in (1.20), we may always estimate \tilde{v} above by ignoring the Dirichlet boundary condition and using the fact that $e^{-t}\tilde{v}$ solves the heat equation. Thus,

$$\tilde{v}(t, x + 2t + t^\gamma) \lesssim t^{-1/2} \exp\left\{t - \frac{(x + 2t + t^\gamma)^2}{4t}\right\} = t^{-1/2} \exp\left\{-x - \frac{x^2}{4t} - \frac{x}{\sqrt{t}} \frac{t^{\gamma-1/2}}{2} - t^\gamma - \frac{t^{2(\gamma-1/2)}}{4}\right\}.$$

Recalling that $\gamma > 1/2$, we see that on the diffusive scale $x \sim \sqrt{t}$, the Gaussian term $x^2/4t$ and the $t^{-1/2}$ in front are (much) lower order and, thus, negligible, but all other terms are large. Hence, our sub-solution should contain all such terms to be reasonably sharp. In particular, while the $xt^{\gamma-1}$ term appears small at first glance since $\gamma < 1$, it is not negligible in the diffusive scale $x \sim \sqrt{t}$. While the terms depending only on t show up as obvious integrating factors, this term will not. Hence, the key to the proof below is in carefully taking account of this term. Note that here we see the effect of $\gamma > 1/2$.

Proof of Lemma 4.3. We show how to “guess” the form of the sub-solution v . We begin by removing an exponential from \tilde{v} and changing to the moving frame. Define, for $x \in \mathbb{R}^+$,

$$z(t, x) := e^x \tilde{v}(t, 2t + (t+1)^\gamma - 1 + x),$$

so that (4.5) becomes

$$\begin{aligned} z_t &\leq z_{xx} + \gamma(t+1)^{\gamma-1}(z_x - z), \quad t > 0, x > 0, \\ z(t, 0) &= 0, \\ z(0, x) &= e^x w_0(x). \end{aligned} \tag{5.9}$$

Turning to self-similar variables,

$$\tau = \log(1+t), \quad y = (t+1)^{-1/2}x, \quad \text{and} \quad \zeta(\tau, y) = z(e^\tau - 1, e^{\tau/2}y),$$

we wish to construct ζ that satisfies the inequality

$$\zeta_\tau \leq \zeta_{yy} + \frac{y}{2}\zeta_y + \gamma e^{(\gamma-1/2)\tau}\zeta_y - \gamma e^{\gamma\tau}\zeta. \tag{5.10}$$

As $\gamma > 1/2$, the drift in (5.10) is not a perturbation anymore. The heuristic discussion preceding this proof indicates that we should consider

$$\zeta(\tau, y) = e^{-\alpha y e^{(\gamma-1/2)\tau}} \psi(\tau, y),$$

with $\alpha \in \mathbb{R}$ to be determined. Then we require

$$\psi_\tau \leq L\psi + (\gamma - 2\alpha)e^{(\gamma-1/2)\tau}\psi_y - \left(1 + \alpha(\gamma - \alpha)e^{(2\gamma-1)\tau} + \gamma e^{\gamma\tau}\right)\psi - \alpha(1 - \gamma)ye^{(\gamma-1/2)\tau}\psi, \tag{5.11}$$

with L as in (5.3). To remove the drift term, we set $\alpha = \gamma/2$. Then (5.11) becomes

$$\psi_\tau - L\psi + \left(1 + \frac{\gamma^2}{4}e^{(2\gamma-1)\tau} + \gamma e^{\gamma\tau}\right)\psi + \frac{\gamma}{2}(1 - \gamma)ye^{(\gamma-1/2)\tau}\psi \leq 0.$$

Further, writing

$$\psi(\tau, y) = \exp\left\{-\tau - e^{\gamma\tau} - \frac{\gamma^2}{4(2\gamma-1)}e^{(2\gamma-1)\tau}\right\}\Psi(\tau, y),$$

we arrive at

$$\Psi_\tau - L\Psi + \frac{1}{2}\gamma(1 - \gamma)ye^{(\gamma-1/2)\tau}\Psi \leq 0. \tag{5.12}$$

To deal with the last term in (5.12), let $a, a', b > 0$ be constants to be determined and define

$$\Psi(\tau, y) = y \exp\left\{-ae^{\tau(2\gamma-1)} - a'\tau - \frac{y^2}{b}\right\}.$$

By a direct computation, we see that

$$\begin{aligned} \Psi_\tau - L\Psi + \frac{\gamma}{2}(1 - \gamma)ye^{(\gamma-1/2)\tau}\Psi \\ = \left[-a' - a(2\gamma-1)e^{\tau(2\gamma-1)} - \frac{y^2}{b}\left(\frac{4}{b} - 1\right) + \left(\frac{6}{b} - \frac{3}{2}\right) + \frac{\gamma(1-\gamma)}{2}ye^{(\gamma-1/2)\tau}\right]\Psi. \end{aligned} \tag{5.13}$$

It is clear that to have (5.12), we must choose $b < 4$. For simplicity, we take $b = 2$, and

$$a' = \frac{6}{b} - \frac{3}{2} = \frac{3}{2},$$

so that (5.13) becomes

$$\Psi_\tau - L\Psi + \frac{\gamma}{2}(1 - \gamma)ye^{(\gamma-1/2)\tau}\Psi = \left[-a(2\gamma-1)e^{\tau(2\gamma-1)} + \frac{\gamma(1-\gamma)}{2}ye^{(\gamma-1/2)\tau} - \frac{y^2}{2}\right]\Psi.$$

The choice

$$a \geq \frac{\gamma^2(1-\gamma)^2}{8(2\gamma-1)}$$

ensures that (5.12) holds. Returning to our original variables, we see that

$$\begin{aligned} v(t, x) &= \tilde{v}(t, 2t + (t+1)^\gamma - 1 + x) = e^{-x} \zeta(\log(1+t), (t+1)^{-1/2}x) \\ &= e^{-x} e^{-\alpha(t+1)^{-1/2}x(1+t)^{\gamma-1/2}} \psi(\log(1+t), (t+1)^{-1/2}x) \\ &= \frac{1}{1+t} \exp \left\{ -x - \frac{\gamma}{2}x(1+t)^{\gamma-1} - (1+t)^\gamma - \frac{\gamma^2(1+t)^{2\gamma-1}}{4(2\gamma-1)} \right\} \Psi(\log(1+t), (t+1)^{-1/2}x) \\ &= \frac{x}{(1+t)^3} \exp \left\{ -x - \frac{\gamma}{2}x(1+t)^{\gamma-1} - (1+t)^\gamma - \left[\frac{\gamma^2}{4(2\gamma-1)} + a \right] (1+t)^{2\gamma-1} - \frac{x^2}{2(1+t)} \right\}. \end{aligned}$$

This concludes the proof. \square

6. The Fisher-KPP equation with a Gompertz non-linearity

A side effect of our analysis gives the asymptotics for a related local equation:

$$u_t - \Delta u = f_r(u). \quad (6.1)$$

Here, we assume that $f_r \in C^1$, $r \in (1, \infty)$, and there exist positive constants θ_f , δ_f , and A_f such that

$$f_r(0) = 0, \quad f_r(u) > 0 \text{ for all } u \in (0, \theta_f), \quad f_r(\theta_f) = 0, \quad f_r(u) = 0 \text{ for all } u \geq \theta_f, \quad (6.2)$$

and

$$u \left(1 - A_f \log \left(\frac{1}{u} \right)^{1-r} \right) \leq f_r(u) \leq u \left(1 - A_f^{-1} \log \left(\frac{1}{u} \right)^{1-r} \right), \quad (6.3)$$

for $u \in (0, \delta_f)$.

Theorem 6.1. *Suppose that the initial condition $u_0(x)$ for (6.1) is as in (1.8). If $r > 3$, then the solution $u(t, x)$ propagates with a logarithmic delay:*

$$\lim_{L \rightarrow \infty} \limsup_{t \rightarrow \infty} \sup_{x \geq L} u \left(t, 2t - \frac{3}{2} \log t + x \right) = 0, \quad (6.4)$$

and

$$\lim_{L \rightarrow \infty} \limsup_{t \rightarrow \infty} \sup_{x \leq -L} \left| u \left(t, 2t - \frac{3}{2} \log t + x \right) - \theta_f \right| = 0. \quad (6.5)$$

If $r = 3$, $u(t, x)$ propagates with a larger logarithmic delay: there exists $S_A > s_A > 3/2$ such that

$$\liminf_{t \rightarrow \infty} \sup_{x \leq 0} \left| u \left(t, 2t - S_A \log t + x \right) - \theta_f \right| = 0, \quad (6.6)$$

and

$$\lim_{t \rightarrow \infty} \sup_{x \leq 0} u \left(t, 2t - s_A \log t + x \right) = 0. \quad (6.7)$$

If $r \in (1, 3)$, then the delay is algebraic: there exist $C_f > c_f > 0$, depending only on f_r , such that

$$\lim_{t \rightarrow \infty} \sup_{x \geq 0} u \left(t, 2t - c_f t^{\frac{3-r}{1+r}} + x \right) = 0, \quad (6.8)$$

and

$$\lim_{t \rightarrow \infty} \sup_{x \leq 0} \left| u \left(t, 2t - C_f t^{\frac{3-r}{1+r}} + x \right) - \theta_f \right| = 0. \quad (6.9)$$

The proof of (6.4) follows directly from Section 3. The proofs of (6.5), (6.6), and (6.9) follow from what was done in Section 4, combined with a standard argument saying that the convergence is necessarily to the steady state θ_f (see, e.g., [17]). The bounds (6.7) and (6.8) need additional ingredients; the work is similar so we only detail the computations for (6.8). Indeed, since our non-linearity is local, we cannot “pull” information from the front as we did above when we used the value of u at the front to bound $\phi \star u$ far ahead of the front. In order to get around this, we state a weak lower bound on u .

Lemma 6.2. *Let the hypotheses of Theorem 6.1 be satisfied. Then there exists $\delta_f > 0$, depending only on f , such that*

$$u(t, x) \geq \exp\{-\delta_f t^\gamma\}$$

for all t sufficiently large and all $x \leq 2t + t^\gamma$, where we again define $\gamma = 2/(1+r)$.

Such a bound follows from the analysis of the lower bound in part (3) of Proposition 3.2 and requires no new ideas. As such, we omit the proof.

The main point in the proof of Theorem 6.1 is to use the lower bound in Lemma 6.2 on u along with the form of the non-linearity to replace the estimate of $\phi \star u$ that we used in the proof of the upper bound in Theorem 1.1 when $r \in (1, 3)$.

Proof of (6.8) assuming Lemma 6.2. We use a super-solution

$$\bar{v}(t, x) := B \exp\left\{-\left(x - 2t + 2c_f t^{2\gamma-1}\right)\right\},$$

with $c_f > 0$ to be determined. Then \bar{v} satisfies

$$\bar{v}_t = \bar{v}_{xx} + \bar{v}\left(1 - 2c_f(2\gamma - 1)t^{2\gamma-2}\right).$$

On the other hand, using the bound on f (6.3) along with Lemma 6.2, we have that, for all t sufficiently large and $x \leq 2t + t^\gamma$,

$$u_t - u_{xx} = f_r(u) \leq u\left(1 - A_f \log\left(\frac{1}{u(t, x)}\right)^{1-r}\right) \leq u\left(1 - A_f \delta_f^{1-r} t^{\gamma(1-r)}\right).$$

Recalling $2\gamma - 2 = \gamma(1 - r)$, and choosing c_f such that $A_f \delta_f^{r-1} \geq 2c_f(2\gamma - 1)$, we see that \bar{v} is a super-solution for u . \square

7. The local-in-time Harnack inequality: Proposition 1.2

Proof of Proposition 1.2. Up to a shift in time, we may assume that $t = 0$. We may also assume that $c \equiv 0$. Indeed, let

$$u_\pm(t, x) = e^{\pm t \|c\|_{L^\infty([0, T] \times \mathbb{R})}} w(t, x),$$

where w solves the heat equation

$$w_t = w_{xx},$$

with the initial condition $w(t = 0, x) = u(t = 0, x)$. Then u_+ is a super-solution to u while u_- is a sub-solution to u . Hence, we have

$$\frac{u(T, x + y)}{\|u_-\|_{L^\infty}^{1-1/p} u(T, x)^{1/p}} \leq \frac{u_+(T, x + y)}{\|u_-\|_{L^\infty}^{1-1/p} u_-(T, x)^{1/p}} \leq e^{2\|c\|_{L^\infty} T} \frac{w(T, x + y)}{\|w\|_{L^\infty}^{1-1/p} w(T, x)^{1/p}}.$$

In view of this inequality, it is enough to prove the claim for w , that is, solutions to the heat equation.

Let G be the one-dimensional heat kernel $G(t, x) = (4\pi t)^{-1/2} e^{-x^2/(4t)}$. Fix $s = (p + 1)/2p$, notice that $s \in (0, 1)$ and $sp > 1$, and let q be the dual exponent of p . Then we have

$$\begin{aligned}
w(T, x + y) &= \int_{\mathbb{R}} w(0, z) G(T, x + y - z) dz \\
&\leq \|w\|_{\infty}^{1-1/p} \int_{\mathbb{R}} w(0, z)^{1/p} G(T, x + y - z)^s G(T, x + y - z)^{1-s} dz \\
&\leq \|w\|_{\infty}^{1-1/p} \left(\int_{\mathbb{R}} w(0, z) G(T, x + y - z)^{sp} dz \right)^{1/p} \|G^{1-s}(T, \cdot)\|_q \\
&\leq C_p T^{\frac{s}{2} - \frac{1}{2p}} \|w\|_{\infty}^{1-1/p} \left(\int_{\mathbb{R}} w(0, z) G(T, x + y - z)^{sp} dz \right)^{1/p}.
\end{aligned} \tag{7.1}$$

Above, we have used that

$$\begin{aligned}
\|G^{1-s}(T, \cdot)\|_q &= (4\pi T)^{-\frac{1}{2}(1-s)} \left(\int_{\mathbb{R}^n} e^{-q(1-s)\frac{x^2}{4T}} dx \right)^{\frac{1}{q}} \\
&= (4\pi T)^{-\frac{1}{2}(1-s)} \left(\left(\frac{4T\pi}{q(1-s)} \right)^{\frac{1}{2}} \right)^{\frac{1}{q}} = C_p T^{\frac{1}{2q} - \frac{1}{2}(1-s)} = C_p T^{\frac{s}{2} - \frac{1}{2p}}.
\end{aligned}$$

We now seek a bound on $G(T, x + y - z)^{sp}$ in terms of $G(T, x - z)$. To this end, we recall that $sp > 1$, let $x' = x - z$ and we compute

$$\begin{aligned}
\frac{G(T, x' + y)^{sp}}{G(T, x')} &= (4\pi T)^{\frac{(1-sp)}{2}} \exp \left\{ -\frac{sp(x' + y)^2}{4T} + \frac{|x'|^2}{4T} \right\} \\
&= (4\pi T)^{\frac{(1-sp)}{2}} \exp \left\{ -\frac{sp|x'|^2}{4T} - \frac{spx'y}{2T} - \frac{s py^2}{4T} + \frac{|x'|^2}{4T} \right\} \\
&= (4\pi T)^{\frac{(1-sp)}{2}} \exp \left\{ -\frac{(sp-1)|x'|^2}{4T} - \frac{spx'y}{2T} - \frac{s py^2}{4T} \right\} \\
&\leq (4\pi T)^{\frac{(1-sp)}{2}} \exp \left\{ -\frac{(sp-1)|x'|^2}{4T} + \frac{(sp-1)|x'|^2}{4T} + \frac{(sp)^2 y^2}{4T(sp-1)} - \frac{s py^2}{4T} \right\} \\
&= (4\pi T)^{\frac{(1-sp)}{2}} \exp \left\{ \frac{(sp)^2 y^2}{4T(sp-1)} - \frac{s py^2}{4T} \right\} = (4\pi T)^{\frac{(1-sp)}{2}} \exp \left\{ \frac{s py^2}{4T(sp-1)} \right\}.
\end{aligned}$$

Define $\beta = \frac{sp}{4(sp-1)}$. Using the above bound in (7.1), we obtain

$$\begin{aligned}
w(t, x + y) &\leq C_p e^{\frac{\beta y^2}{T}} T^{\frac{s}{2} - \frac{1}{2p} + \frac{(1-sp)}{2p}} \|w\|_{\infty}^{1-1/p} \left(\int_{\mathbb{R}} w(0, z) G(T, x - z) dz \right)^{1/p} \\
&= C e^{\beta y^2/T} \|w\|_{\infty}^{1-1/p} w(t, x)^{1/p}.
\end{aligned}$$

In the second line we used the explicit choice of s to simplify the exponent of T . This concludes the proof. \square

Declaration of competing interest

There are no competing interests.

Appendix A. Proofs of Lemmas 2.2, 5.1 and 5.2

Proof of Lemma 5.1. The proof of this lemma is similar to that of a corresponding estimate in [17]. However, the proof there only deals with moving boundary conditions of the form $2t + r \log(t)$. Hence, for completeness, we provide a streamlined proof. Recall that ζ solves

$$\bar{\zeta}_\tau = L\bar{\zeta} + \varepsilon e^{(\gamma-1/2)\tau} \bar{\zeta}_y.$$

To rectify the fact that the operator L is not self-adjoint, we remove a Gaussian term. Let

$$\bar{\zeta}(\tau, y) = e^{-y^2/8} \zeta^*(\tau, y),$$

then ζ^* satisfies

$$\zeta_\tau^* + M\zeta^* = \varepsilon e^{(\gamma-1/2)\tau} \left(\bar{\zeta}_y^* - \frac{y}{4} \bar{\zeta}^* \right), \quad (\text{A.1})$$

where

$$M\zeta^* := -\zeta_{yy}^* + \left(\frac{y^2}{16} - \frac{3}{4} \right) \zeta^*.$$

The principle eigenvalue of M is associated to the eigenfunction

$$\psi(y) := (2\sqrt{\pi})^{-1/2} y e^{-y^2/8}.$$

Define the non-negative quadratic form

$$Q(f) := \langle Mf, f \rangle = \int_{\mathbb{R}} \left(f_y^2 + \left(\frac{y^2}{16} - \frac{3}{4} \right) f^2 \right) dy,$$

for all $f \in H^1(0, \infty)$ such that $yf \in L^2(\mathbb{R}^+)$.

Multiplying (A.1) by ζ^* and integrating, we obtain

$$\partial_\tau \|\zeta^*\|_{L^2(\mathbb{R}^+)}^2 + 2Q(\zeta^*) = -2\varepsilon e^{(\gamma-1/2)\tau} \int_0^\infty \frac{y}{4} (\zeta^*)^2 dy \leq 0.$$

Hence ζ^* is bounded uniformly in L^2 independently of τ . Next, let $\zeta_1^* = \langle \psi, \zeta^* \rangle$. We have

$$|\partial_\tau \zeta_1^*| \leq \varepsilon e^{-(\gamma-1/2)\tau} (|\langle (-\psi_y), \zeta^* \rangle| + |\zeta_1^*|) \lesssim \varepsilon e^{-(\gamma-1/2)\tau} \|\zeta^*\|_2. \quad (\text{A.2})$$

Integrating this inequality in τ and using the L^2 bound above, we obtain

$$|\zeta_1^*(\tau) - \zeta_1^*(0)| \lesssim \varepsilon \|\bar{\zeta}(0, \cdot)\|_2. \quad (\text{A.3})$$

We now show that the component of ζ^* that is orthogonal to ψ decays in time. Let

$$\zeta^{*\perp} := \zeta^* - \zeta_1^* \psi,$$

then

$$Q(\zeta^{*\perp}) \geq \frac{1}{2} \|\zeta^{*\perp}\|_2^2.$$

Using (A.1), we obtain

$$\partial_\tau \|\zeta^{*\perp}\|_{L^2(\mathbb{R}^+)}^2 + 2Q(\zeta^{*\perp}) \lesssim \varepsilon e^{-(\gamma-1/2)\tau} \|\zeta^*(0, \cdot)\|_2 \|\zeta^{*\perp}\|_2,$$

from which we deduce that

$$\|\bar{\zeta}^\perp(\tau, \cdot)\|_2 \lesssim e^{-(\gamma-1/2)\tau} \|\bar{\zeta}(0, \cdot)\|_2.$$

Gathering all estimates concludes the proof. \square

Proof of Lemma 5.2. Recall that $\bar{\zeta}$ solves

$$\bar{\zeta}_\tau = L\bar{\zeta} + \frac{1}{2} \bar{\zeta}_y.$$

To pass to a self-adjoint form, write

$$\bar{\zeta}(\tau, y) = \exp \left\{ -\frac{y^2}{8} - \frac{y}{4} \right\} \zeta^*(\tau, y),$$

so that ζ^* solves

$$\zeta_\tau^* + M_1 \zeta^* = 0,$$

where,

$$M_1 \zeta^* := -\zeta_{yy}^* + \left[\left(\frac{y^2}{16} - \frac{3}{4} \right) + \left(\frac{y}{8} + \frac{1}{16} \right) \right] \zeta^* = M \zeta^* + \left(\frac{y}{8} + \frac{1}{16} \right) \zeta^*.$$

This operator is now self-adjoint with a compact resolvent. Let ψ and λ be the principal eigenfunction and eigenvalue of the operator above satisfying the boundary condition $\psi(0) = 0$ and the normalization $\|\psi\|_{L^2(\mathbb{R}^+)} = 1$.

Observe that

$$Q(f) := \langle Mf, f \rangle = \langle Mf, f \rangle + \left\langle \left(\frac{y}{8} + \frac{1}{16} \right) f, f \right\rangle \geq 0,$$

and thus $\lambda > 0$.

Write

$$\zeta^* := \langle \psi, \zeta^* \rangle \psi + \zeta^{*\perp},$$

so that

$$Q(\zeta^{*\perp}) \geq \mu \|\zeta^{*\perp}\|_2^2, \quad (\text{A.4})$$

where μ is the second eigenvalue of M . After a time differentiation we have

$$\langle \psi_\varepsilon, \zeta^* \rangle(\tau) = \langle \psi, \zeta^* \rangle(0) e^{-\lambda \tau},$$

and as a consequence of (A.4), that

$$\|\zeta^{*\perp}\|_2(\tau) \leq \|\zeta^{*\perp}\|_2(0) e^{-\mu \tau}.$$

Then, locally we have $\|\zeta^{*\perp}\|_\infty(\tau) \lesssim e^{-\mu \tau}$ by parabolic regularity. This yields

$$\bar{\zeta} := \exp \left\{ -\frac{y^2}{8} - \frac{y}{4} \right\} \left(\left(\int_{\mathbb{R}^+} \psi(y) \exp \left\{ \frac{y^2}{8} + \frac{y}{4} \right\} \bar{\zeta}(0, y) dy \right) \psi(y) e^{-\lambda \tau} + \bar{h}(\tau, y) e^{-\mu \tau} \right), \quad (\text{A.5})$$

where \bar{h} is bounded in τ , locally in y . To finish, we simply note that, by elliptic regularity theory, for any compact set $K \subset [0, \infty)$, there exists $C_K > 0$ such that

$$\frac{y}{C_K^{-1}} \leq \psi(y) \leq C_K y \quad \text{for all } y \in K$$

since $\psi(0) = 0$. This concludes the proof. \square

Proof of Lemma 2.2. As before, we pass to the moving frame and remove an exponential: let

$$v(t, x) = e^{-x + (2t - \tilde{s}_\phi \log(t+t_0))} \bar{v}(t, x - (2t - \tilde{s}_\phi \log(t+t_0))).$$

Then, we want

$$\bar{v}_t \geq \bar{v}_{xx} - \bar{v}_x \frac{\tilde{s}_\phi}{t+t_0} + \bar{v} \left(\frac{\tilde{s}_\phi}{t+t_0} - \nu \right).$$

Changing now to self-similar variables and recalling the definition $\varepsilon = t_0^{-1/2}$, let

$$\tau = \log \left(1 + \frac{t}{t_0} \right), \quad y = \frac{x}{\sqrt{t+t_0}}, \quad \zeta(\tau, y) = \bar{v} \left(t_0(e^\tau - 1), t_0^{1/2} e^{\tau/2} y \right),$$

$$\text{and } N(\tau, y) := t_0 e^\tau \bar{v} \left(t_0(e^\tau - 1), t_0^{1/2} e^{\tau/2} y \right) = \frac{1}{C A_\phi \left(y + \varepsilon e^{-\tau/2} ((S_\phi - \tilde{s}_\phi)(\tau + \log(t_0)) + L) \right)^2}.$$

The important point here is that $N(\infty, \cdot)$ is neither infinity nor zero as it would be for another choice of r . This is what induces the larger delay.

Then we must find ζ that satisfies:

$$\zeta_\tau - L\zeta + (1 - \tilde{s}_\phi + N(0, y))\zeta \geq \tilde{s}_\phi \varepsilon e^{-\tau/2} \zeta_y + \zeta (N(0, y) - N(\tau, y)).$$

We have changed the order to emphasize that the right hand side is a small error.

Let $\bar{\zeta} = e^{-\tau/2} e^{y^2/8} \zeta$. The above yields

$$\bar{\zeta}_\tau + M\bar{\zeta} + \left(\frac{3}{2} - \tilde{s}_\phi + N(0, y)\right)\bar{\zeta} \geq \tilde{s}_\phi \varepsilon e^{-\tau/2} \bar{\zeta}_y - \frac{y}{4} \tilde{s}_\phi \varepsilon e^{-\tau/2} \bar{\zeta} + \bar{\zeta} (N(0, y) - N(\tau, y)). \quad (\text{A.6})$$

We now define $\bar{\zeta}$. As above $M + N(0, \cdot)$ is self-adjoint with a compact resolvent. Let $\bar{\zeta}_0$ and λ_0 be its principle eigenlements. Using the Rayleigh quotient and testing with $ye^{-y^2/8}$, we see immediately that $\lambda_0 > 0$. We need only verify that $\bar{\zeta}$ satisfies (A.6); indeed, setting $\tilde{s}_\phi = 3/2 + \lambda_0/2$,

$$\begin{aligned} \bar{\zeta}_\tau + M\bar{\zeta} + \left(\frac{3}{2} - \tilde{s}_\phi + N(0, y)\right)\bar{\zeta} - \tilde{s}_\phi \varepsilon e^{-\tau/2} \bar{\zeta}_y + \frac{y}{4} \tilde{s}_\phi \varepsilon e^{-\tau/2} \bar{\zeta} - \bar{\zeta} (N(0, y) - N(\tau, y)) \\ = \frac{\lambda_0}{2} \bar{\zeta} - \tilde{s}_\phi \varepsilon e^{-\tau/2} \bar{\zeta}_y + \frac{y}{4} \tilde{s}_\phi \varepsilon e^{-\tau/2} \bar{\zeta} - \bar{\zeta} (N(0, y) - N(\tau, y)). \end{aligned}$$

The first, third, and fourth terms are all positive. A simple maximum principle argument yields that $-\bar{\zeta}_y$ is positive for all $y \geq y_0$ for some $y_0 > 0$. The Hopf maximum principle implies $(\lambda_0/2)\bar{\zeta} - \tilde{s}_\phi \varepsilon e^{-\tau/2} \bar{\zeta}_y > 0$ for all $y > e^{-\tau/2}/C$ for some $C > 0$. Thus, $\bar{\zeta}$ satisfies (A.6) on $[e^{-\tau/2}/C, \infty)$, which, after translating back to physical variables, concludes the proof. \square

References

- [1] H. Berestycki, G. Nadin, B. Perthame, L. Ryzhik, The non-local Fisher-KPP equation: travelling waves and steady states, *Nonlinearity* 22 (2009) 2813–2844.
- [2] H. Berestycki, L. Nirenberg, Travelling fronts in cylinders, *Ann. Inst. Henri Poincaré, Anal. Non Linéaire* 9 (5) (1992) 497–572.
- [3] E. Bouin, C. Henderson, Forthcoming.
- [4] E. Bouin, C. Henderson, L. Ryzhik, The Bramson logarithmic delay in the cane toads equations, preprint, <https://arxiv.org/abs/1610.03285>, 2016.
- [5] M. Bramson, Maximal displacement of branching Brownian motion, *Commun. Pure Appl. Math.* 31 (5) (1978) 531–581.
- [6] M. Bramson, Convergence of solutions of the Kolmogorov equation to travelling waves, *Mem. Amer. Math. Soc.* 44 (285) (1983), iv+190.
- [7] N.F. Britton, Spatial structures and periodic travelling waves in an integro-differential reaction-diffusion population model, *SIAM J. Appl. Math.* 50 (6) (1990) 1663–1688.
- [8] B. Dennis, J.M. Ponciano, S.R. Lele, M.L. Taper, D.F. Staples, Estimating density dependence, process noise, and observation error, *Ecol. Monogr.* 76 (3) (2006) 323–341.
- [9] A. Ducrot, On the large time behaviour of the multi-dimensional Fisher-KPP equation with compactly supported initial data, *Nonlinearity* 28 (4) (2015) 1043–1076.
- [10] J. Fang, X.-Q. Zhao, Monotone wavefronts of the nonlocal Fisher-KPP equation, *Nonlinearity* 24 (11) (2011) 3043–3054.
- [11] M. Fang, O. Zeitouni, Slowdown for time inhomogeneous branching Brownian motion, *J. Stat. Phys.* 149 (1) (2012) 1–9.
- [12] G. Faye, M. Holzer, Modulated traveling fronts for a nonlocal Fisher-KPP equation: a dynamical systems approach, *J. Differ. Equ.* 258 (7) (2015) 2257–2289.
- [13] S. Genieys, V. Volpert, P. Auger, Pattern and waves for a model in population dynamics with nonlocal consumption of resources, *Math. Model. Nat. Phenom.* 1 (1) (2006) 65–82.
- [14] S.A. Gourley, Travelling front solutions of a nonlocal Fisher equation, *J. Math. Biol.* 41 (3) (2000) 272–284.
- [15] F. Hamel, Qualitative properties of monostable pulsating fronts: exponential decay and monotonicity, *J. Math. Pures Appl.* (9) 89 (4) (2008) 355–399.
- [16] F. Hamel, C. Henderson, Propagation in a Fisher-KPP equation with non-local advection, preprint, <https://arxiv.org/abs/1707.05710>, 2017.
- [17] F. Hamel, J. Nolen, J.-M. Roquejoffre, L. Ryzhik, A short proof of the logarithmic Bramson correction in Fisher-KPP equations, *Netw. Heterog. Media* 8 (1) (2013) 275–289.
- [18] F. Hamel, J. Nolen, J.-M. Roquejoffre, L. Ryzhik, The logarithmic delay of KPP fronts in a periodic medium, *J. Eur. Math. Soc.* 18 (3) (2016) 465–505.
- [19] F. Hamel, L. Ryzhik, On the nonlocal Fisher-KPP equation: steady states, spreading speed and global bounds, *Nonlinearity* 27 (11) (2014) 2735–2753.
- [20] P. Maillard, O. Zeitouni, Slowdown in branching Brownian motion with inhomogeneous variance, *Ann. Inst. Henri Poincaré Probab. Stat.* 52 (3) (2016) 1144–1160.

- [21] G. Nadin, L. Rossi, L. Ryzhik, B. Perthame, Wave-like solutions for nonlocal reaction-diffusion equations: a toy model, *Math. Model. Nat. Phenom.* 8 (3) (2013) 33–41.
- [22] J. Nolen, J.-M. Roquejoffre, L. Ryzhik, Power-like delay in time inhomogeneous Fisher-KPP equations, *Commun. Partial Differ. Equ.* 40 (3) (2015) 475–505.
- [23] J. Nolen, J.-M. Roquejoffre, L. Ryzhik, Refined long time asymptotics for Fisher-KPP fronts, arXiv preprint, arXiv:1607.08802, 2016.
- [24] J. Nolen, J.-M. Roquejoffre, L. Ryzhik, Convergence to a single wave in the Fisher-KPP equation, *Chin. Ann. Math., Ser. B* 38 (2) (2017) 629–646.
- [25] S. Penington, The spreading speed of solutions of the non-local Fisher-kpp equation, <https://arxiv.org/abs/1708.07965>.
- [26] M.I. Roberts, A simple path to asymptotics for the frontier of a branching Brownian motion, *Ann. Probab.* 41 (5) (2013) 3518–3541.
- [27] K. Uchiyama, The behavior of solutions of some nonlinear diffusion equations for large time, *J. Math. Kyoto Univ.* 18 (3) (1978) 453–508.