

Convex integration solutions to the transport equation with full dimensional concentration $\star, \star\star$

Stefano Modena ^{a,*}, Gabriel Sattig ^b

^a Technische Universität Darmstadt, Fachbereich Mathematik, D-64289 Darmstadt, Germany

^b Universität Leipzig, Mathematisches Institut, D-04109 Leipzig, Germany

Received 16 April 2019; received in revised form 13 January 2020; accepted 12 March 2020

Available online 18 March 2020

Abstract

We construct infinitely many incompressible Sobolev vector fields $u \in C_t W_x^{1, \tilde{p}}$ on the periodic domain \mathbb{T}^d for which uniqueness of solutions to the transport equation fails in the class of densities $\rho \in C_t L_x^p$, provided $1/p + 1/\tilde{p} > 1 + 1/d$. The same result applies to the transport-diffusion equation, if, in addition, $p' < d$.

© 2020 L'Association Publications de l'Institut Henri Poincaré. Published by Elsevier B.V. All rights reserved.

Keywords: Transport equation; Continuity equation; Convex integration; Non-uniqueness; Concentrated Mikado

1. Introduction

This paper deals with the problem of (non)uniqueness of solution to the Cauchy problem for the transport equation

$$\partial_t \rho + \nabla \rho \cdot u = 0, \tag{1.1}$$

on the d -dimensional flat torus $\mathbb{T}^d := \mathbb{R}^d / \mathbb{Z}^d$, where $u : [0, T] \times \mathbb{T}^d \rightarrow \mathbb{R}^d$ is a given (locally integrable) vector field and $\rho : [0, T] \times \mathbb{T}^d \rightarrow \mathbb{R}$ is the unknown density. We will always assume that u is incompressible, i.e.

$$\operatorname{div} u = 0, \tag{1.2}$$

in the sense of distributions. Under this condition, (1.1) is formally equivalent to the continuity equation

$$\partial_t \rho + \operatorname{div}_x(\rho u) = 0. \tag{1.3}$$

We prove the following theorem.

\star This research was supported by the ERC Grant Agreement No. 724298. The authors wish to thank Prof. László Székelyhidi and Jonas Hirsch for several useful discussions on the topic of this paper.

$\star\star$ A part of this paper was written during the visit of the authors to the Hausdorff Research Institute for Mathematics (HIM), University of Bonn, Jan-Apr 2019. This visit was supported by the HIM. Both, this support and the hospitality of HIM, are gratefully acknowledged.

* Corresponding author.

E-mail addresses: modena@mathematik.tu-darmstadt.de (S. Modena), gabriel.sattig@math.uni-leipzig.de (G. Sattig).

Theorem 1.1. *Let $p \in [1, \infty)$, $\tilde{p} \in [1, \infty)$, and assume that*

$$\frac{1}{p} + \frac{1}{\tilde{p}} > 1 + \frac{1}{d}. \tag{1.4}$$

Then there are infinitely many incompressible vector fields satisfying

$$u \in C_t L_x^{p'} \cap C_t W_x^{1, \tilde{p}} \tag{1.5}$$

for which uniqueness of distributional solutions to the Cauchy problem for the transport equation (1.1) fails in the class of densities

$$\rho \in C_t L_x^p.$$

Moreover, if $p = 1$, it holds $u \in C([0, T] \times \mathbb{T}^d)$.

Here and in the following we will use the notation $C_t L_x^p := C([0, T], L^p(\mathbb{T}^d))$, and, similarly, $L_t' L_x^p := L^r((0, T), L^p(\mathbb{T}^d))$.

Remark.

- (i) In the proof we will show that there are non-trivial solutions with zero initial data, thus proving non-uniqueness. However, any smooth function with zero mean value can serve as initial data for our “wild solutions”. For the precise statement see Theorem 1.2.
- (ii) As a matter of fact, one can strengthen condition (1.5) and produce vector fields which satisfy

$$u \in C_t L_x^{p'} \cap \bigcap_{\substack{\tilde{p} \text{ such that} \\ (1.4) \text{ holds}}} C_t W_x^{1, \tilde{p}}$$

and, moreover, $\|u\|_{L^{p'}} \leq \varepsilon$, for any fixed $\varepsilon > 0$. See Theorem 1.2 below.

- (iii) Theorem 1.1 can be extended to cover the case of the transport-diffusion equation and to produce more regular densities and fields, provided more restrictive conditions on the exponents p, \tilde{p} are assumed. See Theorems 1.3 and 1.4 below for the precise statements.

1.1. Background

It is well known that, when u is at least Lipschitz continuous (in the space variable), the solution to (1.1) is given by the implicit formula

$$\rho(t, X(t, x)) = \rho(0, x), \tag{1.6}$$

where $X(t, x)$ is the flow solving the ODE

$$\begin{aligned} \partial_t X(t, x) &= u(t, X(t, x)), \\ X(0, x) &= x. \end{aligned} \tag{1.7}$$

It is in general of great importance, both for theoretical interest and for the applications to many physical models, to study the well posedness of the Cauchy problem (1.1), in the case the vector field u is not smooth, i.e. less than Lipschitz continuous.

There are several ways to state the well posedness problem in the weak setting. The one we propose here is one possibility. We refer to [17] for a more comprehensive discussion. Fix an exponent $p \in [1, \infty]$ and denote by p' its dual Hölder

$$\frac{1}{p} + \frac{1}{p'} = 1.$$

We ask two questions.

(a) Do existence and uniqueness of distributional solutions to (1.1) hold in the class of densities

$$\rho \in L_t^\infty L_x^p \tag{1.8}$$

for a given vector field

$$u \in L_t^1 L_x^{p'}? \tag{1.9}$$

(b) Is the relation (1.6) still valid, in some weak sense? In other words, is there still a connection between the Lagrangian world (1.7) and the Eulerian one (1.1)?

Let us observe that the choice of the class (1.8) is motivated by the fact that, for smooth solutions of (1.1)-(1.2), every L^p norm is constant in time: it is thus reasonable to expect that, for weak solutions, the L^p norm, if not constant, remains, at least, uniformly bounded in time. Once the class for the density (1.8) is fixed, the choice (1.9) for the vector field is natural, because in this way the product $\rho u \in L^1((0, T) \times \mathbb{T}^d)$ and thus the transport equation (1.1), in its equivalent form (1.3), can be considered in distributional sense.

We list now some answers to the questions (a), (b) above, which can be found in the literature. The first consideration is that the *existence* of distributional solutions is a pretty easy task. Indeed, regularizing the vector field and the initial datum, one can use the classical theory for ODE and formula (1.6) to produce a sequence of approximate solutions, which turns out to be uniformly bounded in $L_t^\infty L_x^p$. From such sequence one can then extract a weakly converging subsequence, whose limit is a solution to (1.1), because of the linearity of the equation.

Let us now discuss some *uniqueness* results. In their groundbreaking paper [12], R. DiPerna and P.L. Lions proved that, for every $p \in [1, \infty]$, uniqueness holds in the class of densities (1.8) for a given vector field u as in (1.9), provided, in addition,

$$u \in L_t^1 W_x^{1,p'}. \tag{1.10}$$

Moreover, the incompressibility assumption can be substituted by the weaker requirement $\operatorname{div} u \in L^\infty$ (see also [19] for a further relaxation in the case of the continuity equation). DiPerna and Lions’ proof is based on a regularization argument. Denote by ρ^ε (resp. u^ε) the convolution of ρ (resp. u) with a compactly supported standard mollifier $\eta_\varepsilon = \varepsilon^{-d} \eta(\cdot/\varepsilon)$ and observe that

$$\partial_t \rho^\varepsilon + \operatorname{div}(\rho^\varepsilon u^\varepsilon) = \operatorname{div}(\rho^\varepsilon u^\varepsilon - (\rho u)^\varepsilon) =: r_\varepsilon,$$

i.e. ρ^ε solves (1.3) with smooth vector field u^ε up to some error, the “commutator” r_ε , which arises from the fact that the convolution does not commute with multiplication. By partial integration r_ε takes the form (assuming $\operatorname{div} u = 0$)

$$r_\varepsilon(t, x) = \int_{\mathbb{R}^d} \rho(t, x - \varepsilon z) \frac{u(t, x) - u(t, x - \varepsilon x)}{\varepsilon} \nabla \eta(z) dz.$$

Such expression suggests that the commutator r_ε converges to zero as $\varepsilon \rightarrow 0$ (and thus uniqueness of solutions holds) if the integrability of ρ is Hölder dual to the integrability of ∇u , i.e. if $\rho \in L_t^\infty L_x^p$, and $\nabla u \in L_t^1 L_x^{p'}$, which is exactly DiPerna and Lions’ condition (1.10). In other words, the interplay between the integrability of the density and the integrability of the derivative of the vector field plays a crucial role: very roughly speaking, a Sobolev vector field is “Lipschitz like” on a very large set, and there is just a very small “bad” set, where ∇u can be very large. A density ρ with integrability L^p that “matches” the integrability $L^{p'}$ of ∇u does not see the bad set of u , and this implies uniqueness.

A natural question is now whether it is possible to lower the regularity (1.10) of u and still have uniqueness of solutions in $L_t^\infty L_x^p$.

In the class of *bounded* densities, (i.e. $p = \infty$ in our notation), L. Ambrosio [1] showed in 2004 that uniqueness holds if the vector field $u \in L^1((0, T), BV(\mathbb{T}^d))$ and it has bounded divergence, whereas S. Bianchini and P. Bonicatto in [3] were able to prove uniqueness in the BV framework for the more general class of *nearly incompressible* vector fields.

Concerning question (b) above, it is a general principle in the theory of the transport equation that, whenever existence and uniqueness for the PDE (1.1) holds in the class of *bounded* densities, then existence and uniqueness

holds also for the ODE (1.7), in the sense of the *regular Lagrangian flow* and, moreover, the bridge (1.6) between the Lagrangian world and the Eulerian one still holds true. We refer to [2] for a detailed discussion in this direction.

From the analysis above, it follows that the uniqueness results present in the literature are based essentially on two assumptions on the vector field: on one side, a bound on the derivative Du is needed (e.g. u Sobolev or BV); on the other side, a condition on the divergence of u is required (e.g. $\operatorname{div} u = 0$, or $\operatorname{div} u \in L^\infty$, or u nearly incompressible).

The most part of the counterexamples to uniqueness that can be found in the literature are based on the absence of at least one of those two conditions. There are counterexamples to uniqueness with Sobolev vector field with unbounded divergence (e.g. in DiPerna and Lions’ paper [12]), and there are counterexamples to uniqueness for incompressible vector fields, which do not possess one full derivative (e.g. $u \in W^{s,1}$ for every $s < 1$, but $u \notin W^{1,1}$), see, for instance, [12], [11]. All such counterexamples are based on the failure of uniqueness at a *Lagrangian* level: one constructs a pathological vector field for which the ODE admits two different flows of solutions and then uses such flows to produce non-unique solutions to the PDE: once again, the connection (1.6) is crucial.

1.2. Non-uniqueness for Sobolev vector fields and our contribution

The mentioned counterexamples, therefore, do not answer the question whether uniqueness holds in the class of densities (1.8), if

$$u \text{ is incompressible, } u \in L_t^1 W_x^{1,\tilde{p}}, \text{ but } \tilde{p} < p'. \tag{1.11}$$

In such framework there are two competing mechanisms. On one side, by DiPerna and Lions result, uniqueness holds, at least, in the class of bounded densities, and thus, by the observation made before, uniqueness at the Lagrangian level is satisfied (again in the sense of the regular Lagrangian flow): in other words, the vector field is very well behaved from the ODE point of view. On the other side, the integrability of ρ and the integrability of Du do not “match” anymore and thus, referring the heuristic introduced above, it could happen that an L^p density “sees the bad set” of a $W^{1,\tilde{p}}$ vector field, so that purely Eulerian non-uniqueness phenomena could appear.

The framework (1.11) was considered, for the first time, quite recently in [17] and [18], where the analog of Theorem 1.1 was proven, with assumption (1.4) substituted by the strongest assumption

$$\frac{1}{p} + \frac{1}{\tilde{p}} > 1 + \frac{1}{d-1}, \tag{1.12}$$

using a convex integration approach and exploiting a *concentration* mechanism, in the spirit of the *intermittency* added to the convex integration schemes by T. Buckmaster and V. Vicol in [6].

Our main result, namely Theorem 1.1, shows that such approach can be extended to produce examples of non-uniqueness for the transport equation with *full dimensional concentration*, i.e. with d instead of $d - 1$ in (1.12). Notice that the result in [17,18] and our Theorem 1.1 in particular implies that the duality between Lagrangian and Eulerian world is completely destroyed, even for Sobolev and incompressible (thus, quite “well behaved” vector field): there are many distributional solutions, but only one among them is transported by the regular Lagrangian flow as in (1.6).

It is still an open question whether uniqueness of weak solutions to (1.1) holds if the Sobolev integrability \tilde{p} of the field, $Du \in L_t^1 L_x^{\tilde{p}}$, lies in the range

$$1 < \frac{1}{p} + \frac{1}{\tilde{p}} \leq 1 + \frac{1}{d}, \tag{1.13}$$

and thus whether Theorem 1.1 is or is not optimal. Let us nevertheless observe that, for $p = 1$, Theorem 1.1 provides existence of continuous vector fields

$$u \in C_t W_x^{1,\tilde{p}} \tag{1.14}$$

for every $\tilde{p} < d$, for which uniqueness fails (in the class $\rho \in C_t L_x^1$). On the other side, in a recent result by L. Caravenna and G. Crippa [7,8] uniqueness (for $\rho \in L_{t,x}^1$) is proven, provided (1.14) is satisfied for some $\tilde{p} > d$ (in

particular u is continuous) and u satisfies the additional assumption of “uniqueness of forward-backward characteristics”. We refer to [7,8] for the precise definition. Such result could suggest that, at least in the case $p = 1$, Theorem 1.1 (and in particular condition (1.4)) could be sharp.

A last point is worth mentioning. Contrary to other recent results in convex integration (e.g. [6,9,15,16]) where *concentration* or *intermittency* have been used, in this paper we use a completely physical space based approach and we deliberately avoid any use of Fourier methods and Littlewood-Paley theory. This has, in our opinion, at least two advantages. First, the paper is completely self contained, in particular we do not use any abstract theorem on Fourier multipliers. Secondly, we think that a proof developed in the physical space can provide a better understanding of the structure of the “anomalous” vector fields we are exhibiting and therefore could help in getting an insight on the relation, if any, between the (very well behaved) Lagrangian structure of the vector fields and the non-Lagrangian solutions we construct.

We conclude this section observing that the proof of Theorem 1.1 is an immediate consequence of the following more general theorem, whose proof is the main topic of the paper.

Theorem 1.2 (Solutions for Sobolev vector fields). *Let $\varepsilon > 0$, let $\bar{\rho} \in C^\infty([0, T] \times \mathbb{T}^d)$ with zero mean value in the space variable and let $\bar{u} \in C^\infty([0, T] \times \mathbb{T}^d, \mathbb{R}^d)$ be a divergence-free vector field. Set $E := \{t \in [0, T] : \partial_t \bar{\rho} + \operatorname{div}(\bar{\rho}\bar{u}) = 0\}$. Let $p \in [1, \infty)$ and define $q \in [1, \infty)$ such that*

$$\frac{1}{p} + \frac{1}{q} = 1 + \frac{1}{d}. \tag{1.15}$$

Then there are functions $\rho : [0, T] \times \mathbb{T}^d \rightarrow \mathbb{R}$ and $u : [0, T] \times \mathbb{T}^d \rightarrow \mathbb{R}^d$ such that

- (i) $\rho \in C([0, T], L^p(\mathbb{T}^d))$ and $u \in C([0, T], L^{p'}(\mathbb{T}^d)) \cap \bigcap_{\bar{p} < q} C([0, T], W^{1, \bar{p}}(\mathbb{T}^d))$.
 If $p = 1$ then u is also continuous: $u \in C([0, T] \times \mathbb{T}^d)$;
- (ii) (ρ, u) is a distributional solution of (1.3)–(1.2);
- (iii) $(\rho, u)(t) = (\bar{\rho}, \bar{u})(t)$ for all $t \in E$;
- (iv) $\|\rho(t) - \bar{\rho}(t)\|_{L^p} < \varepsilon$ for all $t \in [0, T]$.

Statement (iv) can be replaced by the similar

$$(iv') \quad \|u(t) - \bar{u}(t)\|_{L^{p'}} < \varepsilon \text{ for all } t \in [0, T].$$

From this theorem, Theorem 1.1, i.e. the non-uniqueness of the transport equation, can be easily deduced.

Proof of Theorem 1.1, assuming Theorem 1.2. Let $\bar{\rho} \in C^\infty(\mathbb{T}^d)$ with zero mean value but not identically zero. Choose $\chi : [0, T] \rightarrow [0, 1]$ smooth such that χ is equal to zero on $[0, T/3]$ and one on $[2T/3, T]$. Then the function $(t, x) \mapsto \chi(t)\bar{\rho}(x)$ is smooth and has zero mean value in x at any time. We can apply Theorem 1.2 on $\chi\bar{\rho}$ and $\bar{u} \equiv 0$ and obtain a solution of the transport equation (ρ, u) with the claimed regularity. As at times $t \in [0, \frac{T}{3}] \cup [\frac{2T}{3}, T]$ the transport equation is solved by $(\chi\bar{\rho}, \bar{u})$ in the strong sense, in particular the initial and final values of ρ are maintained because of statement (iii) of the theorem. Therefore $\rho|_{t=0} \equiv 0$ and $\rho|_{t=T} = \bar{\rho} \neq 0$. \square

1.3. Some comments on the method used in the proof

The proof of Theorem 1.2 is based on a convex integration technique: smooth approximate solutions to the continuity equations are constructed, which in the weak limit produce an exact but only distributional solution. In each iterations step the error is decreased by adding a small oscillating perturbation to both density and velocity field.

In the past years convex integration has been applied very successfully on the Euler equations in order positively prove Onsager’s conjecture (see, for instance [14,5]). However, for obtaining Sobolev vector fields, i.e. fields with one full derivative (in some $L^{\bar{p}}$ space) new ideas are required. Inspired by the *intermittent Beltrami flow* used in the [6] (see also [4] for the related notion of *intermittent jets*), L. Székelyhidi and the first author adopted, as building block of their construction in the mentioned papers [17,18], some stationary solutions to the continuity equation called

concentrated Mikado densities and field, proving the analog of Theorem 1.1 under the less restrictive assumption (1.12). The idea of using “Mikado flows” for the equation of fluid dynamics was introduced for the first time by S. Daneri and L. Székelyhidi in [10]. The “concentrated” Mikado are suitable modifications of the standard Mikado, having different scaling in different L^p norms. The $d - 1$ in (1.12) comes from the fact that Mikado functions depends only on $d - 1$ coordinates and thus only a $(d - 1)$ -dimensional concentration is possible.

In the present paper, we are able to substitute $d - 1$ with d , as we use, as building block of our construction, suitable approximate solutions to the continuity equation, called *space-time Mikado densities and fields*, see Section 4.1 for the precise definition. Adding the time dependence to the building block allows, roughly speaking, to gain one further dimension and thus to pass from (1.12) to (1.4).

1.4. Extension to transport-diffusion and to higher regularity

Similarly to [17,18], Theorem 1.2 (and thus also Theorem 1.1) can be extended to cover the case of the transport-diffusion equation

$$\begin{aligned} \partial_t \rho + \operatorname{div}(\rho u) - \Delta \rho &= 0, \\ \operatorname{div} u &= 0, \end{aligned} \tag{1.16}$$

provided more restrictive conditions on the exponent p, \tilde{p} are assumed. Roughly speaking, the non-uniqueness produced by the transport term $\operatorname{div}(\rho u)$ (i.e. by the interplay between density and field) can be so strong that it can beat the regularizing effect induced by a diffusion operator (see to [17] for a more comprehensive discussion on this subject).

Theorem 1.3 (Analog of Theorem 1.2 for the transport-diffusion equation). *Let $\varepsilon > 0$, let $\bar{\rho} \in C^\infty([0, T] \times \mathbb{T}^d)$ with zero mean value and let $\bar{u} \in C^\infty([0, T] \times \mathbb{T}^d, \mathbb{R}^d)$ be a divergence-free field. Set $E := \{t \in [0, T] : \partial_t \bar{\rho} + \operatorname{div}(\bar{\rho} \bar{u}) - \Delta \bar{\rho} = 0\}$. Let $p \in (1, \infty)$ and $\tilde{p} \in [1, \infty)$ such that*

$$\frac{1}{p} + \frac{1}{\tilde{p}} > 1 + \frac{1}{d}, \quad p' < d. \tag{1.17}$$

Then there are functions $\rho : [0, T] \times \mathbb{T}^d \rightarrow \mathbb{R}$ and $u : [0, T] \times \mathbb{T}^d \rightarrow \mathbb{R}^d$ such that

- (i) $\rho \in C([0, T], L^p(\mathbb{T}^d))$ and $u \in C([0, T], L^{p'}(\mathbb{T}^d)) \cap C([0, T], W^{1, \tilde{p}}(\mathbb{T}^d))$;
- (ii) (ρ, u) is a distributional solution of (1.16);
- (iii) $(\rho, u)(t) = (\bar{\rho}, \bar{u})(t)$ for all $t \in E$;
- (iv) $\|\rho(t) - \bar{\rho}(t)\|_{L^p} < \varepsilon$ for all $t \in [0, T]$.

Statement (iv) can be replaced by the similar

$$(iv') \quad \|u(t) - \bar{u}(t)\|_{L^{p'}} < \varepsilon \text{ for all } t \in [0, T].$$

Remark. Notice that (1.17) in particular requires $d > 2$, so we cannot show non-uniqueness for the dissipative equation for $d = 2$ as in the “inviscid” transport equation.

Theorems 1.2 and 1.3 can be further generalized to cover the generalized transport-diffusion equation

$$\begin{aligned} \partial_t \rho + \operatorname{div}_x(\rho u) + L_k \rho &= 0, \\ \operatorname{div}_x u &= 0, \end{aligned} \tag{1.18}$$

where L_k is any constant-coefficient linear differential operator of grade k (not necessarily elliptic), and to produce more regular densities and vector fields.

Theorem 1.4 (Analog for solutions with higher regularity and higher order diffusion). *Let $\varepsilon > 0$, let $\bar{\rho} \in C^\infty([0, T] \times \mathbb{T}^d)$ with zero mean value and let $\bar{u} \in C^\infty([0, T] \times \mathbb{T}^d, \mathbb{R}^d)$ be a divergence-free field. Let $p, \tilde{p} \in [1, \infty)$ and $m, \tilde{m} \in \mathbb{N}$ such that*

$$\frac{1}{p} + \frac{1}{\tilde{p}} > 1 + \frac{m + \tilde{m}}{d} \text{ and } \tilde{p} < \frac{d}{\tilde{m} + k - 1}. \tag{1.19}$$

Then there are $s \in [p, \infty]$ and functions $\rho : [0, T] \times \mathbb{T}^d \rightarrow \mathbb{R}$ and $u : [0, T] \times \mathbb{T}^d \rightarrow \mathbb{R}^d$ such that

- (i) $\rho \in C([0, T], L^s(\mathbb{T}^d))$, $u \in C([0, T], L^{s'}(\mathbb{T}^d))$ and, moreover, $\rho \in C([0, T], W^{m, p}(\mathbb{T}^d))$, $u \in C([0, T], W^{\tilde{m}, \tilde{p}}(\mathbb{T}^d))$;
- (ii) (ρ, u) is a distributional solution of (1.18);
- (iii) $(\rho, u)(t) = (\bar{\rho}, \bar{u})(t)$ for all $t \in E$ defined as in Theorem 1.2;
- (iv) $\|\rho(t) - \bar{\rho}(t)\|_{L^s} < \varepsilon$ for all $t \in [0, T]$.

Statement (iv) can be replaced by the similar

$$(iv') \quad \|u(t) - \bar{u}(t)\|_{L^{s'}} < \varepsilon \text{ for all } t \in [0, T].$$

Remark. Observe also that, if we choose $m = 0, \tilde{m} = 1, k = 2$ in Theorem 1.4, the first condition in (1.19) reduces to the first condition in (1.17), nevertheless (1.19) is not equivalent to (1.17). Indeed (1.17) implies (1.19), but the viceversa is not true, in general. This can be explained by the fact that Theorem 1.3, for any given p , produces a vector field $u \in C_t L_x^p$, whereas Theorem 1.4 produces $u \in C_t L_x^{s'}$ for some $s' \leq p'$.

Remark. In Section 2 we state the main Proposition of this paper, namely Proposition 2.1, and we show how Theorem 1.2 can be deduced from Proposition 2.1. In Sections 3-6 we give a complete proof of Proposition 2.1, assuming $p > 1$, for the sake of simplicity. In Section 7 we give a sketch of the proof of Proposition 2.1 in the case $p = 1$ as well as a sketch of the proofs of Theorems 1.3 and 1.4.

1.5. Notations

We fix some notations which will be used throughout the paper.

- Integrals, L^p -norms and Sobolev norms of functions defined on $[0, T] \times \mathbb{T}^d$ will always be evaluated on the space \mathbb{T}^d at a single time t , we will write

$$\|\rho(t)\|_{L^p} = \|\rho(t, \cdot)\|_{L^p(\mathbb{T}^d)} \text{ and } \int \rho = \int_{\mathbb{T}^d} \rho(t, x) dx.$$

- Similarly, all differential operators (except ∂_t , of course) apply on the space variable: $\partial_j = \frac{\partial}{\partial x_j}$, $\operatorname{div} = \operatorname{div}_x$, $\Delta = \Delta_x, \dots$
- In contrast, C^k -norms are always evaluated on the space-time $[0, T] \times \mathbb{T}^d$.
- If a function is stated to have zero mean value we always mean ‘in the space variable’. Define C_0^∞ to be the space of smooth functions which have zero mean value:

$$C_0^\infty(\mathbb{T}^d) := \left\{ f : \mathbb{T}^d \rightarrow \mathbb{R} \text{ smooth such that } \int_{\mathbb{T}^d} f(x) dx = 0 \right\}.$$

- If not specified otherwise, for a periodic function $f : \mathbb{T}^d \rightarrow \mathbb{R}$ and $\lambda \in \mathbb{N}_+$, $f_\lambda : \mathbb{T}^d \rightarrow \mathbb{R}$ denotes the dilation $f_\lambda(x) = f(\lambda x)$. Note that

$$\|D^k f_\lambda\|_{L^p(\mathbb{T}^d)} = \lambda^k \|D^k f\|_{L^p(\mathbb{T}^d)}. \tag{1.20}$$

2. Main proposition and proof of the theorem

In this section we state the main proposition of this paper, Proposition 2.1, and we use it in order to prove Theorem 1.2. Proposition 2.1 will be proven in details in Sections 3-6, assuming, for simplicity, $p > 1$. A sketch of the proof in the case $p = 1$ can be found in Section 7.1.

We introduce the (incompressible) continuity-defect equation

$$\begin{aligned} \partial_t \rho + \operatorname{div}(\rho u) &= -\operatorname{div} R \\ \operatorname{div} u &= 0 \end{aligned} \tag{2.1}$$

as an approximation of the transport equation. The iteration step of the Convex Integration scheme deals with solution to this system.

Proposition 2.1. *There is a constant $M > 0$ such that the following holds. Let $p \in [1, \infty)$ and $\tilde{p} \in [1, \infty)$ so that*

$$\frac{1}{p} + \frac{1}{\tilde{p}} > 1 + \frac{1}{d}. \tag{2.2}$$

Then for any $\delta, \eta > 0$ and any smooth solution (ρ_0, u_0, R_0) of the continuity defect equation (2.1) there is another smooth solution (ρ_1, u_1, R_1) which fulfills the estimates

$$\|\rho_1(t) - \rho_0(t)\|_{L^p} \leq M\eta \|R_0(t)\|_{L^1}^{1/p} \tag{2.3a}$$

$$\|u_1(t) - u_0(t)\|_{L^{p'}} \leq \frac{M}{\eta} \|R_0(t)\|_{L^1}^{1/p'} \tag{2.3b}$$

$$\|u_1(t) - u_0(t)\|_{W^{1,\tilde{p}}} \leq \delta \tag{2.3c}$$

$$\|R_1(t)\|_{L^1} \leq \delta \tag{2.3d}$$

for all $t \in [0, T]$. Furthermore the solution is not changed at times where it is a proper solution of (1.3)–(1.2), i.e. if $R_0(t, \cdot) \equiv 0$ for some $t \in [0, T]$ then $R_1(t) \equiv 0$ and $(\rho_1, u_1)(t) \equiv (\rho_0, u_0)(t)$.

Proof of Theorem 1.2, assuming Proposition 2.1. We will use the proposition to construct a sequence $(\rho_n, u_n, R_n)_{n \in \mathbb{N}}$ of smooth solutions to (2.1) bounded in the space

$$C\left([0, T], L^p(\mathbb{T}^d) \times \left(L^{p'} \cap W^{1,\tilde{p}}(\mathbb{T}^d, \mathbb{R}^d)\right) \times L^1(\mathbb{T}^d, \mathbb{R}^d)\right)$$

for any $\tilde{p} < q$ (with q as defined in (1.15)), which in the limit will produce a solution of (1.3)–(1.2).

Set $(\rho_0, u_0) := (\bar{\rho}, \bar{u})$ as given in the statement of the theorem and define

$$R_0(t) := -\nabla \Delta^{-1} [\partial_t \bar{\rho}(t) + \operatorname{div}(\bar{\rho}(t)\bar{u}(t))].$$

Recall that $\partial_t \bar{\rho}$ has zero mean value by assumption and $\operatorname{div}(\bar{\rho}\bar{u})$ also, being a divergence, so the definition is correct. Then clearly (ρ_0, u_0, R_0) is a smooth solution of (2.1).

Set $\delta_0 := \|R_0\|_{C_t L_x^1}$ and choose a sequence of positive numbers $\delta_n, n \geq 1$ such that the sum $\sum_n \delta_n^{1/2}$ converges. (Then in particular $\sum_n \delta_n < \infty$.) Furthermore choose sequences $(\tilde{p}_n)_{n \in \mathbb{N}} \subset [1, q)$ and $(\eta_n)_{n \in \mathbb{N}} \subset (1, \infty)$ such that

$$\tilde{p}_n \xrightarrow{n \rightarrow \infty} q \text{ and } \delta_n^{1/p} \eta_n = \sigma \delta_n^{1/2}$$

for some $\sigma > 0$ to be chosen later and observe that $\delta_n^{1/p'}/\eta_n = \delta_n^{1/2}/\sigma$. By repeated application of Proposition 2.1 we obtain a sequence of smooth solutions (ρ_n, u_n, R_n) fulfilling the bounds (uniformly in time)

$$\|\rho_{n+1}(t) - \rho_n(t)\|_{L^p} \leq M\eta_n \|R_n(t)\|_{L^1}^{1/p} \leq M\sigma \delta_n^{1/2} \tag{2.4a}$$

$$\|u_{n+1}(t) - u_n(t)\|_{L^{p'}} \leq \frac{M}{\eta_n} \|R_n(t)\|_{L^1}^{1/p'} \leq \frac{M}{\sigma} \delta_n^{1/2} \tag{2.4b}$$

$$\|u_{n+1}(t) - u_n(t)\|_{W^{1,\tilde{p}_n}} \leq \delta_{n+1} \tag{2.4c}$$

$$\|R_{n+1}(t)\|_{L^1} \leq \delta_{n+1} \tag{2.4d}$$

$$R_n(t) = 0 \implies R_{n+1}(t) = 0. \tag{2.4e}$$

Clearly there are functions $\rho \in C_t L_x^p$ and $u \in C_t L_x^{p'} \cap C_t W_x^{1, \tilde{p}}$ for any $\tilde{p} < q$ such that $\rho_n \rightarrow \rho$ in $C_t L_x^p$ and $u_n \rightarrow u$ in $C_t L_x^{p'}$ and $C_t W_x^{1, \tilde{p}}$. Moreover, we have $\rho_n u_n \rightarrow \rho u$ and $R_n \rightarrow 0$ in $C_t L_x^1$, which proves statements (i) and (ii) of the theorem. For $t \in E$ by (2.4e) we have $R_n(t) = 0$ for all n and therefore, by (2.4a) and (2.4b)

$$\rho_n(t) = \bar{\rho}(t), \quad u_n(t) = \bar{u}(t) \quad \forall n$$

which implies statement (iii). For the last statement we need to choose a sufficiently small (or large) σ so that $M\sigma \sum_{n=0}^\infty \delta_n^{1/2} < \varepsilon$ (or $M\sigma^{-1} \sum_{n=0}^\infty \delta_n^{1/2} < \varepsilon$). So we can ensure that statement (iv) (or statement (iv)', respectively) holds by our choice of σ . If $p = 1$ (and thus $p' = \infty$), then the continuity in space-time of the limit u follows from (2.4b), observing that, in this case, u is the uniform limit of the smooth vector fields u_n . This concludes the proof of the main theorem. \square

We will only prove Proposition 2.1 in the case $p > 1$, the proof will cover Sections 4 to 6. The case $p = 1$, in which the obtained velocity field is in particular continuous (although continuity via Sobolev embeddings just exactly fails to hold), is more delicate to prove. We refer to [18] for the details and will sketch the strategy and the necessary adaptations in Section 7.

3. Technical tools

In this section we provide some technical tools we will use throughout the paper.

3.1. Improved Hölder inequality for fast oscillations

We recall the following lemma from [17]:

Lemma 3.1. *For $p \in [1, \infty]$ there is a constant C_p such that for all smooth functions f, g on the torus \mathbb{T}^d and $\lambda \in \mathbb{N}$:*

$$|\|fg\|_{L^p} - \|f\|_{L^p} \|g\|_{L^p}| \leq \frac{C_p}{\lambda^{1/p}} \|f\|_{C^1} \|g\|_{L^p}.$$

Remark. In particular this lemma supplies the Hölder-like inequality

$$\|fg\|_{L^p} \leq \|f\|_{L^p} \|g\|_{L^p} + \frac{C_p}{\lambda^{1/p}} \|f\|_{C^1} \|g\|_{L^p}, \tag{3.1}$$

which allows to bound the product by the L^p norm of both functions, plus some error term which is small if one function is fastly oscillating, i.e. λ is large.

3.2. Higher derivatives and antiderivatives

As for smooth f , with $\int_{\mathbb{T}^d} f = 0$, the Poisson equation $\Delta u = f$ has a solution on the flat torus which is unique up to addition of a constant, the inverse Laplacian

$$\Delta^{-1} : C_0^\infty \rightarrow C_0^\infty, \quad f \mapsto u$$

is well-defined as an operator on the space C_0^∞ . We can now use it to define higher order (anti)derivatives with a simple structure.

Definition. For any smooth function $f \in C^\infty(\mathbb{T}^d)$ on the torus and non-negative integers k we define the differential operator \mathcal{D}^k :

$$\mathcal{D}^k f = \begin{cases} \Delta^{k/2} f, & \text{if } k \text{ even,} \\ \nabla \Delta^{\frac{k-1}{2}} f, & \text{if } k \text{ odd,} \end{cases}$$

with the convention that $\mathcal{D}^0 = \Delta^0 = Id$.

For negative k the definition is identical with the additional condition $f \in C_0^\infty(\mathbb{T}^d)$, which is necessary so that negative powers of the Laplacian are meaningful.

Remark. The basic properties of the operators \mathcal{D}^k include

- It commutes with derivatives: $\partial^\alpha \mathcal{D}^k f = \mathcal{D}^k \partial^\alpha f$ for all $k \in \mathbb{Z}$ and any multi-index α .
- Partial Integration: For any $k, n, m \in \mathbb{Z}$ and $f, g \in C_0^\infty(\mathbb{T}^d)$

$$\int_{\mathbb{T}^d} \mathcal{D}^k f \cdot \mathcal{D}^{m+n} g = (-1)^n \int_{\mathbb{T}^d} \mathcal{D}^{k+n} f \cdot \mathcal{D}^m g,$$

where the ‘ \cdot ’ denotes scalar product if both factors are vectors, otherwise standard multiplication.

- Scaling: $\mathcal{D}^k u_\lambda = \lambda^k (\mathcal{D}^k u)_\lambda$ for any $k \in \mathbb{Z}$ and $\lambda \in \mathbb{N}$.

3.3. Calderon-Zygmund estimates

We first recall the usual Calderon-Zygmund inequality in the following form.

Remark (Classical Calderon-Zygmund inequality). Let $p \in (1, \infty)$. There is a constant $C_{d,p}$ such that for any smooth compactly supported function f the following inequality holds:

$$\|f\|_{W^{2,p}(\mathbb{R}^d)} \leq C_{d,p} \|\Delta f\|_{L^p(\mathbb{R}^d)}. \tag{3.2}$$

We refer to [13] for the proof.

It is now a small step to show that the same statement can be transferred to the periodic setting: we include the proof for completeness.

Lemma 3.2 (Calderon-Zygmund on the flat torus). Let $p \in (1, \infty)$. There is a constant $C_{d,p}$ such that for any $f \in C_0^\infty(\mathbb{T}^d)$ the following inequality holds:

$$\|f\|_{W^{2,p}(\mathbb{T}^d)} \leq C_{d,p} \|\Delta f\|_{L^p(\mathbb{T}^d)}. \tag{3.3}$$

Proof. Let $f \in C_0^\infty(\mathbb{T}^d)$ and $N \in \mathbb{N}$. We treat f as a periodic map $f : \mathbb{R}^d \rightarrow \mathbb{R}$ and identify \mathbb{T}^d with the unit cube $(0, 1)^d$. Choose a smooth cut-off function $\chi \in C^\infty(\mathbb{R})$ such that $\chi(x) = 1$ if $x \leq 0$ and $\chi(x) = 0$ if $x \geq 1$. Define the function $f_N \in C_c^\infty(\mathbb{R}^d)$ by

$$f_N(x) := \left(\prod_{i=1}^d \chi(|x_i| - N) \right) f(x).$$

Now the classical Calderon-Zygmund inequality (3.2) and the fact that f_N is supported in the cube $[-N - 1, N + 1]^d$ yield

$$\|f_N\|_{W^{2,p}([-N,N]^d)} \leq \|f_N\|_{W^{2,p}(\mathbb{R}^d)} \leq C_{d,p} \|\Delta f_N\|_{L^p(\mathbb{R}^d)} = C_{d,p} \|\Delta f_N\|_{L^p([-N-1,N+1]^d)}$$

and therefore, using that $\|\chi\|_{C^0} = 1$ and $f_N = f$ on $[-N, N]^d$.

$$(2N)^d \|f\|_{W^{2,p}(\mathbb{T}^d)} \leq C_{d,p} \left[(2N + 2)^d \|\Delta f\|_{L^p(\mathbb{T}^d)} + \left((2N + 2)^d - (2N)^d \right) (\|\chi'\|_{C^0} \|\nabla f\|_{L^p(\mathbb{T}^d)} + \|\chi''\|_{C^0} \|f\|_{L^p(\mathbb{T}^d)}) \right].$$

If $N \rightarrow \infty$ the dominating terms are the ones with the factor $(2N)^d$, and so

$$\|f\|_{W^{2,p}(\mathbb{T}^d)} \leq C_{d,p} \|\Delta f\|_{L^p(\mathbb{T}^d)}$$

holds with the same constant as in the full space setting. \square

Lemma 3.3 (Estimates on antiderivatives). *Let $p \in (1, \infty)$ and $k \in \mathbb{N}$. There is a constant $C_{d,p,k}$ such that*

$$\|\mathcal{D}^{-k} f\|_{W^{k,p}(\mathbb{T}^d)} \leq C_{d,p,k} \|f\|_{L^p(\mathbb{T}^d)} \tag{3.4}$$

holds for any $f \in C_0^\infty(\mathbb{T}^d)$.

Proof. If k is even, the inequality arises simply from iterated application of the Calderon-Zygmund inequality on the torus:

$$\|\mathcal{D}^{-k} f\|_{W^{k,p}} = \|\Delta^{-k/2} f\|_{W^{k,p}} \leq C_{d,p} \|\Delta^{-k/2+1} f\|_{W^{k-2,p}} \leq \dots \leq C_{d,p}^{k/2} \|f\|_{L^p}.$$

For odd numbers k observe that the same iteration leaves us with

$$\|\mathcal{D}^{-k} f\|_{W^{k,p}} \leq C_{d,p}^{(k-1)/2} \|\mathcal{D}^{-1} f\|_{W^{1,p}} = C_{d,p}^{(k-1)/2} \|\nabla \Delta^{-1} f\|_{W^{1,p}}$$

and clearly

$$\|\nabla \Delta^{-1} f\|_{W^{1,p}} \leq \|\Delta^{-1} f\|_{W^{2,p}} \leq C_{d,p} \|f\|_{L^p}$$

so the stated inequality holds with $C_{d,p,k} = C_{p,d}^{\lceil k/2 \rceil}$. \square

Lemma 3.4 (End point estimates on antiderivatives). *Let $p \in [1, \infty]$ and $k \in \mathbb{N}_+$. There is a constant $C_{d,p,k}$ such that*

$$\|\mathcal{D}^{-k} f\|_{W^{k-1,p}(\mathbb{T}^d)} \leq C_{d,p,k} \|f\|_{L^p(\mathbb{T}^d)} \tag{3.5}$$

holds for any $f \in C_0^\infty(\mathbb{T}^d)$.

Proof. In the case $p \in (1, \infty)$ there is nothing to show as the statement is just a weaker form of (3.4).

For $p = \infty$ we use Sobolev embeddings on every derivative of order $k - 1$ and smaller to control the Sobolev norm of a smooth function g : for every multiindex α , with $|\alpha| \leq k - 1$,

$$\|\partial^\alpha g\|_{L^\infty} \leq C_d \|D\partial^\alpha g\|_{L^{d+1}} \implies \|g\|_{W^{k-1,\infty}} \leq C_d \|g\|_{W^{k,d+1}}.$$

If we set $g = \mathcal{D}^{-k} f$ and we use the previous Lemma, we obtain

$$\|\mathcal{D}^{-k} f\|_{W^{k-1,\infty}} \leq C_d \|\mathcal{D}^{-k} f\|_{W^{k,d+1}} \leq C_{d,p,k} \|f\|_{L^{d+1}} \leq C_{d,p,k} \|f\|_{L^\infty}.$$

For $p = 1$ we consider the dual characterization of the L^1 -norm:

$$\begin{aligned} \|g\|_{L^1} &= \max \left\{ \frac{1}{\|\phi\|_{L^\infty}} \int_{\mathbb{T}^d} g\phi : \phi \in L^\infty(\mathbb{T}^d) \setminus \{0\} \right\} \\ &= \sup \left\{ \frac{1}{\|\phi\|_{L^\infty}} \int_{\mathbb{T}^d} g\phi : \phi \in C^\infty(\mathbb{T}^d) \setminus \{0\} \right\}. \end{aligned}$$

If $\int_{\mathbb{T}^d} g = 0$ we can restrict the definition to test functions in $C_0^\infty(\mathbb{T}^d)$, still obtaining the inequalities

$$\frac{1}{2} \|g\|_{L^1} \leq \sup \left\{ \frac{1}{\|\phi\|_{L^\infty}} \int_{\mathbb{T}^d} g\phi : \phi \in C_0^\infty(\mathbb{T}^d) \setminus \{0\} \right\} \leq \|g\|_{L^1} \tag{3.6}$$

where the first inequality comes from the fact that $\int g(\phi - f\phi) = \int g\phi$ and $\|\phi - f\phi\|_{L^\infty} \leq 2\|\phi\|_{L^\infty}$ hold for any ϕ . Using this, we can estimate for any multiindex α of order $k - 1$ or smaller

$$\begin{aligned}
 \|\partial^\alpha \mathcal{D}^{-k} f\|_{L^1} &\leq \sup_{\phi \in C_0^\infty(\mathbb{T}^d)} \frac{2}{\|\phi\|_{L^\infty}} \int_{\mathbb{T}^d} (\partial^\alpha \mathcal{D}^{-k} f) \phi \\
 &= \sup_{\phi \in C_0^\infty(\mathbb{T}^d)} \frac{2}{\|\phi\|_{L^\infty}} \int_{\mathbb{T}^d} f (\partial^\alpha \mathcal{D}^{-k} \phi) \\
 &\leq \sup_{\phi \in C_0^\infty(\mathbb{T}^d)} \frac{2}{\|\phi\|_{L^\infty}} \|f\|_{L^1} \|\partial^\alpha \mathcal{D}^{-k} \phi\|_{L^\infty} \\
 &\leq \|f\|_{L^1} \sup_{\phi \in C_0^\infty(\mathbb{T}^d)} \frac{C_{d,\infty,k}}{\|\phi\|_{L^\infty}} \|\phi\|_{L^\infty} \\
 &= C_{d,\infty,k} \|f\|_{L^1}
 \end{aligned}$$

where in the last inequality (3.5) with $p = \infty$ was applied. Summation over all such α then yields (3.5):

$$\|\mathcal{D}^{-k} f\|_{W^{k-1,1}} = \sum_{|\alpha| \leq k-1} \|\partial^\alpha \mathcal{D}^{-k} f\|_{L^1} \leq \sum_{|\alpha| \leq k-1} C_{d,\infty,k} \|f\|_{L^1} = C_{d,1,k} \|f\|_{L^1}. \quad \square$$

3.4. Improved antidivergence for fast oscillations

The first order antiderivative \mathcal{D}^{-1} is an anti-divergence operator, which we will call standard anti-divergence operator. It will be used in situations when the estimate provided in Lemma 3.4 with $k = 1$ suffices. However, in many steps of the proof of Proposition 2.1 refined estimates on the antidivergence are necessary. We therefore introduce a bilinear operator which is apt to control the antidivergence of a product of functions if one of them is fast oscillating.

Definition (Bilinear anti-divergence operator). Let $N \in \mathbb{N}$. Define the operator

$$\begin{aligned}
 \mathcal{R}_N : C^\infty(\mathbb{T}^d) \times C_0^\infty(\mathbb{T}^d) &\rightarrow C^\infty(\mathbb{T}^d; \mathbb{R}^d) \\
 \mathcal{R}_N(f, g) &:= \sum_{k=0}^{N-1} (-1)^k \mathcal{D}^k f \mathcal{D}^{-k-1} g + \mathcal{D}^{-1} \left((-1)^N \mathcal{D}^N f \cdot \mathcal{D}^{-N} g - \int_{\mathbb{T}^d} fg \right). \tag{3.7}
 \end{aligned}$$

Here the ‘ \cdot ’ indicates the scalar product if needed, i.e. if N is odd, and the standard product otherwise. Note that both arguments must be smooth but only the second argument g is supposed to have zero mean value.

Lemma 3.5 (Properties of \mathcal{R}_N). Let $N \in \mathbb{N}$, $f \in C^\infty(\mathbb{T}^d)$ and $g \in C_0^\infty(\mathbb{T}^d)$.

(i) \mathcal{R}_N is an antidivergence operator in the sense that

$$\operatorname{div}(\mathcal{R}_N(f, g)) = fg - \int_{\mathbb{T}^d} fg.$$

(ii) \mathcal{R}_N satisfies the Leibniz rule:

$$\partial_j(\mathcal{R}_N(f, g)) = \mathcal{R}_N(\partial_j f, g) + \mathcal{R}_N(f, \partial_j g).$$

(iii) If $p, r, s \in [1, \infty]$ such that $\frac{1}{p} = \frac{1}{r} + \frac{1}{s}$, then the following inequality holds:

$$\|\mathcal{R}_N(f, g)\|_{L^p} \leq \sum_{k=0}^{N-1} \|\mathcal{D}^k f\|_{L^r} \|\mathcal{D}^{-k-1} g\|_{L^s} + C_{d,p} \|\mathcal{D}^N f\|_{L^r} \|\mathcal{D}^{-N} g\|_{L^s}. \tag{3.8}$$

Proof. (i) By induction in N . By definition we have

$$\mathcal{R}_0(f, g) = \mathcal{D}^{-1} \left(fg - \int_{\mathbb{T}^d} fg \right)$$

so the statement follows from the remark on standard anti-divergence. Now let $N > 0$ and w.l.o.g. assume N to be even, then

$$\begin{aligned} \operatorname{div}(\mathcal{R}_N(f, g)) - \left(fg - \int_{\mathbb{T}^d} fg \right) &= \overbrace{- \left(fg - \int_{\mathbb{T}^d} fg \right)}^{=0 \text{ by assumption}} + \operatorname{div}(\mathcal{R}_{N-1}(f, g)) \\ &= -(-1)^{N-1} \mathcal{D}^{N-1} f \cdot \mathcal{D}^{-N+1} g + (-1)^N \mathcal{D}^N f \mathcal{D}^{-N} g \\ &\quad + \operatorname{div} \left((-1)^{N-1} \mathcal{D}^{N-1} f \mathcal{D}^{-N} g \right) \\ &= \mathcal{D}^{N-1} f \cdot \mathcal{D}^{-N+1} g + \mathcal{D}^N f \mathcal{D}^{-N} g \\ &\quad - \operatorname{div} \left(\mathcal{D}^{N-1} f \right) \mathcal{D}^{-N} g - \mathcal{D}^{N-1} f \cdot \nabla \mathcal{D}^{-N} g \\ &= 0 \end{aligned}$$

by definition of the operators \mathcal{D}^k .

(ii) is proven by lengthy but straightforward computation which we omit here.

(iii) Use the standard Hölder inequality on each term of the definition of \mathcal{R}_N . For the last summand note that Lemma 3.4 in particular implies $\|\mathcal{D}^{-1}h\|_{L^p} \leq C(d, p)\|h\|_{L^p}$; furthermore $\|h - \int_{\mathbb{T}^d} h\|_{L^p} \leq 2\|h\|_{L^p}$ for any p . \square

Remark. The bilinear anti-divergence and inequality (3.8) are only useful if applied on functions g_λ which are fast oscillating, as then we gain the oscillation parameter λ as small factor. In particular the following two estimates will be used throughout the paper. Let $p \in [1, \infty]$, $\lambda, N \in \mathbb{N}$, $f \in C^\infty(\mathbb{T}^d)$ and $g \in C_0^\infty(\mathbb{T}^d)$. Then:

$$\|\mathcal{R}_N(f, g_\lambda)\|_{L^p} \leq C_{d,p,N} \|g\|_{L^p} \left(\sum_{k=0}^{N-1} \lambda^{-k-1} \|\mathcal{D}^k f\|_{L^\infty} + \lambda^{-N} \|\mathcal{D}^N f\|_{L^\infty} \right), \tag{3.9}$$

$$\|\mathcal{R}_N(f, g_\lambda)\|_{L^p} \leq C_{d,p,N} \|g\|_{L^\infty} \left(\sum_{k=0}^{N-1} \lambda^{-k-1} \|\mathcal{D}^k f\|_{L^p} + \lambda^{-N} \|\mathcal{D}^N f\|_{L^p} \right). \tag{3.10}$$

The proof of (3.9)-(3.10) is direct consequence of (3.8) and Lemma 3.4.

4. The perturbations

In this section we introduce the basic building blocks of our construction, namely the *space-time Mikado densities and field*, which allow us to get a “full dimensional concentration”, i.e. to assume (1.4) instead of (1.12). We then use the Mikado functions to define and estimate ρ_1, u_1 .

4.1. Space-time Mikado densities and fields

For given $\zeta, v \in \mathbb{T}^d$, consider the line on \mathbb{T}^d

$$\mathbb{R} \ni s \mapsto \zeta + sv \in \mathbb{T}^d.$$

Lemma 4.1 (*Space-time Mikado lines*). *There exist $r > 0$ and $\zeta_1, \dots, \zeta_d \in \mathbb{T}^d$ such that the lines*

$$\mathbf{x}_j : \mathbb{R} \rightarrow \mathbb{T}^d, \mathbf{x}_j(s) = \zeta_j + se_j$$

satisfy

$$d_{\mathbb{T}^d}(\mathbf{x}_i(s), \mathbf{x}_j(s)) > 2r \quad \forall s \in \mathbb{R}, \forall i \neq j, \tag{4.1}$$

where $d_{\mathbb{T}^d}$ denotes the Euclidian distance on the torus.

Remark. We can think to the lines x_j as the trajectories of d particles moving on the torus with speed 1 and along different directions. The claim of the Lemma is that such particles have different positions at every time.

Proof. We define

$$\zeta_i := \frac{i}{d}e_i, \quad i = 1, \dots, d,$$

where e_j denotes the j -th standard unit vector in \mathbb{R}^d .

Let $i \neq j$ be fixed. If, for some $s \in \mathbb{R}$,

$$x_i(s) = x_j(s) \text{ in } \mathbb{T}^d,$$

then

$$(\zeta_j + se_j) - (\zeta_i + se_i) \in \mathbb{Z}^d$$

and thus

$$\frac{i}{d} + s \in \mathbb{Z}, \quad \frac{j}{d} + s \in \mathbb{Z},$$

which implies, taking the difference,

$$\frac{i-j}{d} \in \mathbb{Z},$$

a contradiction. Therefore, for every $s \in \mathbb{R}$ and $i \neq j$, $x_i(s) \neq x_j(s)$ and thus there must be $r > 0$ such that (4.1) holds. \square

Let φ be a smooth function on \mathbb{R}^d , with

$$\text{supp } \varphi \subseteq B(P, r) \subseteq (0, 1)^d,$$

where $P = (1/2, \dots, 1/2) \in (0, 1)^d$ and $B(P, r)$ is the ball with radius r centered at P , and assume that

$$\int_{\mathbb{R}^d} \varphi^2 = 1.$$

For a given p (fixed in the statement of Proposition 2.1), and its dual Hölder exponent p' define the constants

$$a := \frac{d}{p}, \quad b := \frac{d}{p'} \quad \text{so that } a + b = d \tag{4.2}$$

and the scaled functions (defined on the whole space \mathbb{R}^d , thus not periodic)

$$\varphi_\mu(x) := \mu^a \varphi(\mu x), \quad \tilde{\varphi}_\mu(x) := \mu^b \varphi(\mu x), \quad \mu \geq 1.$$

Lemma 4.2. For every $\mu \geq 1$, $k \in \mathbb{N}$, $r \in [1, \infty]$,

$$\|D^k \varphi_\mu\|_{L^r(\mathbb{R}^d)} = \mu^{a - \frac{d}{r} + k} \|D^k \varphi\|_{L^r(\mathbb{R}^d)}, \quad \|D^k \tilde{\varphi}_\mu\|_{L^r(\mathbb{R}^d)} = \mu^{b - \frac{d}{r} + k} \|D^k \varphi\|_{L^r(\mathbb{R}^d)}. \tag{4.3}$$

Moreover,

$$\int_{\mathbb{R}^d} \varphi_\mu \tilde{\varphi}_\mu = 1. \tag{4.4}$$

The proof is straightforward and thus it is omitted. Note in particular that the $L^p(\mathbb{R}^d)$ -norm of φ_μ and the $L^{p'}(\mathbb{R}^d)$ -norm of $\tilde{\varphi}_\mu$ are invariant of the scaling. Note also that $\text{supp } \varphi_\mu = \text{supp } \tilde{\varphi}_\mu$ and both are contained in a ball with radius at most r . For any given $y \in \mathbb{T}^d$, we define the translation

$$\tau_y : \mathbb{T}^d \rightarrow \mathbb{T}^d, \quad \tau_y(x) := x - y.$$

Notice that, for every smooth periodic map g

$$\|D^k(g \circ \tau_y)\|_{L^r} = \|D^k g\|_{L^r} \quad \forall k \in \mathbb{N}, \quad \forall r \in [1, \infty].$$

Lemma 4.3. *There are periodic functions*

$$\varphi_\mu^j : \mathbb{T}^d \rightarrow \mathbb{R}, \quad \tilde{\varphi}_\mu^j : \mathbb{T}^d \rightarrow \mathbb{R}, \quad j = 1, \dots, d,$$

such that the same scaling as in (4.3) holds:

$$\|D^k \varphi_\mu^j\|_{L^r} = \mu^{a-\frac{d}{r}+k} \|D^k \varphi\|_{L^r}, \quad \|D^k \tilde{\varphi}_\mu^j\|_{L^r} = \mu^{b-\frac{d}{r}+k} \|D^k \varphi\|_{L^r}. \tag{4.5}$$

Moreover, for every $i = 1, \dots, d$,

$$\int_{\mathbb{T}^d} (\varphi_\mu^i \circ \tau_{se_i})(\tilde{\varphi}_\mu^i \circ \tau_{se_i}) = 1, \tag{4.6}$$

and, for every $i \neq j$ and $s \in \mathbb{R}$,

$$(\varphi_\mu^i \circ \tau_{se_i})(\tilde{\varphi}_\mu^j \circ \tau_{se_j}) = 0. \tag{4.7}$$

Notice that (4.7) means

$$\varphi_\mu^i(x - se_i)\tilde{\varphi}_\mu^j(x - se_j) = 0$$

for every $x \in \mathbb{T}^d$.

Proof. Since $\varphi_\mu, \tilde{\varphi}_\mu$ have support contained in $(0, 1)^d$, we can consider their periodic extensions, still denoted, with a slight abuse of notation, by $\varphi_\mu, \tilde{\varphi}_\mu$, respectively. We define now the periodic maps

$$\varphi_\mu^j := \varphi_\mu \circ \tau_{\zeta_j}, \quad \tilde{\varphi}_\mu^j := \tilde{\varphi}_\mu \circ \tau_{\zeta_j},$$

where ζ_1, \dots, ζ_d are the points given by Lemma 4.1. It is immediate from the definition and from (4.3)-(4.4) that (4.5)-(4.6) holds. Let now $x \in \mathbb{T}^d, s \in \mathbb{R}$. We have

$$\varphi_\mu^i(x - se_i)\tilde{\varphi}_\mu^j(x - se_j) = \varphi_\mu(x - \zeta_i - se_i)\tilde{\varphi}_\mu(x - \zeta_j - se_i) = \varphi_\mu(x - \mathbf{x}_i(s))\tilde{\varphi}_\mu(x - \mathbf{x}_j(s)).$$

Observe that, by Lemma 4.1,

$$d_{\mathbb{T}^d}(x - \mathbf{x}_i(s), x - \mathbf{x}_j(s)) = d_{\mathbb{T}^d}(\mathbf{x}_i(s), \mathbf{x}_j(s)) > 2r.$$

Since the support of φ_μ and $\tilde{\varphi}_\mu$ coincide and are both contained in a ball with radius at most r , it must be

$$\varphi_\mu(x - \mathbf{x}_i(s))\tilde{\varphi}_\mu(x - \mathbf{x}_j(s)) = 0,$$

and thus (4.7) holds. \square

We introduce now the building block of our construction, the space-time Mikado densities and fields. Besides the families of functions $\varphi_\mu^j, \tilde{\varphi}_\mu^j, \mu \geq 1, j = 1, \dots, d$, we fix a smooth periodic function $\psi : \mathbb{T}^{d-1} \rightarrow \mathbb{R}$ satisfying

$$\int_{\mathbb{T}^{d-1}} \psi = 0, \quad \int_{\mathbb{T}^{d-1}} \psi^2 = 1$$

and we define

$$\psi^j : \mathbb{T}^d \rightarrow \mathbb{R}, \quad \psi^j(x) = \psi^j(x_1, \dots, x_d) := \psi(x_1, \dots, x_{j-1}, x_{j+1}, \dots, x_d),$$

for every $j = 1, \dots, d$, so that

$$\int_{\mathbb{T}^d} \psi^j = 0, \quad \int_{\mathbb{T}^d} (\psi^j)^2 = 1. \tag{4.8}$$

Introduce the parameters

- λ ‘fast oscillation’, $\in \mathbb{N}$
- μ ‘concentration’, $\gg \lambda$
- ω ‘phase speed’
- ν ‘very fast oscillation’, $\in \lambda\mathbb{N}, \gg \lambda$

to be chosen in the very end of the proof. Now we can define the Mikado functions, for $j = 1, \dots, d$:

- Mikado density $\Theta_{\lambda, \mu, \omega, \nu}^j(t, x) := \varphi_\mu^j(\lambda(x - \omega t e_j)) \psi^j(\nu x)$,
- Mikado field $W_{\lambda, \mu, \omega, \nu}^j(t, x) := \tilde{\varphi}_\mu^j(\lambda(x - \omega t e_j)) \psi^j(\nu x) e_j$,
- Quadratic corrector $Q_{\lambda, \mu, \omega, \nu}^j(t, x) := \frac{1}{\omega} (\varphi_\mu^j \tilde{\varphi}_\mu^j)(\lambda(x - \omega t e_j)) (\psi^j(\nu x))^2$.

We will use also the shorter notation

$$\begin{aligned} \Theta_{\lambda, \mu, \omega, \nu}^j &= \Theta_{\lambda, \mu, \omega, \nu}^j(t) := ((\varphi_\mu^j)_\lambda \circ \tau_{\omega t e_j}) \psi_\nu^j, \\ W_{\lambda, \mu, \omega, \nu}^j &= W_{\lambda, \mu, \omega, \nu}^j(t) := ((\tilde{\varphi}_\mu^j)_\lambda \circ \tau_{\omega t e_j}) \psi_\nu^j e_j, \\ Q_{\lambda, \mu, \omega, \nu}^j &= Q_{\lambda, \mu, \omega, \nu}^j(t) := \frac{1}{\omega} ((\varphi_\mu^j \tilde{\varphi}_\mu^j)_\lambda \circ \tau_{\omega t e_j}) (\psi_\nu^j)^2, \end{aligned}$$

where we have used the notation $g_\lambda(x) := g(\lambda x)$ (and $g_\nu(x) := g(\nu x)$), for $g : \mathbb{T}^d \rightarrow \mathbb{R}$.

Remark. The Mikados defined here do not form a stationary solution of the incompressible transport equation, in contrast to those used in [17,18]. The ideal cancellation properties $\partial_t \Theta_{\lambda, \mu, \omega, \nu}^j = \operatorname{div}(\Theta_{\lambda, \mu, \omega, \nu}^j W_{\lambda, \mu, \omega, \nu}^j) = 0 = \operatorname{div} W_{\lambda, \mu, \omega, \nu}^j$ for every j (and in particular if summed over all j) cannot hold here because of the time-dependence and compact support in space of the function $\varphi(\lambda(x - \omega t e_j))$. However, ψ is still time-independent and divergence-free so that

$$\partial_t \Theta_{\lambda, \mu, \omega, \nu}^j = -\lambda \omega \left((\partial_j \varphi_\mu^j)_\lambda \circ \tau_{\omega t e_j} \right) \psi_\nu^j, \tag{4.9}$$

$$\operatorname{div} W_{\lambda, \mu, \omega, \nu}^j = \lambda \left((\partial_j \tilde{\varphi}_\mu^j)_\lambda \circ \tau_{\omega t e_j} \right) \psi_\nu^j e_j, \tag{4.10}$$

holds and, because of the fact that $Q^j = \frac{1}{\omega} \Theta^j W^j$, we still have a set of functions similar to a solution to the transport equation, as stated in the following proposition.

Set

$$\epsilon := \frac{d}{p} + \frac{d}{\tilde{p}} - d - 1 = \frac{d}{\tilde{p}} - \frac{d}{p'} - 1 > 0. \tag{4.11}$$

Note that $\epsilon > 0$, because of (2.2).

Proposition 4.4. Define the global constants M (not depending on p, \tilde{p}) by

$$M := 2d \max_{k, k'=0,1} \left\{ \|D^k \varphi\|_{L^\infty} \|D^{k'} \psi\|_{L^\infty}, \|\varphi\|_{L^\infty}^2 \|\psi\|_{L^\infty}^2 \right\}. \tag{4.12}$$

The Mikado functions obey the following bounds:

$$\left\| \Theta_{\lambda, \mu, \omega, v}^j(t) \right\|_{L^p} \leq \frac{M}{2d}, \quad \left\| W_{\lambda, \mu, \omega, v}^j(t) \right\|_{L^{p'}} \leq \frac{M}{2d}, \quad \left\| Q_{\lambda, \mu, \omega, v}^j(t) \right\|_{L^p} \leq \frac{M\mu^b}{\omega}, \tag{4.13a}$$

$$\left\| \Theta_{\lambda, \mu, \omega, v}^j(t) \right\|_{L^1} \leq \frac{M}{\mu^b}, \quad \left\| W_{\lambda, \mu, \omega, v}^j(t) \right\|_{L^1} \leq \frac{M}{\mu^a}, \quad \left\| Q_{\lambda, \mu, \omega, v}^j(t) \right\|_{L^1} \leq \frac{M}{\omega}, \tag{4.13b}$$

$$\left\| W_{\lambda, \mu, \omega, v}^j(t) \right\|_{C^0} \leq M\mu^b, \tag{4.13c}$$

$$\left\| W_{\lambda, \mu, \omega, v}^j(t) \right\|_{W^{1, \tilde{p}}} \leq M \frac{\lambda\mu + v}{\mu^{1+\epsilon}}. \tag{4.13d}$$

Furthermore, for every $i \neq j$,

$$\Theta_{\lambda, \mu, \omega, v}^i W_{\lambda, \mu, \omega, v}^j = 0 \tag{4.14}$$

and the Mikado functions ‘solve the continuity equation’ in the sense that

$$\partial_t Q_{\lambda, \mu, \omega, v}^j + \operatorname{div} \left(\Theta_{\lambda, \mu, \omega, v}^j W_{\lambda, \mu, \omega, v}^j \right) = 0 \tag{4.15}$$

on $[0, T] \times \mathbb{T}^d$.

Proof. The inequalities in (4.13a)-(4.13b)-(4.13c) are immediate consequence of (4.5). We show only the first inequality in (4.13a), the other ones being completely similar:

$$\begin{aligned} \left\| \Theta_{\lambda, \mu, \omega, v}^j(t) \right\|_{L^p} &\leq \left\| (\varphi_\mu^j)_\lambda \circ \tau_{\omega t e_j} \right\|_{L^p} \left\| \psi_v^j \right\|_{L^\infty} \\ &= \left\| \varphi_\mu^j \right\|_{L^p} \left\| \psi^j \right\|_{L^\infty} \\ &= \left\| \varphi \right\|_{L^p} \left\| \psi \right\|_{L^\infty} \\ &\leq \left\| \varphi \right\|_{L^\infty} \left\| \psi \right\|_{L^\infty} \\ &\leq \frac{M}{2d}. \end{aligned}$$

Inequality (4.13d) requires direct calculation: using (1.20), we get

$$\begin{aligned} \left\| W_{\lambda, \mu, \omega, v}^j(t) \right\|_{W^{1, \tilde{p}}} &\leq \left\| (\tilde{\varphi}_\mu^j)_\lambda \circ \tau_{\omega t e_j} \right\|_{L^{\tilde{p}}} \left\| \psi_v^j \right\|_{L^\infty} + \left\| D \left((\tilde{\varphi}_\mu^j)_\lambda \circ \tau_{\omega t e_j} \right) \right\|_{L^{\tilde{p}}} \left\| \psi_v^j \right\|_{L^\infty} \\ &\quad + \left\| (\tilde{\varphi}_\mu^j)_\lambda \circ \tau_{\omega t e_j} \right\|_{L^{\tilde{p}}} \left\| D(\psi_v^j) \right\|_{L^\infty} \\ &\leq \left\| \tilde{\varphi}_\mu^j \right\|_{L^{\tilde{p}}} \left\| \psi^j \right\|_{L^\infty} + \lambda \left\| D\tilde{\varphi}_\mu^j \right\|_{L^{\tilde{p}}} \left\| \psi^j \right\|_{L^\infty} + v \left\| \tilde{\varphi}_\mu^j \right\|_{L^{\tilde{p}}} \left\| D\psi^j \right\|_{L^\infty} \\ \text{(by (4.5))} &\leq \mu^{d/p'-d/\tilde{p}} \left\| \varphi \right\|_{L^{\tilde{p}}} \left\| \psi \right\|_{L^\infty} + \lambda \mu^{d/p'-d/\tilde{p}+1} \left\| D\varphi \right\|_{L^{\tilde{p}}} \left\| \psi \right\|_{L^\infty} \\ &\quad + v \mu^{d/p'-d/\tilde{p}} \left\| \varphi \right\|_{L^{\tilde{p}}} \left\| D\psi \right\|_{L^\infty} \\ &\leq M \left(\frac{\lambda}{\mu^\epsilon} + \frac{v}{\mu^{1+\epsilon}} \right). \end{aligned}$$

Equality (4.14) is an immediate consequence of (4.7). To prove (4.15), we observe that

$$\Theta_{\lambda, \mu, \omega, v}^j(t, x) W_{\lambda, \mu, \omega, v}^j(t, x) = \omega Q_{\lambda, \mu, \omega, v}^j(t, x) e_j = F(x - \omega t e_j) \psi_v^j(x) e_j,$$

for some $F : \mathbb{T}^d \rightarrow \mathbb{R}$, whose precise form is not important. Since $\psi_v^j e_j$ is time independent and divergence free, we get

$$\begin{aligned} \operatorname{div} \left(\Theta_{\lambda, \mu, \omega, v}^j W_{\lambda, \mu, \omega, v}^j \right) &= \nabla F \cdot \psi_v^j e_j, \\ \partial_t Q_{\lambda, \mu, \omega, v}^j &= -\nabla F \cdot \psi_v^j e_j, \end{aligned}$$

and thus (4.15) holds. \square

4.2. Definition of perturbations

Given (ρ_0, u_0, R_0) as in Proposition 2.1, we denote by $R_0^j(t, x)$ the components of the vector $R_0(t, x)$, i.e.

$$R_0(t, x) = \sum_{j=1}^d R_0^j(t, x) e_j.$$

We now define the new density and velocity field as

$$\begin{aligned} \rho_1(t, x) &:= \rho_0(t, x) + \vartheta(t, x) + \vartheta_c(t) + q(t, x) + q_c(t) \\ u_1(t, x) &:= u_0(t, x) + w(t, x) + w_c(t, x) \end{aligned}$$

where ϑ, q and w are the Mikado density, quadratic corrector term and Mikado flow weighted by the defect field R_0 , defined as follows:

$$\begin{aligned} \vartheta(t, x) &:= \eta \sum_{j=1}^d \chi_j(t, x) \operatorname{sgn} \left(R_0^j(t, x) \right) \left| R_0^j(t, x) \right|^{1/p} \Theta_{\lambda, \mu, \omega, \nu}^j(t, x), \\ w(t, x) &:= \frac{1}{\eta} \sum_{j=1}^d \chi_j(t, x) \left| R_0^j(t, x) \right|^{1/p'} W_{\lambda, \mu, \omega, \nu}^j(t, x), \\ q(t, x) &:= \sum_{j=1}^d \chi_j^2(t, x) R_0^j(t, x) Q_{\lambda, \mu, \omega, \nu}^j(t, x). \end{aligned}$$

Here $\lambda, \mu, \omega, \nu$ will be chosen in Section 6 to conclude the proof of Proposition 2.1, the $\chi_j : [0, T] \times \mathbb{T}^d \rightarrow [0, 1]$ are cut-off functions which ensure the smoothness of the perturbations at the zero set of R_0^j :

$$\chi_j(t, x) = \begin{cases} 0 & \text{if } |R_0^j(t, x)| \leq \frac{\delta}{4d}, \\ 1 & \text{if } |R_0^j(t, x)| \geq \frac{\delta}{2d}, \end{cases}$$

and η and δ are the strictly positive numbers which appear in the statement of Proposition 2.1.

The parameters $\lambda, \mu, \omega, \nu \gg 1$ will be fixed in Section 6. We will however use the shorter notation

$$\vartheta(t) := \sum_j a_j(t) \Theta^j(t), w(t) := \sum_j b_j(t) W^j(t), q(t) := \sum_j a_j(t) b_j(t) Q^j(t),$$

where a_j, b_j are defined as

$$a_j(t) := \eta \chi_j(t) \operatorname{sgn} \left(R_0^j(t) \right) \left| R_0^j(t) \right|^{1/p}, b_j(t) := \frac{1}{\eta} \chi_j(t) \left| R_0^j(t) \right|^{1/p'}.$$

Notice that

$$a_j(t) b_j(t) = \chi_j^2(t) R_0^j(t),$$

and the following estimates hold true:

$$\|a_j(t)\|_{L^p} \leq \eta \|R_0(t)\|_{L^1}^{1/p}, \quad \|b_j(t)\|_{L^{p'}} \leq \eta^{-1} \|R_0(t)\|_{L^1}^{1/p'} \tag{4.16a}$$

and, for every $k \in \mathbb{N}$,

$$\|a_j\|_{C^k}, \|b_j\|_{C^k} \leq C(\eta, \delta, \|R_0\|_{C^k}). \tag{4.16b}$$

The corrector terms ϑ_c, q_c are needed for ρ_1 to have zero mean value:

$$\vartheta_c(t) := - \int_{\mathbb{T}^d} \vartheta(t, x) dx$$

$$q_c(t) := - \int_{\mathbb{T}^d} q(t, x) dx.$$

The corrector term w_c is needed for u_1 to be divergence-free. We first compute

$$\begin{aligned} \operatorname{div} w(t) &= \sum_j \operatorname{div}(b_j(t) W^j(t)) \\ &= \sum_j \operatorname{div} \left(b_j(t) \left((\tilde{\varphi}_\mu^j)_\lambda \circ \tau_{\omega t e_j} \right) \psi_v^j e_j \right) \\ &= \sum_j \nabla \left(b_j(t) \left((\tilde{\varphi}_\mu^j)_\lambda \circ \tau_{\omega t e_j} \right) \cdot e_j \right) \psi_v^j. \end{aligned}$$

We thus define

$$w_c(t) := - \sum_j \mathcal{R}_N \left(f_j(t), \psi_v^j \right), \tag{4.17}$$

where we set for simplicity

$$f_j(t) := \nabla \left(a_j(t) \left((\tilde{\varphi}_\mu^j)_\lambda \circ \tau_{\omega t e_j} \right) \right) \tag{4.18}$$

and N is some large integer, which will be chosen in Section 6 together with the parameters $\lambda, \mu, \omega, \nu$. Notice that this definition of the corrector w_c really cancels the divergence of w .

4.3. Estimates on the perturbations

In this section we will formulate and prove all the necessary estimates on the perturbations, beginning with the density terms.

Remark. In this and in the next two sections, Sections 5 and 6, we will denote by C any constant which can depend on the constant M defined in (4.12), on all the parameters in the statement of Proposition 2.1, i.e.

$$p, \tilde{p}, \delta, \eta, \rho_0, u_0, R_0,$$

on the parameter N to be fixed in Section 6 (and on the properties of the functions ϕ, ψ fixed in Section 4.1, in particular their derivatives and antiderivatives up to order N as in the definition of w_c), but not on

$$\lambda, \mu, \omega, \nu.$$

Lemma 4.5 (ϑ in L^p comparable to R_0). *It holds*

$$\|\vartheta(t)\|_{L^p} \leq \frac{M\eta}{2} \|R_0(t)\|_{L^1}^{1/p} + \frac{C}{\lambda^{1/p}}. \tag{4.19}$$

Proof. Applying the improved Hölder inequality (3.1) with $f = a_j(t)$ and $g_\lambda = \Theta^j(t)$ (recall that $\Theta^j(t)$ is $1/\lambda$ -periodic, as ν is an integer multiple of λ) we obtain

$$\begin{aligned} \|\vartheta(t)\|_{L^p} &\leq \sum_j \|a_j(t)\|_{L^p} \left\| \Theta^j(t) \right\|_{L^p} + \frac{C_p}{\lambda^{1/p}} \|a_j\|_{C^1} \left\| \Theta^j(t) \right\|_{L^p} \\ \text{(by (4.13a) and (4.16))} &\leq \frac{M\eta}{2d} \|R_0(t)\|_{L^1}^{1/p} + \frac{C}{\lambda^{1/p}}. \quad \square \end{aligned}$$

Lemma 4.6 (*q small in L^p*). *It holds*

$$\|q(t)\|_{L^p} \leq \frac{C\mu^b}{\omega}. \tag{4.20}$$

Proof. We obtain (4.20) simply from the Hölder inequality, using (4.13a) and (4.16b):

$$\|q(t)\|_{L^p} \leq \sum_j \|a_j b_j\|_{C^0} \|Q^j(t)\|_{L^p} \leq C \frac{\mu^b}{\omega}. \quad \square$$

Lemma 4.7 (*ϑ_c and q_c small as numbers*). *It holds*

$$|\vartheta_c(t)| \leq C\mu^{-b}, \tag{4.21}$$

$$|q_c(t)| \leq C\omega^{-1}. \tag{4.22}$$

Proof. Clearly the correctors are bounded by the L^1 -norm of $\vartheta(t)$ and $q(t)$, so (4.21) and (4.22) follow immediately from (4.13b) and (4.16b):

$$|\vartheta_c(t)| \leq \|\vartheta(t)\|_{L^1} \leq C\mu^{-b}, \quad |q_c(t)| \leq \|q(t)\|_{L^1} \leq C\omega^{-1}. \quad \square$$

Lemma 4.8 (*w in $L^{p'}$ comparable to R_0*). *It holds*

$$\|w(t)\|_{L^{p'}} \leq \frac{M}{2\eta} \|R_0(t)\|_{L^1}^{1/p'} + \frac{C}{\lambda^{1/p'}}. \tag{4.23}$$

Proof. The proof is completely analog to the one of (4.19) and is thus omitted. \square

Lemma 4.9 (*w small in $W^{1,\tilde{p}}$*). *It holds*

$$\|w(t)\|_{W^{1,\tilde{p}}} \leq \frac{C(\lambda\mu + \nu)}{\mu^{1+\epsilon}}. \tag{4.24}$$

Proof. We only use Hölder together with (4.13d) and (4.16b) and obtain

$$\begin{aligned} \|w(t)\|_{W^{1,\tilde{p}}} &\leq \sum_j \|b_j(t)W^j(t)\|_{W^{1,\tilde{p}}} \\ &\leq \sum_j \|b_j\|_{C^1} \|W^j(t)\|_{W^{1,\tilde{p}}} \\ &\leq \frac{C(\lambda\mu + \nu)}{\mu^{1+\epsilon}}. \quad \square \end{aligned}$$

Lemma 4.10 (*Estimates on f_j*). *For every $k, h \in \mathbb{N}$ and $r \in [1, \infty]$*

$$\|\mathcal{D}^k \mathcal{D}^h f_j(t)\|_{L^r} \leq C(\lambda\mu)^{k+h+1} \mu^{b-d/r}.$$

Proof. Recalling the definition of f_j in (4.18), we have

$$\begin{aligned} \|\mathcal{D}^k \mathcal{D}^h f_j(t)\|_{L^r} &\leq \|f_j(t)\|_{W^{k+h,r}} \\ &\leq \|a_j(t) \left((\tilde{\varphi}_\mu^j)_\lambda \circ \tau_{\omega t e_j} \right)\|_{W^{k+h+1,r}} \\ &\leq C \|a_j\|_{C^{k+h+1}} \|(\tilde{\varphi}_\mu^j)_\lambda\|_{W^{k+h+1,r}} \\ \text{(by (4.16b))} &\leq C \lambda^{k+h+1} \|\tilde{\varphi}_\mu^j\|_{W^{k+h+1,r}} \\ \text{(by (4.5))} &\leq C(\lambda\mu)^{k+h+1} \mu^{b-d/r}. \quad \square \end{aligned}$$

Lemma 4.11 (w_c small in $L^{p'}$). *It holds*

$$\|w_c(t)\|_{L^{p'}} \leq C \left(\sum_{k=1}^N \left(\frac{\lambda\mu}{\nu}\right)^k + \frac{(\lambda\mu)^{N+1}}{\nu^N} \right). \tag{4.25}$$

Proof. Applying (3.10) to the definition (4.17) of w_c we immediately obtain

$$\|w_c(t)\|_{L^{p'}} \leq \sum_j C \|\psi\|_{L^\infty} \left(\sum_{k=0}^{N-1} \frac{\|\mathcal{D}^k f_j(t)\|_{L^{p'}}}{\nu^{k+1}} + \frac{\|\mathcal{D}^N f_j(t)\|_{L^{p'}}}{\nu^N} \right).$$

The conclusion follows applying Lemma 4.10 with $h = 0$, $r = p'$ and recalling that $b = d/p'$. \square

Lemma 4.12 (w_c small in $W^{1,\tilde{p}}$). *It holds*

$$\|w_c(t)\|_{W^{1,\tilde{p}}} \leq C \frac{\lambda\mu + \nu}{\mu^{1+\epsilon}} \left(\sum_{k=1}^N \left(\frac{\lambda\mu}{\nu}\right)^k + \frac{(\lambda\mu)^{N+1}}{\nu^N} \right). \tag{4.26}$$

Proof. We will only estimate $\|Dw_c(t)\|_{L^{\tilde{p}}}$ as the estimate on $\|w_c(t)\|_{L^{\tilde{p}}}$ is very similar to the proof of the previous lemma (we just gain a factor of $\mu^{-(1+\epsilon)}$ because of the integrability of $\tilde{\varphi}_\mu$). By statement (ii) of Lemma 3.5 we can split Dw_c into:

$$Dw_c(t) = - \sum_j D\mathcal{R}_N \left(f_j(t), \psi_\nu^j \right) = - \sum_j \mathcal{R}_N \left(Df_j(t), \psi_\nu^j \right) + \mathcal{R}_N \left(f_j, D \left(\psi_\nu^j \right) \right).$$

Both terms can now be estimated analog to the previous lemma by application of (3.10), resulting in (the constant may change from line to line)

$$\begin{aligned} \|Dw_c(t)\|_{L^{\tilde{p}}} &\leq C \sum_j \left[\|\psi\|_{L^\infty} \left(\sum_{k=0}^{N-1} \frac{\|\mathcal{D}^k Df_j(t)\|_{L^{\tilde{p}}}}{\nu^{k+1}} + \frac{\|\mathcal{D}^N Df_j(t)\|_{L^{\tilde{p}}}}{\nu^N} \right) \right. \\ &\quad \left. + \|D\psi_\nu\|_{L^\infty} \left(\sum_{k=0}^{N-1} \frac{\|\mathcal{D}^k f_j(t)\|_{L^{\tilde{p}}}}{\nu^{k+1}} + \frac{\|\mathcal{D}^N f_j(t)\|_{L^{\tilde{p}}}}{\nu^N} \right) \right] \\ \text{(by Lemma 4.10)} &\leq C \left[\mu^{d/p'-d/\tilde{p}} \left(\sum_{k=1}^N \frac{(\lambda\mu)^{k+1}}{\nu^k} + \frac{(\lambda\mu)^{N+2}}{\nu^N} \right) \right. \\ &\quad \left. + \nu \mu^{d/p'-d/\tilde{p}} \left(\sum_{k=1}^N \frac{(\lambda\mu)^k}{\nu^k} + \frac{(\lambda\mu)^{N+1}}{\nu^N} \right) \right] \\ &= C \frac{\lambda\mu + \nu}{\mu^{1+\epsilon}} \left(\sum_{k=1}^N \left(\frac{\lambda\mu}{\nu}\right)^k + \frac{(\lambda\mu)^{N+1}}{\nu^N} \right). \quad \square \end{aligned}$$

5. The new defect field

5.1. Definition of R_1

Given the perturbations defined in the previous section we now have to find a vector field R_1 so that (ρ_1, u_1, R_1) solve (2.1) on $[0, T] \times \mathbb{T}^d$. This is achieved basically by taking the anti-divergence of the left hand side of (2.1), but as we want to show that R_1 can be chosen arbitrarily small in L^1 in order to prove (2.3d), we need to be careful about the exact form of the anti-divergence. Therefore, decompose the left hand side of (2.1) as

$$\begin{aligned}
 -\operatorname{div} R_1 &= \partial_t \rho_1 + \operatorname{div}(\rho_1 u_1) \\
 &= \underbrace{\partial_t \rho_0 + \operatorname{div}(\rho_0 u_0)}_{=-\operatorname{div} R_0} + \partial_t(\vartheta + \vartheta_c + q + q_c) + \operatorname{div}(\rho_0(w + w_c)) + \operatorname{div}((\vartheta + q)u_0) \\
 &\quad + \operatorname{div}((\vartheta + q)(w + w_c)) + \underbrace{(\vartheta_c + q_c) \operatorname{div}((u_0 + w + w_c))}_{=0 \text{ by def. of } w_c} \\
 &= \partial_t(q + q_c) + \operatorname{div}(\vartheta w - R_0) \\
 &\quad + \partial_t(\vartheta + \vartheta_c) + \operatorname{div}(\rho_0 w + \vartheta u_0) \\
 &\quad + \operatorname{div}(q(u_0 + w)) \\
 &\quad + \operatorname{div}((\rho_0 + \vartheta + q)w_c).
 \end{aligned} \tag{5.1}$$

In the next sections we analyze each line in (5.1) separately. In particular we will define and estimate R^χ (in (5.2)), $R^{\text{time},1}$ (in (5.5)), R^{quadr} (in (5.7)), R^{lin} (in (5.11)), $R^{\text{time},2}$ (in (5.12)), R^q (in (5.15)), R^{corr} (in (5.17)), so that

$$\begin{aligned}
 \partial_t(q + q_c) + \operatorname{div}(\vartheta w - R_0) &= \operatorname{div} R^{\text{time},1} + \operatorname{div} R^{\text{quadr}} + \operatorname{div} R^\chi, \\
 \partial_t(\vartheta + \vartheta_c) + \operatorname{div}(\rho_0 w + \vartheta u_0) &= \operatorname{div} R^{\text{time},2} + \operatorname{div} R^{\text{lin}}, \\
 \operatorname{div}(q(u_0 + w)) &= \operatorname{div} R^q, \\
 \operatorname{div}((\rho_0 + \vartheta + q)w_c) &= \operatorname{div} R^{\text{corr}},
 \end{aligned}$$

and thus

$$-\operatorname{div} R_1 = \partial_t \rho_1 + \operatorname{div}(\rho_1 u_1)$$

for

$$-R_1 := R^{\text{time},1} + R^{\text{quadr}} + R^\chi + R^{\text{time},2} + R^{\text{lin}} + R^q + R^{\text{corr}}.$$

5.2. Analysis of the first line in (5.1)

We write

$$R_0 = \sum_j R_0^j e_j = \sum_j (1 - \chi_j^2) R_0^j e_j + \sum_j \chi_j^2 R_0^j e_j$$

and thus

$$\begin{aligned}
 -\operatorname{div} R_0 &= \operatorname{div} \left(R^\chi - \sum_j \chi_j^2 R_0^j e_j \right) \\
 &= \operatorname{div} R^\chi - \sum_j \nabla(\chi_j^2 R_0^j) \cdot e_j \\
 &= \operatorname{div} R^\chi - \sum_j \nabla(a_j b_j) \cdot e_j
 \end{aligned}$$

where we set

$$R^\chi := - \sum_j (1 - \chi_j^2) R_0^j e_j. \tag{5.2}$$

Observe now that, because of (4.14),

$$\vartheta w = \sum_j a_j b_j \Theta^j W^j = \sum_j \chi_j^2 R_0^j \Theta^j W^j.$$

Therefore

$$\operatorname{div}(\vartheta w) = \sum_j a_j b_j \operatorname{div}(\Theta^j W^j) + \nabla(a_j b_j) \cdot \Theta^j W^j$$

and thus

$$\begin{aligned}
 \operatorname{div}(\vartheta w - R_0) &= \sum_j a_j b_j \operatorname{div}(\Theta^j W^j) + \nabla(a_j b_j) \cdot \Theta^j W^j + \operatorname{div} R^\chi - \sum_j \nabla(a_j b_j) \cdot e_j \\
 &= \sum_j a_j b_j \operatorname{div}(\Theta^j W^j) + \nabla(a_j b_j) \cdot [\Theta^j W^j - e_j] + \operatorname{div} R^\chi \\
 &= \sum_j a_j b_j \operatorname{div}(\Theta^j W^j) \\
 &\quad + \left(\nabla(a_j b_j) \cdot [\Theta^j W^j - e_j] - \int \nabla(a_j b_j) \cdot [\Theta^j W^j - e_j] \right) \\
 &\quad + \int \nabla(a_j b_j) \cdot [\Theta^j W^j - e_j] \\
 &\quad + \operatorname{div} R^\chi.
 \end{aligned} \tag{5.3}$$

On the other side

$$\begin{aligned}
 \partial_t(q + q_c) &= \sum_j a_j b_j \partial_t Q^j + \partial_t(a_j b_j) Q^j + q'_c \\
 &= \sum_j a_j b_j \partial_t Q^j + \left(\partial_t(a_j b_j) Q^j - \int \partial_t(a_j b_j) Q^j \right) + \left(\int \partial_t(a_j b_j) Q^j + q'_c \right).
 \end{aligned} \tag{5.4}$$

Putting together (5.3) and (5.4) we get

$$\begin{aligned}
 \partial_t(q + q_c) + \operatorname{div}(\vartheta w - R_0) &= \sum_j a_j b_j \underbrace{[\partial_t Q^j + \operatorname{div}(\Theta^j W^j)]}_{=0 \text{ by (4.15)}} \\
 &\quad + \left(\partial_t(a_j b_j) Q^j - \int \partial_t(a_j b_j) Q^j \right) \\
 &\quad + \left(\nabla(a_j b_j) \cdot [\Theta^j W^j - e_j] - \int \nabla(a_j b_j) \cdot [\Theta^j W^j - e_j] \right) \\
 &\quad + \operatorname{div} R^\chi \\
 &\quad + \underbrace{\int \partial_t(a_j b_j) Q^j + q'_c + \int \nabla(a_j b_j) \cdot [\Theta^j W^j - e_j]}_{=0, \text{ as the l.h.s. has zero mean value}} \\
 &\quad \quad \quad \text{and each other line in the r.h.s. has zero mean value} \\
 &= \sum_j \left(\partial_t(a_j b_j) Q^j - \int \partial_t(a_j b_j) Q^j \right) \\
 &\quad + \left(\nabla(a_j b_j) \cdot [\Theta^j W^j - e_j] - \int \nabla(a_j b_j) \cdot [\Theta^j W^j - e_j] \right) \\
 &\quad + \operatorname{div} R^\chi \\
 &= \operatorname{div} R^{\text{time},1} + \operatorname{div} R^{\text{quadr}} + \operatorname{div} R^\chi,
 \end{aligned}$$

where $R^{\text{time},1}$ is defined by

$$R^{\text{time},1} := \sum_j \left\{ \mathcal{D}^{-1} \left(\partial_t(a_j b_j) Q^j - \int_{\mathbb{T}^d} \partial_t(a_j b_j) Q^j \right) \right\}, \tag{5.5}$$

and R^{quadr} is defined in such a way that

$$\operatorname{div} R^{\text{quadr}} = \sum_j \left\{ \nabla(a_j b_j) \cdot [\Theta^j W^j - e_j] - \int \nabla R_0^j \cdot [\Theta^j W^j - e_j] \right\}, \tag{5.6}$$

as follows. We first compute

$$\begin{aligned} \nabla(a_j b_j) \cdot [\Theta^j W^j - e_j] &= \nabla(a_j b_j) \cdot \left[(\varphi_\mu^j \tilde{\varphi}_\mu^j)_\lambda \circ \tau_{\omega t e_j} (\psi^j)^2 - 1 \right] e_j \\ &= \nabla(a_j b_j) \cdot \left[(\varphi_\mu^j \tilde{\varphi}_\mu^j)_\lambda \circ \tau_{\omega t e_j} \left((\psi^j)^2 - 1 \right) + ((\varphi_\mu^j \tilde{\varphi}_\mu^j)_\lambda \circ \tau_{\omega t e_j} - 1) \right] e_j \\ &= \nabla(a_j b_j) \cdot \left[(\varphi_\mu^j \tilde{\varphi}_\mu^j)_\lambda \circ \tau_{\omega t e_j} \left((\psi^j)^2 - 1 \right)_\nu + (\varphi_\mu^j \tilde{\varphi}_\mu^j - 1)_\lambda \circ \tau_{\omega t e_j} \right] e_j \\ &= \partial_j(a_j b_j) \left[(\varphi_\mu^j \tilde{\varphi}_\mu^j)_\lambda \circ \tau_{\omega t e_j} \left((\psi^j)^2 - 1 \right)_\nu + (\varphi_\mu^j \tilde{\varphi}_\mu^j - 1)_\lambda \circ \tau_{\omega t e_j} \right]. \end{aligned}$$

We then define

$$\begin{aligned} R^{\text{quadr},1} &:= \sum_j \mathcal{R}_1 \left(\partial_j(a_j b_j) (\varphi_\mu^j \tilde{\varphi}_\mu^j)_\lambda \circ \tau_{\omega t e_j}, \left((\psi^j)^2 - 1 \right)_\nu \right), \\ R^{\text{quadr},2} &:= \sum_j \mathcal{R}_1 \left(\partial_j(a_j b_j), (\varphi_\mu^j \tilde{\varphi}_\mu^j - 1)_\lambda \circ \tau_{\omega t e_j} \right), \end{aligned}$$

and

$$R^{\text{quadr}} := R^{\text{quadr},1} + R^{\text{quadr},2}, \tag{5.7}$$

so that (5.6) holds. Notice that the definitions of $R^{\text{quadr},1}$ and $R^{\text{quadr},2}$ are well posed, as

$$\int_{\mathbb{T}^d} \left((\psi^j)^2 - 1 \right)_\nu = 0, \quad \int_{\mathbb{T}^d} (\varphi_\mu^j \tilde{\varphi}_\mu^j - 1)_\lambda \circ \tau_{\omega t e_j} = 0,$$

because of (4.6) and (4.8). We now estimate $R^\chi, R^{\text{time},1}, R^{\text{quadr}}$.

Lemma 5.1 (Bound on R^χ). *It holds*

$$\|R^\chi(t)\|_{L^1} \leq \frac{\delta}{2}. \tag{5.8}$$

Proof. From the definition of χ_j it is obvious that $|R_0^j(t, x)| \leq \frac{\delta}{2d}$ on the support of $(1 - \chi_j^2(t, x))$, so

$$\|R^\chi(t)\|_{L^1} \leq \sum_j \int_{\operatorname{spt}(1 - \chi_j^2(t))} |R_0^j(t, x)| dx \leq d \int_{\mathbb{T}^d} \frac{\delta}{2d} \leq \frac{\delta}{2}. \quad \square$$

Lemma 5.2 (Bound on $R^{\text{time},1}$). *It holds*

$$\|R^{\text{time},1}(t)\|_{L^1} \leq C \frac{1}{\omega}. \tag{5.9}$$

Proof. Using the definition of $R^{\text{time},1}$ in (5.5) and applying Lemma 3.4, we get

$$\begin{aligned} \|R^{\text{time},1}(t)\|_{L^1} &\leq C \sum_j \|\partial_t(a_j(t) b_j(t)) Q^j(t)\|_{L^1} \\ &\leq C \sum_j \|\partial_t(a_j b_j)\|_{C^0} \|Q^j(t)\|_{L^1} \\ &\text{(by (4.13b))} \leq C \frac{1}{\omega}. \quad \square \end{aligned}$$

Lemma 5.3 (Bound on R^{quadr}). *It holds*

$$\|R^{\text{quadr}}(t)\|_{L^1} \leq C \left(\frac{\lambda\mu}{\nu} + \frac{1}{\lambda} \right). \tag{5.10}$$

Proof. First observe that both terms in the definition of R^{quadr} need to be handled separately as the fast oscillation term of $R^{\text{quadr},1}$ is $(1/\nu)$ -periodic whereas in $R^{\text{quadr},2}$ there is only $(1/\lambda)$ -periodicity. For $R^{\text{quadr},1}$, (3.10) (with $N = 1$) and standard Hölder gives us

$$\begin{aligned} \|R^{\text{quadr},1}(t)\|_{L^1} &\leq \frac{C}{\nu} \|\psi^2 - 1\|_{C^0} \left(\left\| \partial_j (a_j(t)b_j(t)) \left(\varphi_\mu^j \tilde{\varphi}_\mu^j \right)_\lambda \circ \tau_{\omega t e_j} \right\|_{L^1} \right. \\ &\quad \left. + \left\| \mathcal{D}^1 \left(\partial_j (a_j(t)b_j(t)) \left(\varphi_\mu^j \tilde{\varphi}_\mu^j \right)_\lambda \circ \tau_{\omega t e_j} \right) \right\|_{L^1} \right) \\ &\leq \frac{C}{\nu} \left(\|\partial_j (a_j(t)b_j(t))\|_{C^0} \left\| \left(\varphi_\mu^j \tilde{\varphi}_\mu^j \right)_\lambda \circ \tau_{\omega t e_j} \right\|_{L^1} \right. \\ &\quad \left. + \|\partial_j (a_j(t)b_j(t))\|_{C^1} \left\| \left(\varphi_\mu^j \tilde{\varphi}_\mu^j \right)_\lambda \circ \tau_{\omega t e_j} \right\|_{W^{1,1}} \right) \\ &\leq \frac{C}{\nu} \|a_j b_j\|_{C^2} \left(\|\varphi_\mu^j \tilde{\varphi}_\mu^j\|_{L^1} + \lambda \|\varphi_\mu^j \tilde{\varphi}_\mu^j\|_{W^{1,1}} \right) \\ &\leq \frac{C\lambda\mu}{\nu}, \end{aligned}$$

where in the last step we used (4.5). For $R^{\text{quadr},2}$ we apply (3.9) (again with $N = 1$) and obtain

$$\begin{aligned} \|R^{\text{quadr},2}(t)\|_{L^1} &\leq C \|\varphi_\mu^j \tilde{\varphi}_\mu^j - 1\|_{L^1} \left(\frac{1}{\lambda} \|\partial_j (\chi_j^2 R_0^j)\|_{C^0} + \frac{1}{\lambda} \|\partial_j (\chi_j^2 R_0^j)\|_{C^1} \right) \\ &\leq C \frac{1}{\lambda}, \end{aligned}$$

as $\|\varphi_\mu^j \tilde{\varphi}_\mu^j\|_{L^1} = 1$, by (4.5). Together these two estimates supply the required bound. \square

5.3. Analysis of the second line in (5.1)

We have

$$\begin{aligned} \partial_t (\vartheta + \vartheta_c) + \text{div}(\vartheta u_0 + \rho_0 w) &= \sum_j a_j \partial_t \Theta^j + (\partial_t a_j) \Theta^j + \text{div}(\vartheta u_0 + \rho_0 w) + \vartheta'_c \\ &= \sum_j \left(a_j \partial_t \Theta^j - \int a_j \partial_t \Theta^j \right) \\ &\quad + \left((\partial_t a_j) \Theta^j - \int (\partial_t a_j) \Theta^j \right) + \text{div}(\vartheta u_0 + \rho_0 w) \\ &\quad + \underbrace{\int a_j \partial_t \Theta^j + \int (\partial_t a_j) \Theta^j + \vartheta'_c}_{=0 \text{ as the l.h.s. and each other line in the r.h.s. has zero mean value}} \\ &= \text{div} R^{\text{time},2} + \text{div} R^{\text{lin}}, \end{aligned}$$

where

$$R^{\text{lin}} := \mathcal{D}^{-1} \left((\partial_t a_j) \Theta^j - \mathcal{F} (\partial_t a_j) \Theta^j \right) + \vartheta u_0 + \rho_0 w \tag{5.11}$$

and $R^{\text{time},2}$ is defined in such a way that

$$\text{div } R^{\text{time},2} = \sum_j \left(a_j \partial_t \Theta^j - \mathcal{F} a_j \partial_t \Theta^j \right),$$

as follows. Using (4.9), we get

$$a_j \partial_t \Theta^j = -\lambda \omega a_j \left((\partial_j \varphi_\mu^j)_\lambda \circ \tau_{\omega t e_j} \right) \psi_v^j$$

and thus we can define

$$R^{\text{time},2} := -\lambda \omega \sum_j \mathcal{R}_N \left(a_j \left((\partial_j \varphi_\mu^j)_\lambda \circ \tau_{\omega t e_j}, \psi_v^j \right), \right) \tag{5.12}$$

where N will be fixed in Section 6, as we have already stressed.

Lemma 5.4 (Bound on R^{lin}). *It holds*

$$\| R^{\text{lin}}(t) \|_{L^1} \leq C \left(\frac{1}{\mu^a} + \frac{1}{\mu^b} \right). \tag{5.13}$$

Proof. For the first term in the definition (5.11) of R^{lin} , Lemma 3.4 yields

$$\begin{aligned} \left\| \mathcal{D}^{-1} \left(\partial_t a_j(t) \Theta^j(t) - \mathcal{F} \partial_t a_j(t) \Theta^j(t) \right) \right\|_{L^1} &\leq C \| \partial_t a_j(t) \Theta^j(t) \|_{L^1} \\ &\leq C \| \partial_t a_j \|_{C^0} \| \Theta^j(t) \|_{L^1} \\ &\text{(by (4.13b))} \leq \frac{C}{\mu^b}. \end{aligned}$$

For the second term in the definition (5.11) of R^{lin} , simply apply Hölder’s inequality

$$\begin{aligned} \| \rho_0(t) w(t) \|_{L^1} &\leq \| \rho_0 \|_{C^0} \| a_j \|_{C^0} \| W^j(t) \|_{L^1} \\ &\text{(by (4.13b))} \leq \frac{C}{\mu^a}. \end{aligned}$$

The third term is handled completely analog, resulting in

$$\| \vartheta(t) u_0(t) \|_{L^1} \leq \frac{C}{\mu^b}.$$

By adding the three terms we obtain the required bound. \square

Lemma 5.5 (Bound on $R^{\text{time},2}$). *It holds*

$$\| R^{\text{time},2}(t) \| \leq C \frac{\omega}{\mu^b} \left(\sum_{k=1}^N \left(\frac{\lambda \mu}{v} \right)^k + \frac{(\lambda \mu)^{N+1}}{v^N} \right). \tag{5.14}$$

Proof. $R^{\text{time},2}$ is defined in (5.12) by application of the bilinear anti-divergence operator \mathcal{R}_N of Section 3.4 to the product of $a_j (\partial_j \varphi_\mu^j)_\lambda \circ \tau_{\omega t e_j}$ and ψ_v^j , so (3.10) yields

$$\begin{aligned}
 \|R^{\text{time},2}(t)\|_{L^1} &\leq C\lambda\omega\left(\sum_{k=0}^{N-1}\frac{1}{\nu^{k+1}}\left\|\mathcal{D}^k\left(a_j(t)(\partial_j\varphi_\mu^j)_\lambda\circ\tau_{\omega t e_j}\right)\right\|_{L^1}+\frac{1}{\nu^N}\left\|\mathcal{D}^N\left(a_j(t)(\partial_j\varphi_\mu^j)_\lambda\circ\tau_{\omega t e_j}\right)\right\|_{L^1}\right) \\
 &\leq C\lambda\omega\left(\sum_{k=0}^{N-1}\frac{1}{\nu^{k+1}}\left\|a_j(t)(\partial_j\varphi_\mu^j)_\lambda\circ\tau_{\omega t e_j}\right\|_{W^{k,1}}+\frac{1}{\nu^N}\left\|a_j(t)(\partial_j\varphi_\mu^j)_\lambda\circ\tau_{\omega t e_j}\right\|_{W^{N,1}}\right) \\
 &\leq C\lambda\omega\left(\sum_{k=0}^{N-1}\frac{1}{\nu^{k+1}}\|a_j\|_{C^k}\left\|(\partial_j\varphi_\mu^j)_\lambda\right\|_{W^{k,1}}+\frac{1}{\nu^N}\|a_j\|_{C^N}\left\|(\partial_j\varphi_\mu^j)_\lambda\right\|_{W^{N,1}}\right) \\
 &\leq C\lambda\omega\left(\sum_{k=0}^{N-1}\frac{\left\|(\partial_j\varphi_\mu^j)_\lambda\right\|_{W^{k,1}}}{\nu^{k+1}}+\frac{\left\|(\partial_j\varphi_\mu^j)_\lambda\right\|_{W^{N,1}}}{\nu^N}\right) \\
 \text{(by (4.5))} &\leq C\lambda\mu^{1-b}\omega\left(\sum_{k=0}^{N-1}\frac{(\lambda\mu)^k}{\nu^{k+1}}+\frac{(\lambda\mu)^N}{\nu^N}\right) \\
 &\leq C\frac{\omega}{\mu^b}\left(\sum_{k=1}^N\left(\frac{\lambda\mu}{\nu}\right)^k+\frac{(\lambda\mu)^{N+1}}{\nu^N}\right),
 \end{aligned}$$

which is exactly the desired inequality. \square

5.4. Analysis of the third line in (5.1)

We simply define

$$R^q := q(u_0 + w). \tag{5.15}$$

Lemma 5.6 (Bound on R^q). *It holds*

$$\|R^q(t)\|_{L^1} \leq C\frac{\mu^b}{\omega}. \tag{5.16}$$

Proof. From the definitions of q and w we immediately get

$$\begin{aligned}
 \|R^q(t)\|_{L^1} &\leq \|q(t)\|_{L^1}\left(\|u_0(t)\|_{C^0}+\|w(t)\|_{C^0}\right) \\
 &\leq \sum_j\|a_j b_j\|_{C^0}\|Q^j(t)\|_{L^1}\left(\|u_0\|_{C^0}+\sum_i\|b_i\|_{C^0}\|W^i(t)\|_{C^0}\right) \\
 &\leq C\sum_j\|Q^j(t)\|_{L^1}\left(1+\sum_i\|W^i(t)\|_{C^0}\right)
 \end{aligned}$$

$$\text{(by (4.13b) and (4.13c))} \leq \frac{C}{\omega}(1+\mu^b),$$

which implies the desired inequality. \square

5.5. Analysis of the fourth line in (5.1)

We simply define

$$R^{\text{corr}} := (\rho_0 + \vartheta + q)w_c. \tag{5.17}$$

Lemma 5.7 (Bound on R^{corr}). *It holds*

$$\|R^{\text{corr}}(t)\|_{L^1} \leq C\left(1+\frac{1}{\lambda^{1/p}}+\frac{\mu^b}{\omega}\right)\left(\sum_{k=1}^N\left(\frac{\lambda\mu}{\nu}\right)^k+\frac{(\lambda\mu)^{N+1}}{\nu^N}\right).$$

Proof. The inequality is easier to prove than to state as it is an immediate consequence of Lemmata 4.5, 4.6 and 4.11. We omit the details. \square

6. Proof of the main proposition

Given the estimates proven in Sections 4 and 5 we are now able to prove Proposition 2.1. Let $p \in (1, \infty)$ and $\tilde{p} \in [1, \infty)$ so that (2.2) holds. Let $\delta, \eta > 0$ and let

$$(\rho_0, u_0, R_0) : [0, T] \times \mathbb{T}^d \rightarrow \mathbb{R} \times \mathbb{R}^d \times \mathbb{R}^d$$

be a smooth solution of the incompressible continuity-defect equation (2.1).

6.1. Choice of parameters

Recall that M was defined in (4.12). Let ϵ be as in (4.11) and note that $\epsilon > 0$ by (2.2). Recall that $a = d/p > 0$ and $b = d/p' > 0$. For some large positive integer λ to be defined later:

- (1) Set $\mu := \lambda^\alpha$ for some $\alpha(\epsilon) > 2\epsilon^{-1} > \epsilon^{-1}$.
- (2) Set $\nu := \lambda^\gamma$ for a natural number $\gamma(\alpha, \epsilon)$ chosen such that

$$\alpha + 1 < \gamma < \alpha(1 + \epsilon)$$

which is possible by the choice of α . In this way, ν is a multiple of λ and the Mikado functions defined in Section 4.1 are λ -periodic.

- (3) Choose $\beta(b, \alpha, \gamma)$ such that

$$b\alpha < \beta < b\alpha + \gamma - (\alpha + 1)$$

which is possible by the first condition on γ , and set $\omega := \lambda^\beta$.

- (4) Finally, choose an integer $N(\alpha, \gamma)$ which is large enough so that

$$\frac{N}{N-1} < \frac{\gamma}{1+\alpha}$$

which is also possible by the first condition on γ .

Let us summarize the conditions imposed by our choice of the parameters α, β, γ and N :

$$1 < \alpha\epsilon \tag{6.1a}$$

$$\alpha + 1 < \gamma \tag{6.1b}$$

$$\gamma < \alpha(1 + \epsilon) \tag{6.1c}$$

$$b\alpha < \beta \tag{6.1d}$$

$$\beta + 1 + \alpha < b\alpha + \gamma \tag{6.1e}$$

$$N(1 + \alpha) < (N - 1)\gamma. \tag{6.1f}$$

6.2. Definition of the new solution

Let (ρ_1, u_1) be as defined in Section 4 and R_1 as in Section 5. Then (ρ_1, u_1, R_1) is a solution of (2.1) as stated in the construction of R_1 . Clearly the solution is smooth in time and space (ensured by the cut-offs χ_j) and it is equal to $(\rho_0, u_0, R_0)(t)$ if $R_0(t) \equiv 0$ holds, as the construction is completely local in time apart from the definition of R^{lin} and $R^{\text{time},1}$, which contain the time derivative of R_0 . However, by the definition of the cut-off functions χ_j it is clear that

$$R_0(t) \equiv 0 \implies \partial_t a_j = \partial_t \left(\chi_j(t, \cdot) |R_0^j(t, \cdot)|^{1/p} \right) \equiv 0$$

and analog for $\partial_t(a_j b_j)$, so also $R^{\text{lin}}(t), R^{\text{time},1}(t) \equiv 0$ holds.

We need to show (2.3a)–(2.3d), which is equivalent to

$$\|\vartheta(t) + q(t) + \vartheta_c(t)\|_{L^p} \leq M\eta \|R_0(t)\|_{L^1}^{1/p} \tag{6.2a}$$

$$\|w(t) + w_c(t)\|_{L^{p'}} \leq \frac{M}{\eta} \|R_0(t)\|_{L^1}^{1/p'} \tag{6.2b}$$

$$\|w(t) + w_c(t)\|_{W^{1,\bar{p}}} \leq \delta \tag{6.2c}$$

$$\left\| \left(R^{\text{time},1} + R^{\text{quadr}} + \text{div } R^\chi + R^{\text{time},2} + R^{\text{lin}} + R^q + R^{\text{corr}} \right) (t) \right\|_{L^1} \leq \delta. \tag{6.2d}$$

Remark. In all these definitions the oscillation parameter $\lambda \in \mathbb{N}$ is still to be fixed. It will be chosen sufficiently large in the following estimates. Note that this is possible as there is no upper bound on λ here.

6.3. Estimates on the perturbations

Set

$$A := \{t \in [0, T] : \|R_0(t)\|_{L^\infty} < \delta/4d\}, \quad B := [0, T] \setminus A.$$

Since R_0 is a smooth function, A is open in $[0, T]$ and thus B is compact. It must then hold

$$\inf_{t \in B} \|R_0(t)\|_{L^1} = \min_{t \in B} \|R_0(t)\|_{L^1} > 0.$$

If $t \in A$, then $\chi_j(t) \equiv 0$ for every j and thus, by definition, $\vartheta(t) = q(t) = \vartheta_c(t) = w(t) = w_c(t) = 0$. Hence, (6.2a) trivially holds. If $t \in B$, Lemmata 4.5, 4.6 and 4.7 provide the desired bound on the density perturbation:

$$\begin{aligned} \|\vartheta(t) + q(t) + \vartheta_c(t) + q_c(t)\|_{L^p} &\leq \|\vartheta(t)\|_{L^p} + \|q(t)\|_{L^p} + |\vartheta_c(t)| + |q_c(t)| \\ &\leq \frac{M\eta}{2} \|R_0(t)\|_{L^1}^{1/p} + C \left(\frac{1}{\lambda^{1/p}} + \frac{\mu^b}{\omega} + \frac{1}{\mu^b} + \frac{1}{\omega} \right) \\ &= \frac{M\eta}{2} \|R_0(t)\|_{L^1}^{1/p} + C \left(\lambda^{-1/p} + \lambda^{b\alpha-\beta} + \lambda^{-b\alpha} + \lambda^{-\beta} \right). \end{aligned}$$

Because of (6.1d) and the facts $p < \infty$ and $b > 0$ the second summand can be made arbitrarily small by choosing λ sufficiently large. More precisely, we can choose λ so that

$$C \left(\lambda^{-1/p} + \lambda^{b\alpha-\beta} + \lambda^{-b\alpha} + \lambda^{-\beta} \right) < \frac{M\eta}{2} \min_{t \in B} \|R_0(t)\|_{L^1}^{1/p},$$

which, in particular, proves (6.2a). Notice that, taking the minimum of the $\|R_0(t)\|_{L^1}$, we ensure that λ can be chosen independent of t .

For the L^p -bound on the velocity perturbation we need Lemmata 4.8 and 4.11.

$$\begin{aligned} \|w + w_c\|_{L^{p'}} &\leq \frac{M}{2\eta} \|R_0\|_{L^1}^{1/p'} + C \left(\frac{1}{\lambda^{1/p'}} + \sum_{k=1}^N \left(\frac{\mu\lambda}{\nu} \right)^k + \frac{(\mu\lambda)^{N+1}}{\nu^N} \right) \\ &= \frac{M}{2\eta} \|R_0\|_{L^1}^{1/p'} + C \left(\lambda^{-1/p'} + \sum_{k=1}^N \left(\lambda^{1+\alpha-\gamma} \right)^k + \lambda^{(N+1)(1+\alpha)-N\gamma} \right). \end{aligned}$$

Because of (6.1b) we have $\lambda^{1+\alpha-\gamma} < 1$, so the sum inside the parentheses is bounded by $N\lambda^{1+\alpha-\gamma}$. Furthermore

$$(N + 1)(1 + \alpha) - N\gamma < N(1 + \alpha) - (N - 1)\gamma < 0$$

holds by (6.1f). Observe also that $p' < \infty$, so all the exponents of λ in the parentheses are negative so the term can be made arbitrarily small by choosing λ sufficiently large, which proves (6.2b).

For (6.2c) we apply Lemmata 4.9 and 4.12 and obtain

$$\begin{aligned} \|w + w_c\|_{W^{1,\tilde{p}}} &\leq C \left(\frac{\lambda\mu + \nu}{\mu^{1+\epsilon}} \right) \left(1 + \sum_{k=1}^N \left(\frac{\lambda\mu}{\nu} \right)^k + \frac{(\lambda\mu)^{N+1}}{\nu^N} \right) \\ &= C \left(\lambda^{1-\alpha\epsilon} + \lambda^{\gamma-\alpha(1-\epsilon)} \right) \left(\sum_{k=0}^N \lambda^{k(1+\alpha-\gamma)} + \lambda^{(N+1)(1+\alpha)-N\gamma} \right). \end{aligned}$$

Again because of (6.1b) and (6.1f) each summand inside the second parentheses is bounded by 1, so the inequality boils down to

$$\|w + w_c\|_{W^{1,\tilde{p}}} \leq C(N + 2) \left(\lambda^{1-\alpha\epsilon} + \lambda^{\gamma-\alpha(1-\epsilon)} \right).$$

Both exponents of λ in this expression are negative: The first one is by condition (6.1a) and the second by (6.1c). Therefore, if λ is large enough, (6.2c) holds.

6.4. Estimates on the new error

By Lemma 5.1 the smoothness corrector term R^χ is bounded in L^1 by $\frac{\delta}{2}$ so in order to prove (6.2d) we need to show that the sum of all other components of the defect field R_1 is smaller than $\frac{\delta}{2}$ in L^1 . Most of the terms are bounded analog to the density and velocity perturbations, by Lemmata 5.3, 5.4 and 5.6:

$$\begin{aligned} \|R^{\text{quadr}}\|_{L^1} &\leq C \left(\lambda^{1+\alpha-\gamma} + \lambda^{-1} \right), \\ \|R^{\text{lin}}\|_{L^1} &\leq C \left(\lambda^{-a\alpha} + \lambda^{-b\alpha} \right), \\ \|R^q\|_{L^1} &\leq C \lambda^{b\alpha-\beta}, \\ \|R^{\text{time},1}\|_{L^1} &\leq C \lambda^{-\beta}. \end{aligned}$$

These terms are small for large λ because of (6.1b) (first line), as $a, b > 0$ (second line) and by (6.1d) (third and fourth line).

The two remaining terms require more attention. Lemma 5.7 provides the following bound on R^{corr} :

$$\|R^{\text{corr}}\|_{L^1} \leq C \left(1 + \lambda^{-1/p} + \lambda^{b\alpha-\beta} \right) \left(\sum_{k=1}^N \lambda^{k(1+\alpha-\gamma)} + \lambda^{(N+1)(1+\alpha)-N\gamma} \right).$$

By (6.1d) the term in the first parentheses is bounded by 3, the second one is small for large λ because of (6.1b) and (6.1f) by the same argument as above in the estimate of the velocity perturbation. The last remaining term is $R^{\text{time},2}$, which is taken care of in Lemma 5.5:

$$\begin{aligned} \|R^{\text{time},2}\|_{L^1} &\leq C \lambda^{\beta-b\alpha} \left(\sum_{k=1}^N \lambda^{k(1+\alpha-\gamma)} + \lambda^{(N+1)(1+\alpha)-N\gamma} \right) \\ &= C \lambda^{\beta+1+\alpha-(b\alpha+\gamma)} \left(\sum_{k=0}^{N-1} \lambda^{k(1+\alpha-\gamma)} + \lambda^{N(1+\alpha)-(N-1)\gamma} \right). \end{aligned}$$

Now (6.1b) and (6.1f) implies that the parentheses is bounded by $N + 1$. Moreover the exponent $\beta + 1 + \alpha - (b\alpha + \gamma)$ is negative because of condition (6.1e), so the term is arbitrarily small if λ is chosen sufficiently large. This concludes the proof of (6.2d) and thus the proof of the proposition.

7. Sketch of the proof of Proposition 2.1 for $p = 1$ and of Theorems 1.3 and 1.4

7.1. The case of continuous vector fields

The proof of Proposition 2.1 at some points requires an integrability of the density perturbation ϑ which is strictly better than L^1 , most crucially in Lemma 5.4: smallness for the term $\|\vartheta u_0\|_{L^1}$ is impossible in the construction of the perturbation as presented in the previous sections.

In [18] the same problem was solved by letting the Mikados “deform with the flow” so that the transport term in the linear part of R_1 ,

$$\operatorname{div} R^{\text{transport}} = (\partial_t + u_0 \cdot \nabla) \left(\vartheta - \int_{\mathbb{T}^d} \vartheta \right)$$

is sufficiently small because of a cancellation in the Mikado function.

More precisely, since u_0 is smooth, there exists the “inverse flow map”, a smooth function $\Phi : [0, T] \times \mathbb{T}^d \rightarrow \mathbb{T}^d$ which solves

$$\partial_t \Phi + u_0 \cdot \nabla \Phi = 0, \quad \Phi(t_0, x) = x.$$

Moreover $\Phi(t, \cdot) : \mathbb{T}^d \rightarrow \mathbb{T}^d$ is close to the identity if t is close to t_0 . In [18] the perturbations are now defined using the pushforward of the Mikado density and flow. Ignoring corrector and cut-offs and using our notation the density perturbations locally in time has the representation

$$\vartheta(t, x) = \eta \sum_j R_0^j(t, x) \Theta_{\lambda, \mu}^j(\Phi(t, x)).$$

It is easy to see that from this definition the transport term in the new defect field reduces to

$$(\partial_t + u_0(t, x) \cdot \nabla) \vartheta(t, x) = \eta \sum_j \Theta_{\lambda, \mu}^j(\Phi(t, x)) (\partial_t + u_0(t, x) \cdot \nabla) R_0^j(t, x),$$

whose anti-divergence is of order $1/\lambda$ in L^1 -norm, because of the fast oscillating Mikado $\Theta_{\lambda, \mu}^j$.

In the construction presented in Section 4 it is advantageous to apply the pushforward only on the fast oscillating factor $\psi(vx)$ and not on the space-time Mikado functions $\varphi^j(t, x)$, which ensure the disjoint support where necessary. The density perturbation then takes the form

$$\vartheta(t, x) = \eta \sum_j R_0^j(t, x) \varphi_\mu^j(\lambda(x - \omega t e_j)) \psi^j(v\Phi(t, x)).$$

On the one hand the transport term also contains derivatives of $(\varphi_\mu)_\lambda$, which excludes the possibility of a cheap L^1 -estimate. However, the term is almost identical to $\partial_t \vartheta$, so it is possible to estimate its anti-divergence analog to Lemma 5.5. On the other hand, since the definition of the space-time Mikado functions $\varphi^j(t, x)$ remains untouched, we still have disjoint support of Mikados in different directions, so there will not be any nontrivial interactions (“Third issue” in Section 2 of [18]) which need to be controlled.

All the other estimates in Sections 4 to 6 remain valid under this redefinition, so Proposition 2.1 can be proved with $p = 1$. For the technical details see [18].

7.2. Handling the diffusion term

In order to prove Theorem 1.3 we only need to add minor adjustments and one more estimate to the proof presented in Sections 3 to 6. The cheapest way to prove that (ρ_n, u_n, R_n) converges to a solution of (1.16) is by showing that $\nabla \rho_n$ converges in L^1 . This way we can keep the construction of the perturbations untouched and just add $\nabla(\rho_n - \rho_{n-1})$ to the new defect field R_n . Then clearly

$$\partial_t \rho_1 + \operatorname{div}(\rho_1 u_1) - \Delta \rho_1 = -\operatorname{div}(R_1) - \Delta \rho_1 = -\operatorname{div}(R_1 + \nabla \rho_1)$$

holds and it suffices to show that $\nabla(\rho_1 - \rho_0)$ is small in L^1 . This estimate is straightforward: with the notation introduced in Section 4 we obtain

$$\begin{aligned} \|\nabla\vartheta\|_{L^1} &\leq C \frac{1 + \lambda\mu + \nu}{\mu^b} = C \frac{1 + \lambda^{1+\alpha} + \lambda^\gamma}{\lambda^{b\alpha}}, \\ \|\nabla q\|_{L^1} &\leq C \frac{1 + \lambda\mu + \nu}{\omega} = C \frac{1 + \lambda^{1+\alpha} + \lambda^\gamma}{\lambda^\beta} \end{aligned}$$

(and trivially $\nabla\vartheta_c = 0$). We need to redefine ϵ so that

$$0 < \epsilon < \min \left\{ \frac{d}{\tilde{p}} - \frac{d}{p'} - 1, \frac{d}{p'} - 1 \right\},$$

which is always possible by the additional condition (1.17) in the statement of the theorem. Choose the parameters α, β, γ exactly as before and observe that

$$b > 1 + \epsilon \implies b\alpha > \alpha(1 + \epsilon) > \gamma > 1 + \alpha$$

by conditions (6.1c) and (6.1a) and therefore $\|\nabla\vartheta\|_{L^1}$ is small for large λ . Similarly, $\|\nabla q\|_{L^1}$ is also small as by (6.1d) in particular $\beta > \gamma, 1 + \alpha$. This concludes the proof of an analog of Proposition 2.1 in the viscous case and thus Theorem 1.3.

7.3. Solutions of higher regularity

Also for Theorem 1.4 the already existing proof requires only some adjustments and more estimates. For the sake of completeness and in order to motivate the extra conditions in the statement we state the analog of the main proposition.

Proposition 7.1. *There is constant $M > 0$ such that the following holds. Let $p, \tilde{p} \in [1, \infty)$ and $m, \tilde{m} \in \mathbb{N}$ such that (1.19) holds. There is $s \in (1, \infty)$ such that for any $\delta > 0$ and any smooth solution (ρ_0, u_0, R_0) of*

$$\begin{aligned} \partial_t \rho + \operatorname{div}(\rho u) + L_k \rho &= -\operatorname{div} R, \\ \operatorname{div} u &= 0, \end{aligned}$$

there is another smooth solution (ρ_1, u_1, R_1) which fulfills for any $t \in [0, T]$

$$\|\rho_1(t) - \rho_0(t)\|_{L^s} \leq M \|\mathbf{R}_0(t)\|^{1/s} \tag{7.1a}$$

$$\|u_1(t) - u_0(t)\|_{L^{s'}} \leq M \|\mathbf{R}_0(t)\|^{1/s'} \tag{7.1b}$$

$$\|\rho_1(t) - \rho_0(t)\|_{W^{m,p}} \leq \delta \tag{7.1c}$$

$$\|u_1(t) - u_0(t)\|_{W^{\tilde{m},\tilde{p}}} \leq \delta \tag{7.1d}$$

$$\|\rho_1(t) - \rho_0(t)\|_{W^{k-1,1}} \leq \delta \tag{7.1e}$$

$$\|\mathbf{R}_1(t)\|_{L^1} \leq \delta \tag{7.1f}$$

$$R_0(t) \equiv 0 \implies R_1(t) \equiv 0. \tag{7.1g}$$

Proof of Theorem 1.4. For the order k differential operator of L_k there is an operator \tilde{L}_k such that

$$L_k f = \operatorname{div} \tilde{L}_k f \text{ for any smooth } f.$$

Observe that $\|\tilde{L}f\|_{L^r} \lesssim \|f\|_{W^{k-1,r}}$, so (7.1e) in particular implies

$$\left\| \tilde{L}_k(\rho_1 - \rho_0) \right\|_{L^1} \leq \delta.$$

This guarantees that $R_n(t) \rightarrow 0$ in L^1 , uniformly in time. Completely analog to the proof of Theorem 1.2 we construct a sequence (ρ_n, u_n, R_n) of smooth solutions satisfying the bounds

$$\begin{aligned} \|\rho_{n+1}(t) - \rho_n(t)\|_{L^s} &\leq M \|R_n(t)\|^{1/s} \leq M \delta_{n-1}^{1/s} \\ \|u_{n+1}(t) - u_n(t)\|_{L^{s'}} &\leq M \|R_n(t)\|^{1/s'} \leq M \delta_{n-1}^{1/s'} \\ \|\rho_{n+1}(t) - \rho_n(t)\|_{W^{m,p}} &\leq \delta_n \\ \|u_{n+1}(t) - u_n(t)\|_{W^{\tilde{m},\tilde{p}}} &\leq \delta_n \\ \|\rho_{n+1}(t) - \rho_n(t)\|_{W^{k-1,1}} &\leq \delta_n \\ \|R_{n+1}(t)\|_{L^1} &\leq \delta_n \\ R_n(t) \equiv 0 &\implies R_{n+1}(t) \equiv 0 \end{aligned}$$

for $(\rho_0, u_0) = (\bar{\rho}, \bar{u})$ and a sequence of positive numbers $(\delta_n)_{n \in \mathbb{N}}$ chosen such that

$$\sum_{n \in \mathbb{N}} \delta_n^{1/s} < \infty, \quad \sum_{n \in \mathbb{N}} \delta_n^{1/s'} < \infty, \quad \sum_{n \in \mathbb{N}} \delta_n < \infty,$$

and, in addition,

$$M \sum_{n \in \mathbb{N}} \delta_n^{1/s} < \varepsilon$$

if we want to show (iv) or

$$M \sum_{n \in \mathbb{N}} \delta_n^{1/s'} < \varepsilon$$

if we want to show (iv'). Then the limit

$$\rho_n \xrightarrow{n \rightarrow \infty} \rho \text{ in } C([0, T], W^{m,p}(\mathbb{T}^d)), \quad u_n \xrightarrow{n \rightarrow \infty} u \text{ in } C([0, T], W^{\tilde{m},\tilde{p}}(\mathbb{T}^d))$$

fulfills statements (i)–(iv) of the theorem. \square

We only give a sketch of the proof of Proposition 7.1, as it is mostly analog to the proof of Proposition 2.1. The only important difference is that in general $u_1 \in L^{p'}$ does not hold, which is needed for the L^1 -convergence of the product $\rho_n u_n$ and we want the density perturbation to be small in the Sobolev space $W^{m,p}$, which was not necessary before. We address both issues by defining the Mikados in a slightly different way: the ‘‘concentration scaling’’ of Mikado density Θ_λ and Mikado field W_λ is now given by

$$\varphi_\mu(x) = \mu^a \varphi(\mu x), \quad \tilde{\varphi}_\mu(x) = \mu^b \varphi(\mu x) \text{ where } a = \frac{d}{s}, \quad b = \frac{d}{s'}$$

for $s \in (1, \infty)$ chosen such that

$$\frac{1}{p} - \frac{m}{d} > \frac{1}{s} = 1 - \frac{1}{s'} > 1 + \frac{\tilde{m}}{d} - \frac{1}{\tilde{p}} \text{ and } \frac{1}{s'} > \frac{k-1}{d}.$$

Note that such an s must exist because of (1.19).

With a suitable M and positive numbers $\epsilon_1, \epsilon_2, \epsilon_3$ defined as

$$\begin{aligned} \epsilon_1 &:= \frac{d}{m} \left(\frac{1}{p} - \frac{1}{s} \right) - 1 \\ \epsilon_2 &:= \frac{d}{\tilde{m}} \min \left\{ \frac{1}{\tilde{p}} - \frac{1}{s'}, \frac{1}{\tilde{p}} - \frac{k-1}{d} \right\} - 1 \\ \epsilon_3 &:= \frac{d}{s'(k-1)} - 1 \end{aligned}$$

the scaling of the Mikados implies

$$\begin{aligned} \|\Theta_{\lambda,\mu,\omega,\nu}\|_{L^s}, \|W_{\lambda,\mu,\omega,\nu}\|_{L^{s'}} &\leq M \\ \|Q_{\lambda,\mu,\omega,\nu}\|_{L^s} &\lesssim \frac{\mu^b}{\omega} \\ \|\Theta_{\lambda,\mu,\omega,\nu}\|_{W^{m,p}} &\lesssim \left(\frac{\lambda\mu + \nu}{\mu^{1+\epsilon_1}}\right)^m \\ \|Q_{\lambda,\mu,\omega,\nu}\|_{W^{m,p}} &\lesssim \frac{(\lambda\mu + \nu)^m \mu^{d/p'}}{\omega} \\ \|W_{\lambda,\mu,\omega,\nu}\|_{W^{\tilde{m},\tilde{p}}} &\lesssim \left(\frac{\lambda\mu + \nu}{\mu^{1+\epsilon_2}}\right)^{\tilde{m}} \\ \|\Theta_{\lambda,\mu,\omega,\nu}\|_{W^{k-1,1}} &\lesssim \left(\frac{\lambda\mu + \nu}{\mu^{1+\epsilon_3}}\right)^{k-1} \\ \|Q_{\lambda,\mu,\omega,\nu}\|_{W^{k-1,1}} &\lesssim \frac{(\lambda\mu + \nu)^{k-1}}{\omega}. \end{aligned}$$

Choosing the parameters $\mu = \lambda^\alpha$, $\omega = \mu^\beta$ and $\nu = \lambda^\gamma$ dependent of b and $\epsilon := \min\{\epsilon_1, \epsilon_2, \epsilon_3\}$ according to (6.1a) the proof of all necessary estimates is analog to those in Sections 4, 5 and 6.

Declaration of competing interest

There is no competing interest.

References

- [1] L. Ambrosio, Transport equation and Cauchy problem for BV vector fields, *Invent. Math.* 158 (2) (2004) 227–260.
- [2] L. Ambrosio, Well posedness of ODE's and continuity equations with nonsmooth vector fields, and applications, *Rev. Mat. Complut.* 30 (3) (2017) 427–450.
- [3] S. Bianchini, P. Bonicatto, A uniqueness result for the decomposition of vector fields in \mathbb{R}^d , *Invent. Math.* (2019).
- [4] T. Buckmaster, M. Colombo, V. Vicol, Wild solutions of the Navier-Stokes equations whose singular sets in time have Hausdorff dimension strictly less than 1, arXiv:1809.00600, 2018.
- [5] T. Buckmaster, C. De Lellis, L. Székelyhidi Jr., V. Vicol, Onsager's conjecture for admissible weak solutions, *Commun. Pure Appl. Math.* 72 (2) (2019) 229–274.
- [6] T. Buckmaster, V. Vicol, Nonuniqueness of weak solutions to the Navier-Stokes equation, *Ann. Math.* (2019).
- [7] L. Caravenna, G. Crippa, Uniqueness and Lagrangianity for solutions with lack of integrability of the continuity equation, *C. R. Math. Acad. Sci. Paris* 354 (12) (2016) 1168–1173.
- [8] L. Caravenna, G. Crippa, A directional Lipschitz extension lemma, with applications to uniqueness and Lagrangianity for the continuity equation, arXiv:1812.06817, 2018.
- [9] A. Cheskidov, X. Luo, Stationary and discontinuous weak solutions of the Navier-Stokes equations, arXiv:1901.07485, 2019.
- [10] S. Daneri, L. Székelyhidi Jr., Non-uniqueness and h-principle for Hölder-continuous weak solutions of the Euler equations, *Arch. Ration. Mech. Anal.* 224 (2) (2017) 471–514.
- [11] N. Depauw, Non unicité des solutions bornées pour un champ de vecteurs BV en dehors d'un hyperplan, *C. R. Math. Acad. Sci. Paris* 337 (4) (2003) 249–252.
- [12] R.J. DiPerna, P.-L. Lions, Ordinary differential equations, transport theory and Sobolev spaces, *Invent. Math.* 98 (3) (1989) 511–547.
- [13] D. Gilbarg, N.S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer-Verlag, Berlin Heidelberg, 2001.
- [14] P. Isett, A Proof of Onsager's Conjecture, *Ann. Math.* 188 (3) (2018) 871–963.
- [15] T. Luo, E.S. Titi, Non-uniqueness of Weak Solutions to Hyperviscous Navier-Stokes Equations – On Sharpness of J.-L. Lions Exponent, arXiv:1808.07595, 2018.
- [16] X. Luo, Stationary solutions and nonuniqueness of weak solutions for the Navier-Stokes equations in high dimensions, arXiv:1807.09318, 2018.
- [17] S. Modena, L. Székelyhidi Jr., Non-uniqueness for the transport equation with Sobolev vector fields, *Ann. PDE* 4 (2) (2018) 18.
- [18] S. Modena, L. Székelyhidi Jr., Non-renormalized solutions to the continuity equation, *Calc. Var. Partial Differ. Equ.* 58 (2019) 208.
- [19] C. Seis, A quantitative theory for the continuity equation, *Ann. Inst. Henri Poincaré C, Anal. Non Linéaire* 34 (7) (2017) 1837–1850.