

ANNALES SCIENTIFIQUES
DE L'UNIVERSITÉ DE CLERMONT-FERRAND 2
Série Probabilités et applications

R. ZAHAROPOL

**A zero-two theorem for a certain class of positive contractions
in finite dimensional L^p -spaces** ($1 \leq p < +\infty$)

Annales scientifiques de l'Université de Clermont-Ferrand 2, tome 78, série *Probabilités et applications*, n° 2 (1984), p. 9-13

http://www.numdam.org/item?id=ASCFPA_1984__78_2_9_0

© Université de Clermont-Ferrand 2, 1984, tous droits réservés.

L'accès aux archives de la revue « Annales scientifiques de l'Université de Clermont-Ferrand 2 » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

A Zero-Two Theorem for a certain class of positive
Contractions in Finite Dimensional L^p -spaces ($1 \leq p < +\infty$)

R. ZAHAROPOL

Summary

Our goal here is to extend Theorem 1.1 from [2] (which is sometimes called the zero-two law for positive contractions in L^1 -spaces) to a class of positive contractions in finite dimensional L^p -spaces ($1 \leq p < +\infty$).

1. A General Lemma

Let (X, Σ, m) be a measure space and $L^p(X, \Sigma, m)$ ($1 \leq p < +\infty$) the usual Banach spaces. By a positive contraction $T: L^p(X, \Sigma, m) \rightarrow L^p(X, \Sigma, m)$ we mean that T is a linear bounded operator which transforms non-negative functions into non-negative functions and its norm is not more than one.

Lemma 1. Let $1 \leq p < +\infty$ and let $T: L^p(X, \Sigma, m) \rightarrow L^p(X, \Sigma, m)$ be a positive contraction. Suppose that there exist $\epsilon > 0$ and $n_0 \in \mathbb{N} \cup \{0\}$ such that $\|T^{n_0+1} - T^{n_0}\|_p^p < 2(1 - \epsilon)$. Let $f \in L^p(X, \Sigma, m)$ be such that for every $n \in \mathbb{N} \cup \{0\}$ $T^n f \cdot T^{n+1} f = 0$. Then $\lim_{n \rightarrow +\infty} \|T^n f\|_p = 0$.

Proof. Clearly it is enough to prove the lemma for $\|f\|_p = 1$.

1. This paper is part of the author's Ph.D. Thesis at the Hebrew University of Jerusalem. I wish to express my deepest gratitude to Professor Harry Furstenberg, my supervisor, for his valuable support and guidance.

If f is as above then

$$\|T^{n_0+1}f - T^{n_0}f\|_p^p = \|T^{n_0+1}f\|_p^p + \|T^{n_0}f\|_p^p < 2(1 - \epsilon) .$$

Using the fact that $(\|T^n f\|_p^p)_n$ is a decreasing sequence we obtain that

$$\|T^{n_0+1}f\|_p < (1 - \epsilon)^{1/p} .$$

It follows that if we note $\rho = (1 - \epsilon)^{1/p}$ then

$$\|(T^{n_0+1} - T^{n_0})T^{n_0+1}f\|_p^p = \|T^{n_0+1}T^{n_0+1}f\|_p^p + \|T^{n_0}T^{n_0+1}f\|_p^p < 2(1 - \epsilon)\rho^p$$

and we obtain that

$$\|T^{2(n_0+1)}f\|_p < \rho^2 .$$

By induction it follows that for every $h \in \mathbb{N}$ $\|T^{h(n_0+1)}f\|_p < \rho^h$ and using the fact that $(\|T^n f\|_p)_n$ is a decreasing sequence it follows that $\lim_n \|T^n f\|_p = 0$.

Remark. In Lemma 1 we may drop the assumption of T being positive. However we will need this assumption later on.

2. The Finite Dimensional L^p -spaces and the Theorem

We will now consider the following case:

Let $k \in \mathbb{N}$, $k \geq 2$ be and we note $X = \{1, 2, \dots, k\}$, $\Sigma = \mathcal{P}(X)$. Let m_1, \dots, m_k be k non-zero positive real numbers. We will denote by m the measure generated by m_1, \dots, m_k (that is $m(\{i\}) = m_i$, $i = 1, \dots, k$). We will call the space $L^p(X, \Sigma, m)$ a finite dimensional L^p -space and we will note

$\mathcal{L}_p(k, m) = L^p(X, \Sigma, m)$. A positive contraction $T: \mathcal{L}_p(k, m) \rightarrow \mathcal{L}_p(k, m)$ is generated by a matrix $(a_{ij})_{i,j=1,2,\dots,k}$ and the resulting positive contraction $T^n (n \in \mathbb{N})$ is generated by $(a_{ij}^{(n)})_{i,j=1,\dots,k}$.

If $x \in \mathcal{L}_p(k, m)$, $x = (x_1, \dots, x_k)$ then $Tx = \left(\sum_{i=1}^k x_i a_{ij} \right)_{j=1,\dots,k}$ and

$$T^n x = \left(\sum_{i=1}^k x_i a_{ij}^{(n)} \right)_{j=1,\dots,k}.$$

If T is a positive contraction on $\mathcal{L}_1(k, m)$ we will note

$$\Omega = \{i \in X \mid \text{for every } n \in \mathbb{N} \cup \{0\} \text{ and } j = 1, \dots, k \quad \sum_{j=1}^k a_{ij}^{(n)} m_j = m_i$$

$$\text{and } a_{i\ell}^{(n)} \cdot a_{i\ell}^{(n+1)} = 0\}.$$

Lemma 2. Let $T: \mathcal{L}_1(k, m) \rightarrow \mathcal{L}_1(k, m)$ be a positive contraction. Then the following are equivalent:

- a) for every $n \in \mathbb{N} \cup \{0\}$, $\|T^{n+1} - T^n\|_1 = 2$
- b) $\Omega \neq \emptyset$.

Proof. a) \Rightarrow b) Suppose $\Omega = \emptyset$. It follows that for every $i \in \{1, 2, \dots, k\}$

there exists $n_i \in \mathbb{N}$ such that $\sum_{j=1}^k a_{ij}^{(n_i)} m_j < m_i$ or there exists j_0 such that $a_{ij_0}^{(n_i)} \cdot a_{ij_0}^{(n_i+1)} \neq 0$. In other words for every $i \in \{1, 2, \dots, k\}$ there exists n_i such that $\|(T^{n_i+1} - T^{n_i})1_{\{i\}}\|_1 < 2m_i = 2\|1_{\{i\}}\|_1$.

If we note $n_0 = \max\{n_1, \dots, n_k\}$ it follows that for every $n \geq n_0$ and for every $i \in \{1, 2, \dots, k\}$ $\|(T^{n+1} - T^n)1_{\{i\}}\|_1 < 2m_i$ (as for every $i \in \{1, 2, \dots, k\}$

the sequence $(\| (T^{n+1} - T^n) 1_{\{i\}} \|_1)_n$ is a decreasing one).

It follows that for $n \geq n_0$ $\|T^{n+1} - T^n\|_1 < 2$.

b) \Rightarrow a) If $\Omega \neq \emptyset$ then for every $n \in \mathbb{N} \cup \{0\}$ and $i \in \Omega$ $\|T^{n+1} 1_{\{i\}} - T^n 1_{\{i\}}\|_1 = 2\|1_{\{i\}}\|_1$ and as T is a positive contraction it follows that for every $n \in \mathbb{N} \cup \{0\}$ $\|T^{n+1} - T^n\|_1 = 2$.

Now we are able to prove the desired result:

Theorem 3. Let p be such that $1 \leq p < +\infty$ and let T be simultaneously a positive contraction of $L^1(k,m)$ and $L^p(k,m)$. If there exists $n_0 \in \mathbb{N} \cup \{0\}$ such that $\|T^{n_0+1} - T^{n_0}\|_p < 2^{1/p}$ then $\lim_n \|T^{n+1} - T^n\|_p = 0$.

Proof. If $\lim_n \|T^{n+1} - T^n\|_p \neq 0$ then $\lim_n \|T^{n+1} - T^n\|_1 \neq 0$ (as every two norms in R^{k^2} are equivalent). Using the zero-two law for positive contractions in L^1 -spaces (Theorem 1.1 from [2]) it follows that for every $n \in \mathbb{N} \cup \{0\}$ $\|T^{n+1} - T^n\|_1 = 2$ and by Lemma 2 it follows that $\Omega \neq \emptyset$. If $i \in \Omega$ then the characteristic function $1_{\{i\}}$ satisfies the conditions of Lemma 1 and it follows that $\lim_n \|T^{n+1} 1_{\{i\}} - T^n 1_{\{i\}}\|_p = 0$. We obtain that $\lim_n \|T^{n+1} 1_{\{i\}} - T^n 1_{\{i\}}\|_1 = 0$ which contradicts the fact that $i \in \Omega$.

References

1. Neveu, J.: "Mathematical foundations of the calculus of probability", San Francisco, London, Amsterdam: Holden Day 1965.
2. Ornstein, D. and Sucheston, L.: "An operator Theorem on L_1 convergence to zero with applications to Markov kernels". Ann. Math. Statist. 1970 vol. 41, no. 5, 1631-1639.

R. ZAHAROPOL
Département de Mathématiques
Université Hébraïque

GIVAT-RAM
Jérusalem
Israël

Reçu en Septembre 1983