

ANNALI DELLA SCUOLA NORMALE SUPERIORE DI PISA *Classe di Scienze*

D. R. ADAMS

Traces of potentials arising from translation invariant operators

Annali della Scuola Normale Superiore di Pisa, Classe di Scienze 3^e série, tome 25, n° 1 (1971), p. 203-217

http://www.numdam.org/item?id=ASNSP_1971_3_25_1_203_0

© Scuola Normale Superiore, Pisa, 1971, tous droits réservés.

L'accès aux archives de la revue « Annali della Scuola Normale Superiore di Pisa, Classe di Scienze » (<http://www.sns.it/it/edizioni/riviste/annaliscienze/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

TRACES OF POTENTIALS ARISING FROM TRANSLATION INVARIANT OPERATORS

by D. R. ADAMS

For a function u in the space $W^{\alpha,p}(E_n)$ (the usual Sobolev space of L_p functions on Euclidean n -space E_n with distribution derivatives of orders $\leq \alpha$ in L_p), it is possible to characterize the «restriction» or trace of u (call it u^*) to certain lower dimensional manifolds M provided their dimension d , satisfies $d > n - \alpha p$.

In this paper the characterization is given in terms of the Lebesgue or L_p class of u^* , where the norm of u^* is taken with respect to an appropriate measure μ concentrated on M . In particular, it is known that if $n - d < \alpha p < n$ and $p > 1$, then $u^* \in L_r(M)$, $1 \leq r \leq dp/(n - \alpha p)$, when M is «smooth» and μ is the surface area measure on M . The usual procedure for proving this is to first obtain the result for a subset of a d -dimensional hyperplane in E_n and then extend via a change of variables to manifolds which are diffeomorphic images of d -dimensional coordinate patches. In such a method, the essence is to work coordinate wise, from E_n down to the hyperplane. This paper presents a new method for achieving this, which in addition allows an extension of the trace result to sets M of fractional Hausdorff dimension d , $0 < d \leq n$.

Since every $u \in W^{\alpha,p}$ can be represented as a Bessel potential of an L_p function (see [1]), we will consider functions u in the form of potentials $T(f)$, where $f \in L_p$ and $T \in S_\alpha$, S_α being the class of translation invariant operators of smoothness α , $\alpha > 0$ (see section 1 for the definitions). Theorem 1 then states that for each class S_α there is a corresponding class of «appropriate» measures $\mu(\mathcal{L}_{1,d}^+)$ for which $T(f) \in L_{p^*}(\mu)$, $p^* = dp/(n - \alpha p)$,

Pervenuto alla Redazione il 9 Giugno 1970.

This paper was written while the Author was visiting the Istituto per le Applicazioni del Calcolo of Rome, supported by a C.N.R. fellowship.

$n - d < \alpha p < n$. Here the set M is the support of μ . When T is the Riesz potential operator, Theorem 1 can be improved (Theorem 2). In this case, the condition $\mu \in \mathcal{L}_{1;d}^+$ is both necessary and sufficient for the map $T: L_p \rightarrow L_{p^*}(\mu)$ to be continuous.

The class of measures $\mathcal{L}_{1;d}^+$ is closely related to the Hausdorff d -dimensional measure H_d by a well known theorem of O. Frostman (see [3]). In particular if $H_d(M) > 0$, then there is a measure μ concentrated on M such that $\mu \in \mathcal{L}_{1;d}^+$ and $\mu \not\equiv 0$.

Theorem 1 can also be viewed another way: if μ_0 is given then the condition — μ_0 restricted to M belongs to $\mathcal{L}_{1;d}^+$ — will be a sufficient condition on M to insure that $T(f)$ has a $L_{p^*}(\mu_0)$ trace on M . For example, if $\mu_0 = H_d$, d a positive integer, then any smooth compact manifold in E_n satisfies this condition.

I am grateful to G. Stampacchia for pointing out the result appearing in the appendix of [9]. It is the forerunner of lemmas 1 and 2.

Section 1 contains the preliminaries and section 2, the statements and proofs of the main results. In section 3, I have attempted to point out relationships between Theorem 1 and various results appearing in the literature.

1. Preliminaries.

1.1. Let $\mathcal{A}_\gamma$ denote the usual Banach space of all bounded Hölder continuous functions of exponent $\gamma > 0$ defined on E_n (see for example [8] for a precise definition). A linear transformation T which maps $\mathcal{A}_\gamma$ into $\mathcal{A}_{\gamma+\alpha}$, $\alpha > 0$, boundedly and which commutes with translations will be termed a linear translation invariant operator of smoothness α and the class of all such T will be denoted by S_α . Here we will be content to list the various properties of $T \in S_\alpha$ needed for this paper.

If $T \in S_\alpha$, then it is known (see [8]) that for $0 < \alpha < 1$, T applied to any smooth function f is given by $T(f)(x) = \int k(x-y)f(y)dy$ where k , the kernel of T , satisfies

$$(1) \quad \int |k(x)| dx < \infty,$$

$$(2) \quad \int |k(x-y) - k(x)| dx \leq Q|y|^\alpha,$$

Q a constant independent of y . Here the symbol $\int \dots dx$ denotes integration over E_n with respect to n -dimensional Lebesgue measure m_n .

Our main interest lies with the Riesz potential operator whose kernel is $h^\alpha(x) = |x|^{\alpha-n}$ and the Bessel potential operator J^α . In general, J^α is defined for all real α by: J^α is the mapping given by the convolution with a tempered distribution and whose Fourier Transform is $(2\pi)^{-n/2} \cdot (1 + |x|^2)^{-\alpha/2}$. When $\alpha > 0$, the kernel of J^α is denoted by g_α and satisfies, in addition to (1) and (2)

$$(3) \quad 0 < g_\alpha(x) \leq Q h_\alpha(x), \quad \text{for all } x \in E_n.$$

Here Q is a constant independent of x . For additional properties of g_α see [1].

A basic feature of the map J^β is the fact that it is a bicontinuous isomorphism of $\mathcal{A}_\gamma$ to $\mathcal{A}_{\gamma+\beta}$ as long as $\gamma + \beta > 0$. From this, it easily follows that for any $T \in \mathcal{S}_\alpha$, $T = J^\beta T J^{-\beta}$, i. e. T commutes with J^β .

1.2. By $\mathcal{M}$ we will understand the collection of all completions of Borel measures on E_n and by $\mathcal{L}_1$ those $\mu \in \mathcal{M}$ for which $\|\mu\|_1 = \text{total variation of } \mu < \infty$. We will use the Morrey space notation $\mathcal{L}_{1;d}$ to denote those $\mu \in \mathcal{M}$ for which

$$(4) \quad V_x(\mu, r) = |\mu|(\{y : |x - y| < r\}) \leq A r^d,$$

for all $x \in E_n$ and all $r \geq 0$. Here $0 < d \leq n$ and $|\mu|$ denotes the sum of the positive and negative parts of μ . A is a constant independent of x and r .

The notation $\|\cdot\|_p$ will represent the usual Lebesgue p -norm, $1 \leq p \leq \infty$, with respect to m_n . For any other measure $\mu \in \mathcal{M}^+$, the symbol $\|\cdot\|_{p,\mu}$ will be used. L_p and $L_p(\mu)$ will denote the corresponding Lebesgue function spaces. $\mathcal{M}_0$ denotes the measures with compact support.

The superscript « $+$ » is used to indicate the subclass of non-negative elements. The letter Q will denote various constants, possibly not the same constant in any one proof, whereas A, A_1 , etc. will denote specific constants.

2. The main results.

2.1. The results of principal interest are Theorems 1 and 2 below

THEOREM 1: For $T \in \mathcal{S}_\alpha$ and $\mu \in \mathcal{L}_{1;d}^+$, there exists a constant Q such that for all f in L_p ,

$$\|T(f)\|_{p^*,\mu} \leq Q \|f\|_p$$

provided $n - d < \alpha p < n$, $0 < d \leq n$, $p > 1$. Here $p^* = dp/(n - \alpha p)$ and Q is independent of f .

REMARK 1: Special cases of Theorem 1 are known, e. g. when $\mu = m_n$, it is the Theorem of Stein-Zygmund (see Section 2.4); when $\mu = m_d$, d an integer, and T the Riesz potential operator, it becomes the imbedding result of II' in [5].

The program for proving Theorem 1 will be: (a) to establish necessary and sufficient conditions on a non-negative Borel measure μ in order that the above inequality holds for the Riesz potential operator, (b) to show then that Theorem 1 holds for the Bessel potential operator (using (3)), and finally (c), to establish Theorem 1 for general T (using (b) and the Theorem of Stein-Zygmund). Thus the main burden of the proof of Theorem 1 is in establishing (a). This can be stated as follows:

THEOREM 2: The necessary and sufficient condition for

$$\|h_\alpha * f\|_{p^*, \mu} \leq Q \|f\|_p$$

to hold for all $f \in L_p$, Q a constant independent of f , with $\mu \in \mathcal{M}^+$ ($p^* = dp/n - \alpha p$, $0 < d \leq n$, $n - d < \alpha p < n$, $p > 1$) is that $\mu \in \mathcal{L}_{1;d}^+$.

2.2. The proof of the sufficiency for Theorem 2 involves an estimate on the number $\|h_\alpha * \mu^K\|_{p'}$, where $p' = p/(p-1)$ and μ^K denotes μ restricted to K , K a compact set in E_n of positive μ measure. To obtain the desired estimate, two main cases are considered, namely $1 < p \leq 2$ and $p > 2$. In the first case, $h_{\alpha p} * \mu^K$ is estimated in the L_∞ norm and then $h_\alpha * \mu^K$ in the $L_{p'}$ norm (lemmas 1 and 2).

For the second case we note that

$$(5) \quad \|h_\alpha * \mu^K\|_{p'}^{p'} = \int h_\alpha * (h_\alpha * \mu^K)^{1/(p-1)} d\mu^K = \int u^K d\mu^K$$

where $u^K(x) = h_\alpha * f^K(x)$, $f^K(y) = [h_\alpha * \mu^K(y)]^{1/(p-1)}$. Hence it suffices to estimate u^K in the L_∞ norm. To do this, observe that

$$(6) \quad u^K(x) = \int h_\alpha(x-y) f^K(y) dy = \int_0^\infty h_\alpha(r) dV_x(f^K, r).$$

Here $V_x(f^K, r)$ is given by (4), for a measure with density f^K . In lemmas 3-5, estimates for the functions $f^K(y)$ and $V_x(f^K, r)$ are obtained. Finally, lemma 6 is the desired estimate on u^K .

LEMMA 1 : $h_{\alpha p} * \mu^K(x) \leq A_1 \mu(K)^{(\alpha p - n + d)/d}$, for all $x \in E_n$, $A_1 = 1 + A(n - \alpha p)/(\alpha p - n + d)$.

PROOF : $h_{\alpha p} * \mu^K(x) = \int_0^\infty h_{\alpha p}(r) dV_x(\mu^K, r)$ and altho the function $V_x(\mu^K, r)$ is not in general continuous in r (for each fixed x), it is non decreasing and left continuous. This formula follows from the definitions of the integrals involved.

Integrating by parts we get

$$\begin{aligned} \int_0^\infty h_{\alpha p}(r) dV_x(\mu^K, r) &= - \int_0^\infty V_x(\mu^K, r) dh_{\alpha p}(r) \\ &= (n - \alpha p) \int_0^\infty V_x(\mu^K, r) r^{\alpha p - n - 1} dr, \end{aligned}$$

since as $r \rightarrow 0$, $h_{\alpha p}(r) \cdot V_x(\mu^K, r) \leq A r^{\alpha p - n + d}$, $\alpha p - n + d > 0$ whereas $h_{\alpha p}(r) \cdot V_x(\mu^K, r) \leq \mu(K) r^{\alpha p - n}$, as $r \rightarrow \infty$, $\alpha p - n < 0$. Thus

$$h_{\alpha p} * \mu^K(x) \leq (n - \alpha p) \left(\int_0^\sigma + \int_\sigma^\infty \right) V_x(\mu^K, r) \cdot r^{\alpha p - n - 1} dr = (n - \alpha p) (I_1 + I_2).$$

$$I_1 \leq A \int_0^\sigma r^{d + \alpha p - n - 1} dr = \frac{A}{\alpha p - n + d} \cdot \sigma^{\alpha p - n + d},$$

$$I_2 \leq \mu(K) \int_\sigma^\infty r^{\alpha p - n - 1} dr = \frac{1}{n - \alpha p} \cdot \mu(K) \cdot \sigma^{\alpha p - n}.$$

The result now follows by choosing $\sigma = \mu(K)^{1/d}$.

LEMMA 2 : For $1 < p \leq 2$,

$$\|h_\alpha * \mu^K\|_{p'} \leq A_2 \cdot \mu(K)^{(\alpha p - n + dp)/dp},$$

where $A_2 = C\left(\frac{\alpha p}{2}, \frac{\alpha p}{2}\right)^{1/p'} \cdot A_1^{1/p}$; $C(\alpha, \beta)$ is the Riesz convolution constant, i. e. $h_\alpha * h_\beta = C(\alpha, \beta) \cdot h_{\alpha + \beta}$ for $\alpha + \beta < n$.

PROOF: For $1 < p < 2$ (the case $p = 2$ can be handled by a trivial modification) choose $\theta : 0 < \theta < 1$ and $1 = \theta p/2 + (1 - \theta)p$, then

$$h_\alpha(x) = h_{\alpha p/2}(x)^\theta h_{\alpha p}(x)^{1-\theta}.$$

Thus from Hölder's inequality, we have

$$h_\alpha * \mu^K(y) \leq [h_{\alpha p/2} * \mu^K(y)]^\theta \cdot [h_{\alpha p} * \mu^K(y)]^{1-\theta},$$

and

$$(7) \quad \|h_\alpha * \mu^K\|_{p'} \leq \|h_{\alpha p} * \mu^K\|_\infty^{1-2/p'} \cdot \|h_{\alpha p/2} * \mu^K\|_2^{2/p'}$$

by the choice of θ , i. e. $\theta p' = 2$. But

$$(8) \quad \begin{aligned} \|h_{\alpha p/2} * \mu^K\|_2^2 &= C \left(\frac{\alpha p}{2}, \frac{\alpha p}{2} \right) \int h_{\alpha p} * \mu^K d\mu^K \\ &\leq C \left(\frac{\alpha p}{2}, \frac{\alpha p}{2} \right) \|h_{\alpha p} * \mu^K\|_\infty \cdot \mu(K). \end{aligned}$$

(7) and (8), together with lemma 1, now give the desired result.

LEMMA 3 : For $p > 2$,

$$V_x(f^K, r) \leq A_3 \mu(K)^{1/(p-1)} \cdot r^{n-(n-\alpha)/(p-1)}$$

for all $x \in E_n$ and all $r \geq 0$. $A_3 = \omega_n (1 + 3^\alpha/\alpha)^{1/(p-1)}$, ω_n = area of the unit sphere in E_n .

PROOF: Since $p > 2$, Hölder's inequality gives

$$(9) \quad \begin{aligned} V_x(f^K, r) &\leq (\omega_n r^n)^{1-1/(p-1)} \cdot \{V_x(h_\alpha * \mu^K, r)\}^{1/(p-1)} \\ &= (\omega_n r^n)^{1-1/(p-1)} \cdot \{I_3 + I_4\}^{1/(p-1)}, \end{aligned}$$

where

$$I_3 = \int_{K \cap \{|x-z| > 2r\}} d\mu(z) \int_{|x-y| < r} h_\alpha(y-z) dy$$

which never exceeds

$$\int_{K \cap \{|x-z| > 2r\}} d\mu(z) \int_{|x-y| < r} h_\alpha(r) dy$$

since $|y-z| \geq r$. Thus $I_3 \leq r^{\alpha-n} \cdot \omega_n r^n \cdot \mu(K)$.

And

$$\begin{aligned} I_4 &= \int_{K \cap [|x-z| \leq 2r]} d\mu(z) \int_{|x-y| < r} h_\alpha(y-z) dy \\ &\leq \int_{K \cap [|x-z| \leq 2r]} d\mu(z) \int_{|y-z| \leq 3r} h_\alpha(y-z) dy, \end{aligned}$$

Since now $|y-z| \leq 3r$. Thus $I_4 \leq \omega_n \mu(K) \cdot (3r)^\alpha / \alpha$.

LEMMA 4: For $p > 2$ and $0 < \alpha < n-d$,

$$V_x(f^K, r) \leq A_4 r^{n-(n-\alpha-d)/(p-1)}$$

for all $x \in E_n$ and all $r \geq 0$. $A_4 = \omega_n A^{1/(p-1)} \cdot \left(\frac{n-\alpha}{n-\alpha-d} + \frac{2^d 3^\alpha}{\alpha} \right)^{1/(p-1)}$.

PROOF: Equivalently this lemma asserts that f^K belongs to the Morrey class $\mathcal{L}_{1,b}$, with $b = n - (n - \alpha - d)/(p - 1)$ when $\alpha < n - d$ (compare this to lemma 5).

For fixed x , we consider y such that $|x-y| < r$, then

$$h_\alpha * \mu^K(y) \leq \left(\int_{|x-z| \geq 2r} + \int_{|x-z| < 2r} \right) h_\alpha(y-z) d\mu(z) = I_5 + I_6.$$

Let

$$\varphi_y(\varrho) = \int_{[|x-z| \geq 2r] \cap [|y-z| < \varrho]} d\mu(z)$$

and note :

- (i) $\varphi_y(\varrho) = 0$, when $0 \leq \varrho \leq r$;
- (ii) $\varphi_y(\varrho) \leq V_y(\mu, \varrho) \leq A\varrho^d$, for all $y \in E_n$ and $\varrho \geq 0$;
- (iii) For each fixed y , $\varphi_y(\varrho)$ is non-decreasing in ϱ and left continuous.

$$I_5 = \int_r^\infty h_\alpha(\varrho) d\varphi_y(\varrho) = (n-\alpha) \int_r^\infty \varphi_y(\varrho) \varrho^{\alpha-n-1} d\varrho$$

using (i) and then integrating by parts.

Also note that $h_\alpha(\varrho) \varphi_y(\varrho) \rightarrow 0$, as $\varrho \rightarrow \infty$ by (ii). Thus

$$I_b \leq (n - \alpha) A \int_r^\infty \varrho^{\alpha-n+d-1} d\varrho = \frac{(n - \alpha)}{(n - \alpha - d)} A r^{\alpha-n+d}.$$

$$\int_{|x-y| < r} I_6 dy \leq \int_{|x-z| < 2r} d\mu(z) \int_{|y-z| < 3r} h_\alpha(y-z) dy = \omega_n (3r)^{\alpha/\alpha} \cdot V_x(\mu, 2r).$$

With these estimates and (9) of lemma 3, the result follows.

LEMMA 5: For $p > 2$ and $0 \leq n - d < \alpha$,

$$f^K(y) \leq A_5 \cdot \mu(K)^{(d+\alpha-n)/d(p-1)}$$

for all $y \in E_n$. $A_5 = [A(1 + (n - \alpha)/(\alpha - n + d)) + 1]^{1/(p-1)}$.

PROOF: In contrast to lemma 4, f^K is no longer in a Morrey class, but in a Hölder class with exponent $(d + \alpha - n)/(p - 1)$.

$$\begin{aligned} [f^K(y)]^{p-1} &= \int h_\alpha(y-z) d\mu^K(z) \\ &= \left(\int_{|y-z| > \sigma} + \int_{|y-z| \leq \sigma} \right) h_\alpha(y-z) d\mu^K(z) = I_7 + I_8. \end{aligned}$$

Again with $\sigma = \mu(K)^{1/d}$,

$$I_7 \leq \sigma^{\alpha-n} \mu(K),$$

$$\begin{aligned} I_8 &\leq \int_0^\sigma h_\alpha(\varrho) dV_y(\mu, \varrho) \\ &\leq h_\alpha(\sigma) V_y(\mu, \sigma) + (n - \alpha) \int_0^\sigma V_y(\mu, \varrho) \varrho^{\alpha-n-1} d\varrho \\ &\leq \left(A + \frac{(n - \alpha)}{(\alpha - n + d)} A \right) \sigma^{\alpha-n+d}, \end{aligned}$$

the result now follows easily.

LEMMA 6 : For $p > 2$, there is a constant A_6 independent of the set K such that

$$u^K(x) \leq A_6 \mu(K)^{(\alpha p - n + d)/d(p-1)}.$$

Hence by (5), $\|h_a * \mu^K\|_{p'} \leq A_6^{1/p'} \mu(K)^{(\alpha p - n + dp)/dp}$.

PROOF : case (1) $0 < \alpha < n - d$: Integrating by parts in (6),

$$u^K(x) = (n - \alpha) \int_0^\infty V_x(f^K, r) r^{\alpha - n - 1} dr$$

since by lemma 4, $V_x(f^K, r) \cdot h_a(r)$ is $O(r^{(\alpha p - n + d)/(p-1)})$ as $r \rightarrow 0$, and is $O(r^{(\alpha p - n)/(p-1)})$ as $r \rightarrow \infty$, by lemma 3. Thus

$$u^K(x) = (n - \alpha) \left(\int_0^\sigma + \int_\sigma^\infty \right) V_x(f^K, r) r^{\alpha - n - 1} dr = (n - \alpha) (I_9 + I_{10}).$$

Applying lemma 4 to I_9 and lemma 3 to I_{10} , we have

$$I_9 \leq \frac{A_4(p-1)}{(\alpha p - n + d)} \cdot \sigma^{(\alpha p - n + d)/(p-1)},$$

and

$$I_{10} \leq \frac{A_3(p-1)}{(n - \alpha p)} \cdot \mu(K)^{1/(p-1)} \cdot \sigma^{(\alpha p - n)/(p-1)}.$$

The result follows taking $\sigma = \mu(K)^{1/d}$.

case (2) $0 \leq n - d < \alpha$:

$$u^K(x) = \left(\int_{|x-y| \leq \sigma} + \int_{|x-y| > \sigma} \right) h_a(x-y) f^K(y) dy = I_{11} + I_{12}.$$

Applying lemma 5 to I_{11} and lemma 3 to I_{12} , we have

$$I_{11} \leq \frac{A_5 \omega_n}{\alpha} \mu(K)^{(d + \alpha - n)/d(p-1)} \cdot \sigma^\alpha,$$

$$I_{12} \leq \frac{(n - \alpha)(p-1)}{(n - \alpha p)} A_3 \mu(K)^{1/(p-1)} \cdot \sigma^{(\alpha p - n)/(p-1)},$$

with the same choice of σ .

case (3) $0 < \alpha = n - d$: This case is resolved by interpolating between cases (1) and (2) as follows: choose pairs (α_i, p) with $n - d < \alpha_i p < n$, $i = 0, 1$ but $0 < \alpha_0 < n - d < \alpha_1 < n$. Let $\alpha = \theta \alpha_0 + (1 - \theta) \alpha_1$, $0 < \theta < 1$.

As before $h_\alpha(x) = h_{\alpha_0}(x)^\theta \cdot h_{\alpha_1}(x)^{1-\theta}$ and upon applying Hölder's inequality, we have

$$f^K(y) \leq [f_0^K(y)]^\theta \cdot [f_1^K(y)]^{1-\theta}$$

where $f_i^K(y) = [h_{\alpha_i} * \mu^K(y)]^{1/(p-1)}$, $i = 0, 1$. Thus

$$u^K(x) \leq [h_{\alpha_0} * f_0^K(x)]^\theta \cdot [h_{\alpha_1} * f_1^K(x)]^{1-\theta} = [u_0^K(x)]^\theta \cdot [u_1^K(x)]^{1-\theta}.$$

Case (1) gives $u_0^K(x) \leq A_6' \mu(K)^{(\alpha_0 p - n + d)/d(p-1)}$ and case (2) gives $u_1^K(x) \leq A_6'' \mu(K)^{(\alpha_1 p - n + d)/d(p-1)}$. Hence it is now clear that a finite constant A_6 may be chosen with the required properties.

2.3. PROOF OF THEOREM 2: For the sufficiency, lemmas 2 and 6 are used to show that for fixed α , the Riesz potential operator is of weak type $(L_p, L_{p^*}(\mu))$, when $n - d < \alpha p < n$.

Let $E_t = \{x : |h_\alpha * f(x)| > t\}$, $t > 0$ and $f \in L_p$.

$$t \mu(E_t) \leq \int h_\alpha * |f|(x) d\mu^{E_t}(x) = \int h_\alpha * \mu^{E_t}(x) |f(x)| dx \leq \|h_\alpha * \mu^{E_t}\|_{p'} \|f\|_p.$$

Now since lemmas 2 and 6 hold for all compact sets K , and the constants A_2 and A_6 are independent to K , these estimates must also hold for K replaced by E_t , a G_δ -set, since μ is a Borel measure. Thus

$$t \mu(E_t) \leq Q' \|f\|_p \mu(E_t)^{(\alpha p - n + dp)/dp}$$

or

$$\mu(E_t) \leq \left(\frac{Q' \|f\|_p}{t} \right)^{p^*}, \quad p^* = dp/(n - \alpha p).$$

We now apply the well known interpolation theorem of Marcinkiewicz to deduce the strong type estimate required.

To prove the necessity, we choose a particular L_p function, namely the characteristic function of the ball $B_r(x_0) = \{x : |x - x_0| < r\}$, $r > 0$ and $x_0 \in E_n$ arbitrary. Denote this function by $\chi_r(x)$. $\|\chi_r\|_p = (\omega_n r^n)^{1/p}$. On the other hand

$$\begin{aligned} \|h_\alpha * \chi_r\|_{p^*, \mu} &\geq \left\{ \int_{|x-x_0| < r} \left(\int_{|y-x_0| < r} h_\alpha(x-y) dy \right)^{p^*} d\mu(x) \right\}^{1/p^*} \\ &\geq h_\alpha(2r) \omega_n r^n [V_{x_0}(\mu, r)]^{1/p^*}. \end{aligned}$$

Thus if $h_a : L_p \rightarrow L_{p^*}(\mu)$ is continuous, we immediately get

$$V_{x_0}(\mu, r) \leq A' r^d.$$

The proof of Theorem 2 is now complete.

REMARK 2: It is interesting to note that the region in the (p, α) plane $1 < p < \infty$, $0 < \alpha < n$, for which the above result holds is the region between the two hyperbolas $\alpha p = n - d$ and $\alpha p = n$. It is possible to «shift» this region to obtain a result of additional interest (see Remark 6).

Making the changes: $\mu \rightarrow \mu_2$, $m_n \rightarrow \mu_1$ and $h_a \rightarrow h_{a+d_1}$, where $0 \leq d_1 < d_2 \leq n$, $\mu_1 \in \mathcal{L}_{1; n-d_1}^+$ and $\mu_2 \in \mathcal{L}_{1; d_2-d_1}^+$, we easily get

$$(10) \quad \|h_{a+d_1} * f \mu_1\|_{p^*, \mu_2} \leq Q \|f\|_{p, \mu_1}$$

where $p^* = (d_2 - d_1)p / (n - \alpha p - d_1)$, $n - d_2 < \alpha p < n - d_1$, $p > 1$. Here $f \mu_1$ denotes a measure with density f .

2.4. To see that Theorem 1 holds for h_a replaced by g_a , it is only necessary to combine (3) with Theorem 2.

Necessary and sufficient conditions on μ for g_a are possible only if the variation of μ is allowed to grow more rapidly at infinity.

PROOF OF THEOREM 1: This extension of Theorem 2 can now be established by applying the theorem of Stein-Zygmund [8]. This result may be stated as follows:

THEOREM: If $T \in S_a$, then there is a constant Q such that

$$\|T(f)\|_q \leq Q \|f\|_p, \quad q = np / (n - \alpha p)$$

for all $f \in L_p$; Q is independent of f . $1 < p < \infty$, $0 < \alpha p < n$.

For $T \in S_a$, choose β satisfying $\frac{(n-d)(n-\alpha p)}{dp} < \beta < \alpha$. Note that this is always possible since $\alpha p > n - d$. From 1.1, $T \cdot J^{-\beta} \in S_{a-\beta}$, thus the theorem of Stein-Zygmund yields

$$\|T \cdot J^{-\beta}(f)\|_q \leq Q_1 \|f\|_p$$

where $q = np / [n - (\alpha - \beta)p]$. Using Theorem 2 (which we now know is true for the Bessel potential operators),

$$\|T(f)\|_{q^*, \mu} = \|J^\beta(T \cdot J^{-\beta}(f))\|_{q^*, \mu} \leq Q_2 \|T \cdot J^{-\beta}(f)\|_q$$

where $q^* = dq/(n - \beta q) = dp/(n - \alpha p) = p^*$. Hence

$$\|T(f)\|_{p^*, \mu} \leq Q_2 \cdot Q_1 \|f\|_p.$$

Note that the conditions $1 < q < q^* < \infty$ are satisfied when $1 < p < \infty$, $n - d < \alpha p < n$, and by the choice of β .

REMARK 3: It might be noted that if the more general interpolation theorem of R. Hunt [4] had been used in place of the theorem of Marcinkiewicz, it would be possible to deduce that any $T \in S_\alpha$ maps the Lorentz space $L(p, q)$ continuously into $L(p^*, s)(\mu)$, with $q \leq s$, the usual restrictions on α , p and d . In particular when $q = s = \infty$, T maps weak- L_p continuously into weak- $L_{p^*}(\mu)$.

3. Related results.

3.1. We begin by giving potential versions of two classical trace theorems.

THEOREM 3: Let $T \in S_\alpha$ and $\mu \in \mathcal{L}_1^+; d$, then there is a constant Q such that for all $f \in L_p$

$$\|\Delta_t T(f)\|_{r, \mu} \leq Q |t|^{a-n/p+d/r} \|f\|_p$$

where $\max\{dp/[n - (\alpha - 1)p], p\} < r < p^*$, $n - d < \alpha p < n$. Here Δ_t denotes the first difference; Q is to be independent of f and t .

PROOF: The restrictions on r insure that the exponent of $|t|$ is always positive and less than 1. It is easy to see that $\Delta_t T(f)(x) = (\Delta_t k) * f(x)$, where k is the kernel of T .

We write $T = (J^{-\theta} \cdot T) \cdot J^\theta = k_{\alpha-\theta} * g_\theta$ where $k_{\alpha-\theta}$ is the kernel of $J^{-\theta}$. T , θ chosen to satisfy initially $(n - d)/p < \theta < \alpha$. Then

$$\Delta_t T(f)(x) = \int \Delta_t k_{\alpha-\theta}(z) \cdot g_\theta * f_z(x) dz$$

since $k = k_{\alpha-\theta} * g_\theta$. Here $f_z(y)$ denotes $f(y - z)$. By the inequality of Minkowski and Theorem 1, we have

$$\begin{aligned} \|\Delta_t T(f)\|_{r, \mu} &\leq \int \|\Delta_t k_{\alpha-\theta}(z) \cdot \|g_\theta * f_z\|_{r, \mu} dz \\ &\leq \int \|\Delta_t k_{\alpha-\theta}(z) \cdot Q \|f_z\|_p dz \end{aligned}$$

where $r = dp/(n - \theta p)$. Since $\|f_z\|_p = \|f\|_p$, we have, using (2)

$$\| \Delta_t T(f) \|_{r, \mu} \leq Q' \|f\|_p |t|^{\alpha - \theta}, \quad 0 < \alpha - \theta < 1.$$

But $\theta = n/p - d/r$, hence the theorem follows.

THEOREM 4: For $T \in S_\alpha$, $f \rightarrow T(f)$ is a compact mapping of L_p into $L_r(\mu)$ for any $\mu \in \mathcal{M}_1^+ \cap \mathcal{L}_{1; d}^+$, $1 \leq r < p^*$, $n - d < \alpha p < n$.

PROOF: Let $\{f_k\}$ be a bounded sequence in L_p , then there exists a weakly convergent subsequence $f'_k \rightarrow f, f \in L_p$. By (1) and (2) $k_{\alpha - \theta} * f'_k \rightarrow k_{\alpha - \theta} * f$ strongly in L_p locally, by the familiar Riesz compactness criterion. But since $\mu \in \mathcal{M}_0$, $T(f'_k) \rightarrow T(f)$ in μ measure and thus using Theorem 1 the result follows by a standard argument.

3.2. We now consider a « dual » to Theorem 1 and then apply it to obtain an extension of a theorem of Campanato [2].

THEOREM 5: Let $T \in S_\alpha$ and $\mu \in \mathcal{L}_{1; d}^+$, then there is a constant Q such that for all $g \in L_q(\mu)$,

$$\|T(g\mu)\|_{\bar{q}} \leq \|g\|_{q, \mu} Q$$

where $\bar{q} = nq/[d + q(n - \alpha - d)]$ and $1 < q < \bar{q} < \infty$. Here Q is independent of g .

PROOF: Let $g \in L_{p^{**}}(\mu)$ and $f \in L_p$, $p^* = dp/(n - \alpha p)$, then

$$\int T(g\mu)f dx = \int T(f)g d\mu \leq \|T(f)\|_{p^*, \mu} \|g\|_{p^{**}, \mu} \leq Q \|f\|_p \|g\|_{p^{**}, \mu},$$

the last inequality following from Theorem 1. The result now follows by taking $q = p^{**}$ and $\bar{q} = p'$.

Let $\mathcal{L}_{t; n-\lambda}$, $1 \leq t < \infty$, $0 < \lambda < n$, denote the class of measures $\mu \in \mathcal{L}_{1; n-\lambda}$ which are absolutely continuous with respect to m_n with density f satisfying $|f|^t \cdot m_n \in \mathcal{L}_{1; n-\lambda}^+$.

THEOREM 6: If $\mu \in \mathcal{L}_{1; n-\lambda}$, then $h_\alpha * \mu \in \mathcal{L}_{t; n-t(\lambda-\alpha)}$ where $1 \leq t < \lambda/(\lambda - \alpha)$, $0 < \alpha < \lambda$.

PROOF: In Theorem 5, take $g(x) = \chi_{2r}(x)$, the characteristic function of the ball $B_{2r}(x_0)$, and $t = \bar{q}$, $\bar{q}/q = (n - t(\lambda - \alpha))/(n - \lambda)$, $d = n - \lambda$, then

$$\int_{|x-x_0| < r} |h_\alpha * \mu_{2r}|^t dx \leq Q' r^{n-t(\lambda-\alpha)}$$

where μ_{2r} is μ restricted to $B_{2r}(x_0)$. The condition $1 < q < \bar{q} < \infty$ is equivalent to $n/(n - \alpha) < t < \lambda/(\lambda - \alpha)$.

It now remains to estimate the t -power of the variation over $B_r(x_0)$ of $h_\alpha * (\mu - \mu_{2r})$. However, this quantity is just the variation of $|I_5|^t$ over $B_r(x_0)$ (I_5 as in lemma 4) with x_0 playing the role of x and $\alpha < \lambda$.

Next, if $1 \leq t < n/(n - \alpha)$, then

$$\begin{aligned} \left\{ \int_{|x-x_0| < r} |h_\alpha * \mu_{2r}|^t dx \right\}^{1/t} &\leq \int_{|y-x_0| < 2r} \left\{ \int_{|x-x_0| < r} h_\alpha(x-y)^t dx \right\}^{1/t} d|\mu|(y) \\ &\leq Q r^{[n-t(\lambda-\alpha)]/t}. \end{aligned}$$

The remaining integral is handled as before.

Finally, for any t in the interval $[1, \lambda/(\lambda - \alpha)]$, a simple interpolation argument gives the result.

REMARK 4: The result of Campanato [2] can be stated as follows: If $f \in \mathcal{L}_{p; n-\lambda}$, then $h_\alpha * f \in \mathcal{L}_{t; n-\sigma}$ where $\alpha p < \lambda$, $1 \leq t < \lambda p/(\lambda - \alpha p)$, and $\lambda > \sigma > t(\lambda - \alpha p)/p$, $1 < p < \infty$.

His proof fails when $p = 1$, which Theorem 6 now treats. Note, when $p = 1$, it is possible to take $\sigma = t(\lambda - \alpha)$.

REMARK 5: It might be interesting to find conditions on μ for which $t = \lambda/(\lambda - \alpha)$ is allowed in Theorem 6, for in general, it is known that $\mu \in \mathcal{L}_{1; n-\lambda}$ is *not* sufficient. Indeed, if $\lambda = \alpha p$, then $\lambda/(\lambda - \alpha) = p'$ and such a μ does not even insure that $h_\alpha * \mu \in L_{p'}$ locally (for this see [6]). From the proof of Theorem 6, it appears that any condition on μ which insures $\|h_\alpha * \mu_{2r}\|_t^t \leq Q r^{n-\lambda}$, $t = \lambda/(\lambda - \alpha)$ will be sufficient. Two such conditions are:

- (i) $h_\alpha : L_{\lambda/\alpha} \rightarrow L_{\lambda/\alpha}(\mu)$ continuously, $0 < \alpha < \lambda$,
and
(ii) $h_\alpha * (h_\alpha * \mu)^{\alpha/(\lambda-\alpha)}$ is bounded on E_n .

The proofs require no new ideas.

REMARK 6: Finally, we observe that (10) is an extension of a theorem of Stein and Weiss [7] — see in particular their Theorem B^* in the case $p < q$, which corresponds to our case $p < p^*$. The example referred to in the above remark easily shows that no extension of this generality is possible when $p = p^*$.

REFERENCES

- [1] A. P. CALDERON, « *Lebesgue spaces of differential functions and distributions* », Proc Symp. Pre Math. Vol. 4, pp. 33-49, Amer. Math. Soc., Providence, R. I. 1961.
- [2] S. CAMPANATO, « *Proprietà di inclusione per spazi di Morrey* », Ricerche di Matematica, 12 (1963) pp. 67-86.
- [3] L. CARLSON, « *Selected problems in exceptional sets* », Van Nostrand 1967.
- [4] R. HUNT, « *An extension of the Marcinkiewicz interpolation theorem to Lorentz spaces* », Bull. Amer. Math. Soc. 70 (1964), pp. 803-807.
- [5] V. IL'IN, « *Generalization of an integral inequality* », Uspehi Mat. Nauk vol. II (1956) n. 4 (70) pp. 131-138.
- [6] N. MEYERS, « *A theory of capacities for potentials of functions in Lebesgue classes* », to appear.
- [7] E. M. STEIN and G. WEISS, « *Fractional integrals on n -dimensional Euclidean space* », J. Math. Mech. 7 (1958), pp. 503-514.
- [8] E. M. STEIN and A. ZYGMUND, « *Boundedness of translation invariant operators on Hölder spaces and L_p -spaces* », Ann. Math. 85 (1967) pp. 337-349.
- [9] G. STAMPACCHIA, « *Contributi alla regolarizzazione delle soluzioni dei problemi al contorno per equazioni del secondo ordine ellittiche* », Ann. Sc. Norm. Sup. Pisa, 12 (1958) pp. 223-245.