Astérisque

S. GOLDSTEIN

Spectrum of measurable flows

Astérisque, tome 40 (1976), p. 95-98

<http://www.numdam.org/item?id=AST_1976__40__95_0>

© Société mathématique de France, 1976, tous droits réservés.

L'accès aux archives de la collection « Astérisque » (http://smf4.emath.fr/Publications/Asterisque/) implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/conditions). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
http://www.numdam.org/

Société Mathématique de France
Astérisque 40 (1976) p.95-98

SPECTRUM OF MEASURABLE FLOWS

S. GOLDSTEIN

Let $\{T_t\}$ be a measurable flow (one parameter group of measure preserving transformations satisfying standard measurability condition) on the (Lebesgue) probability space (X,Σ,μ). Denote by H the generator of the group $U_t = e^{-itH}$ of induced unitaries and let $sp(H)$ be its spectrum.

Let $P(x)$ be the period of $x \in X$ (if x is aperiodic, $P(x) = \infty$). We have the

Proposition.

If $||P||_\infty = \infty$, $sp\ (H) = \mathbb{R}$

Corollary. If $\{T_t\}$ has an aperiodic component and in particular, if T^t is ergodic and $T^t = 1$ only if $t = 0$, then $sp\ (H) = \mathbb{R}$.

Proof: We first give an a argument which only establishes the latter half of the corollary. The argument, which is a generalization of the standard proof that the discrete spectrum of an ergodic transformation forms a group, may be applied to prove the appropriate results for automorphisms of C^*-algebras.

Let $sp\ \psi$ denote the support of $\psi \in L^2(X,\mu)$ in the spectral representation determined by H . Then

Lemma 1. If $\psi, \phi \in L^\infty(X,\mu)$ then $sp(\psi\phi) \subset sp\psi + sp\phi$.

Proof: For $\psi \in L^2(X,\mu)$ let

$$\hat{\psi}^\tau(E) = \frac{1}{2\pi}\int_{-\infty}^{\infty} e^{iEt}\, e^{-\frac{\tau}{2}t^2}(U_t\psi)\, dt \quad \text{(Bochner Integral)}$$

By the spectral theorem :

$$\hat{\psi}^\tau(E) = \delta_E^\tau(H)\psi$$

where

$$\delta_E^\tau(x) = \frac{1}{\sqrt{2\pi\tau}}\, e^{-(x-E)^2/2\tau} .$$

We claim that if $\lim_{t\to 0} ||\hat{\psi}^\tau(E)||_2 = 0$ for all E in an open set Θ, then $sp(\psi)$ and Θ are disjoint. [By the spectral theorem :

$$\int_{\Theta'} dE\, ||\hat{\psi}^\tau(E)||_2^2 = \frac{1}{2\pi\tau}\int_{\Theta'} dE \int d\mu_\psi(E')\, e^{-(E'-E)^2/\tau}$$

$$= \frac{1}{2\pi\tau}\int d\mu_\psi(E') \int_{\Theta'} dE\, e^{-\frac{(E'-E)^2}{\tau}} \sim \frac{1}{\sqrt{4\pi\tau}}\, \mu_\psi(\Theta') .$$

Therefore if $||\hat{\psi}^\tau(E)||_2 \to 0$ uniformly on Θ', $\mu_\psi(\Theta') = 0$. Hence, by Egoroff's theorem, $\mu_\psi(\Theta) = 0$ (here μ_ψ is the spectral measure belonging to ψ)] .

Now $\widehat{(\psi\phi)}^\tau(E_o) = \hat{\psi}^{\tau/2} * \hat{\phi}^{\tau/2}(E_o) = \int \hat{\psi}^{\tau/2}(E_o - E)\, \hat{\phi}^{\tau/2}(E)\, dE$,

and

$$||\widehat{(\psi\phi)}^\tau(E_o)||_2 \le \int ||\hat{\psi}^{\tau/2}(E_o-E)\, \hat{\phi}^{\tau/2}(E)||_2\, dE \le \frac{\text{const}}{\tau^2}\, e^{\delta^2/2\tau}\int e^{-\text{const}'E^2} dE \xrightarrow[t\to 0]{} 0$$

if $E_o \notin sp\psi + sp\phi$, where

$$\delta = \inf_E \text{dist}(E, sp\phi) \vee \text{dist}(E_o - E, sp\psi) = \frac{1}{2}\,\text{dist}(E_o, sp\psi + sp\phi).$$

In the above we have used these estimates :

$$||\hat{\psi}^\tau(E)||_\infty \le \int \left(\frac{1}{2\pi}\, e^{-\tau\tau^2/2}\, dt\right) ||\psi||_\infty = \frac{1}{\sqrt{2\pi\tau}}\, ||\psi||_\infty$$

$$||\hat{\psi}^\tau(E)||_2 \le \frac{1}{\sqrt{2\pi\tau}}\, ||\psi||_2\, e^{-[\text{dist}(E,sp\psi)]^2/2\tau}$$

(spectral theorem) and

$$||\psi\phi||_2 \leq ||\psi||_\infty \, ||\phi||_\infty \, ||\psi||_2 \, ||\phi||_2$$

Theorem.

Let T_t *be ergodic. Then sp(H) forms a group.*

Proof. Let $C\psi(x) = \bar{\psi}(x)$. Then $C\, U_t = U_t\, C$, i.e. $CH = -HC$. Therefore if $\lambda \in sp(H)$, $-\lambda \in sp(H)$. Let

$$E_1 \in \sigma(H) \ , \ sp\ \psi \quad (E_1 - \tfrac{1}{n},\ E_1 + \tfrac{1}{n}) , ||\psi||_2 = 1$$

$$E_2 \in \sigma(H) \ , \ sp\ \phi \quad (E_2 - \tfrac{1}{n},\ E_2 + \tfrac{1}{n}) , ||\phi||_2 = 1 \quad .$$

By ergodicity,

$$\frac{1}{T}\int_0^T dt \int d\mu\, |\psi|\,|\phi(T_t x)| \xrightarrow[T \to \infty]{} \left(\int d\mu\, |\psi|\right)\left(\int d\mu\, |\phi|\right) > 0 \ ,$$

so we may assume that $\psi\phi \neq 0$. By lemma 1, $\psi\phi$ is therefore an approximate eigenvector of H of eigenvalue $E_1 + E_2$, so that $E_1 + E_2 \in sp(H)$.

Now if $sp(H)$ is discrete, it must be of the form

$$\{n\, E_o \mid n = \ldots, -2, -1, 0, 1, 2, \ldots\}$$

and $T_\tau = \mathbf{1}$ for $\tau = 2\pi/E_o$. If $sp(H)$ is not discrete, it must be $\mathbb{R}$, since it is closed.

The proposition itself should follow from the decomposition of T_t into its ergodic components. We instead prove it directly, using an entirely different approach.

Assume first that T_t has a nontrivial aperiodic component. This may be represented as a special flow [ ], i.e. as a flow built under a function $f \geq \delta > 0$ and aperiodic automorphism (X_o, μ_o, T_o) . (X, μ) is identified with the "region under the function f on (X_o, μ_o)" with measure given by $\mu \times$ Lebesgue measure.

For $t > 0$, $T_t(x,s) = (x,s+t)$ for $s+t \leqslant f(x)$ and

$$T_t(x,s) = (T_o x, s+t-f(x)) \text{ otherwise.}$$

Since T_o is aperiodic, for any $n > 0$ there exists a set A with $\mu_o(A) > 0$ and $A, T_o(A), \ldots T_o^{n-1}(A)$ disjoint [ ]. Therefore there exist in X "rectangular tubes" of arbitrarily large length in which T_t induces a uniform translation. Consequently we may construct approximate eigenvectors (approximate plane waves e^{its}) corresponding to any $\lambda \in \mathbb{R}$. Finally, if T_t has no aperiodic component, we may construct approximate eigenfunctions is above by employing the representation of T_t, as the flow built under the function $\pi(x)$ and the identity automorphism. Here $\pi(x)$ is the period of $x \in X_o$. Since $||P(x)||_\infty = \infty$, we obviously have "tubes" of arbitrarily large length on which the motion is a uniform translation.

Dept. of Mathematics
White Hall
Cornell University
Ithaca, New York 14853
U.S.A.