Astérisque

FRANCESCO GUERRA

Introduction to stochastic field theory (abstract)

Astérisque, tome 40 (1976), p. 99-102

<http://www.numdam.org/item?id=AST_1976__40__99_0>

© Société mathématique de France, 1976, tous droits réservés.

L'accès aux archives de la collection « Astérisque » (http://smf4.emath.fr/Publications/Asterisque/) implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/conditions). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
http://www.numdam.org/

Société Mathématique de France
Astérisque 40 (1976) p.99-102

INTRODUCTION TO STOCHASTIC FIELD THEORY *

(Abstract)

Francesco Guerra

We consider classical dynamical systems with a finite or infinite number of degrees of freedom and with Lagrangian of canonical type which arise in many areas of physics. Typical examples are the harmonic and anharmonic oscillator, the free Klein-Gordon scalar field, the vector meson Proca field, the electromagnetic Maxwell field, the Klein-Gordon field with polynomial self-interaction, etc. To each of these classical dynamical systems we associate some stochastic processes following the prescriptions of Nelson stochastic mechanics [9,10] extended to general Lagrangian systems [7,4]. In particular we focus our attention on the ground state stochastic process of lowest energy and study its properties.

As a rather unexpected and surprising result [7] we find that the ground state stochastic process so constructed coincides with the stochastic process introduced in the so-called Euclidean formulation of quantum field theory. We recall here that the Euclidean theory, initiated by Schwinger [14] and Nakano [8], after the work of Symanzik [17] and Nelson [11,12] has become an essential tool [18,15] for the constructive quantum field theory program of Glimm and Jaffe and

* Research sponsored by the National Science Foundation, Grant No. GP-40768X.

their followers [3, 18]. In the usual interpretation the Euclidean theory is introduced starting from quantum field theory in the Wightman formulation[16] and then performing an analytic continuation in the time variables from real Minkowski time to the imaginary "Euclidean" time [14, 17]. The inverse analytic continuation is necessary in order to recover Wightman field theory from the Euclidean theory [11,13,15,1]. According to the proposed new interpretation [7] we can introduce "Euclidean"-Markov fields directly inside the framework of physical space-time and no analytic continuation is necessary both for the mathematical construction and the physical interpretation.

An attractive feature of our way of introducing Euclidean fields via stochastic mechanics is a simple derivation [7] of the Feynman-Kac-Nelson formula (see [6]) from the theory of stochastic differential equations [2].

The role of Euclidean field theory and stochastic mechanics as possible tools for the quantization of general relativity is stressed in [5].

REFERENCES

[1] J. Fröhlich, Verification of Axioms for Euclidean and Relativistic Fields and Haag's Theorem in a Class of $P(\phi)_2$-Models, Ann. Inst. Henri Poincaré XXI, 271 (1974).

[2] I.I. Gihman and A.V. Skorohod, Stochastic Differential Equations, Springer-Verlag, Berlin (1972).

[3] J. Glimm and A. Jaffe, Quantum Field Theory Models, In Statistical Mechanics and Quantum Field Theory, C. de Witt, R. Stora, Eds., Gordon and Breach, New York (1973)

[4] F. Guerra, On Stochastic Field theory, Suppl. J. de Physique 34, C1-95 (1973).

[5] F. Guerra, report at the Conference on Quantum Dynamics: Models and Mathematics, Bielefeld DBR, September 8-12, 1975.

[6] F. Guerra, L. Rosen and B. Simon, The $P(\phi)_2$ Euclidean Quantum Field Theory as Classical Statistical Mechanics, Annals of Mathmatics 101, 111 -(1975).

[7] F. Guerra and P. Ruggiero, New interpretation of the Euclidean-Markov Field in the Framework of Physical Minkowski Space-Time, Phys.Rev. Lett. 31, 1022 (1973), and paper in preparation.

[8] T. Nakano, Quantum Field Theory in Terms of Euclidean Parameters, Prog.Theor.Phys. 21, 241 (1959).

[9] E. Nelson, Derivation of the Schrödinger Equation from Newtonian Mechanics, Phys.Rev. 150, 1079 (1966).

[10] E. Nelson, Dynamical Theories of Brownian Motion, Princeton University Press, 1967.

[11] E. Nelson, Construction of Quantum Fields from Markoff Fields, J.F.A.12, 94 (1973) **The Free Markoff Field**, ibidem 12.211 (1973).

[12] E. Nelson, Probability Theory and Euclidean Field Theory, contribution to [18].

[13] K. Osterwalder and R Schrader, Axioms for Euclidean Green's Functions, Comm.Math. Phys. 31, 83 (1973), 42, 281 (1975).

[14] J. Schwinger, On the Euclidean Structure of Relativistic Field Theory, Proc. Nat. Acad. Sc. U.S. 44, 956 (1958).

[15] B. Simon, The $P(\phi)_2$ Euclidean (Quantum) Field Theory, Princeton Series in Physics, 1974.

[16] R. Streater and A.S. Wightman, PCT, Spin and Statistic and All That, Benjamin, New York, 1964.

[17] K. Symanzik, Euclidean Quantum Field Theory, In Local Quantum Theory, R. Jost, Ed., Academic Press, New York, 1969.

[18] G. Velo and A.S. Wightman, Eds., Constructive Quantum Field Theory, Springer Verlag, Berlin, 1973.

Via A. Falcone 70,
80127 Napoli
Italy.