

Astérisque

HARALD NIEDERREITER

ROBERT F. TICHY

**Metric theorems on uniform distribution and
approximation theory**

Astérisque, tome 147-148 (1987), p. 319-323

http://www.numdam.org/item?id=AST_1987__147-148__319_0

© Société mathématique de France, 1987, tous droits réservés.

L'accès aux archives de la collection « Astérisque » (<http://smf4.emath.fr/Publications/Asterisque/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

METRIC THEOREMS ON UNIFORM DISTRIBUTION AND APPROXIMATION THEORY

by

Harald NIEDERREITER and Robert F. TICHY

A sequence (x_n) , $n = 1, 2, \dots$, of points in $[0, 1)^k$ is called uniformly distributed (u.d.) if

$$\lim_{N \rightarrow \infty} N^{-1} \text{card} \{1 \leq n \leq N: x_n \in J\} = \text{vol}(J)$$

for all subintervals J of $[0, 1)^k$. A sequence (x_n) of numbers in $[0, 1)$ is called completely uniformly distributed (c.u.d.) if for all $k \geq 1$ the sequence of points $x_n = (x_n, x_{n+1}, \dots, x_{n+k-1})$, $n=1, 2, \dots$, is u.d., and a sequence (x_n) of real numbers is called c.u.d. mod 1 if the sequence $(\{x_n\})$ of fractional parts is c.u.d.

Franklin [3] proved that the sequence (x^n) is c.u.d. mod 1 for almost all $x > 1$ (in the sense of Lebesgue measure). Knuth [4] defined a sequence (x_n) in $[0, 1)$ to be random if, for every sequence (b_n) of distinct positive integers obtained by an effective algorithm, the sequence (x_{b_n}) is c.u.d. and raised the problem of whether $(\{x^n\})$ satisfies this definition of randomness for almost all $x > 1$. This problem was recently settled in the affirmative by the following result.

THEOREM 1 (Niederreiter and Tichy [7]). - If (a_n) is any sequence of distinct positive integers, then the sequence (x^{a_n}) is c.u.d. mod 1 for almost all $x > 1$.

The following lemmas are the key ingredients in the proof.

LEMMA 1 (Davenport, Erdős and LeVeque [2]). - Let (u_n) be a sequence of Lebesgue-measurable functions on the interval $[a, b]$ and put

$$I(N) = \int_a^b \left| \frac{1}{N} \sum_{n=1}^N e(u_n(x)) \right|^2 dx, \text{ where } e(t) = e^{2\pi it}.$$

If $\sum_{N=1}^{\infty} I(N)/N$ converges, then

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N e(u_n(x)) = 0 \text{ for almost all } x \in [a, b].$$

LEMMA 2 (van der Corput [8]). - Let the function u on [a,b] have a continuous and monotone first derivative that satisfies $|u'(x)| \geq L > 0$ for all $x \in [a,b]$. Then

$$\left| \int_a^b e(u(x)) dx \right| \leq \frac{1}{L}.$$

LEMMA 3. - If P is a monic polynomial of degree $d \geq 1$, then for the supremum norm $\|P\|_\infty$ on [a,b] we have

$$(1) \quad \|P\|_\infty \geq 2^{1-2d} (b-a)^d.$$

Lemma 3 is a standard result in approximation theory (see [6,p.39]). The lower bound (1) is in fact best possible, as it is attained by the normalized Chebyshev polynomial on [a,b] of degree d. The question of whether the integer exponents in Theorem 1 can be replaced by real exponents leads to another problem in approximation theory, namely to establish a lower bound for $\|P\|_\infty$ when

$$(2) \quad P(x) = x^{\lambda_s} + \sum_{j=1}^{s-1} c_j x^{\lambda_j}$$

is a monic Müntz polynomial with real coefficients c_j and real exponents λ_j satisfying $0 \leq \lambda_1 < \lambda_2 < \dots < \lambda_s$, $s \geq 2$. The lower bound should not depend on the c_j . For $[a,b] = [0,1]$ this is a classical problem considered by Müntz and Szász. By applying the transformation $x \rightarrow bx$, intervals $[0,b]$ can also be handled. For $[a,b]$ with $a > 0$ we cannot use the transformation trick since $P(a + (b - a)x)$ is not necessarily a Müntz polynomial. We now have the following result. We are grateful to Prof. J. Korevaar for providing the important reference [5].

LEMMA 4. - For any monic Müntz polynomial P given by (2) and any interval [a,b] with $0 < a < b$ we have

$$\|P\|_\infty \geq C_s a^{\lambda_s} \prod_{j=1}^{s-1} \frac{\lambda_s - \lambda_j}{\lambda_j - \lambda_1 + (1/\tau)},$$

where $\tau = \frac{1}{2} \log \frac{b}{a}$ and the constant $C_s > 0$ depends only on s.

Proof.- We first assume that $\lambda_1 = 0$. By a result in [5] there exists a sequence (γ_n) , $n = 0, 1, \dots$, with $\gamma_n > 0$ and

$\sum_{n=0}^{\infty} \gamma_n = 1$ such that $\prod_{n=0}^{\infty} \cos(\gamma_n z)$ is entire and

$$(3) \quad \prod_{n=0}^{\infty} |\cos(\gamma_n u)| \leq C e^{-\sqrt{u}} \text{ for all } u \geq 0,$$

where C is an absolute constant. Then the complex function

$$F(z) = \prod_{j=1}^{s-1} (z - i\lambda_j) \prod_{n=0}^{\infty} \cos(\gamma_n \tau z)$$

is entire and of exponential type τ , i.e. $|F(z)| = O_{\varepsilon}(e^{(\tau+\varepsilon)|z|})$ for all $\varepsilon > 0$, and the restriction of F to \mathbb{R} is in $L^2(\mathbb{R})$. By the Paley-Wiener theorem (see [1,p.103]) we have

$$F(z) = \int_{-\tau}^{\tau} f(t)e^{-izt} dt$$

for some $f \in L^2[-\tau, \tau]$. With $\sigma = \frac{1}{2} \log(ab)$ we get

$$G(z) := e^{-i\sigma z} F(z) = \int_{-\tau+\sigma}^{\tau+\sigma} f(t-\sigma)e^{-izt} dt = \int_{\log a}^{\log b} g(t)e^{-izt} dt$$

with $g \in L^2[\log a, \log b]$. By Fourier inversion we obtain

$$(4) \quad g(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} G(u)e^{itu} du \text{ for } t \in [\log a, \log b].$$

Put $h(x) = x^{-1}g(\log x)$ for $x \in [a, b]$. Then with $x = e^t$,

$$(5) \quad \int_a^b x^{\lambda_j} h(x) dx = \int_{\log a}^{\log b} g(t)e^{\lambda_j t} dt = G(i\lambda_j) \text{ for } 1 \leq j \leq s.$$

For P in (2) we use (5) and $G(i\lambda_j) = 0$ for $1 \leq j \leq s-1$ to get

$$(6) \quad \left| \int_a^b x^{\lambda_s} h(x) dx \right| = \left| \int_a^b P(x)h(x) dx \right| \leq \|P\|_{\infty} \int_a^b |h(x)| dx.$$

Then by (4), (5) and (6),

$$(7) \quad |G(i\lambda_s)| \leq \|P\|_{\infty} \int_a^b \frac{dx}{x} \cdot \frac{1}{2\pi} \int_{-\infty}^{\infty} |G(u)| du = \frac{\tau}{\pi} \|P\|_{\infty} \int_{-\infty}^{\infty} |G(u)| du.$$

Now

$$\begin{aligned} |G(i\lambda_s)| &= e^{\sigma\lambda_s} |F(i\lambda_s)| = e^{\sigma\lambda_s} \prod_{j=1}^{s-1} (\lambda_s - \lambda_j) \prod_{n=0}^{\infty} \cosh(\gamma_n \tau \lambda_s) \\ (8) &= b^{\lambda_s} \prod_{j=1}^{s-1} (\lambda_s - \lambda_j) \prod_{n=0}^{\infty} \frac{1}{2} (1 + e^{-2\gamma_n \tau \lambda_s}) \geq b^{\lambda_s} \prod_{j=1}^{s-1} (\lambda_s - \lambda_j) \prod_{n=0}^{\infty} e^{-2\gamma_n \tau \lambda_s} \\ &= a^{\lambda_s} \prod_{j=1}^{s-1} (\lambda_s - \lambda_j). \end{aligned}$$

Using (3) we get

$$\int_{-\infty}^{\infty} |G(u)| du \leq 2C \int_0^{\infty} \prod_{j=1}^{s-1} (u + \lambda_j) e^{-\sqrt{\tau}u} du$$

$$= \frac{2C}{\tau} \int_0^\infty \prod_{j=1}^{s-1} \left(\frac{v}{\tau} + \lambda_j\right) e^{-\sqrt{v}} dv \leq \frac{C}{\tau} \prod_{j=1}^{s-1} \left(\frac{1}{\tau} + \lambda_j\right),$$

and together with (7) and (8) the result for $\lambda_1 = 0$ follows. In the general case the lemma follows from

$$\|P\|_\infty \geq a^{\lambda_1} \|x^{\lambda_s - \lambda_1} + \sum_{j=1}^{s-1} c_j x^{j - \lambda_1}\|_\infty. \quad \square$$

Using Hölder's inequality in (6), the method above yields for the L^p -norm ($1 < p < \infty$)

$$\|P\|_p \geq C_s \tau a^{\lambda_s} \frac{(q-1)^{1/q}}{(a^{1-q} - b^{1-q})^{1/q}} \prod_{j=1}^{s-1} \frac{\lambda_s - \lambda_j}{\lambda_j - \lambda_1 + (1/\tau)},$$

where $p^{-1} + q^{-1} = 1$. For the L^1 -norm we get

$$\|P\|_1 \geq C_s \tau a^{\lambda_s + 1} \prod_{j=1}^{s-1} \frac{\lambda_s - \lambda_j}{\lambda_j - \lambda_1 + (1/\tau)}.$$

THEOREM 2.- If $\gamma \neq 0$ and (δ_n) is any sequence of reals with $\inf \delta_n > -\infty$ and $\inf_{m \neq n} |\delta_m - \delta_n| > 0$, then the sequence (γx^{δ_n}) is c.u.d. mod 1 for almost all $x > 1$.

Proof. We take $\gamma = 1$ to simplify the writing. We indicate the changes that are necessary in the proof of Theorem 1 (see [7]) to obtain Theorem 2. Dropping finitely many initial terms, if necessary, we can assume that all $\delta_n \geq 1$. As in [7] it suffices to show that if the lattice point $\underline{h} = (h_0, \dots, h_{k-1}) \neq \underline{0}$ and $1 < \alpha < \beta$ are fixed, then with

$$I(N) = \int_\alpha^\beta \left| \frac{1}{N} \sum_{n=1}^N e\left(\sum_{j=0}^{k-1} h_j x^{\delta_n + j}\right) \right|^2 dx$$

the series $\sum_{N=1}^\infty I(N)/N$ converges. We take $N \geq 3$. The cases (i) and (ii)

in [7] require no changes, if we note that the number of zeros of a Müntz polynomial in $[\alpha, \beta]$ can be bounded by [6, p.49, Hilfssatz].

In case (iii) the function $R_s(x)$ in [7, equ. (2.12)] is now a Müntz polynomial with $\lambda_s = d \leq (\log N)^{\sigma(s)} \leq \log N$, $\lambda_1 = 0$, $2 \leq s \leq k$, and

$$|\lambda_i - \lambda_j| \geq \delta := \inf_{m \neq n} |\delta_m - \delta_n| \text{ for } i \neq j.$$

Instead of the choice $\epsilon = (\log N)^{-2d}$ in [7] we take $\epsilon = (\log N)^{-2(s-1)-d}$.

For a subinterval $[a, b]$ of $[\alpha, \beta]$ with $|R_s(x)| \leq \epsilon$ for all $x \in [a, b]$ we have then by Lemma 4,

$$\epsilon \geq C_s a^{\lambda_s} \prod_{j=1}^{s-1} \frac{\lambda_s - \lambda_j}{\lambda_j + (1/\tau)} \geq C_s \delta^{s-1} \left(\lambda_s + \frac{1}{\tau}\right)^{1-s},$$

hence

$$\frac{1}{\tau} \geq C'_S \delta (\log N)^2 - \lambda_S \geq C(\underline{h}, \delta) (\log N)^2 \text{ for } N \geq N_0(\underline{h}, \delta).$$

From the definition of τ it follows that

$$\log \frac{b}{a} \leq C_1(\underline{h}, \delta) (\log N)^{-2} \text{ for } N \geq N_0(\underline{h}, \delta),$$

and so

$$\begin{aligned} b - a &\leq [\exp(C_1(\underline{h}, \delta) (\log N)^{-2}) - 1] \beta \\ &\leq C(\underline{h}, \delta, \beta) (\log N)^{-2} \text{ for } N \geq N_1(\underline{h}, \delta). \end{aligned}$$

Therefore, the Lebesgue measure of $V = \{x \in [\alpha, \beta] : |R_S(x)| \leq \epsilon\}$ is $O((\log N)^{-2})$. If W is the closure of the complement of V in $[\alpha, \beta]$, then for $x \in W$ we have

$$|R_S(x)| \geq (\log N)^{-2(s-1)-d} \geq (\log N)^{-2} (\log N)^{\sigma(s)} \text{ for } N \geq N_2(\underline{h}),$$

and so the argument following (2.13) in [7] can be applied. This yields $I(N) = O((\log N)^{-2})$, and the proof is complete. \square

R E F E R E N C E S

- [1] R.P. BOAS, Jr. - Entire Functions. Academic Press, New York, 1954.
- [2] H.DAVENPORT, P. ERDÖS and W.J. LEVEQUE. - On Weyl's criterion for uniform distribution. Michigan Math. J. 10, 311 - 314(1963).
- [3] J.N. FRANKLIN. - Deterministic simulation of random processes. Math. Comp. 17, 28-59 (1963).
- [4] D.E.KNUTH.- The Art of Computer Programming, Vol. 2: Seminumerical Algorithms. Addison - Wesley, Reading, Mass., 1969.
- [5] W.A.J. LUXEMBURG and J. KOREVAAR. - Entire functions and Müntz - Szász type approximation. Trans.Amer.Math.Soc.157, 23-37(1971).
- [6] I.P. NATANSON. - Konstruktive Funktionentheorie. Akademie-Verlag, Berlin, 1955.
- [7] H.NIEDERREITER and R.F. TICHY. - Solution of a problem of Knuth on complete uniform distribution of sequences. Mathematika 32, 26-32(1985).
- [8] J.G. VAN DER CORPUT. - Zahlentheoretische Abschätzungen. Math. Ann. 84, 53 - 79(1921).

HARALD NIEDERREITER
 Kommission für Mathematik
 Österreichische Akademie der
 Wissenschaften
 Dr. Ignaz-Seipel-Platz 2
 A-1010 WIEN
 Austria

ROBERT F.TICHY
 Abteilung für Technische
 Mathematik
 Technische Universität Wien
 Wiedner Hauptstraße 6-10
 A-1040 WIEN
 Austria