

Astérisque

AST

Summary

Astérisque, tome 215 (1993), p. 205

[<http://www.numdam.org/item?id=AST_1993__215__205_0>](http://www.numdam.org/item?id=AST_1993__215__205_0)

© Société mathématique de France, 1993, tous droits réservés.

L'accès aux archives de la collection « Astérisque » (<http://smf4.emath.fr/Publications/Asterisque/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

Summary

Let G be a real Lie group acting on a manifold M . H. Cartan introduced the equivariant de Rham complex and its cohomology $H_G^*(M)$. For a free action, $H_G^*(M)$ is the cohomology $H^*(G \backslash M)$ of the orbit space and if M is the point $\bullet$, $H_G^*(\bullet)$ is the algebra of invariant polynomial functions on the Lie algebra $\mathfrak{g}$ of G . To G -equivariant vector bundles on M , equipped with a G -invariant connection, are associated equivariant Chern classes. It became necessary to consider more general cohomological objects, like the algebra $H_G^\infty(M)$ of equivariant cohomology with C^∞ -coefficients, which is an algebra over $H_G^\infty(\bullet) = C^\infty(\mathfrak{g})^G$, and also the space $H_G^{-\infty}(M)$ of equivariant cohomology with $C^{-\infty}$ -coefficients, which is a module for $H_G^\infty(M)$, and for which $H_G^{-\infty}(\bullet)$ is the space of invariant generalized functions $C^{-\infty}(\mathfrak{g})^G$.

The first of the two articles of this volume, *Cohomologie équivariante et descente*, by Michel Duflot and Michèle Vergne, studies a generalization, denoted by $\mathcal{K}_G(M)$, of the cohomology $H_G^\infty(M)$. It is an algebra over $\mathcal{K}_G(\bullet) = C^\infty(G)^G$. It can be considered as a global analog of $H_G^\infty(M)$, and also as a de Rham version of the equivariant K -theory of M . The construction of $\mathcal{K}_G(M)$ uses the fixed point sets in M of the elements of G which are contained in compact subgroups of G . At least when G is compact, and under certain orientation assumptions, the “integral” over M of an element of $\mathcal{K}_G(M)$ is a G -invariant function on G .

The second article, *Equivariant cohomology with generalized coefficients*, by Shrawan Kumar and Michèle Vergne, undertakes a systematic study of the spaces $H_G^{-\infty}(M)$. Remarkable classes, with no analogs in the C^∞ -theory, are discovered. In particular, for a free action, the integral over M of an element of $H_G^{-\infty}(M)$ is a generalized function on $\mathfrak{g}$ with support 0. When G is compact, a spectral sequence allows a comparison of $H_G^{-\infty}(M)$ with the equivariant cohomology $H_G^*(M)$.

The two articles have common motivations, but they can be read independently.