

# COMPOSITIO MATHEMATICA

L. A. RUBEL

## **A relation between Fourier and Mellin averages**

*Compositio Mathematica*, tome 17 (1965-1966), p. 146-148

[http://www.numdam.org/item?id=CM\\_1965-1966\\_\\_17\\_\\_146\\_0](http://www.numdam.org/item?id=CM_1965-1966__17__146_0)

© Foundation Compositio Mathematica, 1965-1966, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques*

<http://www.numdam.org/>

# A relation between Fourier and Mellin averages

by

L. A. Rubel

We establish a relation here between additive and multiplicative convolution averages of a bounded function. The real numbers are a locally compact Abelian group, under the operation of addition, with Haar measure  $dt$ . The positive real numbers are a locally compact Abelian group, under the operation of multiplication, with Haar measure  $dt/t$ . Given a bounded Lebesgue measurable function  $g$ , with  $g(x)$  defined for all real numbers  $x$ , we may study its behaviour for large values of  $x$  by forming certain averages. One kind is with respect to integrable functions on  $(-\infty, \infty)$ , the other with respect to integrable functions on  $(0, \infty)$  where we use only the restriction of  $g$  to  $(0, \infty)$ . In each case, integrability is with respect to the appropriate measure, and the average depends only on the behaviour of  $g$  at  $+\infty$ . We call the first kind of average a Fourier average, the second kind a Mellin average, and we establish a connection between them. We shall assume that all our functions are Lebesgue measurable.

**MAIN THEOREM.** If  $g$  is bounded,  $K \geq 0$ ,

$$\int_{-\infty}^{\infty} K(t)dt = 1, H \geq 0, \quad \text{and} \quad \int_0^{\infty} H(t)dt/t = 1,$$

then

$$\limsup_{x \rightarrow \infty} \int_0^{\infty} H(x/t)g(t)dt/t \leq \limsup_{x \rightarrow \infty} \int_{-\infty}^{\infty} K(x-t)g(t)dt.$$

The next result follows from the main theorem on normalizing  $K$  so that  $\int_{-\infty}^{\infty} K(t)dt = 1$  (i.e. replacing  $K(t)$  by  $K(t)/\int_{-\infty}^{\infty} K(s)ds$ ), and writing  $H = H^+ - H^-$ , where  $H^+(t) = \max(H(t), 0)$  and  $H^-(t) = -\min(H(t), 0)$ . Now considering the normalizations of  $H^+$  and  $H^-$ , the main theorem and the corresponding result for  $\liminf$  may be applied.

**TAUBERIAN THEOREM.** Suppose  $g$  is bounded,

$$K \geq 0, \quad 0 < \int_{-\infty}^{\infty} K(t)dt < \infty,$$

and 
$$\lim_{x \rightarrow \infty} \int_{-\infty}^{\infty} K(x-t)g(t)dt = 0.$$

If 
$$\int_0^\infty |H(t)|dt/t < \infty,$$
 then 
$$\lim_{x \rightarrow \infty} \int_0^\infty H(x/t)g(t)dt = 0.$$

**PROOF.** We need the following result from [2, p. 1005].

**LEMMA 1.** Given a bounded function  $g$  and  $0 < \xi < 1$ , let

$$L(\xi) = \limsup_{x \rightarrow \infty} \frac{1}{x - \xi x} \int_{\xi x}^x g(t)dt,$$

and let  $L(1) = \sup_{0 < \xi < 1} L(\xi)$ . There exists a bounded function  $g^*$  such that  $g^* \geq g$  and  $\lim_{x \rightarrow \infty} x^{-1} \int_0^x g^*(t)dt = L(1)$ .

Let us write  $A = \limsup \int_0^\infty H(x/t)g(t)dt/t$  and  $B = \limsup \int_{-\infty}^\infty K(x-t)g(t)dt$ . We must prove  $A \leq B$ , which obviously follows from the next two lemmas.

**LEMMA 2.**  $A \leq L(1)$ .

**LEMMA 3.**  $L(1) \leq B$ .

We prove Lemma 2 via Lemma 1. Since  $g \leq g^*$  and  $K \geq 0$ , we have  $A \leq \limsup \int_0^\infty H(x/t)g^*(t)dt$ . But since  $\lim_{x \rightarrow \infty} x^{-1} \int_0^x g^*(t)dt = L(1)$  (i.e., the Cesaro limit of  $g^*$  is  $L(1)$ ), we may apply the Mellin form of the Wiener Tauberian theorem [1, p. 296] to conclude that  $\lim \int_0^\infty H(x/t)g^*(t)dt/t = L(1)$ , and hence  $A \leq L(1)$ . In more detail, we have  $\lim_{x \rightarrow \infty} x^{-1} \int_1^x g^*(t)dt = L(1)$ , and we may write  $x^{-1} \int_1^x g^*(t)dt = \int_0^\infty g^*(t)C(x/t)dt/t$ , where  $C(s) = 0$  for  $0 < s < 1$ , and  $C(s) = s^{-1}$  for  $s \geq 1$ . Denoting by  $C^\wedge$  the Mellin transform of  $C$ ,  $C^\wedge(r) = \int_0^\infty t^{ir}C(t)dt/t$ , we have  $C^\wedge(r) = (1-ir)^{-1}$ . Since  $C^\wedge(r) \neq 0$  for real  $r$ , we obtain the conclusion.

To prove Lemma 3, it is enough to do it under the special hypothesis that for some  $N$ ,  $K(x) = 0$  for  $|x| \geq N$ . The general case follows on letting

$$K_N(x) = \begin{cases} K(x)/\int_{-N}^N K(t)dt & \text{for } |x| \leq N \\ 0 & \text{for } |x| > N, \end{cases}$$

and then letting  $N \rightarrow \infty$ . Let us write

$$(K * g)(x) = \int_{-\infty}^\infty K(x-t)g(t)dt.$$

We shall prove that for  $\xi < 1$ ,

$$(1) \quad \int_{\xi x}^x (K * g)(y)dy = \int_{\xi x}^x g(t)dt + o(x).$$

If this is done, we get

$$L(\xi) \leq \limsup_{x \rightarrow \infty} \sup_{\xi x \leq y \leq x} (K * g)(y)$$

from which Lemma 3 follows directly. To prove (1), write

$$I(x) = \int_{\xi x}^x (K * g)(y) dy = \int_{-\infty}^{\infty} g(t) \int_{\xi x}^x K(y-t) dy dt.$$

But  $\int_{\xi x}^x K(y-t) dy$  vanishes if  $t < \xi x - N$  or  $t > x + N$ . And  $\int_{\xi x}^x K(y-t) dy = \int_{\xi x-t}^{x-t} K(y) dy$ . Hence

$$I(x) = \int_{\xi x - N}^{x + N} g(t) \int_{\xi x - t}^{x - t} K(y) dy dt.$$

We write  $\int_{\xi x - N}^{x + N} = \int_{\xi x - N}^{\xi x + N} + \int_{\xi x + N}^{x - N} + \int_{x - N}^{x + N}$ . For  $\xi x + N < t < x - N$ ,  $\int_{\xi x - t}^{x - t} K(y) dy = 1$ , and for any  $a$  and  $b$  with  $a < b$ ,

$$0 \leq \int_a^b K(y) dy \leq 1.$$

Hence

$$I(x) = \int_{\xi x + N}^{x - N} g(t) dt + 0(1) = \int_{\xi x}^x g(t) dt + 0(1),$$

and the proof is complete.

#### REFERENCES

G. H. HARDY

[1] *Divergent Series*, Oxford, 1949.

L. A. RUBEL

[2] Maximal means and Tauberian theorems, *Pacific J. Math.*, Vol. 10, No. 3, 1960, pp. 997—1007.

(Oblatum 25-3-63)

University of Illinois