

COMPOSITIO MATHEMATICA

ALLEN ALTMAN

STEVEN L. KLEIMAN

On the purity of the branch locus

Compositio Mathematica, tome 23, n° 4 (1971), p. 461-465

http://www.numdam.org/item?id=CM_1971__23_4_461_0

© Foundation Compositio Mathematica, 1971, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

ON THE PURITY OF THE BRANCH LOCUS

by

Allen Altman and Steven L. Kleiman

Let $f: X \rightarrow Y$ be a quasi-finite morphism of schemes and U the open subset of X where f is étale. The various theorems about the purity of the branch locus give conditions for U to be all of X . We offer a simple elementary proof that $U = X$ in the rather useful case when Y is smooth over a locally noetherian scheme S and U contains every point of depth ≤ 1 and is dense in the fibers over S . The proof is inspired by Zariski's original method [5] for characteristic 0. After the usual sort of reductions, Y becomes the spectrum of the ring of formal power series in a vector T of variables. Zariski proved that the functions on X also form a power series ring by expanding them in Taylor series in T . The appropriate differential operators $(1/i!)(\partial^i g/\partial T^i)$ first lift canonically over U and then extend over all of X because of the depth condition. However, lifting these operators amounts to constructing formal descent data for X (cf. [2] and [4]) and we present the proof from this point of view without mentioning differential operators, characteristics, or descent (and without using deeper results from formal geometry¹). The various standard results we need have been collected in [1] and all our references below are to this source.

THEOREM (Purity of the branch locus; cf. VI, 6.8). *Let S be a locally noetherian scheme, $g: Y \rightarrow S$ a smooth morphism and $f: X \rightarrow Y$ a quasi-finite morphism. Let U be the open subset of X where f is étale. Assume U contains every point x where $\text{depth}(O_x) \leq 1$ and that $(U \cap X(s))$ is dense in the fiber $X(s)$ for all s in S . Then $U = X$.*

NOTE. (i) If $g(Y) = S$ and $f(U)$ contains every point y of Y where $\text{dim}(O_y) \leq \text{dim}(S)$, then automatically $(U \cap X(s))$ is dense in $X(s)$ for all s in S .

(ii) If $(g \circ f)(U) = S$, the following conditions are equivalent:

(a) X satisfies S_2 (resp. X is normal) and U contains every point x where $\text{dim}(O_x) \leq 1$.

¹ However, using such results, Grothendieck [3] has also proved that $U = X$ when Y is locally a complete intersection and U contains every point of dimension ≤ 2 .

(b) U satisfies S_2 (resp. U is normal) and U contains every point x where $\text{depth}(O_x) \leq 1$.

(c) S satisfies S_2 (resp. S is normal) and U contains every point x where $\text{depth}(O_x) \leq 1$.

Indeed, the equivalence of (a) and (b) results directly from the definitions (resp. and Serre’s criterion). The equivalence of (b) and (c) holds by (VII, 4.9) because $U \rightarrow S$ is smooth and surjective.

(iii) In view of (i) and (ii), the theorem (applied to X minus the components of codimension one of the branch locus) implies that if X satisfies S_2 (e.g., X normal) and $\dim(S) \leq 1$, then the branch locus of f has pure codimension 1.

PROOF. By way of contradiction, assume $U \neq X$. Let x be a generic point of an irreducible component of $(X - U)$. We shall prove f is étale at x .

Let $y = f(x)$ and $s = g(y)$. Consider the flat base change $\text{Spec}(k) \rightarrow S$ where $k = O_y$. The hypotheses clearly hold for $f \otimes k$ and $g \otimes k$; by (VII, 5.11), $U \otimes k$ is the open set on which $f \otimes k$ is étale; the depth condition holds by virtue of (VII, 4.2); and clearly $U \otimes k$ is dense in the fibers over $\text{Spec}(k)$. Thus we may assume that S is the spectrum of a local ring k and that there exists a section $h : S \rightarrow Y$ such that $h(s) = y$.

Note that O_x/O_y is étale if (and only if) \hat{O}_x/\hat{O}_y is, that Y is an étale extension of a polynomial ring $k[[T_1, \dots, T_n]]$ with y lying over (T) , that $\text{depth}(\hat{O}_x) = \text{depth}(O_x)$ by (VII, 4.2) and that \hat{O}_x is a localization of $O_x \otimes_{O_y} \hat{O}_y$. Replace X by $\text{Spec}(\hat{O}_x)$, Y by $\text{Spec}(\hat{O}_y)$ and k by \hat{k} . While g is no longer of finite type, now $O_y \cong k[[T_1, \dots, T_n]]$, f is finite and $U = X - \{x\}$. Furthermore, clearly U contains every point of depth ≤ 1 and $(g \circ f)(U) = S$. Let $V = (Y - \{y\})$. Then f is étale over V and since $\text{depth}_{O_y}(B) = \text{depth}_B(B)$ where $B = O_x$ by (III, 3.16), the open set V contains every point $z \in Y$ such that $\text{depth}_{O_z}(B_z) \leq 1$.

Finally, it suffices to construct an isomorphism $X_O \times_S Y \xrightarrow{\sim} X$ where $X_O = X \times_Y S$. For then, by (VII, 5.11), X_O/S is étale because $U = X_O \times_S V$ is étale over V and $V \rightarrow S$ is surjective and flat; whence X/Y is étale because $Y \rightarrow S$ is surjective and flat. Thus it suffices to prove the following theorem (whose proof will be presented after several preliminary lemmas).

THEOREM. *Let k be a noetherian ring and $A = k[[T_1, \dots, T_n]]$ a formal power series ring. Let B be a finite A -algebra which is étale over every prime p of A where $\text{depth}(B_p) \leq 1$. Then there exists a (canonical) isomorphism $A \otimes_k B_O \xrightarrow{\sim} B$ where $B_O = k \otimes_A B$.*

DEFINITION. Let k be a ring, R a k -algebra. The module of m th principal

parts of R over k , denoted $P^m(R)$, is defined as $(R \otimes_k R)/I^{m+1}$ where I is the diagonal ideal. It is naturally filtered by the powers of I .

LEMMA 1. *Let k be a ring, R a noetherian k -algebra and S an étale extension of R .*

(i) *The natural $(R \otimes_k R)$ -algebra homomorphism $v_m : P^m(R) \otimes_R S \rightarrow P^m(S)$ sending $(a_1 \otimes a_2) \otimes s$ to $a_1 \otimes sa_2$ (resp. to $sa_1 \otimes a_2$) is an isomorphism (where $P^m(R)$ is regarded as an R -module from the right (resp. left)).*

(ii) *The induced map $gr^i(P^m(R)) \otimes_R S \rightarrow gr^i(P^m(S))$ is an isomorphism.*

PROOF. In (i), both filtered modules are separated and complete; so it suffices to show that the $gr^i(v_m)$ are isomorphisms. Since S/R is flat, $gr^i(P^m(R) \otimes_R S)$ is isomorphic to $gr^i(P^m(R)) \otimes_R S$. Thus (i) follows from (ii).

Let I (resp. J) be the diagonal ideal of (R/k) (resp. (S/k)), and set $K = \ker(S \otimes_k S \rightarrow S \otimes_R S)$. As in (VI, 4.9 and 4.10), $K \cong I \otimes_{(R \otimes_k R)} (S \otimes_k S)$ since S/R is flat. Hence $(K^i/K^{i+1}) \cong (I^i/I^{i+1}) \otimes_{(R \otimes_k R)} (S \otimes_k S)$. Also, $(K^i/K^{i+1}) \otimes_{(S \otimes_k S)} S \cong (J^i/J^{i+1})$ since S/R is unramified. Therefore, $(I^i/I^{i+1}) \otimes_{(R \otimes_k R)} S \cong (J^i/J^{i+1})$. Since the $(R \otimes_k R)$ -module structure of (I^i/I^{i+1}) coincides with the left (resp. right) R -module structure of (I^i/I^{i+1}) , this isomorphism coincides with the induced map.

LEMMA 2. *Let R be a noetherian local ring; P, N two finite R -modules. If $\text{depth}(N) \geq 2$, then $\text{depth}(\text{Hom}_R(P, N)) \geq 2$.*

PROOF. An N -regular sequence (x_1, x_2) is easily seen to be $\text{Hom}_R(P, N)$ -regular.

LEMMA 3 (cf. VII, 2.10). *Let R be a noetherian ring, M a finite R -module and V an open subset of $\text{Spec}(R)$.*

(i) *Suppose V contains every point p where $\text{depth}(M_p) = 0$; (e.g., V contains every generic point of $\text{Supp}(M)$ and M satisfies S_1). Then the restriction $M \rightarrow \Gamma(V, \tilde{M})$ is injective.*

(ii) *Suppose V contains every point p where $\text{depth}(M_p) \leq 1$; (e.g., V contains every point of codimension ≤ 1 in $\text{Supp}(M)$ and M satisfies S_2). Then $M \rightarrow \Gamma(V, \tilde{M})$ is bijective.*

PROOF. To prove (i), let $x \in M$ go to zero in $\Gamma(V, \tilde{M})$. Assume $x \neq 0$. Then there exists a prime p in $\text{Ass}(Ax)$. Then $pA_p \in \text{Ass}(A_p x) \subset \text{Ass}(M_p)$, so $\text{depth}(M_p) = 0$. Hence $p \in V$, so $A_p x = 0$; this contradicts $pA_p \in \text{Ass}(A_p x)$.

To prove (ii), let $f \in \Gamma(V, \tilde{M})$. Let E be the ideal of elements $s \in A$ such that sf extends to an element x of M . For every prime p in V the image of f in M_p is a fraction x/s , and it follows that $E \nsubseteq p$. By (III, 1.5),

there exists therefore an element s of E not in any prime p where $\text{depth}(M_p) = 0$. Let x be an element of M extending sf .

Since s is M -regular, V contains every prime p where $\text{depth}((M/sM)_p) = 0$. Since the image of x in (M/sM) is zero on V , it is zero by (i). Thus there exists a g in M such that $x = sg$. Then $s(g-f)$ is zero over V . Since s is M -regular, $g = f$ on V , and the proof is complete.

PROOF OF THEOREM. Let $P = \varinjlim (P^m(A))$. It will suffice to construct a P -isomorphism $u : P \otimes_A B \rightarrow B \otimes_A P$ where in $P \otimes_A B$ (resp. $B \otimes_A P$), P is regarded as an A -module via the second (resp. first) factor. Namely, define $w : A \otimes_k A \rightarrow A$ by $w(a_1 \otimes a_2) = a_2(0)a_1$ where $a_2(0)$ denotes the constant term of a_2 . Then $w(I)$ is contained in $m = T_1A + \dots + T_nA$, so w defines an A -homomorphism $\hat{w} : P \rightarrow A$. Since the diagram

$$\begin{array}{ccc}
 & A \xleftarrow{\hat{w}} P & \\
 & \uparrow & \uparrow j_2 \\
 k = (A/m) & \leftarrow & A
 \end{array}$$

is commutative, $A \otimes_P (P \otimes_A B) = A \otimes_k (k \otimes_A B)$. Hence, $(A \otimes_P u) : (A \otimes_k B_0) \xrightarrow{\sim} B$ is the required isomorphism.

Since $A = k[[T_1, \dots, T_n]]$, the $(A \otimes_k A)$ -module $P^m(A)$, regarded as an A -module on the left (resp. right) is isomorphic to $A^{\oplus r}$ for some r . Therefore $(B \otimes_A P^m(A)) \cong B^{\oplus r}$. Thus, the open set V of $\text{Spec}(A)$ over which B is étale, contains all p where $\text{depth}((B \otimes_A P^m(A))_p) \leq 1$.

Regarding the two $(A \otimes_k A)$ -modules $P^m(A) \otimes_A B$ and $B \otimes_A P^m(A)$ as A -modules on the left, consider $M = \text{Hom}_A(P^m(A) \otimes_A B, B \otimes_A P^m(A))$. By lemma 1, M has a natural section over V . By lemma 2, V contains every point p where $\text{depth}(M_p) \leq 1$. So by lemma 3, this section extends to an A -homomorphism $u_m : P^m(A) \otimes_A B \rightarrow B \otimes_A P^m(A)$. In fact, u_m is an $(A \otimes_k A)$ -homomorphism since it is on V and we may apply 3(i). Similarly, we obtain an $(A \otimes_k A)$ -homomorphism $B \otimes_A P^m(A) \rightarrow P^m(A) \otimes_A B$ which is an inverse to u_m on V ; hence, it is a global inverse. The isomorphisms u_m clearly form a compatible system of maps, inducing the required P -isomorphism $u : P \otimes_A B \rightarrow B \otimes_A P$.

REFERENCES

A. ALTMAN AND S. KLEIMAN
 [1] Introduction to Grothendieck duality theory, Lecture Notes in Math. Vol. 146, Springer-Verlag, Berlin-Heidelberg-New York (1970).
 A. GROTHENDIECK
 [2] Appendix to Crystals and the De Rham Coh. of Sch., in Dix exposés sur la cohomologie des schémas, North-Holland Publ. Co., Amsterdam (1968).

A. GROTHENDIECK

[3] "Sem de Geom. Alg. 2", North-Holland Publ. Co., Amsterdam (1968). Exposé X Theorem (3.4).

N. KATZ

[4] Nilpotent connections and the Monodromy Theorem Applications of a Result of Turrittin", in Sem. on Degeneration of Alg. Var., Inst. for Adv. Study, Princeton, New Jersey (1970). Remark 8.9.16.

O. ZARISKI

[5] "On the purity of the branch locus of algebraic functions," Proc. Nat. Acad. Sci. U.S.A. 44 (1958), 791-796.

(Oblatum 00-00-000)

Allen Altman
Department of Mathematics
University of California, San Diego
La Jolla, Calif. 92037 U.S.A.

Steven L. Kleiman
Department of Mathematics
Massachusetts Institute of Technology
Cambridge, Mass. 02139 U.S.A.

15-11-10