

# COMPOSITIO MATHEMATICA

TAMMO TOM DIECK

**A finiteness theorem for the burnside ring  
of a compact Lie group**

*Compositio Mathematica*, tome 35, n° 1 (1977), p. 91-97

[http://www.numdam.org/item?id=CM\\_1977\\_\\_35\\_1\\_91\\_0](http://www.numdam.org/item?id=CM_1977__35_1_91_0)

© Foundation Compositio Mathematica, 1977, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

## A FINITENESS THEOREM FOR THE BURNSIDE RING OF A COMPACT LIE GROUP

Tammo tom Dieck

Let  $G$  be a compact Lie group and let  $A(G)$  be its Burnside ring [6]. We show that after inverting a finite number of primes the ring  $A(G)$  is generated by idempotent elements. The following result (Theorem 1) about compact Lie groups is basic for our investigations.

Let  $H$  be a subgroup of  $G$  (subgroups will always be closed), let  $NH$  be its normalizer in  $G$  and denote  $NH/H$  by  $WH$ . If  $K$  is a compact Lie group let  $K_0$  be its component of the unit element.

**THEOREM 1:** *There exists an integer  $b$  such that for each closed subgroups  $H$  of  $G$  the index  $|WH : (WH)_0|$  is less than  $b$ .*

Let  $A_c(G)$  be the integral closure of  $A(G)$  in its total quotient ring.

**THEOREM 2:** *There exists an integer  $n > 0$  such that  $nA_c(G) \subset A(G)$ . The minimal such  $n$  is the least common multiple of the numbers  $WH$  where  $H$  runs through all subgroups such that  $WH$  is finite.*

The minimal integer  $n(G)$  provided by Theorem 1 replaces the order of the finite group if one extends the general Artin induction theorem (see Dress [7], Theorem 2, p. 204) to compact Lie groups, whence its importance.

### 1. Normalizers

1.1. We prove in this section Theorem 1. The proof proceeds in three steps: We first reduce to the case that  $WH$  is finite; then we reduce to the case that  $H$  is finite; and finally show that for finite  $H$

with finite  $WH$  the order of  $WH$  is uniformly bounded.

Let  $H$  be a closed subgroup of  $G$  (notation:  $H < G$ ). Let  $\text{Aut}(H)$  be the automorphism group of  $H$  and  $\text{In}(H)$  the closed normal subgroup of inner automorphisms. The group  $\text{Aut } H/\text{In } H$  is discrete. Mapping  $n \in NH$  to the conjugation automorphism  $c(n): h \mapsto nhn^{-1}$  of  $H$  induces a homomorphism  $NH \rightarrow \text{Aut } H/\text{In } H$  with kernel  $ZH \cdot H$ , where  $ZH$  denotes the centralizer of  $H$ . Hence  $NH/ZH \cdot H$  being a compact subgroup of the discrete group  $\text{Aut } H/\text{In } H$  is finite. We conclude

LEMMA 1:  $WH$  is finite if and only if  $ZH/(ZH \cap H)$  is finite.

LEMMA 2: A compact Lie group contains only a finite number of conjugacy classes ( $K$ ) where  $K$  is the centralizer of a closed subgroup.

PROOF: Let  $G$  act on  $M = G$  via conjugation  $G \times M \rightarrow M: (g, m) \mapsto gmg^{-1}$ . If  $H < G$  then the fixed point set  $M^H$  is the centralizer  $ZH$ . A compact differentiable  $G$ -manifold has finite orbit type. Hence there exist finitely many conjugacy classes  $(H_1), \dots, (H_k)$  such that for any closed subgroup  $H$   $M^H = M^K$  and  $(K) = (H_i)$  for a suitable  $i$ .

LEMMA 3: For any  $H < G$  the group  $ZH \cdot H$  has finite index in its normalizer.

PROOF: We have  $Z(ZH \cdot H) < ZH < ZH \cdot H$ ; hence the assertion follows from Lemma 1.

If  $n \in G$  normalizes  $H$  then also  $ZH$  and hence  $ZH \cdot H$ . We therefore have

$$NH/ZH \cdot H < N(ZH \cdot H)/ZH \cdot H.$$

Using Lemma 3 and the existence of an upper bound for the set

$$\{|WH| \mid H < G, WH \text{ finite}\} =: F(G)$$

we obtain

LEMMA 4: There exists an integer  $c$  such that for all  $H < G$  we have

$$|NH/ZH \cdot H| < c.$$

Now we are able to obtain the first reduction of our problem. From the exact sequence  $1 \rightarrow ZH/ZH \cap H \rightarrow WH \rightarrow NH/ZH \cdot H \rightarrow 1$  we see

that  $WH/(WH)_0 \rightarrow NH/ZH \cdot H$  has a kernel which is a quotient of  $ZH/(ZH)_0$ . Lemmas 2 and 4 then show that

$$\{|WH/(WH)_0| | H < G\}$$

is bounded. But note that Lemma 4 requires a bound for the set  $F(G)$ .

1.2. We show by induction over  $|G/G_0|$  and  $\dim G$  that  $F(G)$  has an upper bound  $a = a(G/G_0, \dim G)$ . For finite  $G$  we can take  $a = |G|$ .

Suppose that an upper bound  $a(K/K_0, \dim K)$  is given for all  $K$  with  $\dim K < \dim G$ . Let  $\Sigma(G) = \{H < G | WH \text{ finite}\}$ . Suppose  $H \in \Sigma(G)$  is not finite. We consider the projection  $p: NH_0 \rightarrow NH_0/H_0 =: U$ . Let  $V$  be the normalizer of  $H/H_0$  in  $U$ . Then  $WH = V/(H/H_0)$  and therefore  $H/H_0 \in \Sigma(U)$ . Since  $\dim U < \dim G$  we obtain by induction hypothesis

$$|WH| \leq a(U/U_0, \dim U).$$

We show that the possible values for  $|U/U_0|$  are finite in number. The group  $NH_0$  is the normalizer of a connected subgroup. By [8], Ch. VII, Lemma 3.2, there are only a finite number of conjugacy classes of such subgroups. Hence for a given  $G$  the possible  $|U/U_0|$  are bounded, say  $|U/U_0| \leq m(G)$ . We have inequalities

$$\begin{aligned} |U/U_0| &\leq |NH_0/N_0H_0| |N_0H_0/(NH_0)_0| \\ &\leq |G/G_0| m(G), \end{aligned}$$

where  $N_0$  means normalizer in  $G_0$ . By the classification theory of compact connected Lie groups there are only a finite number in each dimension. Hence there exists a bound for  $|U/U_0|$  depending only on  $|G/G_0|$  and  $\dim G$ . This proves the induction step as far as the non-finite  $H$  in  $\Sigma(G)$  are concerned.

1.3. Let  $H \in \Sigma(G)$  be finite. Let  $\sigma(G)$  be the set of finite subgroups of  $G$ . We use the following classical theorem of Jordan.

LEMMA 5: *There exists an integer  $j = j(|G/G_0|, \dim G)$  with the following properties: To each  $H \in \sigma(G)$  there exists an abelian normal subgroup  $A_H$  of  $H$  such that  $|H/A_H| \leq j$ . Moreover the  $A_H$  can be chosen such that  $H < K$  implies  $A_H < A_K$ .*

PROOF: Boothby and Wang [2]. Wolf [9]. In these references only connected groups are considered. The straightforward extension to non-connected groups we leave to the reader.

If  $H \in \Sigma(G)$  is finite then also  $K := NH$  is finite and by Lemma 1  $K \in \Sigma(G)$ . We choose  $j = j(|G/G_0|, \dim G)$  and  $A_H, A_K$  according to Lemma 5. We have

$$|K/H| \leq |K/A_K| \cdot |A_K/H \cap A_K| \leq j|A_K/H \cap A_K|.$$

Hence it is sufficient to find a bound for  $|A_K/H \cap A_K|$ . Consider the exact sequence  $1 \leftarrow S \leftarrow H \leftarrow A_H \leftarrow 1$ . The conjugation  $c(a)$  with  $a \in A_K$  is trivial on  $A_H$ , because  $A_K > A_H$ , and hence  $c(a)$  induces an automorphism of  $S$ . Since  $|S| \leq j$  this automorphism has order at most  $J = j!$ , i.e.  $c(a^r)$  is the identity on  $S$  and  $A_H$  for a suitable  $r \leq J$ . The group of such automorphisms modulo the subgroups of inner automorphisms by elements of  $A_H$  is isomorphic to  $H^1(S; A_H)$ , with  $S$  acting on  $A_H$  by conjugation. Since this group is annihilated by  $|S|$  we see that  $c(a^s)$  is an inner automorphism by an element of  $A_H$  for a suitable  $s \leq J|S| \leq jJ$ . In other words:  $a^s h^{-1} \in ZH$ . Hence it is sufficient to find a bound for the order of  $A_K \cap ZH/H \cap A_K \cap ZH$ .

Let  $U_1 = A_K \cap ZH$ . By [3], Théorème 1,  $U_1$  is contained in the normalizer  $NT$  of a maximal torus of  $G$ . Put  $U = U_1 \cap T$ . Then  $|U_1/U| \leq |G/G_0| |w(G_0)|$  where  $w(G_0)$  denotes the Weyl group of  $G_0$ . We estimate the order of  $U$ . Since  $U$  is abelian we have  $U < ZU$ . Moreover  $H < ZU$  by definition of  $ZH$ . Since  $U$  is contained in  $ZU$  it is contained in the center  $C := CZU$  of  $ZU$ . The inclusion  $H < ZU$  implies  $C < NH$ . Hence  $C$  is finite.

We proceed to show that for the order of a finite center  $C(G)$  of  $G$  there exists a bound depending only on  $|G/G_0|$  and  $\dim G$ . We let  $G/G_0$  act by conjugation on  $C(G_0)$ . Then  $C(G) \cap G_0$  is the fixed point set of this action. We have  $C(G_0) = A \times T_1$ , where  $A$  is a finite abelian group and  $T_1$  a torus. The group  $A$  is the center of a semisimple group and therefore, by the classification theory of these groups,  $|A|$  is bounded by a constant  $c$  depending only on  $\dim G$ . The exact cohomology sequence associated to the universal covering  $0 \rightarrow \pi_1 T_1 \rightarrow V \rightarrow T_1 \rightarrow 0$  shows, that the fixed point set of the action of  $G/G_0$  on  $T_1 = C(G_0)_0$  is isomorphic to  $H^1(G/G_0, \pi_1 T_1)$ , hence its order is bounded by a constant  $d$  depending only on  $|G/G_0|$  and the rank of  $T_1$ . Hence  $|C(G)| \leq |G/G_0|cd$ .

Finally we have to show that for the possible groups  $ZU$  the order  $|ZU/(ZU)_0|$  is bounded.

$U$  is contained in a maximal torus of  $G$ . Therefore  $ZU$  is a subgroup of maximal rank and  $(ZU)_0$  a connected subgroup of maximal rank. By [4] there exist only finitely many conjugacy classes of connected subgroups of maximal rank. We have

$$|ZU/(ZU)_0| \leq |N(ZU)_0/(ZU)_0| \leq |G/G_0| |N_0(ZU)_0/(ZU)_0|.$$

There are only finitely many possibilities for normalizers  $N_0(ZU)$  in  $G_0$  of  $(ZU)_0$ .

This finishes the proof of Theorem 1.

## 2. The integral closure of $A(G)$

Let  $\phi = \text{Spec}(A(G) \otimes Q)$  be the prime ideal spectrum of  $A(G) \otimes_Z Q =: A_0$  with Zariski topology. By [6], Theorem 4, the prime ideals of  $A_0$  correspond bijectively to kernels of ring homomorphisms  $A_0 \rightarrow Q$ . Therefore  $\phi$  is a totally disconnected compact Hausdorff space ([5], §4. Ex. 16; [1], Ch. 3. Ex. 11).

LEMMA 6: (1) For each  $a \in A_0$  the map  $\varphi_a: \phi \rightarrow Q: \varphi \mapsto \varphi(a)$  is locally constant. If  $a \in A(G)$  then  $\varphi(a) \in Z$ . (2) Let  $C(\phi, Q)$  be the ring of locally constant functions  $\phi \rightarrow Q$ . The ring homomorphism  $\alpha: A_0 \rightarrow C(\phi, Q): a \mapsto \varphi_a$  is an isomorphism. The image  $\alpha A(G)$  is contained in  $C(\phi, Z)$ . (3) The map  $A(G) \rightarrow A_0: a \mapsto a \otimes 1$  is the inclusion of  $A$  into its total quotient ring. The map  $\alpha: A(G) \rightarrow C(\phi, Z)$  is the inclusion of  $A(G)$  into the integral closure of  $A(G)$  in  $A_0 \cong C(\phi, Q)$ .

PROOF: (1) For  $k \in Q$  the set  $\varphi_a^{-1}(k)$  is closed in  $\phi$ , by definition of the Zariski topology. Since  $A_0$  is an absolutely flat ring, this set is also open by [5], §4. Ex. 16.b. Hence  $\varphi_a$  is continuous and there exist only a finite number of non-empty sets  $\varphi_a^{-1}(k)$ , because  $\phi$  is compact. A homomorphism  $A_0 \rightarrow Q$  is induced from a homomorphism  $A(G) \rightarrow Z$ , by [6] Theorem 4.

(2) Since the localizations of  $A_0$  at its prime ideals are canonically isomorphic to  $Q$ , we can identify  $C(\phi, Q)$  with the ring of sections of the structure sheaf of  $A_0$ . Then  $\alpha$  corresponds to the canonical map of  $A_0$  into this ring, hence  $\alpha$  is an isomorphism.

(3) To form the total quotient ring we have to invert the elements which are not zero divisors. Hence  $A_0$  is contained in the total quotient ring. If  $x \in C(\phi, Q) \cong A_0$  is not a zero divisor then it is a locally constant function without zeros, hence a unit. Therefore  $C(\phi, Q)$  is its own total quotient ring. Since a locally constant function  $\phi \rightarrow Z$  takes only finitely many values the ring  $C(\phi, Z)$  is generated by idempotent elements hence integral over any subring. Under the isomorphism  $\alpha$  the ring  $C(\phi, Z)$  corresponds to  $\{a \in A_0 \mid \varphi \in \phi \Rightarrow \varphi(a) \in Z\}$ . If  $x \in A_0$  is integral over  $A(G)$  then  $\varphi(x)$  is integral over  $Z$ , hence  $\varphi(x) \in Z$  and  $\alpha(x) \in C(\phi, Z)$ .

Let  $A_c(G)$  denote the pre-image of  $C(\phi, Z)$  under  $\alpha$ . We recall that  $A(G)$  is additively the free abelian group on homogeneous spaces  $G/H$  where  $H$  runs through a complete system of non-conjugate subgroups  $H$  of  $G$  with finite index in their normalizer ([6], Theorem 1). Let  $\Sigma(G)$  denote the set of conjugacy classes  $(H)$  of subgroups  $H$  of  $G$  with finite  $WH = NH/H$ .

LEMMA 7:  $A_c(G)$  is additively the free abelian group with basis  $x_{(H)} := |WH|^{-1}G/H$ ,  $(H) \in \Sigma(G)$ .

PROOF: The elements  $x_{(H)}$  are contained in  $A_c(G)$ : Suppose  $\varphi \in \phi$  is given. Then there exists  $(K) \in \Sigma(G)$  such that  $\varphi(x \otimes r) = r\chi(x^K)$  where  $\chi$  denotes the Euler characteristic and  $x^K$  the  $K$  fixed point set of any manifold representing  $x$ . But  $WH$  acts freely as a  $G$ -automorphisms group on  $G/H$ , hence also on  $G/H^K$ . Therefore  $\chi(G/H^K)$  is divisible by  $|WH|$  and we see that  $\varphi(x_{(H)}) \in Z$  for all  $\varphi$ . By definition of  $A_c(G)$  this means  $x_{(H)} \in A_c(G)$ .

The elements  $x_{(H)}$  are obviously linearly independent over  $Z$ . We have to show that any  $x \in A_c(G)$  is an integral linear combination of the  $x_{(H)}$ . In any case we have an expression  $x = \sum r_H x_{(H)}$  with rational  $r_H$ . Take  $(L) \in \Sigma(G)$  maximal with respect to inclusion such that  $r_L \neq 0$ . Then  $\sum r_H |WH|^{-1} \chi(G/H^L) \in Z$ . But  $G/H^L = \phi$  for  $(H) \neq (L)$ ,  $r_H \neq 0$ ; and  $\chi(G/L^L) = |WH|$ . Therefore  $r_L \in Z$ . We apply the same argument to  $x - r_L x_{(L)}$  and complete the proof by induction.

Lemma 7 and Theorem 1 give a proof of Theorem 2.

#### REFERENCES

- [1] M. F. ATIYAH and I. G. MACDONALD: *Introduction to Commutative Algebra*. Addison-Wesley Publ. Comp. 1969.
- [2] W. BOOTHBY and H.-C. WANG: On the finite subgroups of connected Lie groups. *Comment. Math. Helv.* 39 (1964) 281-294.
- [3] A. BOREL et J.-P. SERRE: Sur certain sous groupes des groupes de Lie compacts. *Comment. Math. Helv.* 27 (1953) 128-139.
- [4] A. BOREL et J. DE SIEBENTHAL: Les sous-groupes fermés de rang maximum des groupes de Lie clos. *Comment. Math. Helv.* 23 (1949) 200-221.
- [5] N. BOURBAKI: *Algèbre commutative, Chapitre 2*. Hermann, Paris 1961.
- [6] T. TOM DIECK: The Burnside Ring of a Compact Lie Group I. *Math. Ann.* 215 (1975) 235-250.
- [7] A. DRESS: Contributions to the theory of induced representations. *Springer Lecture Notes* 342, (1973) 183-240.

[8] *Seminar on Transformation Groups*. Ed. A. Borel et al. Princeton University Press 1960.

[9] J. A. WOLF: *Spaces of Constant Curvature*. McGraw-Hill, New York 1967.

(Oblatum 26-V-1976)

Math. Institut  
34 Göttingen  
Bunsenstrasse 3-5  
BRD