

COMPOSITIO MATHEMATICA

RUISHI KUWABARA

On isospectral deformations of riemannian metrics

Compositio Mathematica, tome 40, n° 3 (1980), p. 319-324

http://www.numdam.org/item?id=CM_1980_40_3_319_0

© Foundation Compositio Mathematica, 1980, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
http://www.numdam.org/*

ON ISOSPECTRAL DEFORMATIONS OF RIEMANNIAN METRICS

Ruishi Kuwabara

1. Introduction

Let M be an n -dimensional compact orientable C^∞ manifold, and g be a C^∞ Riemannian metric on M . It is known that the Laplace-Beltrami operator $\Delta_g = -g^{ij}\nabla_i\nabla_j$ acting on C^∞ functions on M has an infinite sequence of eigenvalues (denoted by $\text{Spec}(M, g)$)

$$0 = \lambda_0 < \lambda_1 \leq \lambda_2 \leq \cdots \leq \lambda_k \leq \cdots \uparrow + \infty$$

each eigenvalue being repeated as many as its multiplicity.

Consider the following problem [1, p. 233]: Let $g(t)$ ($-\epsilon < t < \epsilon, \epsilon > 0$) be a 1-parameter C^∞ deformation of a Riemannian metric on M . Then, is there a deformation $g(t)$ such that $\text{Spec}(M, g(t)) = \text{Spec}(M, g(0))$ for every t ? Such a deformation is called an *isospectral deformation*.

First, we give some definitions to state the results of this article. The deformation $g(t)$ is called *trivial* if for each t there is a diffeomorphism $\eta(t)$ such that $g(t) = \eta(t)^*g(0)$ (the pull-back of $g(0)$ by $\eta(t)$). For a deformation $g(t)$ the symmetric covariant 2-tensor $h = g'(0)$ is called the *infinitesimal deformation* (*i-deformation*, for short) [2]. By Berger-Ebin [3] h is decomposed as

$$h = \tilde{h} + L_X g(0),$$

where $\nabla^i \tilde{h}_{ij} = 0$ (∇ being the connection induced by $g(0)$) and L_X is the Lie derivative with respect to X . The i-deformation h is called *trivial* if $\tilde{h} = 0$.

The main result of this article is the following.

THEOREM A: *There is no non-trivial isospectral i-deformation of a metric of flat torus.*

REMARK: Concerning the isospectral deformation of a metric of constant curvature, we can easily get the following: *There is no non-trivial isospectral deformation of a metric of constant curvature K if $2 \leq \dim M \leq 5$, or $\dim M = 6$ and $K > 0$.* This result is obtained by combining the results of spectral geometry [1], [4] and those concerning the non-deformability of a metric of constant curvature, the latter being directly derived from the results of Berger-Ebin [3], Mostow [5] and Koiso [2]. (See Tanaka [6], for the case $\dim M = 2$ and $K < 0$.) In the case of flat metrics Sunada [7] showed that there are only finitely many isometry classes of flat manifolds with a given spectrum.

The author wishes to thank Professor M. Ikeda for carefully reading the manuscript and offering valuable comments. Thanks are also due to Mr. N. Koiso, Osaka University, for helpful discussions.

2. Proof of Theorem A

The main part of the proof is to study the variation of the coefficients of Minakshisundaram's expansion:

$$\sum_{k=0}^{\infty} \exp(-\lambda_k s) \underset{s \rightarrow +0}{\sim} \left(\frac{1}{4\pi s}\right)^{n/2} \sum_{j=0}^{\infty} a_j s^j.$$

If $g(t)$ is an isospectral deformation, the coefficients $a_j(t)$ must be constants.

For a Riemannian metric g on M , $dV(g)$, R_{ikm}^i , R_{ij} and τ denote the volume element, the curvature tensor, the Ricci tensor and the scalar curvature, respectively. The coefficients a_j ($j = 0, 1, 2$) are given by

$$\begin{aligned} a_0 &= \text{vol}(M, g) = \int_M dV(g), \quad a_1 = \frac{1}{6} \int_M \tau dV(g), \\ a_2 &= \frac{1}{360} \int_M (2|R|^2 - 2|\rho|^2 + 5\tau^2) dV(g), \end{aligned}$$

where $|R|^2 = R_{ikm}R^{ikm}$ and $|\rho|^2 = R_{ij}R^{ij}$.

Let $g(t)$ be a deformation and $h(t) \equiv g'(t)$ the i-deformation of $g(t)$

at t . Let $a_i(t)$ be the Minakshisundaram's coefficients for $(M, g(t))$. Then the following formulas are obtained by straightforward calculation.

$$(2.1) \quad a'_0(t) = \frac{1}{2} \int_M h_s^s dV(g(t)),$$

$$(2.2) \quad a'_1(t) = \frac{1}{6} \int_M (\frac{1}{2} \tau h_s^s - R_{ij} h^{ij}) dV(g(t)),$$

$$(2.3) \quad a'_2(t) = \frac{1}{360} \int_M [12(\nabla_j \nabla_i \tau) h^{ji} - 6(\nabla_k \nabla^k R_{ji}) h^{ji} + 8R_{jk} R_i^k h^{ji} - 4R_{kjm} R^{km} h^{ji} - 4R_{jkm} R_i^{kms} h^{ji} + 9(\Delta \tau) h_s^s - 10\tau R_{ji} h^{ji} + |R|^2 h_s^s - |\rho|^2 h_s^s + \frac{5}{2} \tau^2 h_s^s] dV(g(t)).$$

Let (M, g_0) be a flat manifold. Then we have

LEMMA 2.1: *If $g(t)$ is an isospectral deformation with $g(0) = g_0$, and $\nabla^i h_{ij} = 0$ holds at $t = 0$, then*

$$(2.4a, b) \quad h_s^s = 0, \quad \nabla_k h_{ji} = 0$$

hold at $t = 0$.

PROOF: Starting from (2.3), we have by tedious calculation,

$$(2.5) \quad a''_2(0) = \frac{1}{120} \int_M [(\nabla_k \nabla^k h^{ji})(\nabla_m \nabla^m h_{ji}) + 3(\Delta h_s^s)^2] dV(g_0).$$

(Note that $\nabla^i h_{ij} = 0$ and $R_{jkm}^i = 0$ at $t = 0$.) From (2.1) and (2.5), $a'_0(0) = a''_2(0) = 0$ holds if and only if (2.4a, b) hold good. Q.E.D.

LEMMA 2.2: *Let $g(t)$ be as in Lemma 2.1. Then*

$$(2.6) \quad \int_M h_{ij} \phi (\nabla^i \nabla^j \phi) dV(g_0) = 0$$

holds for each eigenfunction ϕ of $\Delta_{g(0)}$.

PROOF: For the eigenvalue $\lambda_k(t)$ of $\Delta_{g(t)}$, the following was obtained by Berger [8]:

$$(2.7) \quad \lambda'_k(0) = \int_M [h_{ij} \phi (\nabla^i \nabla^j \phi) + (\nabla^i h_{ij} - \frac{1}{2} \nabla_j h_s^s) \phi (\nabla^i \phi)] dV(g_0),$$

where ϕ is the eigenfunction for $\lambda_k(0)$. Therefore, (2.6) follows from (2.4). Q.E.D.

Now, let us prove Theorem A. Let (M, g_0) be a flat torus given by R^n/L , where L is a lattice, i.e., a discrete abelian subgroup of the group of Euclidean motions in R^n . Let L^* denote the dual lattice, consisting of all $x \in R^n$ such that $(x, y) = \sum_{i=1}^n x^i y^i$ is an integer for all $y \in L$. Then the sets of eigenfunctions and eigenvalues are given by $\{\phi_x(y) = \cos 2\pi(x, y), \psi_x(y) = \sin 2\pi(x, y); x \in L^*\}$ and $\{4\pi^2(x, x); x \in L^*\}$, respectively. Recalling (2.4b), we see that h_{ij} are constants in the coordinates induced from R^n . Therefore, from (2.6) we have

$$4\pi^2 h_{ij} x^i x^j \int_{R^n/L} \{\cos 2\pi(x, y)\}^2 dy = 0, \quad x \in L^*,$$

hence $h_{ij} x^i x^j = (hx, x) = 0$ for $x \in L^*$. This leads to $h = 0$, because the set $\{x/\|x\|; x \in L^*\}$ is obviously dense in $\{x \in R^n; \|x\| = 1\}$. Q.E.D.

3. Conformal deformations

In this section we restrict our study to the conformal deformation,

$$(3.1) \quad g(t) = e^{2\rho(t)} g_0, \quad \text{with } \rho(0) \equiv 0.$$

Set $\sigma(t) = \rho'(t)$, and straightforward calculation gives

$$(3.2) \quad \begin{aligned} a_1''(t) &= \frac{1}{3}(n-1)(n-2) \int_M \sigma \Delta \sigma dV(g(t)) + \frac{1}{6}(n-2)^2 \int_M \tau \sigma^2 dV(g(t)) \\ &\quad + \frac{1}{6}(n-2) \int_M \tau \frac{\partial \sigma}{\partial t} dV(g(t)). \end{aligned}$$

LEMMA 3.1: *Assume that (M, g_0) has a constant scalar curvature. If $g(t)$ is a volume-preserving deformation, we have*

$$(3.3) \quad a_1''(0) = \frac{1}{3}(n-1)(n-2) \int_M \sigma \Delta \sigma dV(g_0) - \frac{1}{3}(n-2)\tau \int_M \sigma^2 dV(g_0).$$

PROOF: It is easy to see that $a_0''(0) = 0$ is written as

$$\int_M \frac{\partial \sigma}{\partial t} dV(g_0) + n \int_M \sigma^2 dV(g_0) = 0.$$

By this equation and $\tau = \text{const.}$, (3.2) is led to (3.3).

Q.E.D.

From now on, we assume $n = \dim M > 2$.

By virtue of the above lemma we have the following theorem.

THEOREM B: *Let (M, g_0) have a constant scalar curvature and λ_1 be the non-zero first eigenvalue of $\Delta_{g(0)}$. If*

$$(3.4) \quad \lambda_1 > \frac{\tau}{n-1}$$

holds, there is no conformal isospectral deformation of g_0 .

PROOF: Let $g(t)$ be a conformal isospectral deformation of g_0 . Since $a'_0(0) = 0$, we have $\int_M \sigma dV(g_0) = 0$. Therefore,

$$\int_M \sigma \Delta \sigma dV(g_0) \geq \lambda_1 \int_M \sigma^2 dV(g_0)$$

holds (see [1, p. 186], for example). Accordingly, if (3.4) holds, we have $a''_1(0) > 0$ unless $\sigma = 0$, i.e., $h = 0$. Q.E.D.

The condition (3.4) is obviously satisfied if $\tau \leq 0$. Further, as shown by Obata [9], it is also satisfied for an Einstein space not isometric with a sphere. In the case of a sphere, $\lambda_1 = \tau/(n-1)$, hence $a''_1(0) \geq 0$ holds. The equality holds only when σ is the eigenfunction for λ_1 , which is equivalent to $\nabla_i \nabla_j \sigma + \{\tau/n(n-1)\} \sigma g_{ij} = 0$ (see [9]). Therefore,

$$h_{ij} = 2\sigma g_{ij} = -\frac{2n(n-1)}{\tau} \nabla_i \nabla_j \sigma.$$

Thus h is trivial.

As a consequence we have the following theorem.

THEOREM C: *Suppose (M, g_0) is an Einstein space, or a space of non-positive constant scalar curvature. Then there is no non-trivial conformal isospectral i -deformation of g_0 .*

REMARK: In the case of $\dim M = 2$, it follows from the Gauss-Bonnet formula that the coefficient $a_1(t)$ is invariant.

REFERENCES

- [1] M. BERGER, P. GAUDUCHON and E. MAZET: *Le spectre d'une variété riemannienne*. Lecture Notes in Mathematics 194. Springer Verlag, 1971.
- [2] N. KOISO: Non-deformability of Einstein metrics, *Osaka J. Math.* 15 (1978) 419–433.
- [3] M. BERGER and D. EBIN: Some decompositions of the space of symmetric tensors on a Riemannian manifold. *J. Diff. Geom.* 3 (1969) 379–392.
- [4] S. TANNO: Eigenvalues of the Laplacian of Riemannian manifolds. *Tôhoku Math. Journ.* 25 (1973) 391–403.
- [5] G.D. MOSTOW: *Strong rigidity of locally symmetric spaces*. Princeton, 1973.
- [6] S. TANAKA: Selberg's trace formula and spectrum. *Osaka J. Math.* 3 (1966) 205–216.
- [7] T. SUNADA: Spectrum of a compact flat manifold (preprint).
- [8] M. BERGER: Sur les premières valeurs propres des variétés riemanniennes. *Compositio Math.* 26 (1973) 129–149.
- [9] M. OBATA: Certain conditions for a Riemannian manifold to be isometric with a sphere. *J. Math. Soc. Japan* 14 (1962) 333–340.

(Oblatum 21-III-1978 & 12-XII-1978)

Department of Applied Mathematics
and Physics
Faculty of Engineering
Kyoto University
Kyoto 606, Japan