

A-QUASICONVEXITY: RELAXATION AND HOMOGENIZATION

ANDREA BRAIDES¹, IRENE FONSECA² AND GIOVANNI LEONI³

Abstract. Integral representation of relaxed energies and of Γ -limits of functionals

$$(u, v) \mapsto \int_{\Omega} f(x, u(x), v(x)) dx$$

are obtained when sequences of fields v may develop oscillations and are constrained to satisfy a system of first order linear partial differential equations. This framework includes the treatment of divergence-free fields, Maxwell's equations in micromagnetics, and curl-free fields. In the latter case classical relaxation theorems in $W^{1,p}$ are recovered.

AMS Subject Classification. 35D99, 35E99, 49J45.

Received March 16, 2000. Revised September 26, 2000.

1. INTRODUCTION

In a recent paper Fonseca and Müller [22] have proved that \mathcal{A} -quasiconvexity is a necessary and sufficient condition for (sequential) lower semicontinuity of a functional

$$(u, v) \mapsto \int_{\Omega} f(x, u(x), v(x)) dx,$$

whenever $f : \Omega \times \mathbb{R}^m \times \mathbb{R}^d \rightarrow [0, \infty)$ is a Carathéodory integrand satisfying

$$0 \leq f(x, u, v) \leq a(x, u) (1 + |v|^q),$$

for a.e. $x \in \Omega$ and all $(u, v) \in \mathbb{R}^m \times \mathbb{R}^d$, where $1 \leq q < \infty$, $a \in L_{\text{loc}}^{\infty}(\Omega \times \mathbb{R}; [0, \infty))$, $\Omega \subset \mathbb{R}^N$ is open, bounded, $u_n \rightarrow u$ in measure, $v_n \rightarrow v$ in $L^q(\Omega; \mathbb{R}^d)$ and $\mathcal{A}v_n \rightarrow 0$ in $W^{-1,q}(\Omega; \mathbb{R}^l)$ (see also [14]). Here, and in what follows, following [32]

$$\mathcal{A} : L^q(\Omega; \mathbb{R}^d) \rightarrow W^{-1,q}(\Omega; \mathbb{R}^l), \quad \mathcal{A}v := \sum_{i=1}^N A^{(i)} \frac{\partial v}{\partial x_i},$$

Keywords and phrases: \mathcal{A} -quasiconvexity, equi-integrability, Young measure, relaxation, Γ -convergence, homogenization.

¹ SISSA, Trieste, Italy; e-mail: braides@sisssa.it

² Department of Mathematical Sciences, Carnegie-Mellon University, Pittsburgh, PA, U.S.A.; e-mail: fonseca@andrew.cmu.edu

³ Dipartimento di Scienze e Tecnologie Avanzate, Università del Piemonte Orientale, Alessandria, Italy; e-mail: leoni@al.unipmn.it

is a constant-rank, first order linear partial differential operator, with $A^{(i)} : \mathbb{R}^d \rightarrow \mathbb{R}^l$ linear transformations, $i = 1, \dots, N$. We recall that \mathcal{A} satisfies the *constant-rank* property if there exists $r \in \mathbb{N}$ such that

$$\text{rank } \mathbb{A}w = r \quad \text{for all } w \in S^{N-1}, \tag{1.1}$$

where

$$\mathbb{A}w := \sum_{i=1}^N w_i A^{(i)}, \quad w \in \mathbb{R}^N.$$

A function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is said to be \mathcal{A} -*quasiconvex* if

$$f(v) \leq \int_Q f(v + w(y)) \, dy$$

for all $v \in \mathbb{R}^d$ and all $w \in C_{1\text{-per}}^\infty(\mathbb{R}^N; \mathbb{R}^d)$ such that $\mathcal{A}w = 0$ and $\int_Q w(y) \, dy = 0$. Here Q denotes the unit cube in \mathbb{R}^N , and the space $C_{1\text{-per}}^\infty(\mathbb{R}^N; \mathbb{R}^d)$ is introduced in Section 2.

The relevance of this general framework, as emphasized by Tartar (see [32, 34–39]), lies on the fact that in continuum mechanics and electromagnetism PDEs other than $\text{curl } v = 0$ arise naturally, and this calls for a relaxation theory which encompasses PDE constraints of the type $\mathcal{A}v = 0$. Some important examples included in this general setting are given by:

(a) [Unconstrained Fields]

$$\mathcal{A}v \equiv 0.$$

Here, due to Jensen’s inequality \mathcal{A} -quasiconvexity reduces to convexity.

(b) [Divergence Free Fields]

$$\mathcal{A}v = 0 \quad \text{if and only if } \text{div } v = 0,$$

where $v : \Omega \subset \mathbb{R}^N \rightarrow \mathbb{R}^N$ (see [33]).

(c) [Magnetostatics Equations]

$$\mathcal{A} \begin{pmatrix} m \\ h \end{pmatrix} := \begin{pmatrix} \text{div}(m + h) \\ \text{curl } h \end{pmatrix} = 0,$$

where $m : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is the *magnetization* and $h : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is the *induced magnetic field* (see [17, 38]); often these are also called Maxwell’s Equations in the micromagnetics literature.

(d) [Gradients]

$$\mathcal{A}v = 0 \quad \text{if and only if } \text{curl } v = 0.$$

Note that $w \in C_{1\text{-per}}^\infty(\mathbb{R}^N; \mathbb{R}^d)$ is such that $\text{curl } w = 0$ and $\int_Q w(y) \, dy = 0$ if and only if there exists $\varphi \in C_{1\text{-per}}^\infty(\mathbb{R}^N; \mathbb{R}^n)$ such that $\nabla \varphi = v$, where $d = n \times N$. Thus in this case we recover the well-known notion of *quasiconvexity* introduced by Morrey [30].

(e) [Higher Order Gradients]

Replacing the target space \mathbb{R}^d by an appropriate finite dimensional vector space E_s^n , it is possible to find a first order linear partial differential operator \mathcal{A} such that $v \in L^p(\Omega; E_s^n)$ and $\mathcal{A}v = 0$ if and only if there exists $\varphi \in W^{s,q}(\Omega; \mathbb{R}^n)$ such that $v = \nabla^s \varphi$ (see Th. 1.3).

This paper is divided into two parts. In the first part we give an integral representation formula for the relaxed energy in the context of \mathcal{A} -quasiconvexity. Precisely, let $1 \leq p < \infty$ and $1 < q < \infty$, and consider the functional

$$F : L^p(\Omega; \mathbb{R}^m) \times L^q(\Omega; \mathbb{R}^d) \times \mathcal{O}(\Omega) \rightarrow [0, \infty)$$

defined by

$$F((u, v); D) := \int_D f(x, u(x), v(x)) \, dx,$$

where $\mathcal{O}(\Omega)$ is the collection of all open subsets of Ω , and the density f satisfies the following hypothesis:

(H) $f : \Omega \times \mathbb{R}^m \times \mathbb{R}^d \rightarrow [0, \infty)$ is Carathéodory function satisfying

$$0 \leq f(x, u, v) \leq C(1 + |u|^p + |v|^q)$$

for a.e. $x \in \Omega$ and all $(u, v) \in \mathbb{R}^m \times \mathbb{R}^d$, and where $C > 0$.

For $D \in \mathcal{O}(\Omega)$ and $(u, v) \in L^p(\Omega; \mathbb{R}^m) \times (L^q(\Omega; \mathbb{R}^d) \cap \ker \mathcal{A})$ define

$$\mathcal{F}((u, v); D) := \inf \left\{ \liminf_{n \rightarrow \infty} F((u_n, v_n); D) : (u_n, v_n) \in L^p(D; \mathbb{R}^m) \times L^q(D; \mathbb{R}^d), \right. \\ \left. u_n \rightarrow u \text{ in } L^p(D; \mathbb{R}^m), \quad v_n \rightharpoonup v \text{ in } L^q(D; \mathbb{R}^d), \quad \mathcal{A}v_n \rightarrow 0 \text{ in } W^{-1,q}(D; \mathbb{R}^l) \right\}. \tag{1.2}$$

It turns out that the condition $\mathcal{A}v_n \rightarrow 0$ imposed in (1.2) may be replaced by requiring that v_n do satisfy the homogeneous PDE $\mathcal{A}v = 0$. Precisely, and in view of Lemma 3.1 and Corollary 3.2 below, it can be shown that

$$\mathcal{F}((u, v); D) = \inf \left\{ \liminf_{n \rightarrow \infty} F((u, v_n); D) : v_n \in L^q(D; \mathbb{R}^d), v_n \rightharpoonup v \text{ in } L^q(D; \mathbb{R}^d), \mathcal{A}v_n = 0 \right\},$$

and thus

$$\mathcal{F}((u, v); D) = \inf \left\{ \liminf_{n \rightarrow \infty} F((u_n, v_n); D) : (u_n, v_n) \in L^p(D; \mathbb{R}^m) \times L^q(D; \mathbb{R}^d), \right. \\ \left. u_n \rightarrow u \text{ in } L^p(D; \mathbb{R}^m), \quad v_n \rightharpoonup v \text{ in } L^q(D; \mathbb{R}^d), \quad \mathcal{A}v_n = 0 \right\} =: \mathcal{F}_0((u, v); D). \tag{1.3}$$

The first main result of the paper is given by the following theorem:

Theorem 1.1. *Under condition (H) and the constant-rank hypothesis (1.1), for all $D \in \mathcal{O}(\Omega)$, $u \in L^p(\Omega; \mathbb{R}^m)$, and $v \in L^q(\Omega; \mathbb{R}^d) \cap \ker \mathcal{A}$, we have*

$$\mathcal{F}((u, v); D) = \int_D \mathcal{Q}_{\mathcal{A}}f(x, u(x), v(x)) dx$$

where, for each fixed $(x, u) \in \Omega \times \mathbb{R}^m$, the function $\mathcal{Q}_{\mathcal{A}}f(x, u, \cdot)$ is the \mathcal{A} -quasiconvexification of $f(x, u, \cdot)$, namely

$$\mathcal{Q}_{\mathcal{A}}f(x, u, v) := \inf \left\{ \int_Q f(x, u, v + w(y)) dy : w \in C_{1\text{-per}}^\infty(\mathbb{R}^N; \mathbb{R}^d) \cap \ker \mathcal{A}, \int_Q w(y) dy = 0 \right\}$$

for all $v \in \mathbb{R}^d$.

Remarks 1.2. (i) Note that in the degenerate case where $\mathcal{A} = 0$, \mathcal{A} -quasiconvex functions are convex and Theorem 1.1 together with condition (1.4) yield a convex relaxation result with respect to $L^p \times L^q$ (weak) convergence. See the monograph of Buttazzo [12] for related results in this context.

(ii) If the function f also satisfies a growth condition of order q from below in the variable v , that is

$$f(x, u, v) \geq \frac{1}{C}|v|^q - C \tag{1.4}$$

for a.e. $x \in \Omega$ and all $(u, v) \in \mathbb{R}^m \times \mathbb{R}^d$, then a simple diagonalization argument shows that $(u, v) \mapsto \mathcal{F}((u, v); D)$ is $L^p \times (L^q\text{-weak})$ lower semicontinuous, *i.e.*,

$$\int_D \mathcal{Q}_{\mathcal{A}}f(x, u(x), v(x)) \, dx \leq \liminf_{n \rightarrow \infty} \int_D \mathcal{Q}_{\mathcal{A}}f(x, u_n(x), v_n(x)) \, dx \tag{1.5}$$

whenever $u_n \in L^p(\Omega; \mathbb{R}^m)$, $v_n \in L^q(\Omega; \mathbb{R}^d) \cap \ker \mathcal{A}$, $u_n \rightarrow u$ in $L^p(D; \mathbb{R}^m)$, $v_n \rightarrow v$ in $L^q(D; \mathbb{R}^d)$. In particular $\mathcal{Q}_{\mathcal{A}}f$ is \mathcal{A} -quasiconvex if f is continuous and

$$\frac{1}{C}|v|^q - C \leq f(v) \leq C(1 + |v|^q)$$

for some $C > 0$, and all $v \in \mathbb{R}^d$ (see the proof of Cor. 5.7).

The lower semicontinuity result (1.5) is not covered by Theorem 3.7 in [22], where it is assumed that the integrand be \mathcal{A} -quasiconvex and *continuous* in the v variable. However, as remarked in [22], in the realm of general \mathcal{A} -quasiconvexity the function $\mathcal{Q}_{\mathcal{A}}f(x, u, \cdot)$ may not be continuous, even if $f(x, u, \cdot)$ is. Indeed in the degenerate case $\ker \mathcal{A} = \{0\}$ all functions are \mathcal{A} -quasiconvex. Also, when $N = 1$, $d = 2$, and $v = (v_1, v_2)$, consider

$$\mathcal{A}v := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} v'_1 \\ v'_2 \end{pmatrix}.$$

Then for $w \in \mathbb{R}$

$$\mathbb{A}w = \begin{pmatrix} 0 & w \\ w & 0 \end{pmatrix}$$

and thus when $|w| = 1$ the matrix $\mathbb{A}w$ has constant rank 1. For any given function $f(v)$ the \mathcal{A} -quasiconvex envelope of f is obtained by convexification in the first component, so that by considering *e.g.* (cf. [22, 28])

$$f_1(v) := e^{-|v_1|v_2^2}, \quad f_2(v) := (1 + |v_1|)^{|v_2|},$$

one gets

$$\mathcal{Q}_{\mathcal{A}}f_1(v) = \begin{cases} 0 & \text{if } v_2 \neq 0 \\ 1 & \text{if } v_2 = 0 \end{cases}, \quad \mathcal{Q}_{\mathcal{A}}f_2(v) = \begin{cases} (1 + |v_1|)^{|v_2|} & \text{if } |v_2| \geq 1 \\ 1 & \text{if } |v_2| < 1. \end{cases}$$

(iii) The continuity of f with respect to v is essential to ensure the representation of \mathcal{F} provided in Theorem 1.1, in contrast with the case where $\mathcal{A}v = 0$ if and only if $\text{curl } v = 0$. In fact, if $f : \mathbb{R}^{n \times N} \rightarrow [0, \infty)$ is a Borel function satisfying the growth condition

$$0 \leq f(v) \leq C(1 + |v|^q)$$

for $C > 0$, $1 \leq q < \infty$, $v \in \mathbb{R}^{n \times N}$, then it can be shown easily that

$$\mathcal{F}(w; D) = \int_D \mathcal{Q}f(\nabla w(x)) \, dx \tag{1.6}$$

for all $D \in \mathcal{O}(\Omega)$, $w \in W^{1,q}(\Omega; \mathbb{R}^n)$, where $\mathcal{Q}f$ is the quasiconvex envelope of f . Indeed, $\mathcal{Q}f$ is a (continuous) quasiconvex function satisfying (H) (see [18], [8] Th. 4.3); therefore by Theorem 1.1

$$w \mapsto \int_D \mathcal{Q}f(\nabla w(x)) \, dx$$

is $W^{1,q}$ -sequentially weakly lower semicontinuous, and so

$$\int_D \mathcal{Q}f(\nabla w(x)) \, dx \leq \mathcal{F}(w; D).$$

Conversely, under hypothesis (H) it is known that $\mathcal{F}(v; \cdot)$ admits an integral representation (see Th. 9.1 in [10], Th. 20.1 in [15])

$$\mathcal{F}(w; D) = \int_D \varphi(\nabla w(x)) \, dx,$$

where φ is a quasiconvex function, and $\varphi(v) \leq f(v)$ for all $v \in \mathbb{R}^{n \times N}$. Hence $\varphi \leq Qf$ and we conclude that (1.6) holds.

For general constant-rank operators \mathcal{A} , and if f is not continuous with respect to v , it may happen that $\mathcal{F}_0((u, v); \cdot)$ is not even the trace of a Radon measure in $\mathcal{O}(\Omega)$ and thus (1.3) fails. As an example, consider $d = 2$, $N = 1$, $\Omega := (0, 1)$, $v = (v_1, v_2)$, and let $\mathcal{A}(v) = 0$ if and only if $v_2' = 0$ as in (ii) above. Let

$$f(v) := \begin{cases} (v_1 - 1)^2 + v_2^2, & \text{if } v_2 \in \mathbb{Q} \\ (v_1 + 1)^2 + v_2^2, & \text{if } v_2 \notin \mathbb{Q}. \end{cases}$$

Although f satisfies a quadratic growth condition of the type (H), and (A_3) holds with $q = 2$, it is easy to see that for all intervals $(a, b) \subset (0, 1)$,

$$\mathcal{F}_0((u, v); (a, b)) = \mathcal{F}_0(v; (a, b)) = \min \left\{ \int_a^b ((v_1 - 1)^2 + v_2^2) dx, \int_a^b ((v_1 + 1)^2 + v_2^2) dx \right\}$$

which is not the trace of a Radon measure on $\mathcal{O}(\Omega)$. On the other hand, it may be shown that (see the Appendix below for a proof)

$$\mathcal{F}((u, v); (a, b)) = \mathcal{F}(v; (a, b)) = \int_a^b (\psi^{**}(v_1) + v_2^2) \, dx,$$

where $\psi^{**}(v_1)$ is the convex envelope of

$$\psi(v_1) := \min \{ (v_1 - 1)^2, (v_1 + 1)^2 \}.$$

(iv) Using the growth condition (H), a mollification argument, and the linearity of \mathcal{A} , it can be shown that (see Rem. 3.3 in [22])

$$\mathcal{Q}_{\mathcal{A}} f(x, u, v) = \inf \left\{ \int_Q f(x, u, v + w(y)) \, dy : w \in L^q_{1\text{-per}}(\mathbb{R}^N; \mathbb{R}^d) \cap \ker \mathcal{A}, \int_Q w(y) \, dy = 0 \right\}.$$

We write $w \in L^q_{1\text{-per}}(\mathbb{R}^N; \mathbb{R}^d) \cap \ker \mathcal{A}$ when $w \in L^q_{1\text{-per}}(\mathbb{R}^N; \mathbb{R}^d)$ and $\mathcal{A}w = 0$ in $W^{-1,q}(Q; \mathbb{R}^l)$.

(v) We may also treat the cases $q = 1, \infty$ and $p = \infty$. See Theorem 3.6 below.

The proof of Theorem 1.1 relies heavily on the use of Young measures (see [5, 40]). However, instead of applying directly the arguments of Fonseca and Müller [22] (based on Balder’s [4] and Kristensen’s [26] approach in the curl-free case), we use these together with the blow-up method introduced by Fonseca and Müller in [20].

Although in Theorem 1.1 the functions u and v are not related to each other, the arguments of the proof work equally well when u and v are not independent. Indeed as a corollary, we can prove the following two theorems:

Theorem 1.3. *Let $1 \leq p \leq \infty$, $s \in \mathbb{N}$, and suppose that $f : \Omega \times E^n_{[s-1]} \times E^n_s \rightarrow [0, \infty)$ is a Carathéodory function satisfying*

$$0 \leq f(x, \mathbf{u}, v) \leq C(1 + |\mathbf{u}|^p + |v|^p), \quad 1 \leq p < \infty,$$

for a.e. $x \in \Omega$ and all $(\mathbf{u}, v) \in E^n_{[s-1]} \times E^n_s$, where $C > 0$, and

$$\chi_{\Omega} f \in L^{\infty}_{\text{loc}}(\mathbb{R}^N \times E^n_{[s-1]} \times E^n_s; [0, \infty)) \quad \text{if } p = \infty.$$

Then for every $u \in W^{s,p}(\Omega; \mathbb{R}^n)$ we have

$$\int_{\Omega} \mathcal{Q}^s f(x, u, \dots, \nabla^s u) dx = \inf \left\{ \liminf_{k \rightarrow \infty} \int_{\Omega} f(x, u_k, \dots, \nabla^s u_k) dx : \{u_k\} \subset W^{s,p}(\Omega; \mathbb{R}^n), \right. \\ \left. u_k \rightharpoonup u \text{ in } W^{s,p}(\Omega; \mathbb{R}^n) \quad (\star \text{ if } p = \infty) \right\},$$

where, for a.e. $x \in \Omega$ and all $(\mathbf{u}, v) \in E_{[s-1]}^n \times E_s^n$,

$$\mathcal{Q}^s f(x, \mathbf{u}, v) := \inf \left\{ \int_Q f(x, \mathbf{u}, v + \nabla^s w(y)) dy : w \in C_{1\text{-per}}^\infty(\mathbb{R}^N; \mathbb{R}^n) \right\}.$$

Remarks 1.4. (i) Here E_s^n stands for the space of n -tuples of symmetric s -linear maps on \mathbb{R}^N ,

$$E_{[s-1]}^n := \mathbb{R}^n \times E_1^n \times \dots \times E_{s-1}^n,$$

and

$$\nabla^l u := \left(\frac{\partial^l u}{\partial x_1^{\alpha_1} \dots \partial x_N^{\alpha_N}} \right)_{\alpha_1 + \dots + \alpha_N = l}, \quad l \geq 1.$$

(ii) When $s = 1$ we recover classical relaxation results (see *e.g.* the work of Acerbi and Fusco [1], Dacorogna [13], Marcellini and Sbordone [28] and the references contained therein).

When $s > 1$ lower semicontinuity results related to Theorem 1.3 are due to Meyers [29], Fusco [23] and Guidorzi and Poggiolini [25], while we are not aware of any integral representation formula for the relaxed energy, when the integrand depends on the full set of variables, that is $f = f(x, u, \dots, \nabla^s u)$. This is due to the fact that classical truncation methods for $s = 1$ cannot be extended in a simple way to truncate higher order derivatives. The results of Fonseca and Müller (see the proof of Lem. 2.15 in [22]), where the truncation is only on the highest order derivative $\nabla^s u$, and Corollary 3.2 below, allows us to overcome this difficulty. Note however that this technique relies heavily on p -equi-integrability, and thus cannot work in the case $p = 1$, if one replaces weak convergence in $W^{s,1}(\Omega; \mathbb{R}^n)$ with the natural convergence, which is strong convergence in $W^{s-1,1}(\Omega; \mathbb{R}^n)$. In this context, a relaxation result has been given by Amar and De Cicco [2], but only when $f = f(\nabla^s u)$, so that truncation is not needed. The general case where f depends also on lower order derivatives has been addressed by Fonseca *et al.* [19].

Theorem 1.5. Let $1 \leq p \leq \infty$, let $\Omega \subset \mathbb{R}^N$ be an open, bounded, connected set, and suppose that $f : \Omega \times \mathbb{R}^N \times \mathbb{R}^{N^2} \rightarrow [0, \infty)$ is a Carathéodory function satisfying

$$0 \leq f(x, u, v) \leq C(1 + |u|^p + |v|^p), \quad 1 \leq p < \infty,$$

for a.e. $x \in \Omega$ and all $(u, v) \in \mathbb{R}^N \times \mathbb{R}^{N^2}$, where $C > 0$, and

$$\chi_\Omega f \in L_{\text{loc}}^\infty(\mathbb{R}^N \times \mathbb{R}^N \times \mathbb{R}^{N^2}; [0, \infty)) \quad \text{if } p = \infty.$$

Then for every $u \in W^{1,p}(\Omega; \mathbb{R}^N)$ such that $\text{div } u = 0$, we have

$$\int_{\Omega} \bar{f}(x, u(x), \nabla u(x)) dx = \inf \left\{ \liminf_{n \rightarrow \infty} \int_{\Omega} f(x, u_n(x), \nabla u_n(x)) dx : \{u_n\} \subset W^{1,p}(\Omega; \mathbb{R}^N), \right. \\ \left. \text{div } u_n = 0, \quad u_n \rightharpoonup u \text{ in } W^{1,p}(\Omega; \mathbb{R}^N) \quad (\star \text{ if } p = \infty) \right\}, \tag{1.7}$$

where, for a.e. $x \in \Omega$ and all $(u, v) \in \mathbb{R}^N \times \mathbb{R}^{N^2}$,

$$\bar{f}(x, u, v) := \inf \left\{ \int_Q f(x, u, v + \nabla w(y)) \, dy : w \in C_{1\text{-per}}^\infty(\mathbb{R}^N; \mathbb{R}^N), \operatorname{div} w = 0 \right\}.$$

Remark 1.6. To the authors' knowledge, this result is new in this generality (for a different proof, with additional smoothness assumptions, see [9]). A related problem was addressed by Dal Maso *et al.* in [16], where it was shown that the Γ -limit of a family of functionals of the type (1.7) may be non local if (H) is violated.

In the second part of the paper we present (Γ -convergence) homogenization results for periodic integrands in the context of \mathcal{A} -quasiconvexity. Let $\varepsilon > 0$ and $1 < q < \infty$, and consider a family of functionals

$$\mathcal{F}_\varepsilon : (L^q(\Omega; \mathbb{R}^d) \cap \ker \mathcal{A}) \times \mathcal{O}(\Omega) \rightarrow [0, \infty)$$

defined by

$$\mathcal{F}_\varepsilon(v; D) := \int_D f\left(\frac{x}{\varepsilon}, v(x)\right) \, dx,$$

where the density f satisfies the following hypotheses:

(A₁) $f : \mathbb{R}^N \times \mathbb{R}^d \rightarrow [0, \infty)$ is a continuous function, Q -periodic in the first argument, that is $f(x + e_i, v) = f(x, v)$ for every $i = 1, \dots, N$, where e_i are the elements of the canonical basis of \mathbb{R}^N ;

(A₂) there exists $C > 0$ such that

$$0 \leq f(x, v) \leq C(1 + |v|^q)$$

for all $(x, v) \in \mathbb{R}^N \times \mathbb{R}^d$;

(A₃) there exists $C > 0$ such that

$$f(x, v) \geq \frac{1}{C}|v|^q - C$$

for all $(x, v) \in \mathbb{R}^N \times \mathbb{R}^d$.

Let $\varepsilon_n \rightarrow 0^+$. We say that a functional

$$\mathcal{J} : (L^q(\Omega; \mathbb{R}^d) \cap \ker \mathcal{A}) \times \mathcal{O}(\Omega) \rightarrow [0, +\infty]$$

is the Γ -lim inf (resp. Γ -lim sup) of the sequence of functionals $\{\mathcal{F}_{\varepsilon_n}\}$ with respect to the weak convergence in $L^q(\Omega; \mathbb{R}^d)$ if for every $v \in L^q(\Omega; \mathbb{R}^d) \cap \ker \mathcal{A}$

$$\mathcal{J}(v; \Omega) = \inf \left\{ \liminf_{n \rightarrow \infty} \text{ (resp. } \limsup_{n \rightarrow \infty} \text{)} \mathcal{F}_{\varepsilon_n}(v_n; \Omega) : v_n \in L^q(\Omega; \mathbb{R}^d) \cap \ker \mathcal{A}, \right. \\ \left. v_n \rightharpoonup v \text{ in } L^q(\Omega; \mathbb{R}^d) \right\}, \tag{1.8}$$

and we write

$$\mathcal{J} = \Gamma - \liminf_{n \rightarrow \infty} \mathcal{F}_{\varepsilon_n} \left(\text{resp. } \mathcal{J} = \Gamma - \limsup_{n \rightarrow \infty} \mathcal{F}_{\varepsilon_n} \right).$$

When finite energy sequences are L^q -equibounded then the infimum in the definition of Γ -lim inf (resp. Γ -lim sup) is attained. We say that the sequence $\{\mathcal{F}_{\varepsilon_n}\}$ Γ -converges to \mathcal{J} if the Γ -lim inf and Γ -lim sup coincide, and we write

$$\mathcal{J} = \Gamma - \lim_{n \rightarrow \infty} \mathcal{F}_{\varepsilon_n}.$$

The functional \mathcal{J} is said to be the $\Gamma - \lim \inf$ (resp. $\Gamma - \lim \sup$) of the *family* of functionals $\{\mathcal{F}_\varepsilon\}$ with respect to the weak convergence in $L^q(\Omega; \mathbb{R}^d)$ if for *every* sequence $\varepsilon_n \rightarrow 0^+$ we have that

$$\mathcal{J} = \Gamma - \lim \inf_{n \rightarrow \infty} \mathcal{F}_{\varepsilon_n} \quad \left(\text{resp. } \mathcal{J} = \Gamma - \lim \sup_{n \rightarrow \infty} \mathcal{F}_{\varepsilon_n} \right),$$

and we write

$$\mathcal{J} = \Gamma - \lim \inf_{\varepsilon \rightarrow 0} \mathcal{F}_\varepsilon \quad \left(\text{resp. } \mathcal{J} = \Gamma - \lim \sup_{\varepsilon \rightarrow 0} \mathcal{F}_\varepsilon \right).$$

Finally, we say that \mathcal{J} is the Γ -*limit* of the *family* of functionals $\{\mathcal{F}_\varepsilon\}$, and we write

$$\mathcal{J} = \Gamma - \lim_{n \rightarrow \infty} \mathcal{F}_{\varepsilon_n},$$

if $\Gamma - \lim \inf$ and $\Gamma - \lim \sup$ coincide.

In the sequel we will also consider functionals \mathcal{J} given by (1.8) where we replace the weak convergence $v_n \rightharpoonup v$ with the convergence $v_n \rightarrow v$ with respect to some metric d . In order to highlight this dependence on the metric d these functionals will be denoted as

$$\mathcal{J} = \Gamma(d) - \lim \inf_{n \rightarrow \infty} \mathcal{F}_{\varepsilon_n} \quad \left(\text{resp. } \mathcal{J} = \Gamma(d) - \lim \sup_{n \rightarrow \infty} \mathcal{F}_{\varepsilon_n} \right),$$

as it is customary (see [10, 15]).

Theorem 1.7. *Under hypotheses $(A_1) - (A_2)$ and the constant-rank hypothesis (1.1),*

$$\mathcal{F}_{\text{hom}} = \Gamma - \lim \inf_{\varepsilon \rightarrow 0} \mathcal{F}_\varepsilon,$$

where

$$\mathcal{F}_{\text{hom}}(v; D) := \int_D f_{\text{hom}}(v) \, dx$$

for all $v \in L^q(\Omega; \mathbb{R}^d) \cap \ker \mathcal{A}$ and $D \in \mathcal{O}(\Omega)$, and

$$f_{\text{hom}}(v) := \inf_{k \in \mathbb{N}} \frac{1}{k^N} \inf \left\{ \int_{kQ} f(x, v + w(x)) \, dx : w \in L^q_{k\text{-per}}(\mathbb{R}^N; \mathbb{R}^d) \cap \ker \mathcal{A}, \int_{kQ} w(x) \, dx = 0 \right\} \quad (1.9)$$

for all $v \in \mathbb{R}^d$. Moreover, if (A_3) holds then

$$\mathcal{F}_{\text{hom}} = \Gamma - \lim_{\varepsilon \rightarrow 0} \mathcal{F}_\varepsilon.$$

For the definition of the space $L^q_{k\text{-per}}(\mathbb{R}^N; \mathbb{R}^d)$, we direct the reader to Section 2.

Remarks 1.8. (i) Using the growth condition (A_2) , a mollification argument, and the linearity of \mathcal{A} , it can be shown that

$$f_{\text{hom}}(v) = \inf_{k \in \mathbb{N}} \frac{1}{k^N} \inf \left\{ \int_{kQ} f(x, v + w(x)) \, dx : w \in L^\infty_{k\text{-per}}(\mathbb{R}^N; \mathbb{R}^d) \cap \ker \mathcal{A}, \int_{kQ} w(x) \, dx = 0 \right\}.$$

See also Corollary 5.7 below.

(ii) When f satisfies the q -Lipschitz condition

$$|f(x, v_1) - f(x, v_2)| \leq C(|v_1|^{q-1} + |v_2|^{q-1} + 1)|v_1 - v_2| \quad (1.10)$$

for all $x \in \mathbb{R}^N$, $v_1, v_2 \in \mathbb{R}^d$, and for some $C > 0$, then the continuity of $f(\cdot, v)$ can be weakened to measurability, namely f can be assumed to be simply Carathéodory. Note that (1.10) is not restrictive when $\mathcal{A} = \text{curl}$, that is when $v = \nabla u$ for some $u \in W^{1,q}(\Omega; \mathbb{R}^m)$, $d = N \times m$. Indeed, in this case in the definition of Γ -convergence we may replace the weak convergence of the gradients in $L^q(\Omega; \mathbb{R}^d)$ with the strong convergence in $L^q(\Omega; \mathbb{R}^m)$ of the potentials normalized to have zero average over Ω , and thus

$$\begin{aligned} \Gamma - \lim_{\varepsilon \rightarrow 0} \int_{\Omega} f\left(\frac{x}{\varepsilon}, v(x)\right) dx &= \Gamma(L^q(\Omega; \mathbb{R}^m)) - \lim_{\varepsilon \rightarrow 0} \int_{\Omega} f\left(\frac{x}{\varepsilon}, \nabla u(x)\right) dx \\ &= \Gamma(L^q(\Omega; \mathbb{R}^m)) - \lim_{\varepsilon \rightarrow 0} \int_{\Omega} \mathcal{Q}f\left(\frac{x}{\varepsilon}, \nabla u(x)\right) dx, \end{aligned}$$

by Proposition 7.13 in [10]. As shown in [27], if $f(x, v)$ is a Borel function which satisfies the growth condition (A_2) then its quasiconvex envelope $\mathcal{Q}f$ satisfies (1.10).

A similar argument fails for general \mathcal{A} -quasiconvexity, since the function $\mathcal{Q}_{\mathcal{A}}f(x, \cdot)$ may not even be continuous, see Remark 1.2(i) above.

In Section 2 we collect preliminary results on Young measures and Γ -convergence. The general relaxation results (see Th. 1.1 and its extension Th. 3.6) are proved in Section 3, and Section 4 is devoted to the applications of the general relaxation principle to Theorems 1.3 and 1.5. Finally, in Section 5 we address homogenization of functionals of \mathcal{A} -constrained vector fields.

2. PRELIMINARIES

We start with some notation. Here Ω is an open, bounded subset of \mathbb{R}^N , \mathcal{L}^N is the N dimensional Lebesgue measure, $S^{N-1} := \{x \in \mathbb{R}^N : |x| = 1\}$ is the unit sphere, and $Q := (-1/2, 1/2)^N$ the unit cube centered at the origin. We set $Q(x_0, \varepsilon) := x_0 + \varepsilon Q$ for $\varepsilon > 0$ and $x_0 \in \mathbb{R}^N$. A function $w \in L^q_{\text{loc}}(\mathbb{R}^N; \mathbb{R}^d)$ is said to be Q -periodic if $w(x + e_i) = w(x)$ for a.e. all $x \in \mathbb{R}^N$ and every $i = 1, \dots, N$, where (e_1, \dots, e_N) is the canonical basis of \mathbb{R}^N . We write $w \in L^q_{1\text{-per}}(\mathbb{R}^N; \mathbb{R}^d)$. More generally, $w \in L^q_{\text{loc}}(\mathbb{R}^N; \mathbb{R}^d)$ is said to be kQ -periodic, $k \in \mathbb{N}$, if $w(k \cdot)$ is Q -periodic. We write $w \in L^q_{k\text{-per}}(\mathbb{R}^N; \mathbb{R}^d)$. Also $C^\infty_{1\text{-per}}(\mathbb{R}^N; \mathbb{R}^d)$ will stand for the space of Q -periodic functions in $C^\infty(\mathbb{R}^N; \mathbb{R}^d)$.

We recall briefly some facts about Young measures which will be useful in the sequel (see e.g. [5, 33]). If D is an open set (not necessarily bounded), we denote by $C_c(D; \mathbb{R}^d)$ the set of continuous functions with compact support in D , endowed with the supremum norm. The dual of the closure of $C_c(D; \mathbb{R}^d)$ may be identified with the set of \mathbb{R}^d -valued Radon measures with finite mass $\mathcal{M}(D; \mathbb{R}^d)$, through the duality

$$\langle \nu, f \rangle := \int_D f(y) d\nu(y), \quad \nu \in \mathcal{M}(D; \mathbb{R}^d), \quad f \in C_c(D; \mathbb{R}^d).$$

A map $\nu : \Omega \rightarrow \mathcal{M}(D; \mathbb{R}^d)$ is said to be *weak-* measurable* if $x \mapsto \langle \nu_x, f \rangle$ are measurable for all $f \in C_c(D; \mathbb{R}^d)$.

The following result is a corollary of the Fundamental Theorem on Young Measures (see [5, 7, 34])

Theorem 2.1. *Let $z_n : \Omega \rightarrow \mathbb{R}^d$ be measurable functions such that*

$$\sup_{n \in \mathbb{N}} \int_{\Omega} |z_n|^q dx < \infty,$$

for some $q > 0$. Then there exists a subsequence $\{z_{n_k}\}$ of $\{z_n\}$ and a weak- measurable map $\nu : \Omega \rightarrow \mathcal{M}(\mathbb{R}^d; \mathbb{R}^d)$ such that*

- (i) $\nu_x \geq 0$, $\|\nu_x\|_{\mathcal{M}} = \int_{\mathbb{R}^d} d\nu_x = 1$ for a.e. $x \in \Omega$;

(ii) if $f : \Omega \times \mathbb{R}^d \rightarrow \mathbb{R}$ is a normal function bounded from below then

$$\liminf_{k \rightarrow \infty} \int_{\Omega} f(x, z_{n_k}(x)) dx \geq \int_{\Omega} \bar{f}(x) dx < \infty,$$

where

$$\bar{f}(x) := \langle \nu_x, f(x, \cdot) \rangle = \int_{\mathbb{R}^d} f(x, y) d\nu_x(y);$$

(iii) for any Carathéodory function $f : \Omega \times \mathbb{R}^d \rightarrow \mathbb{R}$ bounded from below one has

$$\lim_{k \rightarrow \infty} \int_{\Omega} f(x, z_{n_k}(x)) dx = \int_{\Omega} \bar{f}(x) dx < \infty$$

if and only if $\{f(\cdot, z_{n_k}(\cdot))\}$ is equi-integrable.

The map $\nu : \Omega \rightarrow \mathcal{M}(\mathbb{R}^d; \mathbb{R}^d)$ is called the *Young measure* generated by the sequence $\{z_{n_k}\}$.

Proposition 2.2. *If $\{z_n\}$ generates a Young measure ν and $v_n \rightarrow 0$ in measure, then $\{z_n + v_n\}$ still generates the Young measure ν .*

If $1 < q \leq \infty$ then $W^{-1,q}(\Omega; \mathbb{R}^l)$ is the dual of $W_0^{1,q'}(\Omega; \mathbb{R}^l)$, where q' is the Hölder conjugate exponent of q , that is $1/q + 1/q' = 1$. It is well known that $F \in W^{-1,q}(\Omega; \mathbb{R}^l)$ if and only if there exist $g_1, \dots, g_N \in L^q(\Omega; \mathbb{R}^l)$ such that

$$\langle F, w \rangle = \sum_{i=1}^N \int_{\Omega} g_i \cdot \frac{\partial w}{\partial x_i} dx \quad \text{for all } w \in W_0^{1,q}(\Omega; \mathbb{R}^l).$$

Consider a collection of linear operators $A^{(i)} : \mathbb{R}^d \rightarrow \mathbb{R}^l, i = 1, \dots, N$, and define the differential operator

$$\begin{aligned} \mathcal{A} : L^q(\Omega; \mathbb{R}^d) &\longrightarrow W^{-1,q}(\Omega; \mathbb{R}^l) \\ v &\longmapsto \mathcal{A}v \end{aligned}$$

as follows:

$$\langle \mathcal{A}v, w \rangle := \left\langle \sum_{i=1}^N A^{(i)} \frac{\partial v}{\partial x_i}, w \right\rangle = - \sum_{i=1}^N \int_{\Omega} A^{(i)} v \frac{\partial w}{\partial x_i} dx \quad \text{for all } w \in W_0^{1,q}(\Omega; \mathbb{R}^l).$$

Even though the operator \mathcal{A} so defined depends on Ω , we will omit reference to the underlying domain whenever it is clear from the context. In particular, if $v \in L^q_{1\text{-per}}(\mathbb{R}^N; \mathbb{R}^d)$ then we will say that $v \in \ker \mathcal{A}$ if $\mathcal{A}v = 0$ in $W^{-1,q}(Q; \mathbb{R}^l)$.

Throughout the paper we assume that \mathcal{A} satisfies the constant-rank property (1.1).

The following proposition is due to Fonseca and Müller [22].

Proposition 2.3. (i) $(1 < q < +\infty)$ *Let $1 < q < +\infty$, let $\{V_n\}$ be a bounded sequence in $L^q(\Omega; \mathbb{R}^d)$ such that $\mathcal{A}V_n \rightarrow 0$ in $W^{-1,q}(\Omega; \mathbb{R}^l)$, $V_n \rightharpoonup V$ in $L^q(\Omega; \mathbb{R}^d)$, and assume that $\{V_n\}$ generates a Young measure ν . Then there exists a q -equi-integrable sequence $\{v_n\} \subset L^q(\Omega; \mathbb{R}^d) \cap \ker \mathcal{A}$ such that*

$$\int_{\Omega} v_n dx = \int_{\Omega} V dx, \quad \|v_n - V_n\|_{L^s(\Omega)} \rightarrow 0 \quad \text{for all } 1 \leq s < q,$$

and, in particular, $\{v_n\}$ still generates ν . Moreover, if $\Omega = Q$ then $v_n - V \in L^q_{1\text{-per}}(\mathbb{R}^N; \mathbb{R}^d) \cap \ker \mathcal{A}$.

(ii) ($q = 1$) Let $\{V_n\}$ be a sequence converging weakly in $L^1(\Omega; \mathbb{R}^d)$ to a function V , $\mathcal{A}V_n \rightarrow 0$ in $W^{-1,r}(\Omega; \mathbb{R}^l)$ for some $r \in (1, N/(N - 1))$, and assume that $\{V_n\}$ generates a Young measure ν . Then there exists an equi-integrable sequence $\{v_n\} \subset L^1(\Omega; \mathbb{R}^d) \cap \ker \mathcal{A}$ such that

$$\int_{\Omega} v_n \, dx = \int_{\Omega} V \, dx, \quad \|v_n - V_n\|_{L^1(\Omega)} \rightarrow 0,$$

and, in particular, $\{v_n\}$ still generates ν . Moreover, if $\Omega = Q$ then $v_n - V \in L^1_{1\text{-per}}(\mathbb{R}^N; \mathbb{R}^d) \cap \ker \mathcal{A}$.

(iii) ($q = +\infty$) Let $\{V_n\}$ be a sequence that satisfies $V_n \overset{*}{\rightharpoonup} V$ in $L^\infty(\Omega; \mathbb{R}^d)$, $\mathcal{A}V_n \rightarrow 0$ in $L^r(\Omega)$ for some $r > N$, and assume that $\{V_n\}$ generates a Young measure ν . Then there exists a sequence $\{v_n\} \subset L^\infty(\Omega; \mathbb{R}^d) \cap \ker \mathcal{A}$ such that

$$\int_{\Omega} v_n \, dx = \int_{\Omega} V \, dx, \quad \|v_n - V_n\|_{L^\infty(\Omega)} \rightarrow 0,$$

and, in particular, $\{v_n\}$ still generates ν . Moreover, if $\Omega = Q$ then $v_n - V \in L^\infty_{1\text{-per}}(\mathbb{R}^N; \mathbb{R}^d) \cap \ker \mathcal{A}$.

In the second part of the paper we will need the following classical results from Γ -convergence. For a proof see [10].

Proposition 2.4. Let (X, d) be a separable metric space and let $f_n : X \rightarrow [-\infty, \infty]$. Then

(i) there exists an increasing sequence of integers $\{n_k\}$ such that

$$\Gamma(d) - \lim_{k \rightarrow \infty} f_{n_k}(x) \quad \text{exists for all } x \in X.$$

(ii) Moreover

$$f_\infty = \Gamma(d) - \lim_{n \rightarrow \infty} f_n$$

if and only if for every subsequence $\{f_{n_k}\}$ there exists a further subsequence $\{f_{n_{k_j}}\}$ which $\Gamma(d)$ -converges to f_∞ .

3. RELAXATION

In this section we prove Theorem 1.1 and its generalization to the case where $q \in \{1, \infty\}$ and $p = \infty$ (see Th. 3.6).

Lemma 3.1. Let $f : \Omega \times \mathbb{R}^m \times \mathbb{R}^d \rightarrow [0, \infty)$ be a Carathéodory function satisfying (H), with $1 \leq p < \infty$ and $1 < q < \infty$. Let $(u, v) \in L^p(D; \mathbb{R}^m) \times (L^q(D; \mathbb{R}^d) \cap \ker \mathcal{A})$, where $D \in \mathcal{O}(\Omega)$, and consider a sequence of functions $\{(u_k, \hat{v}_k)\} \subset L^p(D; \mathbb{R}^m) \times L^q(D; \mathbb{R}^d)$ such that

$$\begin{aligned} u_k &\rightharpoonup u \quad \text{in } L^p(D; \mathbb{R}^m), & \hat{v}_k &\rightharpoonup v \quad \text{in } L^q(D; \mathbb{R}^d) \\ \mathcal{A}\hat{v}_k &\rightarrow 0 \quad \text{in } W^{-1,q}(D; \mathbb{R}^l). \end{aligned} \tag{3.1}$$

Then we can find a q -equi-integrable sequence $\{v_k\} \subset L^q(D; \mathbb{R}^d) \cap \ker \mathcal{A}$ such that

$$v_k \rightharpoonup v \quad \text{in } L^q(D; \mathbb{R}^d), \quad \int_D v_k \, dx = \int_D v \, dx,$$

and

$$\liminf_{k \rightarrow \infty} \int_D f(x, u(x), v_k(x)) \, dx \leq \liminf_{k \rightarrow \infty} \int_D f(x, u_k(x), \hat{v}_k(x)) \, dx.$$

Proof of Lemma 3.1. Consider a subsequence $\{(u_n, \hat{v}_n)\}$ of $\{(u_k, \hat{v}_k)\}$ such that

$$\lim_{n \rightarrow \infty} \int_D f(x, u_n(x), \hat{v}_n(x)) \, dx = \liminf_{k \rightarrow \infty} \int_D f(x, u_k(x), \hat{v}_k(x)) \, dx$$

and $\{(u_n, \hat{v}_n)\}$ generates the Young measure $\{\delta_{u(x)} \otimes \nu_x\}_{x \in D}$. For $i \in \mathbb{N}$ let

$$F_i := \left\{ x \in D : \text{dist}(x, \partial D) < \frac{1}{i} \right\},$$

and consider cut-off functions θ_i with compact support in D and such that $\theta_i \equiv 1$ in $D \setminus F_i$. Set $w_{i,n} := \theta_i(\hat{v}_n - v) \in L^q(D; \mathbb{R}^d)$ and fix $\varphi \in L^{q'}(D; \mathbb{R}^d)$, where q' is the Hölder conjugate exponent of q . Then

$$\lim_{i \rightarrow \infty} \lim_{n \rightarrow \infty} \int_D \varphi(x) w_{i,n}(x) \, dx = \lim_{i \rightarrow \infty} \lim_{n \rightarrow \infty} \int_D \varphi(x) \theta_i(x) (\hat{v}_n(x) - v(x)) \, dx = 0, \tag{3.2}$$

where we have used the fact that $\hat{v}_k \rightharpoonup v$ in $L^q(D; \mathbb{R}^d)$. Hence $w_{i,n} \rightharpoonup 0$ in $L^q(D; \mathbb{R}^d)$ as $n \rightarrow \infty$ and $i \rightarrow \infty$. Moreover, in view of the compact embedding

$$L^q(D; \mathbb{R}^l) \hookrightarrow W^{-1,q}(D; \mathbb{R}^l)$$

and the assumption that $\mathcal{A}\hat{v}_k \rightarrow 0$ in $W^{-1,q}(D; \mathbb{R}^l)$, we have that

$$\lim_{i \rightarrow \infty} \lim_{n \rightarrow \infty} \mathcal{A}w_{i,n} = 0 \quad \text{in } W^{-1,q}(D; \mathbb{R}^l).$$

Let \mathbb{G} be a countable dense subset of $L^q(D; \mathbb{R}^d)$. By means of a diagonalization process we obtain subsequences $\{u_i := u_{n_i}\}$ and $\{\hat{w}_i := w_{i,n_i} = \theta_i(\hat{v}_{n_i} - v)\}$ such that $\|u_i - u\|_{L^p} \rightarrow 0$, (3.2) holds for each $\varphi \in \mathbb{G}$, and

$$\mathcal{A}\hat{w}_i \rightarrow 0 \quad \text{in } W^{-1,q}(D; \mathbb{R}^l).$$

Hence $\hat{w}_i \rightharpoonup 0$ in $L^q(D; \mathbb{R}^d)$, by the density of \mathbb{G} in $L^q(D; \mathbb{R}^d)$. By Proposition 2.3(i) there exists a q -equi-integrable sequence $\{w_i\} \subset L^q(D; \mathbb{R}^d) \cap \ker \mathcal{A}$ such that $w_i \rightharpoonup 0$ in $L^q(D; \mathbb{R}^d)$, and

$$\int_D w_i \, dx = 0, \quad \|\hat{w}_i - w_i\|_{L^s(D)} \rightarrow 0 \quad \text{for all } 1 \leq s < q. \tag{3.3}$$

Set $v_i := v + w_i$. Then $\int_D v_i \, dx = \int_D v \, dx$, $v_i \rightharpoonup v$ in $L^q(D; \mathbb{R}^d)$. By Hölder's inequality and by (3.3), for $1 \leq s < q$

$$\begin{aligned} \|\hat{v}_{n_i} - v_i\|_{L^s(D)} &\leq \|\hat{v}_{n_i} - v - \hat{w}_i\|_{L^s(D)} + \|\hat{w}_i - w_i\|_{L^s(D)} \\ &\leq \|(1 - \theta_i)(\hat{v}_{n_i} - v)\|_{L^s(D)} + \|\hat{w}_i - w_i\|_{L^s(D)} \\ &\leq \|\hat{v}_{n_i} - v\|_{L^q(D)} |F_i|^r + \|\hat{w}_i - w_i\|_{L^s(D)} \rightarrow 0 \end{aligned} \tag{3.4}$$

as $i \rightarrow \infty$ and where $r := (q - s)/sq$. By (3.4) and Proposition 2.2, the two sequences

$$\{(u(x), v_i(x))\} \quad \text{and} \quad \{(u_i(x), \hat{v}_i(x))\}$$

generate the same Young measure $\{\delta_{u(x)} \otimes \nu_x\}_{x \in D}$. Hence by Theorems 2.1(ii) and (iii)

$$\begin{aligned} \lim_{i \rightarrow \infty} \int_D f(x, u(x), v_i(x)) \, dx &= \int_D \int_{\mathbb{R}^d} f(x, u(x), V) \, d\nu_x(V) \, dx \leq \liminf_{i \rightarrow \infty} \int_D f(x, u_i(x), \hat{v}_i(x)) \, dx \\ &= \liminf_{k \rightarrow \infty} \int_D f(x, u_k(x), \hat{v}_k(x)) \, dx, \end{aligned} \tag{3.5}$$

where we have used the fact that $\{f(x, u(x), v_i(x))\}$ is equi-integrable over D , which follows from (H) and the q -equi-integrability of $\{v_i\}$ over D . \square

It follows immediately from Lemma 3.1 that under its assumptions on f it holds:

Corollary 3.2. For $D \in \mathcal{O}(\Omega)$ and $(u, v) \in L^p(\Omega; \mathbb{R}^m) \times (L^q(\Omega; \mathbb{R}^d) \cap \ker \mathcal{A})$

$$\mathcal{F}((u, v); D) = \inf \left\{ \liminf_{n \rightarrow \infty} \int_D g(x, v_n(x)) \, dx : \{v_n\} \subset L^q(D; \mathbb{R}^d) \cap \ker \mathcal{A} \text{ is } q\text{-equi-integrable} \right. \\ \left. \text{and } v_n \rightarrow v \text{ in } L^q(D; \mathbb{R}^d) \right\},$$

where g is the Carathéodory function defined by

$$g(x, v) := f(x, u(x), v).$$

Note that, by (H), the function g satisfies the growth condition

$$0 \leq g(x, v) \leq C(1 + |u(x)|^p + |v|^q) \tag{3.6}$$

for a.e. $x \in \Omega$ and all $v \in \mathbb{R}^d$. Moreover, since g is a Carathéodory function, by the Scorza-Dragoni theorem for each $j \in \mathbb{N}$ there exists a compact set $K_j \subset \Omega$, with $|\Omega \setminus K_j| \leq 1/j$, such that $g : K_j \times \mathbb{R}^d \rightarrow [0, \infty)$ is continuous. Let K_j^* be the set of Lebesgue points of χ_{K_j} , and set

$$\omega := \bigcup_j (K_j \cap K_j^*) \cap L(u, v), \tag{3.7}$$

where $L(u, v)$ is the set of Lebesgue points of (u, v) . Then

$$|\Omega \setminus \omega| \leq |\Omega \setminus K_j| \leq \frac{1}{j} \rightarrow 0 \text{ as } j \rightarrow \infty.$$

Corollary 3.3. Assume that $x_0 \in \omega$, let $v \in L^q(Q; \mathbb{R}^d) \cap \ker \mathcal{A}$, and consider $r_k \rightarrow 0^+$ and a sequence of functions

$$\{\hat{v}_k\} \subset L^q(Q; \mathbb{R}^d) \cap \ker \mathcal{A}$$

such that

$$\hat{v}_k \rightarrow v \text{ in } L^q(Q; \mathbb{R}^d).$$

Then we can find a q -equi-integrable sequence $\{w_k\} \subset L^q_{1\text{-per}}(\mathbb{R}^N; \mathbb{R}^d) \cap \ker \mathcal{A}$ such that

$$w_k \rightarrow 0 \text{ in } L^q(Q; \mathbb{R}^d), \quad \int_Q w_k \, dx = 0,$$

and

$$\liminf_{k \rightarrow \infty} \int_Q g(x_0, v(y) + w_k(y)) \, dy \leq \liminf_{k \rightarrow \infty} \int_Q g(x_0 + r_k y, \hat{v}_k(y)) \, dy.$$

Proof of Corollary 3.3. We proceed as in the proof of Lemma 3.1 up to (3.4). Since the sequence $\{v_i\}$ is q -equi-integrable, for any $\eta > 0$ there exists $\delta > 0$ such that

$$\sup_i \int_D C(1 + |u(x_0)|^p + |v_i(y)|^q) dy < \eta \tag{3.8}$$

for any measurable set $D \subset Q$, with $|D| < \delta$, and where C is the constant given in (H). Fix $\eta > 0$ and let $\delta > 0$ be given according to (3.8). By the Biting Lemma (see [6]) we may find a further subsequence $\{\hat{v}_{n_j}\} \subset \{\hat{v}_{n_i}\}$ and a set $E \subset Q$ such that $|Q \setminus E| < \delta$ and $\{\hat{v}_{n_j}\}$ is q -equi-integrable over E . Hence there exists $0 < \delta_1 < \delta$ such that

$$\sup_j \int_D C(1 + |u(x_0)|^p + |\hat{v}_{n_j}(y)|^q) dy < \eta \tag{3.9}$$

for any measurable set $D \subset E$, with $|D| < \delta_1$. Moreover, as $\{\hat{v}_{n_j}\}, \{v_j\}$ are bounded in $L^q(Q; \mathbb{R}^d)$, we may find $L > 0$ such that

$$|E \setminus E_j| \leq \delta_1, \quad \text{where } E_j := \{y \in E : |\hat{v}_{n_j}(y)| \leq L, |v_j(y)| \leq L\}. \tag{3.10}$$

Note that by construction of v_i and by Proposition 2.3, $v_i = v + w_i$ where $w_i \in L^q_{1\text{-per}}(\mathbb{R}^N; \mathbb{R}^d) \cap \ker \mathcal{A}$. From the definition of the set ω there exists an integer j_0 such that $x_0 \in K_{j_0} \cap K_{j_0}^*$. Since

$$g : K_{j_0} \times \overline{B_d(0, L)} \rightarrow [0, \infty)$$

is uniformly continuous, there exists $\rho > 0$ such that

$$|g(x, v) - g(x_1, v)| \leq \eta \tag{3.11}$$

for all $(x, v), (x_1, v) \in K_{j_0} \times \overline{B_d(0, L)}$, with $|x - x_1| \leq \rho$. By (3.10) and (3.11)

$$\begin{aligned} \lim_{i \rightarrow \infty} \int_Q g(x_0 + r_{n_i}y, \hat{v}_{n_i}(y)) dy &\geq \liminf_{j \rightarrow \infty} \int_{E_j} g(x_0 + r_{n_j}y, \hat{v}_{n_j}(y)) dy \\ &\geq \liminf_{j \rightarrow \infty} \frac{1}{r_{n_j}^N} \int_{(x_0 + r_{n_j}E_j) \cap K_{j_0}} g(x, \hat{v}_{n_j}((x - x_0)/r_{n_j})) dx \\ &\geq -\eta + \liminf_{j \rightarrow \infty} \frac{1}{r_{n_j}^N} \int_{(x_0 + r_{n_j}E_j) \cap K_{j_0}} g(x_0, \hat{v}_{n_j}((x - x_0)/r_{n_j})) dx. \end{aligned} \tag{3.12}$$

Using, once again, the fact that $|\hat{v}_{n_j}(y)| \leq L$ for $y \in E_j$, by (3.6) we have that

$$\frac{1}{r_{n_j}^N} \int_{(x_0 + r_{n_j}E_j) \setminus K_{j_0}} g(x_0, \hat{v}_{n_j}((x - x_0)/r_{n_j})) dx \leq C(1 + |u(x_0)|^p + L^q) \frac{|Q(x_0, r_{n_j}) \setminus K_{j_0}|}{r_{n_j}^N} \rightarrow 0$$

as $j \rightarrow \infty$, because x_0 is a Lebesgue point of $\chi_{K_{j_0}}$. Consequently, from (3.12) we get

$$\begin{aligned} \lim_{i \rightarrow \infty} \int_Q g(x_0 + r_{n_i}y, \hat{v}_{n_i}(y)) dy &\geq -\eta + \liminf_{j \rightarrow \infty} \frac{1}{r_{n_j}^N} \int_{x_0 + r_{n_j}E_j} g(x_0, \hat{v}_{n_j}((x - x_0)/r_{n_j})) dx \\ &= -\eta + \liminf_{j \rightarrow \infty} \int_{E_j} g(x_0, \hat{v}_{n_j}(y)) dy \\ &\geq -2\eta + \liminf_{j \rightarrow \infty} \int_E g(x_0, \hat{v}_{n_j}(y)) dy, \end{aligned}$$

where we have used (3.6, 3.9) and the fact that $|E \setminus E_j| \leq \delta_1$. We may now proceed as in the previous lemma, using the Carathéory function $h(x, v) := \chi_E(x)g(x_0, v)$, to obtain

$$\lim_{i \rightarrow \infty} \int_Q g(x_0 + r_{n_i}y, \hat{v}_{n_i}(y)) dy \geq -2\eta + \liminf_{j \rightarrow \infty} \int_E g(x_0, v_j(y)) dy \geq -3\eta + \liminf_{j \rightarrow \infty} \int_Q g(x_0, v_j(y)) dy$$

by (3.8). It now suffices to let $\eta \rightarrow 0^+$. □

Theorem 1.1 follows from Lemmas 3.4 and 3.5 below. We will use the notation $\mu|_A$ to denote the restriction of a Radon measure μ to the Borel set A , i.e., $\mu|_A(X) := \mu(X \cap A)$ where X is an arbitrary Borel set in the domain of μ .

Lemma 3.4. $\mathcal{F}((u, v); \cdot)$ is the trace of a Radon measure absolutely continuous with respect to $\mathcal{L}^N|_\Omega$.

Proof of Lemma 3.4. As it is usual, it suffices to prove subadditivity (see e.g. [3, 21]), i.e.

$$\mathcal{F}((u, v); D) \leq \mathcal{F}((u, v); D \setminus \overline{B}) + \mathcal{F}((u, v); C)$$

if $B \subset\subset C \subset\subset D$. Fix $\eta > 0$. By Corollary 3.2 there exist two q -equi-integrable sequences

$$\{v_k\} \subset L^q(D \setminus \overline{B}; \mathbb{R}^d) \cap \ker \mathcal{A}, \quad \{w_k\} \subset L^q(C; \mathbb{R}^d) \cap \ker \mathcal{A},$$

such that

$$v_k \rightharpoonup v \text{ in } L^q(D \setminus \overline{B}; \mathbb{R}^d), \quad w_k \rightharpoonup v \text{ in } L^q(C; \mathbb{R}^d),$$

and

$$\begin{aligned} \lim_{k \rightarrow \infty} \int_{D \setminus \overline{B}} g(x, v_k(x)) dx &\leq \mathcal{F}((u, v); D \setminus \overline{B}) + \eta, \\ \lim_{k \rightarrow \infty} \int_C g(x, w_k(x)) dx &\leq \mathcal{F}((u, v); C) + \eta. \end{aligned}$$

Let θ_j be smooth cut-off functions, $\theta_j \in C_c^\infty(C; [0, 1])$, $\theta_j(x) = 1$ for all $x \in B$, and $|\{0 < \theta_j < 1\}| \rightarrow 0$ as $j \rightarrow \infty$. Set

$$\hat{V}_{j,k} := (1 - \theta_j)v_k + \theta_j w_k.$$

Then, for j fixed,

$$\mathcal{A}\hat{V}_{j,k} = (1 - \theta_j)\mathcal{A}v_k + \theta_j\mathcal{A}w_k - \sum_{i=1}^N A^{(i)}v_k \frac{\partial \theta_j}{\partial x_i} + \sum_{i=1}^N A^{(i)}w_k \frac{\partial \theta_j}{\partial x_i} \rightarrow 0$$

as $k \rightarrow \infty$ in $W^{-1,q}(D; \mathbb{R}^l)$ strong. Using a diagonalization procedure such as that adopted in the proof of Lemma 3.1, we get

$$\hat{V}_j \rightharpoonup v \text{ in } L^q(D; \mathbb{R}^d), \quad \mathcal{A}\hat{V}_j \rightarrow 0 \text{ in } W^{-1,q}(D; \mathbb{R}^l),$$

where $\hat{V}_j := \hat{V}_{j,k_j}$. By Lemma 3.1 we can find a q -equi-integrable sequence $\{V_j\} \subset L^q(D; \mathbb{R}^d) \cap \ker \mathcal{A}$ such that $V_j \rightharpoonup v$ in $L^q(D; \mathbb{R}^d)$ and

$$\liminf_{j \rightarrow \infty} \int_D g(x, V_j(x)) dx \leq \liminf_{j \rightarrow \infty} \int_D g(x, \hat{V}_j(x)) dx.$$

Consequently, in view of Corollary 3.2

$$\begin{aligned} \mathcal{F}((u, v); D) &\leq \liminf_{j \rightarrow \infty} \int_D g(x, V_j(x)) \, dx \leq \liminf_{j \rightarrow \infty} \int_D g(x, \hat{V}_j(x)) \, dx \\ &\leq \limsup_{j \rightarrow \infty} \int_{\{\theta_j=0\}} g(x, v_{k_j}(x)) \, dx + \limsup_{j \rightarrow \infty} \int_{\{\theta_j=1\}} g(x, w_{k_j}(x)) \, dx \\ &\quad + \limsup_{j \rightarrow \infty} \int_{\{0 < \theta_j < 1\}} C(1 + |u(x)|^p + |w_{k_j}(x)|^q + |v_{k_j}(x)|^q) \, dx \\ &\leq 2\eta + \mathcal{F}((u, v); D \setminus \overline{B}) + \mathcal{F}((u, v); C). \end{aligned}$$

It suffices to let $\eta \rightarrow 0^+$. Finally, note that by (H) we have that

$$\mathcal{F}((u, v), \cdot) \leq C(1 + |u|^p + |v|^q) \mathcal{L}^N \llcorner \Omega.$$

□

Lemma 3.5. *For \mathcal{L}^N a.e. $x_0 \in \Omega$ we have*

$$\frac{d\mathcal{F}((u, v); \cdot)}{d\mathcal{L}^N}(x_0) = \mathcal{Q}A f(x_0, u(x_0), v(x_0)).$$

Proof of Lemma 3.5. Fix $x_0 \in \omega$, where ω is defined as in (3.7), and such that

$$\lim_{r \rightarrow 0^+} \frac{1}{r^N} \int_{Q(x_0, r)} |u(x) - u(x_0)|^p \, dx = \lim_{r \rightarrow 0^+} \frac{1}{r^N} \int_{Q(x_0, r)} |v(x) - v(x_0)|^q \, dx = 0 \tag{3.13}$$

and

$$\frac{d\mathcal{F}((u, v); \cdot)}{d\mathcal{L}^N}(x_0) = \lim_{r \rightarrow 0^+} \frac{\mathcal{F}((u, v); Q(x_0, r))}{r^N} < \infty,$$

where, by virtue of Lemma 3.4, we have chosen the radii $r \rightarrow 0^+$ such that

$$\mathcal{F}((u, v); \partial(Q(x_0, r))) = 0.$$

By Corollary 3.2 and for $r > 0$ fixed, let $\{v_{n,r}\} \subset L^q(Q(x_0, r); \mathbb{R}^d) \cap \ker \mathcal{A}$ be such that $v_{n,r} \rightharpoonup v$ in $L^q(Q(x_0, r); \mathbb{R}^d)$ as $n \rightarrow \infty$ and

$$\lim_{n \rightarrow \infty} \int_{Q(x_0, r)} g(x, v_{n,r}(x)) \, dx \leq \mathcal{F}((u, v); Q(x_0, r)) + r^{N+1}.$$

Then

$$\frac{d\mathcal{F}((u, v); \cdot)}{d\mathcal{L}^N}(x_0) \geq \liminf_{r \rightarrow 0^+} \lim_{n \rightarrow \infty} \frac{1}{r^N} \int_{Q(x_0, r)} g(x, v_{n,r}(x)) \, dx = \liminf_{r \rightarrow 0^+} \lim_{n \rightarrow \infty} \int_Q g(x_0 + ry, v(x_0) + w_{n,r}(y)) \, dy$$

where $w_{n,r}(y) := v_{n,r}(x_0 + ry) - v(x_0)$. We claim that $w_{n,r} \rightarrow 0$ in $L^q(Q; \mathbb{R}^d)$ if we first let $n \rightarrow \infty$ and then $r \rightarrow 0^+$. Indeed let $\varphi \in L^{q'}(Q; \mathbb{R}^d)$, where q' is the Hölder conjugate exponent of q . Using Hölder's inequality

and then making a change of variables, we get

$$\begin{aligned} \left| \int_Q \varphi(y) w_{r,n}(y) dy \right| &\leq \left| \int_Q \varphi(y)(v_{n,r}(x_0 + ry) - v(x_0 + ry)) dy \right| + \left| \int_Q \varphi(y)(v(x_0 + ry) - v(x_0)) dy \right| \\ &\leq \left| \frac{1}{r^N} \int_{Q(x_0,r)} \varphi((x - x_0)/r)(v_{n,r}(x) - v(x)) dx \right| \\ &\quad + \|\varphi\|_{L^{q'}(Q)} \left(\frac{1}{r^N} \int_{Q(x_0,r)} |v(x) - v(x_0)|^q dx \right)^{1/q}. \end{aligned}$$

If we now let $n \rightarrow \infty$ the first integral tends to zero, since $v_{n,r} \rightharpoonup v$ in $L^q(Q(x_0, r); \mathbb{R}^d)$. The claim then follows by letting $r \rightarrow 0^+$ and by using (3.13). Diagonalize to get $\hat{w}_k \in L^q(Q; \mathbb{R}^d) \cap \ker \mathcal{A}$ such that $\hat{w}_k \rightharpoonup 0$ in $L^q(Q; \mathbb{R}^d)$ and

$$\frac{d\mathcal{F}((u, v); \cdot)}{d\mathcal{L}^N}(x_0) \geq \lim_{k \rightarrow \infty} \int_Q g(x_0 + r_k y, v(x_0) + \hat{w}_k(y)) dy$$

where $r_k \rightarrow 0$. By Corollary 3.3 there is a q -equi-integrable sequence $\{w_k\} \subset L^q_{1\text{-per}}(\mathbb{R}^N; \mathbb{R}^d) \cap \ker \mathcal{A}$ such that

$$w_k \rightharpoonup 0 \quad \text{in } L^q(Q; \mathbb{R}^d), \quad \int_Q w_k dy = 0,$$

and

$$\begin{aligned} \frac{d\mathcal{F}((u, v); \cdot)}{d\mathcal{L}^N}(x_0) &\geq \lim_{k \rightarrow \infty} \int_Q g(x_0 + r_k y, v(x_0) + \hat{w}_k(y)) dy \\ &\geq \liminf_{k \rightarrow \infty} \int_Q f(x_0, u(x_0), v(x_0) + w_k(y)) dy \geq \mathcal{Q}Af(x_0, u(x_0), v(x_0)). \end{aligned}$$

To conclude the proof of the lemma it remains to show that

$$\frac{d\mathcal{F}((u, v); \cdot)}{d\mathcal{L}^N}(x_0) \leq \mathcal{Q}Af(x_0, u(x_0), v(x_0)) \quad \text{for } \mathcal{L}^N \text{ a.e. } x_0 \in \Omega.$$

Fix $\eta > 0$ and let $w \in C^\infty_{1\text{-per}}(\mathbb{R}^N; \mathbb{R}^d) \cap \ker \mathcal{A}$ be such that $\int_Q w dy = 0$ and

$$\int_Q f(x_0, u(x_0), v(x_0) + w(y)) dy \leq \mathcal{Q}Af(x_0, u(x_0), v(x_0)) + \eta. \tag{3.14}$$

For any fixed $r > 0$ set $w_{n,r}(x) := w(n(x - x_0)/r)$. Then $w_{n,r} \rightharpoonup 0$ in $L^q(Q(x_0, r); \mathbb{R}^d)$ as $n \rightarrow \infty$. Hence, by Corollary 3.2,

$$\frac{d\mathcal{F}((u, v); \cdot)}{d\mathcal{L}^N}(x_0) = \lim_{r \rightarrow 0^+} \frac{\mathcal{F}((u, v); Q(x_0, r))}{r^N} \leq \liminf_{r \rightarrow 0^+} \liminf_{n \rightarrow \infty} \frac{1}{r^N} \int_{Q(x_0,r)} g(x, v(x) + w_{n,r}(x)) dx. \tag{3.15}$$

Fix $L > |v(x_0)| + \|w\|_{L^\infty} + 1$, and let j be such that $x_0 \in K_j \cap K_j^* \cap L(u, v)$, where we are using the notation introduced in (3.7). Since

$$g : K_j \times \overline{B_d(0, L)} \rightarrow [0, \infty)$$

is uniformly continuous, there exists $0 < \rho < 1$ such that

$$|g(x, v) - g(x_1, v_1)| \leq \eta \tag{3.16}$$

for all $(x, v), (x_1, v_1) \in K_j \times \overline{B_d(v(x_0), L)}$, with $|x - x_1| \leq \rho$ and $|v - v_1| \leq \rho$. Let

$$E_{r,\rho} := \{x \in Q(x_0, r) : |v(x) - v(x_0)| \leq \rho\}.$$

We claim that

$$\limsup_{r \rightarrow 0^+} \frac{1}{r^N} \int_{Q(x_0, r) \setminus (E_{r,\rho} \cap K_j)} C(1 + |u(x)|^p + |v(x)|^q + \|w\|_{L^\infty}^q) dx = 0. \tag{3.17}$$

Since $|v(x) - v(x_0)| \geq \rho$ for $x \in Q(x_0, r) \setminus E_{r,\rho}$, we have

$$\begin{aligned} \frac{1}{r^N} \int_{Q(x_0, r) \setminus (E_{r,\rho} \cap K_j)} C(1 + |u(x)|^p + |v(x)|^q + \|w\|_{L^\infty}^q) dx &\leq C \frac{|Q(x_0, r) \setminus (E_{r,\rho} \cap K_j)|}{r^N} \\ &\quad + \frac{C}{r^N} \int_{Q(x_0, r)} (|u(x) - u(x_0)|^p + |v(x) - v(x_0)|^q) dx \end{aligned}$$

and

$$\begin{aligned} \frac{|Q(x_0, r) \setminus (E_{r,\rho} \cap K_j)|}{r^N} &\leq \frac{|Q(x_0, r) \setminus K_j|}{r^N} + \frac{|Q(x_0, r) \setminus E_{r,\rho}|}{r^N} \leq \frac{|Q(x_0, r) \setminus K_j|}{r^N} \\ &\quad + \frac{1}{\rho^q} \frac{C}{r^N} \int_{Q(x_0, r)} |v(x) - v(x_0)|^q dx \rightarrow 0 \quad \text{as } r \rightarrow 0^+, \end{aligned}$$

where we have used (3.13) and the fact that x_0 is a Lebesgue point of χ_{K_j} . Then by (3.6, 3.15–3.17) and (3.14),

$$\begin{aligned} \frac{d\mathcal{F}((u, v); \cdot)}{d\mathcal{L}^N}(x_0) &\leq \liminf_{r \rightarrow 0^+} \liminf_{n \rightarrow \infty} \frac{1}{r^N} \int_{E_{r,\rho} \cap K_j} g(x, v(x) + w_{n,r}(x)) dx \\ &\quad + \limsup_{r \rightarrow 0^+} \frac{1}{r^N} \int_{Q(x_0, r) \setminus (E_{r,\rho} \cap K_j)} C(1 + |u(x)|^p + |v(x)|^q + \|w\|_{L^\infty}^q) dx \\ &\leq \eta + \liminf_{r \rightarrow 0^+} \liminf_{n \rightarrow \infty} \frac{1}{r^N} \int_{Q(x_0, r)} g(x_0, v(x_0) + w_{n,r}(x)) dx \\ &= \eta + \liminf_{n \rightarrow \infty} \int_Q g(x_0, v(x_0) + w(ny)) dy \\ &= \eta + \int_Q g(x_0, v(x_0) + w(y)) dy \leq 2\eta + \mathcal{Q}_A f(x_0, u(x_0), v(x_0)), \end{aligned}$$

by virtue of the equality

$$\liminf_{n \rightarrow \infty} \int_Q g(x_0, v(x_0) + w(ny)) dx = \int_Q g(x_0, v(x_0) + w(y)) dy,$$

which follows from the Q -periodicity of the function $g(x_0, v(x_0) + w(\cdot))$. It now suffices to let $\eta \rightarrow 0^+$. □

As mentioned in the introduction, Theorem 1.1 continues to hold when $q \in \{1, \infty\}$ and $p = \infty$. Indeed, let $1 \leq p, q \leq \infty$ and assume that

(A₄) $f : \Omega \times \mathbb{R}^m \times \mathbb{R}^d \rightarrow [0, \infty)$ is a Carathéodory function satisfying the following growth conditions for a.e. $x \in \Omega$ and all $(u, v) \in \mathbb{R}^m \times \mathbb{R}^d$:

$$0 \leq f(x, u, v) \leq C(1 + |u|^p + |v|^q) \quad \text{if } 1 \leq p, q < \infty,$$

where $C > 0$;

$$0 \leq f(x, u, v) \leq a(x, u) (1 + |v|^q) \quad \text{if } p = \infty \text{ and } 1 \leq q < \infty, \tag{3.18}$$

where $\chi_\Omega a \in L^\infty_{\text{loc}}(\mathbb{R}^N \times \mathbb{R}^d; [0, \infty))$;

$$0 \leq f(x, u, v) \leq b(x, v) (1 + |u|^p) \quad \text{if } 1 \leq p < \infty \text{ and } q = \infty,$$

where $\chi_\Omega b \in L^\infty_{\text{loc}}(\mathbb{R}^N \times \mathbb{R}^m; [0, \infty))$;

$$\chi_\Omega f \in L^\infty_{\text{loc}}(\mathbb{R}^N \times \mathbb{R}^m \times \mathbb{R}^d; [0, \infty)) \quad \text{if } p = q = \infty.$$

For $D \in \mathcal{O}(\Omega)$ and $(u, v) \in L^p(\Omega; \mathbb{R}^m) \times (L^q(\Omega; \mathbb{R}^d) \cap \ker \mathcal{A})$ define

$$\mathcal{F}((u, v); D) := \inf \left\{ \liminf_{n \rightarrow \infty} F((u_n, v_n); D) : (u_n, v_n) \in L^p(D; \mathbb{R}^m) \times L^1(D; \mathbb{R}^d), \right. \\ \left. u_n \rightarrow u \text{ in } L^p(D; \mathbb{R}^m), \quad v_n \rightarrow v \text{ in } L^1(D; \mathbb{R}^d), \quad \mathcal{A}v_n \rightarrow 0 \text{ in } W^{-1,r}(D; \mathbb{R}^l) \right\}$$

if $q = 1$ and for some $r \in (1, N/(N - 1))$; as in (1.2), we set

$$\mathcal{F}((u, v); D) := \inf \left\{ \liminf_{n \rightarrow \infty} F((u_n, v_n); D) : (u_n, v_n) \in L^p(D; \mathbb{R}^m) \times L^q(D; \mathbb{R}^d) \right. \\ \left. u_n \rightarrow u \text{ in } L^p(D; \mathbb{R}^m), \quad v_n \rightarrow v \text{ in } L^q(D; \mathbb{R}^d), \quad \mathcal{A}v_n \rightarrow 0 \text{ in } W^{-1,q}(D; \mathbb{R}^l) \right\}$$

if $1 < q < \infty$;

$$\mathcal{F}((u, v); D) := \inf \left\{ \liminf_{n \rightarrow \infty} F((u_n, v_n); D) : (u_n, v_n) \in L^p(D; \mathbb{R}^m) \times L^\infty(D; \mathbb{R}^d), \right. \\ \left. u_n \rightarrow u \text{ in } L^p(D; \mathbb{R}^m), \quad v_n \overset{*}{\rightharpoonup} v \text{ in } L^\infty(D; \mathbb{R}^d), \quad \mathcal{A}v_n \rightarrow 0 \text{ in } L^r(D; \mathbb{R}^l) \right\}$$

if $q = \infty$ and for some $r > N$.

We can prove the following theorem:

Theorem 3.6. *Under condition (A_4) and the constant-rank hypothesis (1.1), for all $D \in \mathcal{O}(\Omega)$, $u \in L^p(\Omega; \mathbb{R}^m)$ and $v \in L^q(\Omega; \mathbb{R}^d) \cap \ker \mathcal{A}$, we have*

$$\mathcal{F}((u, v); D) = \int_D \mathcal{Q}_\mathcal{A} f(x, u(x), v(x)) \, dx.$$

Proof of Theorem 3.6.

Step 1. Assume first that $1 \leq p < \infty$ and $q = 1$. The proof is similar to the one of Theorem 1.1, with the exceptions that in Lemma 3.1 condition (3.1) should be replaced by

$$u_k \rightarrow u \text{ in } L^p(D; \mathbb{R}^m), \quad \hat{v}_k \rightarrow v \text{ in } L^1(D; \mathbb{R}^d), \\ \mathcal{A}\hat{v}_k \rightarrow 0 \text{ in } W^{-1,r}(D; \mathbb{R}^l) \text{ for some } r \in (1, N/(N - 1)),$$

that we use the compact embedding

$$L^1(D; \mathbb{R}^l) \hookrightarrow W^{-1,r}(D; \mathbb{R}^l), \quad r \in (0, N/(N-1)),$$

to diagonalize $\{w_{i,n}\}$, and (3.3, 3.4) are replaced, respectively, by

$$\int_D w_i \, dx = 0, \quad \|\hat{w}_i - w_i\|_{L^1(D)} \rightarrow 0,$$

$$\begin{aligned} \|\hat{v}_{n_i} - v_i\|_{L^1(D)} &\leq \|\hat{v}_{n_i} - v - \hat{w}_i\|_{L^1(D)} + \|\hat{w}_i - w_i\|_{L^1(D)} \leq \|(1 - \theta_i)(\hat{v}_{n_i} - v)\|_{L^1(D)} + \|\hat{w}_i - w_i\|_{L^1(D)} \\ &\leq \|\hat{v}_{n_i} - v\|_{L^1(F_i)} + \|\hat{w}_i - w_i\|_{L^1(D)} \rightarrow 0, \end{aligned}$$

where we have used the fact that $\|\hat{v}_{n_i} - v\|_{L^1(F_i)} \rightarrow 0$ as $i \rightarrow \infty$, which is due to the equi-integrability of the original sequence $\{\hat{v}_k - v\}$ and the fact that $|F_i| \rightarrow 0$.

Step 2. If $p = \infty$ and $1 \leq q < \infty$ then in Lemma 3.1 the only change needed is in deriving (3.5), which now follows from the fact that, by (3),

$$0 \leq f(x, u(x), v_i(x)) \leq A_\infty (1 + |v_i(x)|^q),$$

where $A_\infty := \text{esssup} \{a(x, u) : x \in \Omega, |u| \leq \|u\|_\infty\} < \infty$, and thus equi-integrability of $\{f(x, u, v_i)\}$ follows from the q -equi-integrability of $\{v_i\}$ over D . Moreover in the remaining of the proof of Theorem 1.1, the growth condition (3.6) should be replaced by

$$0 \leq g(x, v) \leq A_\infty (1 + |v|^q) \tag{3.19}$$

for a.e. $x \in \Omega$ and all $v \in \mathbb{R}^d$.

Step 3. If $1 \leq p \leq \infty$ and $q = \infty$ then in Lemma 3.1 the hypothesis (3.1) should be replaced by

$$\begin{aligned} u_k \rightarrow u \quad \text{in } L^p(D; \mathbb{R}^m), \quad \hat{v}_k \xrightarrow{*} v \quad \text{in } L^\infty(D; \mathbb{R}^d), \\ A\hat{v}_n \rightarrow 0 \quad \text{in } L^r(D; \mathbb{R}^l) \text{ for some } r > N, \end{aligned}$$

the growth condition should be replaced by (3.19) if $1 \leq p < \infty$, $q = \infty$, and by $\chi_\Omega g \in L^\infty_{\text{loc}}(\mathbb{R}^N \times \mathbb{R}^d; [0, \infty))$ if $p = q = \infty$, and we can proceed similarly to the proof of Lemma 3.1 to show that $w_{i,n} \xrightarrow{*} 0$ in $L^\infty(D; \mathbb{R}^d)$ and $\mathcal{A}w_{i,n} \rightarrow 0$ in $L^r(D; \mathbb{R}^l)$, and use Proposition 2.3(iii) to get

$$\|v_i - \hat{v}_{n_i}\|_\infty \rightarrow 0.$$

We omit the details. □

4. PROOFS OF THEOREMS 1.3 AND 1.5

Proof of Theorem 1.3. We present the proof for $1 \leq p < \infty$, the case $p = \infty$ being very similar. Fix $u \in W^{s,p}(\Omega; \mathbb{R}^n)$, and for $D \in \mathcal{O}(\Omega)$ define

$$\mathcal{F}(u; D) := \inf \left\{ \liminf_{k \rightarrow \infty} \int_D f(x, u_k, \dots, \nabla^s u_k) \, dx : \{u_k\} \subset W^{s,p}(D; \mathbb{R}^n), \right. \\ \left. u_k \rightarrow u \text{ in } W^{s,p}(D; \mathbb{R}^n) \right\},$$

and let g be the Carathéodory function

$$g(x, v) := f(x, u(x), \dots, \nabla^{s-1}u(x), v).$$

Reasoning as in Lemma 3.4, it is easy to show that $\mathcal{F}(u; \cdot)$ is the trace of a Radon measure absolutely continuous with respect to $\mathcal{L}^N \llcorner \Omega$.

For any function $v \in L^p(\Omega; E_s^n)$ set

$$\mathcal{G}(v; D) := \inf \left\{ \liminf_{k \rightarrow \infty} \int_D g(x, V_k(x)) dx : \{V_k\} \subset L^p(D; E_s^n) \cap \ker \mathcal{A} \text{ is } p\text{-equi-integrable,} \right. \\ \left. \text{and } V_k \rightharpoonup v \text{ in } L^p(D; E_s^n) \right\},$$

where the differential operator \mathcal{A} is given by

$$\mathcal{A}v := \left(\frac{\partial}{\partial x_i} v_{i_1 \dots i_h j i_{h+2} \dots i_s} - \frac{\partial}{\partial x_j} v_{i_1 \dots i_h i i_{h+2} \dots i_s} \right)_{0 \leq h \leq s-1, 1 \leq i, j, i_1 \dots i_s \leq N}.$$

Here $h = 0$ and $h = s - 1$ correspond to the multi-indices $j i_2 \dots i_s$ and $i_1 \dots i_{s-1} j$. By Theorem 3.6 (and Cor. 3.2), and where the target space \mathbb{R}^d is being replaced by the finite dimensional vector space E_s^n , for any $D \in \mathcal{O}(\Omega)$

$$\mathcal{G}(v; D) = \int_D \mathcal{Q}_{\mathcal{A}}g(x, v(x)) dx,$$

where for a.e. $x \in \Omega$ and for all $v \in E_s^n$,

$$\mathcal{Q}_{\mathcal{A}}g(x, v) := \inf \left\{ \int_Q g(x, v + w(y)) dy : w \in C_{1\text{-per}}^\infty(\mathbb{R}^N; E_s^n) \cap \ker \mathcal{A}, \int_Q w(y) dy = 0 \right\}.$$

As shown in [22],

$$\left\{ w \in C_{1\text{-per}}^\infty(\mathbb{R}^N; E_s^n) : \mathcal{A}w = 0, \int_Q w dx = 0 \right\} = \{ \nabla^s \varphi : \varphi \in C_{1\text{-per}}^\infty(\mathbb{R}^N; \mathbb{R}^N) \}. \tag{4.1}$$

Hence

$$\mathcal{Q}_{\mathcal{A}}g(x, v) = \inf \left\{ \int_Q g(x, v + \nabla^s \varphi(y)) dy : \varphi \in C_{1\text{-per}}^\infty(\mathbb{R}^N; \mathbb{R}^N) \right\}.$$

In particular

$$\mathcal{G}(\nabla^s u; D) = \int_D \mathcal{Q}^s f(x, u, \dots, \nabla^s u) dx. \tag{4.2}$$

Let $\{u_k\} \subset W^{s,p}(\Omega; \mathbb{R}^n)$ be any sequence such that $u_k \rightharpoonup u$ in $W^{s,p}(\Omega; \mathbb{R}^n)$. Extracting a subsequence, if necessary, we may assume that

$$\mathbf{u}_k := (u_k, \dots, \nabla^{s-1}u_k) \rightarrow \mathbf{u} := (u, \dots, \nabla^{s-1}u) \text{ in } L^p(D; E_{[s-1]}^n).$$

Since $\nabla^s u_k \rightharpoonup \nabla^s u$ in $L^p(D; E_s^n)$ and $\mathcal{A}\nabla^s u_k = 0$, by Lemma 3.1 there exists a p -equi-integrable sequence $\{V_k\} \subset L^p(D; E_s^n) \cap \ker \mathcal{A}$ such that $V_k \rightharpoonup \nabla^s u$ in $L^p(D; E_s^n)$ and

$$\liminf_{k \rightarrow \infty} \int_D g(x, V_k(x)) \, dx \leq \liminf_{k \rightarrow \infty} \int_{\Omega} f(x, u_k, \dots, \nabla^s u_k) \, dx.$$

Thus

$$\mathcal{G}(\nabla^s u; D) \leq \mathcal{F}(u; D). \tag{4.3}$$

To prove the converse inequality, fix $x_0 \in \Omega$ and $r > 0$, and consider any p -equi-integrable sequence $\{V_k\} \subset L^p(B(x_0; r); E_s^n) \cap \ker \mathcal{A}$ such that $V_k \rightharpoonup \nabla^s u$ in $L^p(B(x_0; r); E_s^n)$. An induction argument, similar to the one used in [22] to prove (4.1) above, shows that $\mathcal{A}V_k = 0$ if and only if there exists $\varphi_k \in W^{s,p}(B(x_0; r); \mathbb{R}^n)$ such that $\nabla^s \varphi_k = V_k$. By Lemmas 1.1–1.3 in [24], for any $\varphi \in W^{s,p}(B(x_0; r); \mathbb{R}^n)$ we may find a unique function $P \in C^\infty(\mathbb{R}^N; \mathbb{R}^n)$ whose components are polynomials of degree $s - 1$ such that

$$\int_{B(x_0, r)} \nabla^l (\varphi - P) \, dx = 0 \quad 0 \leq l \leq s - 1, \tag{4.4}$$

and a constant $C(n, N, s, p, r) > 0$ such that the following Poincaré type inequality holds

$$\|\varphi - P\|_{W^{s,p}(B(x_0, r); \mathbb{R}^n)} \leq C \|\nabla^s \varphi\|_{L^p(B(x_0, r); E_s^n)}. \tag{4.5}$$

Let P_k and P be the functions associated to φ_k and u , respectively, and satisfying (4.4, 4.5). Since $\nabla^s \varphi_k \rightharpoonup \nabla^s u$ in $L^p(B(x_0; r); E_s^n)$, we have that

$$\varphi_k - P_k \rightharpoonup u - P \text{ in } W^{s,p}(B(x_0; r); \mathbb{R}^n),$$

so

$$u_k := \varphi_k - P_k + P \rightharpoonup u \text{ in } W^{s,p}(B(x_0; r); \mathbb{R}^N).$$

Consider a subsequence of $\{V_k\}$ (not relabelled) such that the two sequences

$$\{(u_k, \dots, \nabla^{s-1} u_k, V_k)\} \quad \text{and} \quad \{(u, \dots, \nabla^{s-1} u, V_k)\}$$

generate the Young measure $\{\delta_{(u(x), \dots, \nabla^{s-1} u)} \otimes \nu_x\}_{x \in B(x_0, r)}$, and

$$(u_k, \dots, \nabla^{s-1} u_k) \rightarrow (u, \dots, \nabla^{s-1} u)$$

pointwise and in $L^p(B(x_0; r); E_{s-1}^n)$. Since $\{V_k\}$ is p -equi-integrable and u_k converge to u strongly in $W^{s-1,p}(\Omega; \mathbb{R}^N)$, it follows from Theorem 2.1 and the growth condition on f that

$$\lim_{k \rightarrow \infty} \int_{B(x_0, r)} f(u_k, \dots, \nabla^{s-1} u_k, V_k) \, dx = \lim_{k \rightarrow \infty} \int_{B(x_0, r)} g(x, V_k(x)) \, dx.$$

Thus

$$\mathcal{G}(\nabla^s u; B(x_0, r)) \geq \mathcal{F}(u; B(x_0, r)),$$

which, together with (4.3), yields

$$\mathcal{G}(\nabla^s u; B(x_0, r)) = \mathcal{F}(u; B(x_0, r)). \tag{4.6}$$

Since $\mathcal{F}(u; \cdot)$ and $\mathcal{G}(\nabla^s u; \cdot)$ are both traces of a Radon measures absolutely continuous with respect to $\mathcal{L}^N \llcorner \Omega$, by (4.2) and (4.6) we immediately obtain that

$$\mathcal{F}(u; D) = \mathcal{G}(\nabla^s u; D) = \int_D \mathcal{Q}f(x, u, \dots, \nabla^s u) dx.$$

□

Proof of Theorem 1.5. We only proof Theorem 1.5 for $1 \leq p < \infty$, the case $p = \infty$ being very similar.

For $v \in \mathbb{R}^{N^2-1}$ let

$$v = (v^{(1)}, \dots, v^{(N)}), \quad \text{where } v^{(i)} \in \mathbb{R}^N, \quad i = 1, \dots, N-1, \quad v^{(N)} \in \mathbb{R}^{N-1}.$$

Given a function $v \in L^p(\Omega; \mathbb{R}^{N^2-1})$ define the differential operator \mathcal{A} as follows

$$\mathcal{A}v := \begin{pmatrix} \text{curl } v^{(1)} \\ \vdots \\ \text{curl } v^{(N-1)} \\ \text{curl } (v^{(N)}, -v_1^{(1)} - \dots - v_{N-1}^{(N-1)}) \end{pmatrix}.$$

A straightforward calculation shows that \mathcal{A} satisfies the *constant-rank* property (1.1).

Given a Carathéodory function $f : \Omega \times \mathbb{R}^N \times \mathbb{R}^{N^2} \rightarrow [0, \infty)$, we define $\hat{f}(x, u, v)$, for $(x, u, v) \in \Omega \times \mathbb{R}^N \times \mathbb{R}^{N^2-1}$, as

$$\hat{f}(x, u, v) = f \left(x, u, \left(v^{(1)}, \dots, v^{(N-1)}, \begin{pmatrix} v^{(N)} \\ -v_1^{(1)} - \dots - v_{N-1}^{(N-1)} \end{pmatrix} \right) \right).$$

Let $u \in W^{1,p}(\Omega; \mathbb{R}^N)$, with $\text{div } u = 0$, and let $\{u_n\} \subset W^{1,p}(\Omega; \mathbb{R}^N)$ be such that $\text{div } u_n = 0$ and $u_n \rightharpoonup u$ in $W^{1,p}(\Omega; \mathbb{R}^N)$. By Lemma 3.1 there exists a p -equi-integrable sequence $\{V_n\} \subset L^p(\Omega; \mathbb{R}^{N^2-1}) \cap \ker \mathcal{A}$ such that $V_n \rightharpoonup v$ in $L^p(D; \mathbb{R}^{N^2-1})$ and

$$\liminf_{n \rightarrow \infty} \int_{\Omega} \hat{f}(x, u, V_n) dx \leq \liminf_{n \rightarrow \infty} \int_{\Omega} \hat{f}(x, u_n, v_n) dx = \liminf_{n \rightarrow \infty} \int_{\Omega} f(x, u_n, \nabla u_n) dx, \tag{4.7}$$

where

$$v_n := \left(\nabla u_n^{(1)}, \dots, \nabla u_n^{(N-1)}, \begin{pmatrix} \frac{\partial u_n^{(N)}}{\partial x_1} \\ \vdots \\ \frac{\partial u_n^{(N)}}{\partial x_{N-1}} \end{pmatrix} \right), \quad v := \left(\nabla u^{(1)}, \dots, \nabla u^{(N-1)}, \begin{pmatrix} \frac{\partial u^{(N)}}{\partial x_1} \\ \vdots \\ \frac{\partial u^{(N)}}{\partial x_{N-1}} \end{pmatrix} \right). \tag{4.8}$$

Define

$$\mathcal{G}(v; D) := \inf \left\{ \liminf_{n \rightarrow \infty} \int_D \hat{g}(x, V_n(x)) dx : \{V_n\} \subset L^p(D; \mathbb{R}^{N^2-1}) \cap \ker \mathcal{A} \text{ is } p\text{-equi-integrable,} \right. \\ \left. \text{and } V_n \rightharpoonup v \text{ in } L^p(D; \mathbb{R}^{N^2-1}) \right\},$$

where \hat{g} is the Carathéodory function defined by $\hat{g}(x, v) := \hat{f}(x, u(x), v)$. By Theorem 3.6 (and Cor. 3.2)

$$\mathcal{G}(v; \Omega) = \int_{\Omega} \mathcal{Q}_{\mathcal{A}} \hat{g}(x, v) \, dx, \tag{4.9}$$

where

$$\mathcal{Q}_{\mathcal{A}} \hat{g}(x, v(x)) := \inf \left\{ \int_Q \hat{f}(x, u(x), v(x) + w(y)) \, dy : w \in C_{1\text{-per}}^{\infty}(\mathbb{R}^N; \mathbb{R}^{N^2-1}) \cap \ker \mathcal{A}, \int_Q w(y) \, dy = 0 \right\}.$$

Now

$$w \in C_{1\text{-per}}^{\infty}(\mathbb{R}^N; \mathbb{R}^{N^2-1}) \cap \ker \mathcal{A} \quad \text{and} \quad \int_Q w(y) \, dy = 0$$

if and only if there exists $\varphi \in C_{1\text{-per}}^{\infty}(\mathbb{R}^N; \mathbb{R}^N)$ such that

$$w = \left(\nabla \varphi^{(1)}, \dots, \nabla \varphi^{(N-1)}, \begin{pmatrix} \frac{\partial \varphi^{(N)}}{\partial x_1} \\ \vdots \\ \frac{\partial \varphi^{(N)}}{\partial x_{N-1}} \end{pmatrix} \right)$$

and $\frac{\partial \varphi^{(N)}}{\partial x_N} = -\frac{\partial \varphi^{(1)}}{\partial x_1} - \dots - \frac{\partial \varphi^{(N-1)}}{\partial x_{N-1}}$. Hence

$$\begin{aligned} \mathcal{Q}_{\mathcal{A}} \hat{g}(x, v(x)) &= \inf \left\{ \int_Q f(x, u(x), \nabla u(x) + \nabla \varphi(y)) \, dy : \varphi \in C_{1\text{-per}}^{\infty}(\mathbb{R}^N; \mathbb{R}^N), \operatorname{div} \varphi = 0 \right\} \\ &= \bar{f}(x, u(x), \nabla u(x)). \end{aligned} \tag{4.10}$$

Thus, by (4.7, 4.9), and (4.10),

$$\int_{\Omega} \bar{f}(x, u(x), \nabla u(x)) \, dx = \mathcal{G}(v; \Omega) \leq \liminf_{n \rightarrow \infty} \int_{\Omega} \hat{f}(x, u, V_n) \, dx \leq \liminf_{n \rightarrow \infty} \int_{\Omega} f(x, u_n(x), \nabla u_n(x)) \, dx,$$

and, in turn,

$$\int_{\Omega} \bar{f}(x, u(x), \nabla u(x)) \, dx \leq \inf \left\{ \liminf_{n \rightarrow \infty} \int_{\Omega} f(x, u_n(x), \nabla u_n(x)) \, dx : \{u_n\} \subset W^{1,p}(\Omega; \mathbb{R}^N), \operatorname{div} u_n = 0, u_n \rightharpoonup u \text{ in } W^{1,p}(\Omega; \mathbb{R}^N) \right\}.$$

To prove the converse inequality, fix $\varepsilon > 0$. By the definition of $\mathcal{G}(v; \Omega)$, there exists a p -equi-integrable sequence $\{V_n\} \subset L^p(D; \mathbb{R}^{N^2-1}) \cap \ker \mathcal{A}$ such that $V_n \rightharpoonup v$ in $L^p(D; \mathbb{R}^{N^2-1})$ and

$$\int_{\Omega} \bar{f}(x, u(x), \nabla u(x)) \, dx + \varepsilon > \liminf_{n \rightarrow \infty} \int_D \hat{g}(x, V_n(x)) \, dx = \liminf_{n \rightarrow \infty} \int_D \hat{f}(x, u(x), V_n(x)) \, dx, \tag{4.11}$$

where we used for v the notation introduced in (4.8). Now $\mathcal{A}V_n = 0$ if and only if there exists $\varphi_n \in W^{1,p}(\Omega; \mathbb{R}^N)$ such that

$$V_n = \left(\nabla \varphi_n^{(1)}, \dots, \nabla \varphi_n^{(N-1)}, \begin{pmatrix} \frac{\partial \varphi_n^{(N)}}{\partial x_1} \\ \vdots \\ \frac{\partial \varphi_n^{(N)}}{\partial x_{N-1}} \end{pmatrix} \right)$$

and $\frac{\partial \varphi_n^{(N)}}{\partial x_N} = -\frac{\partial \varphi_n^{(1)}}{\partial x_1} - \dots - \frac{\partial \varphi_n^{(N-1)}}{\partial x_{N-1}}$. Since $\nabla \varphi_n \rightharpoonup \nabla u$ in $L^p(\Omega; \mathbb{R}^{N^2})$, we have that

$$\varphi_n - \frac{1}{|\Omega|} \int_{\Omega} \varphi_n(x) dx \rightharpoonup U \text{ in } W^{1,p}(\Omega; \mathbb{R}^N),$$

where $U = u + c$ for some constant $c \in \mathbb{R}^N$. So

$$u_n := \varphi_n - \frac{1}{|\Omega|} \int_{\Omega} \varphi_n(x) dx - c \rightharpoonup u \text{ in } W^{1,p}(\Omega; \mathbb{R}^N),$$

and $\operatorname{div} u_n = 0$. Consider a subsequence $\{V_{n_k}\}$ of $\{V_n\}$ such that

$$\lim_{k \rightarrow \infty} \int_{\Omega} \hat{f}(x, u(x), V_{n_k}(x)) dx = \liminf_{n \rightarrow \infty} \int_{\Omega} \hat{f}(x, u(x), V_n(x)) dx$$

and $\{(u_{n_k}, V_{n_k})\}$ and $\{(u, V_{n_k})\}$ generates the Young measure $\{\delta_{u(x)} \otimes \nu_x\}_{x \in \Omega}$. Since $\{V_{n_k}\}$ is p -equi-integrable and u_{n_k} converge to u strongly in $L^p(\Omega; \mathbb{R}^N)$, it follows from Theorem 2.1 and the growth condition on f that

$$\lim_{k \rightarrow \infty} \int_{\Omega} \hat{f}(x, u(x), V_{n_k}(x)) dx = \lim_{k \rightarrow \infty} \int_{\Omega} \hat{f}(x, u_{n_k}(x), V_{n_k}(x)) dx.$$

By (4.11)

$$\begin{aligned} \int_{\Omega} \bar{f}(x, u(x), \nabla u(x)) dx + \varepsilon &> \lim_{k \rightarrow \infty} \int_{\Omega} \hat{f}(x, u_{n_k}(x), V_{n_k}(x)) dx = \lim_{k \rightarrow \infty} \int_{\Omega} f(x, u_{n_k}(x), \nabla u_{n_k}(x)) dx \\ &\geq \inf \left\{ \liminf_{n \rightarrow \infty} \int_{\Omega} f(x, u_n(x), \nabla u_n(x)) dx : \{u_n\} \subset W^{1,p}(\Omega; \mathbb{R}^N), \right. \\ &\quad \left. \operatorname{div} u_n = 0, \quad u_n \rightharpoonup u \text{ in } W^{1,p}(\Omega; \mathbb{R}^N) \right\}. \end{aligned}$$

It now suffices to let $\varepsilon \rightarrow 0^+$. □

5. HOMOGENIZATION

In this section we will limit our analysis to the case where $1 < q < \infty$.

Lemma 5.1. *Let $f : \mathbb{R}^N \times \mathbb{R}^d \rightarrow [0, \infty)$ be a continuous function satisfying (A₁)-(A₂). Let $v \in L^q(D; \mathbb{R}^d) \cap \ker \mathcal{A}$, where $D \in \mathcal{O}(\Omega)$, $\varepsilon_k \rightarrow 0^+$, and let $\{\hat{v}_k\} \subset L^q(D_1; \mathbb{R}^d)$ be a sequence of functions such that*

$$\hat{v}_k \rightharpoonup v \text{ in } L^q(D_1; \mathbb{R}^d), \quad \mathcal{A}\hat{v}_k \rightarrow 0 \text{ in } W^{-1,q}(D_1; \mathbb{R}^l),$$

for some $D_1 \in \mathcal{O}(\Omega)$, with $D_1 \subset D$. Then we can find a q -equi-integrable sequence $\{v_k\} \subset L^q(D; \mathbb{R}^d) \cap \ker \mathcal{A}$ such that $\int_D v_k dx = \int_D v dx$,

$$v_k \rightharpoonup v \text{ in } L^q(D; \mathbb{R}^d), \quad \|\hat{v}_k - v_k\|_{L^s(D_1)} \rightarrow 0 \text{ for all } 1 \leq s < q \tag{5.1}$$

and

$$\begin{aligned} \liminf_{k \rightarrow \infty} \int_{D_1} f(x/\varepsilon_k, v_k(x)) \, dx &\leq \liminf_{k \rightarrow \infty} \int_{D_1} f(x/\varepsilon_k, \hat{v}_k(x)) \, dx, \\ \limsup_{k \rightarrow \infty} \int_{D \setminus D_1} |v_k(x)|^q \, dx &\leq \int_{D \setminus D_1} |v(x)|^q \, dx. \end{aligned} \tag{5.2}$$

Moreover, if $D = Q$, then $v_k = v + w_k$, with $w_k \in L^q_{1\text{-per}}(\mathbb{R}^N; \mathbb{R}^d) \cap \ker \mathcal{A}$.

Remark 5.2. Lemma 5.1 implies, in particular, that for every $v \in L^q(\Omega; \mathbb{R}^d) \cap \ker \mathcal{A}$

$$\Gamma - \liminf_{n \rightarrow \infty} \mathcal{F}_{\varepsilon_n}(v; D) = \inf \left\{ \liminf_{n \rightarrow \infty} \mathcal{F}_{\varepsilon_n}(v_n; D) : v_n \in L^q(D; \mathbb{R}^d) \cap \ker \mathcal{A}, \right. \\ \left. v_n \rightharpoonup v \text{ in } L^q(D; \mathbb{R}^d), \int_D v_n \, dx = \int_D v \, dx \right\},$$

and if $D = Q$ then

$$\Gamma - \liminf_{n \rightarrow \infty} \mathcal{F}_{\varepsilon_n}(v; Q) = \inf \left\{ \liminf_{n \rightarrow \infty} \mathcal{F}_{\varepsilon_n}(v + w_n; Q) : w_n \in L^q_{1\text{-per}}(\mathbb{R}^N; \mathbb{R}^d) \cap \ker \mathcal{A} \right. \\ \left. w_n \rightharpoonup 0 \text{ in } L^q(Q; \mathbb{R}^d), \int_Q w_n \, dx = 0 \right\}.$$

Proof of Lemma 5.1. Let $g(x) := x$ in Q and extend it periodically to \mathbb{R}^N with period 1. Set $g_k(x) := g(x/\varepsilon_k)$. Since $\{g_k\}$ is bounded in L^∞ and $\hat{v}_k \rightharpoonup v$ in $L^q(D_1; \mathbb{R}^d)$, by Theorem 2.1 there exists a subsequence $\{\varepsilon_n\}$ of $\{\varepsilon_k\}$ such that

$$\{(g_n(x), \hat{v}_n(x))\} \text{ generates a Young measure } \{\nu_x\}$$

and

$$\lim_{n \rightarrow \infty} \int_{D_1} f(x/\varepsilon_n, \hat{v}_n(x)) \, dx = \liminf_{k \rightarrow \infty} \int_{D_1} f(x/\varepsilon_k, \hat{v}_k(x)) \, dx.$$

For $i \in \mathbb{N}$ let

$$F_i := \left\{ x \in D_1 : \text{dist}(x, \partial D_1) < \frac{1}{i} \right\}$$

and consider cut-off functions θ_i with compact support in D_1 and such that $\theta_i \equiv 1$ in $D_1 \setminus F_i$. Set $w_{i,n} := \theta_i(\hat{v}_n - v) \in L^q(Q; \mathbb{R}^d)$. Then we can proceed as in the proof of Lemma 3.1 to find a q -equi-integrable sequence $\{v_i := v + w_i\}$, where $\{w_i\}$ satisfies (3.3, 5.1) holds, and the two sequences

$$\{(g_{n_i}(x), v_i(x))\} \quad \text{and} \quad \{(g_{n_i}(x), \hat{v}_{n_i}(x))\}$$

generate the same Young measure $\{\nu_x\}$. Hence by Theorem 2.1

$$\begin{aligned} \lim_{i \rightarrow \infty} \int_{D_1} f(x/\varepsilon_{n_i}, v_i(x)) \, dx &= \int_{D_1} \left(\int_{\mathbb{R}^N \times \mathbb{R}^d} f(X, V) d\nu_x(X, V) \right) dx \leq \lim_{i \rightarrow \infty} \int_{D_1} f(x/\varepsilon_{n_i}, \hat{v}_{n_i}(x)) \, dx \\ &= \liminf_{k \rightarrow \infty} \int_{D_1} f(x/\varepsilon_k, \hat{v}_k(x)) \, dx, \end{aligned}$$

where we have used (A_2) , and the facts that $\{v_i(x)\}$ is q -equi-integrable over D_1 , and that f is a continuous function.

To prove the second inequality in (5.2), we remark that by (3.3) and the fact that $\hat{w}_i = \theta_i(\hat{v}_{n_i} - v) \equiv 0$ outside D_1 , we have for all $1 \leq s < q$

$$\|v_i - v\|_{L^s(D \setminus D_1)} = \|\hat{w}_i - w_i\|_{L^s(D \setminus D_1)} \rightarrow 0.$$

Hence $\{v_i(x)\}$ generates the Young measure $\{\mu_x = \delta_{v(x)}\}$ on $D \setminus D_1$, and since $\{v_i\}$ is q -equi-integrable we have that

$$\limsup_{i \rightarrow \infty} \int_{D \setminus D_1} |v_i(x)|^q dx = \int_{D \setminus D_1} |Y|^q d\mu_x(Y) dx = \int_{D \setminus D_1} |v(x)|^q dx.$$

To complete the proof it suffices to define $v_k := v_{n_i}$ for each $n_i \leq k < n_{i+1}$. Clearly

$$\liminf_{k \rightarrow \infty} \int_{D_1} f(x/\varepsilon_k, v_k(x)) dx \leq \liminf_{i \rightarrow \infty} \int_{D_1} f(x/\varepsilon_{n_i}, v_i(x)) dx.$$

□

Lemma 5.3. *Let $\varepsilon_n \rightarrow 0^+$ and let $\mathcal{R}(\Omega)$ be the family of all finite unions of open cubes contained in Ω and with vertices in \mathbb{Q}^N . Then there exists a subsequence $\{\varepsilon_{n_k}\}$ of $\{\varepsilon_n\}$ such that the Γ -limit*

$$\Gamma - \lim_{k \rightarrow \infty} \mathcal{F}_{\varepsilon_{n_k}}(v; R)$$

exists for all $v \in L^q(R; \mathbb{R}^d) \cap \ker \mathcal{A}$ and for all $R \in \mathcal{R}(\Omega)$.

Proof of Lemma 5.3. Fix $R \in \mathcal{R}(\Omega)$. For simplicity set $\mathcal{F}_n := \mathcal{F}_{\varepsilon_n}$ and let \mathcal{B} denote the closed unit ball of $L^q(R; \mathbb{R}^d)$. For each $l \in \mathbb{N}$ consider

$$l\mathcal{B} := \{v \in L^q(R; \mathbb{R}^d) : \|v\|_{L^q} \leq l\}.$$

Since $q > 1$ the dual of $L^q(R; \mathbb{R}^d)$ is separable, and hence the space $l\mathcal{B}$ endowed with the weak topology is metrizable. Let d_l be any metric which generates the L^q -weak topology. Consider $l = 1$ and apply Proposition 2.4 to the sequence of functionals $\{\mathcal{F}_n(\cdot; R)\}$ restricted to $(\mathcal{B} \cap \ker \mathcal{A}, d_1)$. Then we can find an increasing sequence of integers $\{n_j^1\}$ such that

$$\Gamma(d_1) - \lim_{j \rightarrow \infty} \mathcal{F}_{n_j^1}(v; R)$$

exists for all $v \in \mathcal{B} \cap \ker \mathcal{A}$. We now proceed recursively, so that given $l \in \mathbb{N}$ we apply Proposition 2.4 to the sequence of functionals $\{\mathcal{F}_{n_j^{l-1}}(\cdot; R)\}$ restricted to $(l\mathcal{B} \cap \ker \mathcal{A}, d_l)$ to obtain a subsequence $\{n_j^l\}$ of $\{n_j^{l-1}\}$ such that

$$\Gamma(d_l) - \lim_{j \rightarrow \infty} \mathcal{F}_{n_j^l}(v; R)$$

exists for all $v \in l\mathcal{B} \cap \ker \mathcal{A}$. Let $n_k := n_k^k$. Since $\{n_k\}$ is a subsequence of all $\{n_j^l\}$ we have that for each $l \in \mathbb{N}$

$$\Gamma(d_l) - \lim_{k \rightarrow \infty} \mathcal{F}_{n_k}(v; R)$$

exists for all $v \in l\mathcal{B} \cap \ker \mathcal{A}$.

We claim that the Γ -limit

$$\Gamma - \lim_{k \rightarrow \infty} \mathcal{F}_{n_k}(v; R) \tag{5.3}$$

exists for all $v \in L^q(R; \mathbb{R}^d) \cap \ker \mathcal{A}$. Indeed assume by contradiction that this is not the case. Then there exists $v \in L^q(R; \mathbb{R}^d) \cap \ker \mathcal{A}$ for which

$$\mathcal{F}^-(v; R) := \Gamma - \liminf_{k \rightarrow \infty} \mathcal{F}_{n_k}(v; R) < \mathcal{F}^+(v; R) := \Gamma - \limsup_{k \rightarrow \infty} \mathcal{F}_{n_k}(v; R).$$

Let $v_k \in L^q(R; \mathbb{R}^d) \cap \ker \mathcal{A}$ be such that $v_k \rightharpoonup v$ in $L^q(R; \mathbb{R}^d)$ and

$$\liminf_{k \rightarrow \infty} \mathcal{F}_{n_k}(v_k; R) = \mathcal{F}^-(v; R).$$

Since $v_k \rightharpoonup v$ in $L^q(R; \mathbb{R}^d)$, we may find an integer l_0 such that $v_k, v \in l_0 \mathcal{B} \cap \ker \mathcal{A}$ for all $k \in \mathbb{N}$. Consequently

$$d_{l_0}(v_k, v) \rightarrow 0 \quad \text{as } k \rightarrow \infty,$$

and thus

$$\Gamma(d_{l_0}) - \liminf_{k \rightarrow \infty} \mathcal{F}_{n_k}(v; R) \leq \liminf_{k \rightarrow \infty} \mathcal{F}_{n_k}(v_k; R) = \mathcal{F}^-(v; R) < \mathcal{F}^+(v; R) \leq \Gamma(d_{l_0}) - \limsup_{k \rightarrow \infty} \mathcal{F}_{n_k}(v; R),$$

which contradicts the existence of the Γ -limit $\Gamma(d_{l_0}) - \lim_{k \rightarrow \infty} \mathcal{F}_{n_k}(v; R)$, and where we have used the fact that

$$\begin{aligned} \mathcal{F}^+(v; R) &= \inf \left\{ \limsup_{k \rightarrow \infty} \mathcal{F}_{n_k}(z_k; R) : z_k \in L^q(R; \mathbb{R}^d) \cap \ker \mathcal{A}, \quad z_k \rightharpoonup v \text{ in } L^q(R; \mathbb{R}^d) \right\} \\ &\leq \Gamma(d_{l_0}) - \limsup_{k \rightarrow \infty} \mathcal{F}_{n_k}(v; R) \\ &= \inf \left\{ \limsup_{k \rightarrow \infty} \mathcal{F}_{n_k}(z_k; R) : z_k \in l_0 \mathcal{B} \cap \ker \mathcal{A}, \quad z_k \rightharpoonup v \text{ in } L^q(R; \mathbb{R}^d) \right\}. \end{aligned}$$

Hence (5.3) holds. To conclude the proof of the lemma it suffices to observe that since the family $\mathcal{R}(\Omega)$ is countable, with a diagonal process it is possible to extract a further subsequence for which (5.3) holds for all $R \in \mathcal{R}(\Omega)$. □

Remark 5.4. The previous proof asserts that for any given $D \in \mathcal{O}(\Omega)$ and $\varepsilon_n \rightarrow 0^+$ there exists a subsequence $\{\varepsilon_{n_k}\}$ (depending on the particular set D) of $\{\varepsilon_n\}$ such that such that the Γ -limit

$$\Gamma - \lim_{k \rightarrow \infty} \mathcal{F}_{\varepsilon_{n_k}}(v; D)$$

exists for all $v \in L^q(D; \mathbb{R}^d) \cap \ker \mathcal{A}$.

Lemma 5.5. *Assume that conditions (A₁)-(A₂) hold. Given $\varepsilon_n \rightarrow 0^+$, let $\{\varepsilon_{n_k}\}$ be as in Lemma 5.3, and for any $D \in \mathcal{O}(\Omega)$ set*

$$\mathcal{F}^-(\cdot; D) := \Gamma - \liminf_{k \rightarrow \infty} \mathcal{F}_{\varepsilon_{n_k}}(\cdot; D).$$

Then $\mathcal{F}^-(v; \cdot)$ is the trace of a Radon measure.

Proof of Lemma 5.5. We start by establishing inner regularity. Precisely, we claim that for any $v \in L^q(D; \mathbb{R}^d) \cap \ker \mathcal{A}$ and $D \in \mathcal{O}(\Omega)$

$$\mathcal{F}^-(v; D) = \sup \{ \mathcal{F}^-(v; R) : R \in \mathcal{R}(\Omega), R \subset D \} = \lim_{R \nearrow D} \mathcal{F}^-(v; R), \tag{5.4}$$

where the limit is taken over all finite unions of cubes $R \in \mathcal{R}(\Omega)$ with $R \subset D$. For fixed $\eta > 0$ there exists $\delta > 0$ such that

$$\int_{D_0} C(1 + |v(x)|^q) dx < \eta \tag{5.5}$$

for any measurable set $D_0 \subset D$, with $|D_0| < \delta$, and where C is the constant given in (A_2) . Let $R \in \mathcal{R}(\Omega)$, with $R \subset D$ and $|D \setminus R| < \delta$, and, in light of Lemma 5.3, consider a sequence $\{\hat{v}_k\} \subset L^q(R; \mathbb{R}^d) \cap \ker \mathcal{A}$, with $\hat{v}_k \rightharpoonup v$ in $L^q(R; \mathbb{R}^d)$, and such that

$$\lim_{k \rightarrow \infty} \mathcal{F}_{\varepsilon_{n_k}}(\hat{v}_k; R) = \mathcal{F}^-(v; R).$$

By Lemma 5.1 there exists a q -equi-integrable sequence $\{v_k\} \subset L^q(D; \mathbb{R}^d) \cap \ker \mathcal{A}$ such that

$$v_k \rightharpoonup v \quad \text{in } L^q(D; \mathbb{R}^d), \quad \int_D v_k \, dx = \int_D v \, dx,$$

and

$$\begin{aligned} \liminf_{k \rightarrow \infty} \int_R f(x/\varepsilon_{n_k}, v_k(x)) \, dx &\leq \lim_{k \rightarrow \infty} \int_R f(x/\varepsilon_{n_k}, \hat{v}_k(x)) \, dx, \\ \limsup_{k \rightarrow \infty} \int_{D \setminus R} |v_k(x)|^q \, dx &\leq \int_{D \setminus R} |v(x)|^q \, dx. \end{aligned}$$

Hence

$$\begin{aligned} \mathcal{F}^-(v; D) &\leq \liminf_{k \rightarrow \infty} \int_D f(x/\varepsilon_{n_k}, v_k(x)) \, dx \leq \lim_{k \rightarrow \infty} \int_R f(x/\varepsilon_{n_k}, \hat{v}_k(x)) \, dx + \limsup_{k \rightarrow \infty} \int_{D \setminus R} C(1 + |v_k(x)|^q) \, dx \\ &\leq \mathcal{F}^-(v; R) + \int_{D \setminus R} C(1 + |v(x)|^q) \, dx \leq \mathcal{F}^-(v; R) + \eta, \end{aligned}$$

where we have used (A_2) and (5.5). Consequently

$$\mathcal{F}^-(v; D) \leq \sup \{ \mathcal{F}^-(v; R) : R \in \mathcal{R}(\Omega), R \subset D \} + \eta,$$

and letting $\eta \rightarrow 0^+$ we obtain one inequality in (5.4). To show the opposite inequality, note that if $\{v_k\} \subset L^q(D; \mathbb{R}^d) \cap \ker \mathcal{A}$, with $v_k \rightharpoonup v$ in $L^q(D; \mathbb{R}^d)$, then the restriction of v_k to R belongs to $L^q(R; \mathbb{R}^d) \cap \ker \mathcal{A}$, and $v_k \rightharpoonup v$ in $L^q(R; \mathbb{R}^d)$. Therefore

$$\mathcal{F}^-(v; R) \leq \liminf_{k \rightarrow \infty} \int_R f(x/\varepsilon_{n_k}, v_k(x)) \, dx \leq \liminf_{k \rightarrow \infty} \int_D f(x/\varepsilon_{n_k}, v_k(x)) \, dx$$

and by taking the infimum over all such sequences we get that

$$\mathcal{F}^-(v; R) \leq \mathcal{F}^-(v; D), \tag{5.6}$$

and in turn (5.4) holds.

In order to prove that $\mathcal{F}^-(v; \cdot)$ is the trace of a Radon measure, as it is usual it suffices to prove subadditivity for nested sets (see [3, 21]). Let $B \subset\subset C \subset\subset D$. By (5.4) for fixed $\eta > 0$ we find $R \in \mathcal{R}(\Omega)$ such that $R \subset D$ and

$$\mathcal{F}^-(v; D) \leq \eta + \mathcal{F}^-(v; R).$$

Construct $R_1, R_2 \in \mathcal{R}(\Omega)$ with

$$R \subset R_1 \cup R_2, \quad R_1 \subset D \setminus \overline{B} \quad \text{and} \quad R_2 \subset C.$$

By (5.6) we have

$$\mathcal{F}^-(v; D) \leq \eta + \mathcal{F}^-(v; R) \leq \eta + \mathcal{F}^-(v; R_1 \cup R_2). \tag{5.7}$$

By the definition of Γ -convergence and Lemma 5.1 there exist $v_k \in L^q(R_1; \mathbb{R}^d) \cap \ker \mathcal{A}$ and $w_k \in L^q(R_2; \mathbb{R}^d) \cap \ker \mathcal{A}$, with $v_k \rightharpoonup v$ in $L^q(R_1; \mathbb{R}^d)$ and $w_k \rightharpoonup v$ in $L^q(R_2; \mathbb{R}^d)$, such that

$$\mathcal{F}^-(v; R_1) = \lim_{k \rightarrow \infty} \mathcal{F}_{\tilde{\varepsilon}_{n_k}}(v_k; R_1), \quad \mathcal{F}^-(v; R_2) = \lim_{k \rightarrow \infty} \mathcal{F}_{\tilde{\varepsilon}_{n_k}}(w_k; R_2), \tag{5.8}$$

where $\{\tilde{\varepsilon}_{n_k}\}$ is a subsequence of $\{\varepsilon_{n_k}\}$ and $\{v_k\}, \{w_k\}$ are q -equi-integrable over R_1 and R_2 , respectively. Let θ_j be smooth cut-off functions which are equal to 1 on B and 0 on $D \setminus \overline{C}$, and such that $|\{0 < \theta_j < 1\}| \rightarrow 0$ as $j \rightarrow \infty$. Set

$$\hat{V}_{j,k} := (1 - \theta_j)v_k + \theta_j w_k.$$

For j fixed

$$\mathcal{A}\hat{V}_{j,k} = (1 - \theta_j)\mathcal{A}v_k + \theta_j\mathcal{A}w_k - \sum_{i=1}^N A^{(i)}v_k \frac{\partial \theta_j}{\partial x_i} + \sum_{i=1}^N A^{(i)}w_k \frac{\partial \theta_j}{\partial x_i} \rightarrow 0$$

in $W^{-1,q}(R_1 \cup R_2; \mathbb{R}^l)$ strong, because $\|v_k - w_k\|_{W^{-1,q}(B \setminus \overline{C}; \mathbb{R}^l)} \rightarrow 0$ as $k \rightarrow \infty$. Diagonalize to get $\hat{V}_j := \hat{V}_{j,k_j}$ such that

$$\hat{V}_j \rightharpoonup v \text{ in } L^q(R_1 \cup R_2; \mathbb{R}^d), \quad \mathcal{A}\hat{V}_j \rightarrow 0 \text{ in } W^{-1,q}(R_1 \cup R_2; \mathbb{R}^l).$$

By Lemma 5.1 we can find $V_j \in L^q(R_1 \cup R_2; \mathbb{R}^d) \cap \ker \mathcal{A}$ such that $V_j \rightharpoonup v$ in $L^q(R_1 \cup R_2; \mathbb{R}^d)$ and

$$\liminf_{j \rightarrow \infty} \int_{R_1 \cup R_2} f(x/\tilde{\varepsilon}_{n_{k_j}}, V_j(x)) \, dx \leq \liminf_{j \rightarrow \infty} \int_{R_1 \cup R_2} f(x/\tilde{\varepsilon}_{n_{k_j}}, \hat{V}_j(x)) \, dx.$$

Consequently, by (5.7)

$$\begin{aligned} \mathcal{F}^-(v; D) &\leq \eta + \mathcal{F}^-(R_1 \cup R_2; D) \leq \eta + \liminf_{j \rightarrow \infty} \int_{R_1 \cup R_2} f(x/\tilde{\varepsilon}_{n_{k_j}}, V_j(x)) \, dx \\ &\leq \eta + \liminf_{j \rightarrow \infty} \int_{R_1 \cup R_2} f(x/\tilde{\varepsilon}_{n_{k_j}}, \hat{V}_j(x)) \, dx \leq \eta + \limsup_{j \rightarrow \infty} \int_{R_1} f(x/\tilde{\varepsilon}_{n_{k_j}}, v_{k_j}(x)) \, dx \\ &\quad + \limsup_{j \rightarrow \infty} \int_{R_2} f(x/\tilde{\varepsilon}_{n_{k_j}}, w_{k_j}(x)) \, dx + \limsup_{j \rightarrow \infty} \int_{\{0 < \theta_j < 1\}} C(1 + |w_{k_j}(x)|^q + |v_{k_j}(x)|^q) \, dx \\ &\leq \eta + \mathcal{F}^-(v; R_1) + \mathcal{F}^-(v; R_2) \leq \eta + \mathcal{F}^-(v; D \setminus \overline{B}) + \mathcal{F}^-(v; C), \end{aligned}$$

where we have used (5.6) and the fact that in (5.8) inferior limits are actually limits. It now suffices to let $\eta \rightarrow 0^+$. □

Lemma 5.6. *Under conditions (A₁)-(A₂), for \mathcal{L}^N a.e. $x_0 \in \Omega$ we have*

$$\frac{d\mathcal{F}^-(v; \cdot)}{d\mathcal{L}^N}(x_0) = f_{\text{hom}}(v(x_0)).$$

Proof of Lemma 5.6. We divide the proof in three steps.

Given $\varepsilon_n \rightarrow 0^+$, let $\{\varepsilon_{n_k}\}$ be as in Lemma 5.3. In order to simplify the notations, in the proof of this lemma we will represent $\{\varepsilon_{n_k}\}$ simply by $\{\varepsilon\}$.

Step 1. We claim that

$$\mathcal{F}^-(v(\cdot - x_0); D + x_0) = \mathcal{F}^-(v; D).$$

The proof is similar to the one of Lemma 3.9 in [11]. We present it here for the convenience of the reader. Let $v_\varepsilon \in L^q(D; \mathbb{R}^d) \cap \ker \mathcal{A}$ be such that $v_\varepsilon \rightharpoonup v$ in $L^q(D; \mathbb{R}^d)$ and

$$\mathcal{F}^-(v; D) = \liminf_{\varepsilon \rightarrow 0} \mathcal{F}_\varepsilon(v_\varepsilon; D). \tag{5.9}$$

Consider the sequence $z_\varepsilon := [x_0/\varepsilon] \in \mathbb{Z}^N$, so that $x_\varepsilon := z_\varepsilon \varepsilon$ converges to x_0 . Here $[z] := ([z_1], \dots, [z_N])$, with $[z_i]$ denoting the integer part of $z_i \in \mathbb{R}$. By the periodicity of f ,

$$\mathcal{F}_\varepsilon(v_\varepsilon; D) = \int_D f\left(\frac{x + x_\varepsilon}{\varepsilon}, v_\varepsilon(x)\right) dx = \int_{D+x_\varepsilon} f\left(\frac{y}{\varepsilon}, v_\varepsilon(y - x_\varepsilon)\right) dy.$$

Let $B \subset\subset D$. For ε sufficiently small we have that $D + x_\varepsilon \supset B + x_0$, and thus

$$\mathcal{F}_\varepsilon(v_\varepsilon; D) \geq \int_{B+x_0} f\left(\frac{y}{\varepsilon}, v_\varepsilon(y - x_\varepsilon)\right) dy. \tag{5.10}$$

Since $v_\varepsilon(\cdot - x_\varepsilon) \rightharpoonup v(\cdot - x_0)$ in $L^q(B + x_0; \mathbb{R}^d)$, and $v_\varepsilon(\cdot - x_\varepsilon) \in L^q(B + x_0; \mathbb{R}^d) \cap \ker \mathcal{A}$, by (5.9, 5.10), we obtain

$$\mathcal{F}^-(v; D) \geq \mathcal{F}^-(v(\cdot - x_0); B + x_0).$$

By letting $R \nearrow D + x_0$, $R \in \mathcal{R}(\Omega)$, setting $B := R - x_0$ above, we obtain by (5.4)

$$\mathcal{F}^-(v; D) \geq \mathcal{F}^-(v(\cdot - x_0); D + x_0).$$

The converse inequality follows in a similar way.

Step 2. Next, we show that

$$\frac{d\mathcal{F}^-(v; \cdot)}{d\mathcal{L}^N}(x_0) \geq f_{\text{hom}}(v(x_0)) \quad \text{for } \mathcal{L}^N \text{ a.e. } x_0 \in \Omega.$$

Fix $x_0 \in \Omega$ such that

$$\lim_{r \rightarrow 0^+} \frac{1}{r^N} \int_{Q(x_0, r)} |v(x) - v(x_0)|^q dx = 0 \tag{5.11}$$

and

$$\frac{d\mathcal{F}^-(v; \cdot)}{d\mathcal{L}^N}(x_0) = \lim_{r \rightarrow 0^+} \liminf_{\varepsilon \rightarrow 0^+} \frac{1}{r^N} \int_{rQ} f\left(\frac{x}{\varepsilon}, v_{\varepsilon, r}(x)\right) dx < \infty,$$

where we have used Step 1 and Lemma 5.5, and where we have chosen the radii $r \rightarrow 0^+$ such that $\mathcal{F}^-(v(\cdot + x_0); \partial(rQ)) = 0$. Here $v_{\varepsilon, r} \in L^q(rQ; \mathbb{R}^d) \cap \ker \mathcal{A}$ and $v_{\varepsilon, r} \rightharpoonup v(\cdot + x_0)$ in $L^q(rQ; \mathbb{R}^d)$ as $\varepsilon \rightarrow 0^+$. Then

$$\frac{d\mathcal{F}^-(v; \cdot)}{d\mathcal{L}^N}(x_0) = \lim_{r \rightarrow 0^+} \liminf_{\varepsilon \rightarrow 0^+} \int_Q f\left(\frac{r}{\varepsilon} y, v(x_0) + w_{\varepsilon, r}(y)\right) dy$$

where $w_{\varepsilon, r}(y) := v_{\varepsilon, r}(ry) - v(x_0)$. As in the proof of Lemma 3.5, we have that $w_{\varepsilon, r} \rightharpoonup 0$ in $L^q(Q; \mathbb{R}^d)$ if we first let $\varepsilon \rightarrow 0$ and then $r \rightarrow 0^+$. Diagonalize to get $\hat{w}_k \in L^q(Q; \mathbb{R}^d) \cap \ker \mathcal{A}$ such that $\hat{w}_k \rightharpoonup 0$ in $L^q(Q; \mathbb{R}^d)$,

$$\frac{d\mathcal{F}^-(v; \cdot)}{d\mathcal{L}^N}(x_0) = \lim_{k \rightarrow \infty} \int_Q f(s_k y, v(x_0) + \hat{w}_k(y)) dy,$$

and where $s_k := 1/\varepsilon_k \rightarrow \infty$. By Lemma 5.1, applied to the Carathéodory function $h(x, v) := f(x, v(x_0) + v)$, there exists a q -equi-integrable sequence $\{w_k\} \subset L^q_{1\text{-per}}(\mathbb{R}^N; \mathbb{R}^d) \cap \ker \mathcal{A}$ such that

$$w_k \rightharpoonup 0 \quad \text{in } L^q(Q; \mathbb{R}^d), \quad \int_Q w_k \, dy = 0,$$

and

$$\liminf_{k \rightarrow \infty} \int_Q f(s_k y, v(x_0) + w_k(y)) \, dy \leq \lim_{k \rightarrow \infty} \int_Q f(s_k y, v(x_0) + \hat{w}_k(y)) \, dy.$$

Consequently

$$\frac{d\mathcal{F}^-(v; \cdot)}{d\mathcal{L}^N}(x_0) \geq \liminf_{k \rightarrow \infty} \int_Q f(s_k y, v(x_0) + w_k(y)) \, dy \geq \liminf_{i \rightarrow \infty} \liminf_{k \rightarrow \infty} \int_Q f(s_k y, v(x_0) + \theta_i(y)w_k(y)) \, dy,$$

where $0 \leq \theta_i \leq 1$ are smooth cut-off functions with compact support in Q such that $\theta_i \equiv 1$ in $(1 - 1/i)Q$, and where we used the q -equi-integrability of $\{w_k\}$ and (A_2) . Then $\theta_i w_k \rightharpoonup 0$ in $L^q(Q; \mathbb{R}^d)$ as $k \rightarrow \infty$ and $i \rightarrow \infty$, in this order, and

$$\lim_{i \rightarrow \infty} \lim_{k \rightarrow \infty} \mathcal{A}(\theta_i w_k) = 0 \quad \text{in } W^{-1,q}(Q; \mathbb{R}^l).$$

Diagonalize to get $U_i := \theta_i w_{k_i}$ extended by zero outside Q , such that $U_i \rightharpoonup 0$ in $L^q(Q; \mathbb{R}^d)$, $\mathcal{A}U_i \rightarrow 0$ in $W^{-1,q}(Q; \mathbb{R}^l)$ as $i \rightarrow \infty$, and

$$\liminf_{i \rightarrow \infty} \liminf_{k \rightarrow \infty} \int_Q f(s_k y, v(x_0) + \theta_i(y)w_k(y)) \, dy = \liminf_{i \rightarrow \infty} \int_Q f(s_{k_i}, v(x_0) + U_i(y)) \, dy.$$

Thus

$$\begin{aligned} \frac{d\mathcal{F}^-(v; \cdot)}{d\mathcal{L}^N}(x_0) &\geq \liminf_{i \rightarrow \infty} \frac{1}{s_{k_i}^N} \int_{s_{k_i} Q} f\left(x, v(x_0) + U_i\left(\frac{x}{s_{k_i}}\right)\right) \, dx \\ &\geq \liminf_{i \rightarrow \infty} \frac{1}{s_{k_i}^N} \int_{([s_{k_i}] + 1)Q} f\left(x, v(x_0) + U_i\left(\frac{x}{s_{k_i}}\right)\right) \, dx \\ &\quad - \limsup_{i \rightarrow \infty} \frac{1}{s_{k_i}^N} \int_{([s_{k_i}] + 1)Q \setminus s_{k_i} Q} f\left(x, v(x_0) + U_i\left(\frac{x}{s_{k_i}}\right)\right) \, dx, \end{aligned}$$

where $[s_{k_i}]$ denotes the integer part of s_{k_i} . We claim that the last limit is zero. Indeed

$$\frac{1}{s_{k_i}^N} \int_{([s_{k_i}] + 1)Q \setminus s_{k_i} Q} f\left(x, v(x_0) + U_i\left(\frac{x}{s_{k_i}}\right)\right) \, dx = \int_{\frac{([s_{k_i}] + 1)Q \setminus Q}{s_{k_i}}} f(s_{k_i} y, v(x_0) + U_i(y)) \, dy.$$

Since $([s_{k_i}] + 1)/s_{k_i} \rightarrow 1$, we have that

$$\left| \frac{([s_{k_i}] + 1)Q \setminus Q}{s_{k_i}} \right| = \left(\frac{([s_{k_i}] + 1)}{s_{k_i}} \right)^N - 1 \rightarrow 0,$$

and thus the claim follows from the q -equi-integrability of $\{U_i\}$ and (A_2) . Hence, setting

$$m_i := 1/s_{k_i}, \quad n_i := [s_{k_i}] + 1 \in \mathbb{N},$$

we obtain

$$\frac{d\mathcal{F}^-(v; \cdot)}{d\mathcal{L}^N}(x_0) \geq \liminf_{i \rightarrow \infty} \frac{1}{n_i^N} \int_{n_i Q} f(x, v(x_0) + U_i(m_i x)) \, dx = \liminf_{i \rightarrow \infty} \int_Q f(n_i y, v(x_0) + U_i(n_i m_i y)) \, dy.$$

We claim that

$$U_i(n_i m_i \cdot) \rightarrow 0 \quad \text{in } L^q(Q; \mathbb{R}^d), \quad \mathcal{A}U_i(n_i m_i \cdot) \rightarrow 0 \quad \text{in } W^{-1,q}(Q; \mathbb{R}^l) \tag{5.12}$$

as $i \rightarrow \infty$. Assuming that the claim holds, by Lemma 5.1 there exists a q -equi-integrable sequence $\{V_i\} \subset L^q_{1\text{-per}}(\mathbb{R}^N; \mathbb{R}^d) \cap \ker \mathcal{A}$ such that

$$V_i \rightarrow 0 \quad \text{in } L^q(Q; \mathbb{R}^d), \quad \int_Q V_i \, dy = 0,$$

and

$$\begin{aligned} \frac{d\mathcal{F}^-(v; \cdot)}{d\mathcal{L}^N}(x_0) &\geq \liminf_{i \rightarrow \infty} \int_Q f(n_i y, v(x_0) + U_i(n_i m_i y)) \, dy \geq \liminf_{i \rightarrow \infty} \int_Q f(n_i y, v(x_0) + V_i(y)) \, dy \\ &= \liminf_{i \rightarrow \infty} \frac{1}{n_i^N} \int_{n_i Q} f\left(x, v(x_0) + V_i\left(\frac{x}{n_i}\right)\right) \, dx \geq f_{\text{hom}}(v(x_0)), \end{aligned}$$

and where we have used the facts that

$$V_i\left(\frac{1}{n_i} \cdot\right) \in L^q_{n_i\text{-per}}(\mathbb{R}^N; \mathbb{R}^d) \cap \ker \mathcal{A}, \quad \int_{n_i Q} V_i\left(\frac{1}{n_i} y\right) \, dy = 0.$$

Thus it remains to show (5.12). If $\varphi \in C_c^\infty(Q)$ then

$$\begin{aligned} \int_Q U_i(n_i m_i y) \varphi(y) \, dy &= \frac{1}{(n_i m_i)^N} \int_{n_i m_i Q} U_i(x) \varphi\left(\frac{x}{n_i m_i}\right) \, dx \\ &= \frac{1}{(n_i m_i)^N} \left(\int_Q U_i(x) \varphi(x) \, dx + \int_Q U_i(x) \left(\varphi\left(\frac{x}{n_i m_i}\right) - \varphi(x) \right) \, dx \right), \end{aligned}$$

where we have used the fact that $U_i(x) \equiv 0$ in $n_i m_i Q \setminus Q$. Since $U_i \rightarrow 0$ in $L^q(Q; \mathbb{R}^d)$ and $n_i m_i \rightarrow 1$ the first integral on the right hand side of the previous inequality tends to zero as $i \rightarrow \infty$. By Hölder's inequality

$$\left| \int_Q U_i(x) \left(\varphi\left(\frac{x}{n_i m_i}\right) - \varphi(x) \right) \, dx \right| \leq \left(\sup_l \|U_l\|_{L^q(Q)} \right) \left(\int_Q \left| \varphi\left(\frac{x}{n_i m_i}\right) - \varphi(x) \right|^{q'} \, dx \right)^{1/q'}.$$

Since φ is bounded we can apply Lebesgue Dominated Convergence Theorem to conclude that the right hand side approaches zero as $i \rightarrow \infty$. In a similar way we can show that

$$\mathcal{A}U_i(n_i m_i \cdot) \rightarrow 0 \quad \text{in } W^{-1,q}(Q; \mathbb{R}^l) \text{ as } i \rightarrow \infty.$$

We omit the details.

Step 3. To conclude the proof of the lemma it remains to show that

$$\frac{d\mathcal{F}^-(v; \cdot)}{d\mathcal{L}^N}(x_0) \leq f_{\text{hom}}(v(x_0)) \quad \text{for } \mathcal{L}^N \text{ a.e. } x_0 \in \Omega.$$

By Remark 1.8(i), for any fixed $\eta > 0$ we may find $k \in \mathbb{N}$, $w \in L^\infty_{k\text{-per}}(\mathbb{R}^N; \mathbb{R}^d) \cap \ker \mathcal{A}$ such that $\int_{kQ} w \, dx = 0$ and

$$\frac{1}{k^N} \int_{kQ} f(x, v(x_0) + w(x)) \, dx \leq f_{\text{hom}}(v(x_0)) + \eta. \tag{5.13}$$

For any fixed $r > 0$ and for any $n \in \mathbb{N}$, let $u_{n,r}(x) := w(xnk/r)$. Then $u_{n,r} \in L^\infty(rQ; \mathbb{R}^d) \cap \ker A$, $u_{n,r} \rightharpoonup 0$ in $L^q(rQ; \mathbb{R}^d)$ as $n \rightarrow \infty$, and by Step 1

$$\begin{aligned} \frac{d\mathcal{F}^-(v; \cdot)}{d\mathcal{L}^N}(x_0) &= \lim_{r \rightarrow 0^+} \frac{\mathcal{F}^-(v(\cdot + x_0); rQ)}{r^N} \leq \liminf_{r \rightarrow 0^+} \liminf_{n \rightarrow \infty} \frac{1}{r^N} \int_{rQ} f\left(\frac{x}{\varepsilon_{n,r}}, v(x_0 + x) + u_\varepsilon(x)\right) dx \\ &= \liminf_{r \rightarrow 0^+} \liminf_{n \rightarrow \infty} \int_Q f(nk y, v(x_0 + r y) + w(nk y)) dy, \end{aligned}$$

where $\varepsilon_{n,r} := r/nk$. Since $f(\cdot, v)$ is Q -periodic, there exists $\delta > 0$ such that if $|v - v(x_0)| < \delta$ then

$$\sup_{s \in \mathbb{R}^N} |f(s, v + w(s)) - f(s, v(x_0) + w(s))| < \eta.$$

Setting $E_{r,\delta} := \{y \in Q : |v(x_0 + r y) - v(x_0)| \geq \delta\}$, we deduce that

$$\begin{aligned} \frac{d\mathcal{F}^-(v; \cdot)}{d\mathcal{L}^N}(x_0) &\leq \eta + \limsup_{r \rightarrow 0^+} \limsup_{n \rightarrow \infty} \int_Q f(nk y, v(x_0) + w(nk y)) dy \\ &\quad + \limsup_{r \rightarrow 0^+} \limsup_{n \rightarrow \infty} \int_{E_{r,\delta}} C(1 + |v(x_0 + r y)|^q + |w(nk y)|^q) dy \\ &= \eta + \frac{1}{k^N} \int_{kQ} f(y, v(x_0) + w(y)) dy \leq f_{\text{hom}}(v(x_0)) + 2\eta, \end{aligned}$$

where we have used (5.13), the kQ -periodicity of the function $h(y) := f(y, v(x_0) + w(y))$, the equi-integrability of $\{|u_\varepsilon|^q\}$, and the fact that (5.11) entails

$$\lim_{r \rightarrow 0^+} |\{y \in Q : |v(x_0 + r y) - v(x_0)| \geq \delta\}| = 0.$$

It suffices to let $\eta \rightarrow 0^+$.

□

Proof of Theorem 1.7. We claim that for any $\varepsilon_n \rightarrow 0^+$

$$\mathcal{F}_{\text{hom}}(\cdot; D) = \Gamma - \liminf_{n \rightarrow \infty} \mathcal{F}_{\varepsilon_n}(\cdot; D).$$

By Lemmas 5.5 and 5.6 we always have

$$\mathcal{F}_{\text{hom}}(\cdot; D) \geq \Gamma - \liminf_{k \rightarrow \infty} \mathcal{F}_{\varepsilon_{n_k}} \geq \Gamma - \liminf_{n \rightarrow \infty} \mathcal{F}_{\varepsilon_n}(\cdot; D).$$

Thus assume for contradiction that that there exists $\varepsilon_n \rightarrow 0^+$ and $v \in L^q(D; \mathbb{R}^d) \cap \ker \mathcal{A}$ such that

$$\mathcal{F}_{\text{hom}}(v; D) > \Gamma - \liminf_{n \rightarrow \infty} \mathcal{F}_{\varepsilon_n}(v; D).$$

Let $\{v_n\} \subset L^q(D; \mathbb{R}^d) \cap \ker \mathcal{A}$ be such that $v_n \rightharpoonup v$ in $L^q(D; \mathbb{R}^d)$ and

$$\mathcal{F}_{\text{hom}}(v; D) > \liminf_{n \rightarrow \infty} \mathcal{F}_{\varepsilon_n}(v_n; D),$$

and choose a subsequence $\{\varepsilon_{n_k}\}$ such that

$$\mathcal{F}_{\text{hom}}(v; D) > \lim_{k \rightarrow \infty} \mathcal{F}_{\varepsilon_{n_k}}(v_{n_k}; D).$$

Then, by the previous lemmas, we can extract a further subsequence $\{\varepsilon_{n_{k_j}}\}$ such that

$$\mathcal{F}_{\text{hom}}(v; D) = \Gamma - \liminf_{j \rightarrow \infty} \mathcal{F}_{\varepsilon_{n_{k_j}}}(v; D) \leq \lim_{j \rightarrow \infty} \mathcal{F}_{\varepsilon_{n_{k_j}}}(v_{n_{k_j}}; D) < \mathcal{F}_{\text{hom}}(v; D),$$

which is a contradiction and proves the claim.

Hence it remains to show that, when (A_3) holds, for any $\varepsilon_n \rightarrow 0^+$ and $v \in L^q(D; \mathbb{R}^d) \cap \ker \mathcal{A}$

$$\Gamma - \limsup_{n \rightarrow \infty} \mathcal{F}_{\varepsilon_n}(v; D) \leq \mathcal{F}_{\text{hom}}(v; D).$$

By taking $w_n \equiv v$ and using (A_2) we get

$$\Gamma - \limsup_{n \rightarrow \infty} \mathcal{F}_{\varepsilon_n}(v; D) \leq \limsup_{n \rightarrow \infty} \mathcal{F}_{\varepsilon_n}(w_n; D) \leq C \int_D (|v(x)|^q + 1) dx.$$

Hence for *any* sequence $\{v_n\} \subset L^q(D; \mathbb{R}^d) \cap \ker \mathcal{A}$ such that $v_n \rightharpoonup v$ in $L^q(D; \mathbb{R}^d)$ and

$$\limsup_{n \rightarrow \infty} \mathcal{F}_{\varepsilon_n}(v_n; D) \leq C \int_D (|v(x)|^q + 1) dx,$$

by (A_3) we get that $\sup_n \|v_n\|_{L^q(D)} \leq L < \infty$, where the constant L depends only on the constants in $(A_2), (A_3)$, and on $\|v\|_{L^q(\Omega; \mathbb{R}^d)}$. Using the notation introduced in the proof of Lemma 5.3, we conclude that

$$\Gamma - \limsup_{n \rightarrow \infty} \mathcal{F}_{\varepsilon_n}(v; D) = \Gamma(d_L) - \limsup_{n \rightarrow \infty} \mathcal{F}_{\varepsilon_n}(v; D). \tag{5.14}$$

By Remark 5.4 and by Lemma 5.6, for any subsequence $\{\varepsilon_{n_{k_j}}\}$ of $\{\varepsilon_n\}$ there exists a subsequence $\{\varepsilon_{n_{k_j}}\}$ (depending on D) such that

$$\Gamma - \lim_{j \rightarrow \infty} \mathcal{F}_{\varepsilon_{n_{k_j}}}(v; D) = \mathcal{F}_{\text{hom}}(v; D).$$

By (5.14) this implies that

$$\Gamma(d_L) - \lim_{j \rightarrow \infty} \mathcal{F}_{\varepsilon_{n_{k_j}}}(v; D) = \mathcal{F}_{\text{hom}}(v; D).$$

We can now apply the second part of Proposition 2.4 in the metric space $(L\mathcal{B}, d_L)$ to conclude that

$$\Gamma(d_L) - \lim_{n \rightarrow \infty} \mathcal{F}_{\varepsilon_n}(v; D) = \mathcal{F}_{\text{hom}}(v; D).$$

□

Corollary 5.7. *Under hypotheses (A_1) - (A_3) the function f_{hom} is \mathcal{A} -quasiconvex and the following asymptotic formula holds*

$$f_{\text{hom}}(\xi) = \lim_{T \rightarrow +\infty} \frac{1}{T^N} \inf \left\{ \int_{TQ} f(x, \xi + v(x)) dx : v \in L^q_{T\text{-per}}(\mathbb{R}^N; \mathbb{R}^d) \cap \ker \mathcal{A}, \int_{TQ} v(x) dx = 0 \right\}. \tag{5.15}$$

Proof of Corollary 5.7. It may be shown easily, via a diagonalization procedure and in view of the coercivity condition (A_3) , that $\mathcal{F}_{\text{hom}}(\cdot; Q)$ is L^q -sequentially weakly lower semicontinuous in $\ker \mathcal{A}$. In particular, this entails \mathcal{A} -quasiconvexity for f_{hom} . Indeed, fix $v \in \mathbb{R}^d$ and $w \in L^q_{1\text{-per}}(\mathbb{R}^N; \mathbb{R}^d) \cap \ker \mathcal{A}$, with $\int_Q w(y) dy = 0$, and define $w_n(x) := w(nx)$. Then $w_n \in L^q(Q; \mathbb{R}^d) \cap \ker \mathcal{A}$, $w_n \rightharpoonup 0$ in $L^q(Q; \mathbb{R}^d)$, and so

$$f_{\text{hom}}(v) = \mathcal{F}_{\text{hom}}(v; Q) \leq \liminf_{n \rightarrow \infty} \mathcal{F}_{\text{hom}}(v + w_n; Q) = \liminf_{n \rightarrow \infty} \int_Q f_{\text{hom}}(v + w(nx)) dx = \int_Q f_{\text{hom}}(v + w(x)) dx.$$

Finally, using Theorem 1.7 (A_3), and recalling Remark 5.2, we conclude that

$$\begin{aligned} f_{\text{hom}}(v) &= \min \left\{ \mathcal{F}_{\text{hom}}(v + w; Q) : w \in L^q_{1\text{-per}}(\mathbb{R}^N; \mathbb{R}^d) \cap \ker \mathcal{A}, \int_Q w(x) dx = 0 \right\} \\ &= \lim_{\varepsilon \rightarrow 0} \inf \left\{ \mathcal{F}_\varepsilon(v + w; Q) : w \in L^q_{1\text{-per}}(\mathbb{R}^N; \mathbb{R}^d) \cap \ker \mathcal{A}, \int_Q w(x) dx = 0 \right\}, \end{aligned}$$

and (5.15) follows by setting $T = 1/\varepsilon$ and changing variables in the last expression. □

As a corollary of Theorem 1.7, we obtain the following result via the same choice of the underlying operator \mathcal{A} as in the proof of Theorem 1.5.

Theorem 5.8 (Homogenization with constraint on the divergence). *Assume that conditions $(A_1) - (A_3)$ hold, with $d = N^2$, and let F_ε be defined by*

$$F_\varepsilon(u; D) := \int_D f\left(\frac{x}{\varepsilon}, \nabla u\right) dx$$

on functions $u \in W^{1,q}(\Omega; \mathbb{R}^N)$ such that $\text{div } u = 0$. Then the Γ -limit

$$F(u; D) := \Gamma(L^q) - \lim_{\varepsilon \rightarrow 0} F_\varepsilon(u; D) = \int_D f_{\text{hom}}(\nabla u) dx$$

exists on functions $u \in W^{1,q}(\Omega; \mathbb{R}^N)$ such that $\text{div } u = 0$, where

$$f_{\text{hom}}(v) = \inf_{k \in \mathbb{N}} \frac{1}{k^N} \inf \left\{ \int_{kQ} f(x, v + \nabla w(x)) dx : w \in W^{1,q}_{k\text{-per}}(\mathbb{R}^N; \mathbb{R}^N), \text{div } w = 0 \right\}$$

for all $v \in \mathbb{R}^{N^2}$.

APPENDIX

We prove that in Remark 1.2(iii) in the introduction

$$\mathcal{F}(v; (a, b)) = \mathcal{F}((v_1, v_2); (a, b)) = \int_a^b (\psi^{**}(v_1) + v_2^2) dx, \tag{5.16}$$

where $\psi^{**}(v_1)$ is the convex envelope of

$$\psi(v_1) := \min \{ (v_1 - 1)^2, (v_1 + 1)^2 \} = \begin{cases} (v_1 + 1)^2 & \text{if } v_1 \geq 0, \\ (v_1 - 1)^2 & \text{if } v_1 < 0. \end{cases}$$

Indeed, if $v_1^n \rightharpoonup v_1$ in $L^2(a, b)$, $v_2^n \rightharpoonup v_2$ in $L^2(a, b)$ and $(v_2^n)' \rightarrow 0$ in $H^{-1}(a, b)$ then the function v_2 is constant and Jensen's inequality yields

$$\begin{aligned} \liminf_{n \rightarrow \infty} \int_a^b f(v_1^n, v_2^n) dx &\geq \liminf_{n \rightarrow \infty} \int_a^b [\psi(v_1^n) + (v_2^n)^2] dx \geq \liminf_{n \rightarrow \infty} \int_a^b \psi(v_1^n) dx + \liminf_{n \rightarrow \infty} \int_a^b (v_2^n)^2 dx \\ &\geq \int_a^b \psi^{**}(v_1) dx + (v_2)^2(b - a). \end{aligned}$$

The arbitrariness of the sequence $\{(v_1^n, v_2^n)\}$ allows us to conclude that

$$\mathcal{F}(v; (a, b)) \geq \int_a^b \psi^{**}(v_1) dx + (v_2)^2(b - a).$$

Conversely, suppose that if v_1 is smooth, $v_1 \in L^2(a, b)$, and $|\{x \in (a, b) : v_1(x) = 0\}| = 0$, $v_2 \in \mathbb{R}$, then

$$\mathcal{F}(v; (a, b)) \leq \int_a^b \psi(v_1) dx + (v_2)^2(b - a). \tag{5.17}$$

Then this inequality remains true for $v_1 \in L^2(a, b)$, $v_2 \in \mathbb{R}$ arbitrary, because we may approximate v_1 in L^2 strong by a sequence $\{v_1^n\} \subset L^2(a, b) \cap C^\infty(a, b)$, $|\{x \in (a, b) : v_1^n(x) = 0\}| = 0$, and

$$v_1 \mapsto \mathcal{F}((v_1, v_2); (a, b)), \quad v_1 \mapsto \int_a^b \psi(v_1) dx$$

are, respectively, L^2 -weak lower semicontinuous and L^2 -strong continuous. Once we establish (5.17) for $(v_1, v_2) \in L^2(a, b) \times \mathbb{R}$ then (5.16) follows because, once again, $\mathcal{F}((\cdot, v_2); (a, b))$ is L^2 -weak lower semicontinuous.

Fix now $v_1 \in L^2(a, b) \cap C^\infty(a, b)$, with $|\{x \in (a, b) : v_1(x) = 0\}| = 0$, and let $v_2 \in \mathbb{R}$, $\delta > 0$ be fixed. Set

$$\begin{aligned} A_\delta^- &:= \{x \in (a, b) : (v_1(x) - 1)^2 < (v_1(x) + 1)^2 - \delta\}, \\ A_\delta^+ &:= \{x \in (a, b) : (v_1(x) - 1)^2 > (v_1(x) + 1)^2 + \delta\}. \end{aligned}$$

If for all $\delta > 0$ $A_\delta^- = \emptyset$ then $(v_1(x) - 1)^2 \geq (v_1(x) + 1)^2$ for every $x \in (a, b)$ and we choose $w_n \notin \mathbb{Q}$ with $w_n \rightarrow v_2$. Then

$$\begin{aligned} \mathcal{F}((v_1, v_2); (a, b)) &\leq \liminf_{n \rightarrow \infty} \int_a^b f(v, w_n) dx = \liminf_{n \rightarrow \infty} \int_a^b [(v_1(x) + 1)^2 + (w_n)^2] dx \\ &\leq \int_a^b \psi(v_1) dx + (v_2)^2(b - a). \end{aligned}$$

Similarly (5.17) holds if for all $\delta > 0$ $A_\delta^+ = \emptyset$. Thus assume that for $\delta > 0$ sufficiently small $A_\delta^- \neq \emptyset \neq A_\delta^+$. Choose a cut-off function $\varphi_\delta \in C^\infty((a, b); [0, 1])$ such that $\varphi_\delta \equiv 1$ in A_δ^- , $\varphi_\delta \equiv 0$ in A_δ^+ , and let $z_n \in \mathbb{Q}$, $w_n \notin \mathbb{Q}$, be such that with $w_n, z_n \rightarrow v_2$. Define

$$v_2^{n, \delta}(x) := \varphi_\delta(x) z_n + (1 - \varphi_\delta(x)) w_n.$$

Since $(v_2^{n, \delta})' = \varphi_\delta'(x)(z_n - w_n)$, it is clear that

$$\lim_{\delta \rightarrow 0^+} \lim_{n \rightarrow \infty} \|v_2^{n, \delta} - v_2\|_{L^2} = \lim_{\delta \rightarrow 0^+} \lim_{n \rightarrow \infty} \|(v_2^{n, \delta})'\|_{H^{-1}} = 0.$$

We have

$$\begin{aligned}
 \mathcal{F}((v_1, v_2); (a, b)) &\leq \liminf_{\delta \rightarrow 0^+} \liminf_{n \rightarrow \infty} \int_a^b f(v_1, v_2^{n, \delta}) dx \\
 &= \liminf_{\delta \rightarrow 0^+} \liminf_{n \rightarrow \infty} \left\{ \int_{A_\delta^-} f(v_1, z_n) dx + \int_{A_\delta^+} f(v_1, w_n) dx + \int_{(a, b) \setminus (A_\delta^- \cup A_\delta^+)} f(v_1^n, v_2^{n, \delta}) dx \right\} \\
 &\leq \liminf_{\delta \rightarrow 0^+} \liminf_{n \rightarrow \infty} \left\{ \int_{A_\delta^-} [\psi(v_1) + (z_n)^2] dx + \int_{A_\delta^+} [\psi(v_1) + (w_n)^2] dx + C |(a, b) \setminus (A_\delta^- \cup A_\delta^+)| \right\} \\
 &= \liminf_{\delta \rightarrow 0^+} \left\{ \int_{A_\delta^- \cup A_\delta^+} [\psi(v_1) + (v_2)^2] dx + C |(a, b) \setminus (A_\delta^- \cup A_\delta^+)| \right\} \\
 &= \int_a^b \psi(v_1) dx + (v_2)^2(b - a),
 \end{aligned}$$

where we have used the fact that

$$\begin{aligned}
 |(a, b) \setminus (A_\delta^- \cup A_\delta^+)| &= |\{x \in (a, b) : |(v_1 - 1)^2 - (v_1 + 1)^2| < \delta\}| = |\{x \in (a, b) : |v_1(x)| < \delta/2\}| \\
 &\rightarrow |\{x \in (a, b) : v_1(x) = 0\}| = 0
 \end{aligned}$$

as $\delta \rightarrow 0^+$.

The research of A. Braides was partially supported by the CNR research project “Equazioni Differenziali e Calcolo delle Variazioni”, that of I. Fonseca by the National Science Foundation under Grant No. DMS-9731957 and through the Center for Nonlinear Analysis under Grant No. DMS-9803791, and that of G. Leoni by GNAFA, by MURST, Project “Metodi Variazionali ed Equazioni Differenziali Non Lineari”, by the Italian CNR, through the strategic project “Metodi e modelli per la Matematica e l’Ingegneria”. G. Leoni would like to thank the hospitality and support of the Center for Nonlinear Analysis and of S.I.S.S.A, where part of this work was carried out.

REFERENCES

- [1] E. Acerbi and N. Fusco, Semicontinuity problems in the calculus of variations. *Arch. Rational Mech. Anal.* **86** (1984) 125–145.
- [2] M. Amar and V. De Cicco, Relaxation of quasi-convex integrals of arbitrary order. *Proc. Roy. Soc. Edinburgh Sect. A* **124** (1994) 927–946.
- [3] L. Ambrosio, S. Mortola and V.M. Tortorelli, Functionals with linear growth defined on vector valued *BV* functions. *J. Math. Pures Appl.* **70** (1991) 269–323.
- [4] E.J. Balder, A general approach to lower semicontinuity and lower closure in optimal control theory. *SIAM J. Control Optim.* **22** (1984) 570–598.
- [5] J.M. Ball, A version of the fundamental theorem for Young measures, in *PDE’s and Continuum Models of Phase Transitions*, edited by M. Rascle, D. Serre and M. Slemrod. Springer-Verlag, Berlin, *Lecture Notes in Phys.* **344** (1989) 207–215.
- [6] J.M. Ball and F. Murat, Remarks on Chacon’s biting lemma. *Proc. Amer. Math. Soc.* **107** (1989) 655–663.
- [7] H. Berliocchi and J.M. Lasry, Intégrands normales et mesures paramétrées en calcul des variations. *Bull. Soc. Math. France* **101** (1973) 129–184.
- [8] A. Braides, A homogenization theorem for weakly almost periodic functionals, *Rend. Accad. Naz. Sci. XL Mem. Sci. Fis. Natur. (5)* **104** (1986) 261–281.
- [9] A. Braides, Relaxation of functionals with constraints on the divergence. *Ann. Univ. Ferrara Ser. VII (N.S.)* **33** (1987) 157–177.
- [10] A. Braides and A. Defranceschi, *Homogenization of Multiple Integrals*. Clarendon Press, Oxford (1998).
- [11] A. Braides, A. Defranceschi and E. Vitali, Homogenization of free-discontinuity problems. *Arch. Rational Mech. Anal.* **135** (1996) 297–356.
- [12] G. Buttazzo, *Semicontinuity, relaxation and integral representation problems in the Calculus of Variations*. Longman, Harlow, *Pitman Res. Notes Math. Ser.* **207** (1989).
- [13] B. Dacorogna, *Direct Methods in the Calculus of Variations*. Springer-Verlag, Berlin (1989).
- [14] B. Dacorogna, *Weak Continuity and Weak Lower Semicontinuity for Nonlinear Functionals*. Springer-Verlag, Berlin, *Lecture Notes in Math.* **922** (1982).

- [15] G. Dal Maso, *An Introduction to Γ -Convergence*. Birkhäuser, Boston (1993).
- [16] G. Dal Maso, A. Defranceschi and E. Vitali (private communication).
- [17] A. De Simone, Energy minimizers for large ferromagnetic bodies. *Arch. Rational Mech. Anal.* **125** (1993) 99–143.
- [18] I. Fonseca, The lower quasiconvex envelope of the stored energy function for an elastic crystal. *J. Math. Pures Appl.* **67** (1988) 175–195.
- [19] I. Fonseca, G. Leoni, J. Malý and R. Paroni (in preparation).
- [20] I. Fonseca and S. Müller, Quasiconvex integrands and lower semicontinuity in L^1 . *SIAM J. Math. Anal.* **23** (1992) 1081–1098.
- [21] I. Fonseca and S. Müller, Relaxation of quasiconvex functionals in $BV(\Omega, \mathbb{R}^p)$ for integrands $f(x, u, \nabla u)$. *Arch. Rational Mech. Anal.* **123** (1993) 1–49.
- [22] I. Fonseca and S. Müller, A-quasiconvexity, lower semicontinuity and Young measures. *SIAM J. Math. Anal.* **30** (1999) 1355–1390.
- [23] N. Fusco, Quasi-convessità e semicontinuità per integrali di ordine superiore. *Ricerche Mat.* **29** (1980) 307–323.
- [24] M. Giaquinta and G. Modica, Regularity results for some classes of higher order non linear elliptic systems. *J. reine angew. Math.* **311/312** (1979) 145–169.
- [25] M. Guidorzi and L. Poggiolini, Lower semicontinuity for quasiconvex integrals of higher order. *NoDEA Nonlinear Differential Equations Appl.* **6** (1999) 227–246.
- [26] J. Kristensen, *Finite functionals and Young measures generated by gradients of Sobolev functions*. Mathematical Institute, Technical University of Denmark, Mat-Report No. 1994-34 (1994).
- [27] P. Marcellini, Approximation of quasiconvex functions and semicontinuity of multiple integrals. *Manuscripta Math.* **51** (1985) 1–28.
- [28] P. Marcellini and C. Sbordone, Semicontinuity problems in the Calculus of Variations. *Nonlinear Anal.* **4** (1980) 241–257.
- [29] N.G. Meyers, Quasi-convexity and lower semi-continuity of multiple variational integrals of any order. *Trans. Amer. Math. Soc.* **119** (1965) 125–149.
- [30] C.B. Morrey, *Multiple Integrals in the calculus of Variations*. Springer-Verlag, Berlin (1966).
- [31] S. Müller, Variational models for microstructures and phase transitions, in *Calculus of Variations and Geometric Evolution Problems*, edited by S. Hildebrandt et al. Springer-Verlag, Berlin, *Lecture Notes in Math.* **1713** (1999) 85–210.
- [32] F. Murat, Compacité par compensation : condition nécessaire et suffisante de continuité faible sous une hypothèse de rang constant. *Ann. Scuola Norm. Sup. Pisa Cl. Sc. (4)* **8** (1981) 68–102.
- [33] P. Pedregal, *Parametrized Measures and Variational Principles*. Birkhäuser, Boston (1997).
- [34] L. Tartar, Compensated compactness and applications to partial differential equations, in *Nonlinear Analysis and Mechanics: Heriot-Watt Symposium*, edited by R. Knops. Longman, Harlow, *Pitman Res. Notes Math. Ser.* **39** (1979) 136–212.
- [35] L. Tartar, The compensated compactness method applied to systems of conservation laws, in *Systems of Nonlinear Partial Differential Eq.*, edited by J.M. Ball. Riedel (1983).
- [36] L. Tartar, Étude des oscillations dans les équations aux dérivées partielles nonlinéaires. Springer-Verlag, Berlin, *Lectures Notes in Phys.* **195** (1984) 384–412.
- [37] L. Tartar, H -measures, a new approach for studying homogenisation, oscillations and concentration effects in partial differential equations. *Proc. Roy. Soc. Edinburgh Sect. A* **115** (1990) 193–230.
- [38] L. Tartar, On mathematical tools for studying partial differential equations of continuum physics: H -measures and Young measures, in *Developments in Partial Differential Equations and Applications to Mathematical Physics*, edited by Buttazzo, Galdi and Zanghirati. Plenum, New York (1991).
- [39] L. Tartar, Some remarks on separately convex functions, in *Microstructure and Phase Transitions*, edited by D. Kinderlehrer, R.D. James, M. Luskin and J.L. Ericksen. Springer-Verlag, *IMA J. Math. Appl.* **54** (1993) 191–204.
- [40] L.C. Young, *Lectures on Calculus of Variations and Optimal Control Theory*. W.B. Saunders (1969).