PUBLICATIONS MATHÉMATIQUES ET INFORMATIQUES DE RENNES

NOBUO AOKI

Ergodic Automorphisms of Compact Metric Groups are Isomorphic to Bernoulli Shifts

Publications des séminaires de mathématiques et informatique de Rennes, 1975, fascicule S4
« International Conference on Dynamical Systems in Mathematical Physics », , p. 1-10

<http://www.numdam.org/item?id=PSMIR_1975___S4_A2_0>

© Département de mathématiques et informatique, université de Rennes, 1975, tous droits réservés.

L'accès aux archives de la série « Publications mathématiques et informatiques de Rennes » implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/conditions). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
http://www.numdam.org/

ERGODIC AUTOMORPHISMS OF COMPACT METRIC GROUPS ARE ISOMORPHIC TO BERNOULLI SHIFTS

NOBUO AOKI

I wish to discuss its title. Let X be a compact metric group and μ be its normalized Haar measure. Then (X, μ) is a Lebesgue space. Let σ be an automorphism of X, then σ is an invertible measure preserving transformation of X onto itself. Our problem is concerned with measure theoretic properties of σ.

Throughout this given a transformation of any group, a restriction on a subgroup and an induced transformation on a factor space will be denoted by the same symbol as that of the original transformation, if there is no danger of confusion.

Today we will outline a proof of the following

Theorem 1. An ergodic automorphism of a compact metric abelian group is a Bernoulli shift.

The result has recieved the most attension in the literature. In a two-dimensional torus, Adler and Weiss [1] proved the result using Ornstein's Theorems. In recently, Katznelson [4] showed the result in an n-dimensional torus. Lind [5] gave a proof for the case of an infinite-dimensional torus. The proof which Totoki and the author [2] proved was done independently of Lind's work. The techniques I use are due to Katznelson [4] and Totoki and the author [2].

In order to outline Theorem 1, we prepare the following

Proposition 1. Let X be a compact metric abelian group and σ be an ergodic automorphism of X. Then there exist subgroups X_D, X_A and X_B such that X_D is a σ-invariant totally disconnected subgroup, X_A and X_B are σ-invariant connected subgroups of X and dynamical systems (X_D, σ), (X_A, σ), (X_B, σ) are ergodic, and further (X, σ) is an algebraic factor of $(X_D \otimes X_A \otimes X_B, \sigma \otimes \sigma \otimes \sigma)$.

The proof uses the results of Entropy Theory together with the results of Group Theory.

Proposition 2. The dynamical systems (X_B, σ) and (X_D, σ) have the Bernoulli properties.

We can prove that X_B is locally connected and so by [2] we have (X_B, σ) has the Bernoulli properties. The Bernoulli properties of (X_D, σ) is an indirectapplication of the results of Yuzvinskii [12].

To show the dynamical system (X_A, σ) has the Bernoulli properties, let G_A be the character group of X_A and U be the dual automorphism of G_A induced by $(Ug)(x) = g(\sigma^{-1}x)$ for $g \in G_A$. Each $g \in G_A$ satisfies the following condition

(A) There exist integers $k > 0$, $n_0, n_1, \ldots, n_k$ such that $(n_0, n_1, \ldots, n_k) \neq (0, 0, \ldots, 0)$ and

$$g^{n_0} U g^{n_1} \ldots U^k g^{n_k} = 1 .$$

We denote by $\bar{G}_A$ the minimal divisible extension of G_A and by $\bar{U}$ the automorphism of $\bar{G}_A$ extended by U. If $\bar{X}_A$ is the dual group of $\bar{G}_A$, then $\bar{X}_A$ is a compact connected metric abelian group. If $\bar{\sigma}$ is the dual automorphism of $\bar{X}_A$ induced by $\bar{U}^{-1}$, then as (X_A, σ) has the ergodic properties, it is not hard to see that $(\bar{X}_A, \bar{\sigma})$ has the ergodic properties. Since $G_A \subset \bar{G}_A$, let us define $X_1' = \mathrm{ann}(G_A, \bar{X}_A)$, then the dynamical system $(\bar{X}_A/X_1', \bar{\sigma})$ and (X_A, σ) are isomorphic. And so if $(\bar{X}_A, \bar{\sigma})$ has the Bernoulli properties, then Ornstein's theorems imply that (X_A, σ) has the Bernoulli properties. Therefore, using Propositions 1 and 2 it follows that (X, σ) has the Bernoulli properties.

We resolve this difficult with some lemmas.

Let $\bar{G}_A = \{f_1, f_2, \ldots\}$. We denote by G_n the subgroup of $\bar{G}_A$ generated by $\{\bar{U}^j f_k : -\infty < j < \infty, \ k = 1, 2, \ldots, n\}$ for $n \geq 1$. Then we have $\mathrm{rank}(G_n) < \infty$ for $n \geq 1$. Let $X_n = \mathrm{ann}(G_n, \bar{X}_A)$ for $n \geq 1$, then we have $\bar{\sigma} X_n = X_n$, $n \geq 1$ and $X_1 \supset X_2 \supset \ldots \supset \bigcap_{n=1}^{\infty} X_n = \{e\}$. Since each f_k satisfies the condition (A), we have for each $k \geq 1$

$$f_k^{\ell_k} = \bar{U}^{m_1(k)} f_k \ldots \bar{U}^{m_{p_k}(k)} f_k$$

for some $p_k > 0$, $\ell_k > 0$ and some $(m_1(k), \ldots, m_{p_k}(k)) \neq (0, \ldots, 0)$.

From now on, we fix $\ell_1, \ldots, \ell_n$ and put

(1) $n_o = \ell_1 \dots \ell_n$, $k_o = \max_{1 \leq k \leq n} p_k$.

H denotes the subgroup of G_n generated by $\{f_k, \dots, \bar{U}^{k_o} f_k : 1 \leq k \leq n\}$. Then H is finitely generated, torsionfree, $\bar{H} = \bar{G}_n$ and $\prod_{j=-\infty}^{\infty} \bar{U}^j H = G_n$. Let $X(H) = \text{ann}(H, \bar{X}_A)$, then the character group of $\bar{X}_A/X(H)$ is H and so $\bar{X}_A/X(H)$ is a finite-dimensional torus.

Lemma 1. If $n_o = 1$, then $(\bar{X}_A/X_n, \bar{\sigma})$ has the Bernoulli properties.

The proof is a direct application of the result of Katznelson [4].

Lemma 2. If $n_o > 1$, then $\eta(x) = x^{n_o}$, $x \in \bar{X}_A$, is an automorphism of $\bar{X}_A$ such that $\eta\bar{\sigma} X(H) \subset X(H)$, and the induced factor $\eta\bar{\sigma}$ on $\bar{X}_A/X_n$ is ergodic.

The details of the proof are found in Chapter 1 of [13].

This lemma is essentially utilized in the proof of Lemma 13.

To obtain $(\bar{X}_A/X_n, \bar{\sigma})$ has the Bernoulli properties whenever $n_o > 1$, we construct a sequence $\{\mathcal{P}_n\}$ of weak Bernoulli partitions for the dynamical system $(\bar{X}_A/X_n, \bar{\sigma})$ such that $\mathcal{P}_n < \mathcal{P}_{n+1}$ for $n \geq 1$ and $\bigvee_n \mathcal{P}_n$ is the partition of $\bar{X}_A/X_n$ into single points.

Let M be a positive integer and let $\mathcal{P}$ be a partition of the interval $[0, 2\pi)$ into subintervals of the same lengths $2\pi/M$. The elements of $\mathcal{P}$ will denote successively from the left by $p_j = [a_j, a_{j+1})$, $j = 1, \dots, M$. Let K be an arbitrarily fixed positive integer and set $N_K = n_o^{K^2+K}$ (n_o is the integer satisfying the (1)) For $k > 0$ $K_k(t)$ will denote the Fejér kernel defined on $[0, 2\pi)$.

Lemma 3. Let M be a fixed positive integer. Then there exists a positive integer $\ell = \ell(M)$ such that for each $m \geq 2$ there is a positive number $\delta_m = \delta(m, M)$ satisfying the following :

$$(1 + m^{-2})\left(1 - \frac{8}{(m^\ell - 1)\,\delta_m^2}\right) \geq 1,$$

$$3^{M+1} M^2 \delta_m < 1/m^2 .$$

The proof is elementary.

Lemma 4. Let M and δ_m be as in Lemma 3. Then

$$2\pi + 2M(N_k - 1)\delta_m < 2\pi N_K \quad (2 \leqslant m \leqslant K).$$

The proof is clear from Lemma 3.

From Lemma 4 we can manipulate as follows. For an arbitrary m such that $2 \leqslant m \leqslant K$, set $p_j(m) = [b_j, c_j)$, mod $2\pi N_K$, where $b_j = a_j - (N_K - 1)\delta_m$, $c_j = a_{j+1} + (N_K - 1)\delta_m$ $(1 \leqslant j \leqslant M)$. We translate each $p_j(m)$ by $r'_j(m)$ $(r'_1(m) = 0)$ to the right so that $c_j + r'_j(m) = b_{j+1} + r'_{j+1}(m)$ and denote the translated $p_j(m)$ by $\tilde{p}_j(m)$. Then $\tilde{p}_1(m), \ldots, \tilde{p}_M(m)$ are disjoint and each $\tilde{p}_{j+1}(m)$ borders on $\tilde{p}_j(m)$ from the right and hence we may set $\tilde{p}_j(m) = [a'_j(m), a'_{j+1}(m))$ for $j = 1,2, \ldots, M$.

In the case $M = 3$ we have for instance following figure.



Noting $N_K > 2$, we have clearly

$$(2) \quad 2\delta_m < r'_j(m) < 2(j-1)(N_K - 1)\delta_m + 2\pi j/M$$

$$(2 \leqslant m \leqslant K, \quad 2 \leqslant j \leqslant M).$$

Lemma 5. Let M and δ_m be as in Lemma 3. Then there exists a positive integer $K_o = K_o(M)$ such that for $K > K_o$

$$2\pi + 2M(N_K - 1)\delta_m + 3^{M+1}Mr'_M(m) < 2\pi N_K \quad (2 \leqslant m \leqslant K).$$

The proof is elementary.

We set for m such that $2 \leqslant m \leqslant K$ for $K > K_o$

$$(3) \quad \begin{cases} r_1(m) = r'_1(m) = 0, \\ r_j(m) = r'_j(m) + 3r_{j-1}(m) \quad (2 \leqslant j \leqslant M) \\ p'_1(m) = \tilde{p}_1(m), \\ p'_j(m) = [a'_j(m) + 3r_{j-1}(m),\ a'_{j+1}(m) + 3r_j(m)) \quad (2 \leqslant j \leqslant M) \end{cases}$$

Then $p'_1(m), \ldots, p'_M(m)$ are disjoint and each $p'_{j+1}(m)$ borders on $p'_j(m)$ from the right.

In the case $M = 3$ we have for instance the following figure.



We have for m with $2 \leqslant m \leqslant K$ for $K > K_0$

$$3(r_2(m) + \ldots + r_M(m)) < 3^{M+1}Mr_M'(m).$$

Hence by Lemma 5 we have for $K > K_0$

$$\bigcup_{j=1}^{M} p_j'(m) \subset [0, 2\pi N_K) \qquad (2 \leqslant m \leqslant K).$$

Lemma 6. Let M and δ_m be as in Lemma 3. Then there exists a positive integer $K_1 = K_1(M) > K_0$ such that for $K > K_1$

$$\delta_m + r_j(m)/N_K < 2\pi \qquad (2 \leqslant m \leqslant K, \quad 1 \leqslant j \leqslant M).$$

The proof uses (2), (3) and Lemma 3.

Lemma 7. Let δ_m be as in Lemma 3. Then for m satisfying $m > \max(2, \sqrt{M/\pi})$ and $p_j = [a_j, a_{j+1}) \in \mathfrak{p}$, $j = 1, 2, \ldots, M$,

$$[a_j + \delta_m, a_{j+1} - \delta_m) \neq \phi.$$

The proof is elementary.

Lemma 8. We consider the characteristic function $\chi_{p_j'(m)}$ of $p_j'(m)$ as a $2\pi N_K$-cyclic function on R^1. Then for $K > \max(2, K_1, \sqrt{M/\pi})$ and $N_K t \in [a_j + \delta_m, a_{j+1} - \delta_m)$ for $t \in [0, 2\pi)$,

$$\chi_{p_j'(m)}(N_K t + r_j(m) - s) = 1 \quad (\max(2, \sqrt{M/\pi}) < m \leqslant K, \quad 1 \leqslant j \leqslant M)$$

if $0 < s < N_K \delta_m$ or $2\pi N_K - N_K \delta_m - 2r_j(m) < s < 2\pi N_K$.

The lemma follows from (2), (3) and Lemma 7.

Lemma 9. Let ℓ and δ_m be as in Lemma 3 and let K_1 be as in Lemma 6. Then if $K > \max(2, K_1, \sqrt{M/\pi})$, for each m $(\max(2, \sqrt{M/\pi}) < m \leqslant K)$ and each $p_j \in \mathfrak{p}$ there exists a non-negative function $\tilde{f}_{mp_j}(t)$ on $[0, 2\pi)$ satisfying the following :

$$\tilde{f}_{mp_j}(t) \geqslant 1 \qquad t \in [a_j + \delta_m, a_{j+1} - \delta_m),$$

for some constants c_0, c_k, c_k' $(k = 1, 2, \ldots, m^\ell)$

$$\tilde{f}_{mp_j}(t) = c_0 + \sum_{k=1}^{m^\ell} c_k \tilde{e}^{i(k/N_K, t)} + \sum_{k=1}^{m^\ell} c_k' \tilde{e}^{-i(k/N_K, t)},$$

where $\tilde{e}^{i(k/N_K, t)} = e^{ik(t/N_K)}$ and $\tilde{e}^{-i(k/N_K, t)} = e^{-ik(t/N_K)}$ for k,

$$\sum_{j=1}^{M} \tilde{f}_{mp_j}(t) \leqslant 1 + m^{-2} \qquad t \in [0, 2\pi).$$

The proof uses the results of Lemmas 3 ~ 8 .

We denote by $e^{i(m,t)}$ an exponential function e^{mti} for m .

Lemma 10. Let ℓ and δ_m be as in Lemma 3. For each $p_j \in \mathfrak{P}$ and $K > \max(2,\sqrt{M/\pi})$ we define for $t \in [0, 2\pi)$

$$f_{mp_j}(t) = (1 + m^{-2})\hat{f}_{mp_j}(t) \quad (\max(2,\sqrt{M/\pi}) < m \leqslant K)$$

where

$$\hat{f}_{mp_j}(t) = 1/\pi \int_0^{2\pi} \chi_{p_j}(t - s)K_{m^\ell}(s)\, ds .$$

Then $f_{mp_j}(t)$ is a non-negative function on $[0, 2\pi)$ and we have the following :

$$f_{mp_j}(t) \geqslant 1 \quad t \in [a_j + \delta_m, a_{j+1} - \delta_m) ,$$

for some constants d_o, d_k, d_k' $(k = 1,2, \ldots , m)$

$$f_{mp_j}(t) = d_o + \sum_{k=1}^{m^\ell} d_k e^{i(k, t)} + \sum_{k=1}^{m^\ell} d_k' e^{-i(k, t)} ,$$

$$\sum_{j=1}^{M} f_{mp_j}(t) \leqslant 1 + m^{-2} \qquad t \in [0, 2\pi) .$$

The proof is direct from Katznelson [4].

We can generalize easily Lemmas 9 and 10 on a finite-dimensional torus

We assume that $\bar{X}_A/X(H)$ is r-dimensional. $\bar{X}_A/X(H)$ is algebraically isomorphic to $T^r = [0, 2\pi)^r$ whose character group is the discrete group

$$H_r = \{ e^{i(m, \cdot)} : m \in Z^r \}^{1)}$$

which is algebraically isomorphic to H. $\bar{H}_r$ denotes a multiplicative group $\{ e^{i(q, \cdot)} : q \in Q^r \}^{2)}$ which is a minimal divisible extension of H_r . Now let Y^r denote the dual group of $\bar{H}_r$, and we denote by $(e^{i(q, \cdot)})(y)$, $y \in Y^r$ each character $e^{i(q, \cdot)}$ of Y^r. We note that for $m \in Z^r$

$$(e^{i(m, \cdot)})(Py) = e^{i(m, Py)}$$

where P is the projection from Y^r onto T^r.

From the definitions of G_n and H, we have $\bar{G}_n = \bar{H}$. Thus as H_r and H are algebraically isomorphic, we have the diagram

1) Z^r is the set of all r-dimensional integer vectors.

2) Q^r is the set of all r-dimensional rational vectors.

$$\bar{X}_A/\bar{X}_n \overset{3)}{\cong} Y^r$$
$$\searrow \quad \swarrow P$$
$$T^r$$

Let τ and ξ be automorphisms of Y^r isomorphic to $\bar{\sigma}$ and η of $\bar{X}_A/\bar{X}_n$ respectively. Then $\tau\xi$ induces the endomorphism of T^r (wrtten by the same symbol $\tau\xi$) because $\eta\bar{\sigma}$ induces the endomorphism of $\bar{X}_A/X(H)$ and $\tau\xi$ is given by an $r \times r$ matrix with intger entries.

U_P denotes the linear operator from $\mathcal{C}(T^r)$ into $\mathcal{C}(Y^r)$ defined by $(U_P g)(y) = g(Py)$ for $g \in \mathcal{C}(T^r)$. The adjoint operator $\tau\xi$ on Z^r of the endomorphism $\tau\xi$ on T^r is defined by

$$e^{i(m,\ \tau\xi Py)} = e^{i(\tau\xi m,\ Py)}, \qquad m \in Z^r .$$

We have for $\lambda \in Z^r$

$$(4) \qquad U_\tau^{-1}(U_P\, \tilde{e}^{i(\lambda/N_K, \cdot)})(y) = (U_P\, \tilde{e}^{i(\tau\lambda/N_K, \cdot)})(y), \quad y \in Y^r.$$

Now the eigenvalue of ξ is n_o with multiplicity r . Let the eigenvalues of τ be $\lambda_1, \ \dots \ , \lambda_k$ $(k \leqslant r)$, then the eigenvalues of $\tau\xi$ are $n_o\lambda_1, \ \dots \ , n_o\lambda_k$.

We may consider the matrix τ as operating on R^r and so we decompose

$$R^r = V_{-k} \oplus \ \dots \ \oplus V_o \oplus \ \dots \ \oplus V_q$$

such that each V_j is the τ-invariant subspace of R^r corresponding to the eigenvalues of τ of modulus ρ'_j where $\rho'_{-k} < \ \dots \ < 1 = \rho'_o < \ \dots \ < \rho'_q$. Let $\rho_j = n_o\rho'_j$ and let $V_o = V_{-k} \oplus \ \dots \ \oplus V_{-k'}$ $(k' \geqslant 0)$ be the direct sum of V_j's corresponding to ρ'_j such that $\rho_j \leqslant 1$. Then as V_j is $\tau\xi$-invariant and $\tau\xi$ is ergodic on T^r, we have

$$\tilde{V}_o \cap Z^r = \{0\}.$$

Let M be an arbitrarily fixed positive integer. Now let $\mathcal{P}$ be a partition of T^r $(= [0, 2\pi)^r)$ such that $\mathcal{P} = \otimes_{k=1}^r \mathcal{P}^{(k)}$, each $\mathcal{P}^{(k)}$ being a partition of $[0, 2\pi)$ into subintervals of the same lengths $2\pi/M$.

For arbitrarily fixed $K > 0$ and $N > 0$ we set

$$\mathcal{A}(K) = \bigvee_{m=1}^{K^2} \tau^{-m_P-1}(\mathcal{P}), \qquad \mathcal{B}(K,N) = \bigvee_{m=K^2+K}^{K^2+K+N} \tau^{-m_P-1}(\mathcal{P}).$$

Lemma 11. For a sufficiently large $l = l(\mathcal{P})$ there exists

3) $\bar{X}_n$ is the annihilator of $\bar{G}_n$ in $\bar{X}_A$.

a measurable set E_m with measure $< 1/m^2$ for each $m > \max(2, \sqrt{M/\pi})$ such that for each $p \in \mathcal{P}$ there are non-negative functions $\tilde{f}_{mp}$, f_{mp} on T^r satisfying :

(a) there is a positive integer $K_1 = K_1(\mathcal{P})$ such that if $K > K_1$ and $K \geqslant m$, then

$$\tilde{f}_{mp}(t) \geqslant 1 \quad \text{on} \quad p - E_m ,$$

$$\tilde{f}_{mp}(t) = C_o + \sum_{\substack{k : 1 \leqslant k_j \leqslant m^\ell \\ 1 \leqslant j \leqslant r}} [C_k \tilde{e}^{i(k/N_K,\, t)} + C'_k \tilde{e}^{-i(k/N_K,\, t)}], \quad t \in T^r,$$

where $k = (k_1, \ldots, k_r)$ are vectors of positive integers, C_o, C_k, C_j are some constants and $\tilde{e}^{\pm i(k/N_K,\, t)} = e^{\pm ik(t/N_K)}$,

$$\sum_{p \in \mathcal{P}} \tilde{f}_{mp}(t) \leqslant 1 + m^{-2} \qquad t \in T^r,$$

(b) $$f_{mp}(t) \geqslant 1 \quad \text{on} \quad p - E_m ,$$

$$f_{mp}(t) = D_o + \sum_{\substack{k : 1 \leqslant k_j \leqslant m^\ell \\ 1 \leqslant j \leqslant r}} [D_k e^{i(k,\, t)} + D'_k e^{-i(k,\, t)}] , \quad t \in T^r,$$

where k are as in (a) and D_o, D_k, D'_k are constants,

$$\sum_{p \in \mathcal{P}} f_{mp}(t) \leqslant 1 + m^{-2} \qquad t \in T^r .$$

This lemma is a generalization of Lemmas 9 and 10.

Define the following functions Φ_A and Φ_B on Y^r for $A \in \mathcal{A}(K)$ and $B \in \mathcal{B}(K, N)$ where $K > K_1$,

$$\Phi_A = \prod_{m=1}^{K^2} U_\tau^{-m} U_P \tilde{f}_{Kp_m}(A) , \quad \Phi_B = \prod_{m=K^2+K}^{K^2+K+N} U_\tau^{-m} U_P f_{mp_m}(B) .$$

We denote by μ the normalized Haar measure on Y^r. Then we have the following

Lemma 12. Let $\varepsilon > 0$. Then there exists a positive integer $K_2 = K_2(\mathcal{P}, \varepsilon) > \max(2, K_1, \sqrt{M/\pi})$ and a measurable set E in Y^r such that $\mu(E) < \varepsilon^2$ and for every $K > K_2$ and arbitrary $N > 0$

$$\Phi_A \geqslant 1 \quad \text{on} \quad A - E , \quad \sum_{A \in \mathcal{A}(K)} \int \Phi_A \, d\mu \leqslant 1 + \varepsilon^2$$

and

$$\Phi_B \geqslant 1 \quad \text{on} \quad B - E, \quad \sum_{B \in \mathcal{B}(K,N)} \int \Phi_B \, d\mu \leqslant 1 + \varepsilon^2 .$$

The lemma follows from Lemma 11.

Let ℓ be as in Lemma 11. For arbitrary fixed $K > K_3$ and $N > 0$ we denote by $\tilde{Z}^r$ and $\tilde{Q}^r$ sets of all r-dimensional vectors consisting of $\{1,2, \ldots, (K^2+K+N)^\ell\}$ and $\{1/N_K, 2/N_K, \ldots, K^\ell/N_K\}$ respectively.

Now we define an automorphism κ of Y^r by

$$\kappa y = n_o^{K^2+K+N} N_K y , \quad y \in Y^r .$$

Since for $\lambda \in Z^r$ (4) holds, we have that

$$\{(U_p \tilde{e}^{i(\sum_{m=1}^{K^2} \tau^m \lambda_m, \cdot)})(\kappa y) : \lambda_m \in \tilde{Q}^r \cup [-\tilde{Q}^r]\}$$

$$\cup \{(U_p e^{i(\sum_{m=K^2+K}^{K^2+K+N} \tau^m \lambda_m, \cdot)})(\kappa y) : \lambda_m \in \tilde{Z}^r \cup [-\tilde{Z}^r]\}$$

is a set of characters of Y^r. Further we can prove that the frequency which is common to Φ_A and Φ_B is zero for sufficiently large K. From those facts we have

Lemma 13. There exists a positive integer $K_3 > K_1$ such that for $K > K_3$ and $N > 0$

$$\int \Phi_A \Phi_B \, d\mu = \int \Phi_A \, d\mu \int \Phi_B \, d\mu .$$

Using results of Katznelson [4] and Lemmas 12 and 13, we have

Lemma 14. For $\varepsilon > 0$ there exists $\tilde{K} > \max(K_3, K_2)$ such that $\mathcal{A}(K)$ and $\mathcal{B}(K,N)$ are 11ε-independent for $K > \tilde{K}$ and $N > 0$.

Consequently $P^{-1}(\mathcal{P})$ is an weak Bernoulli partition on Y^r for τ. Let $\mathcal{P}'$ be the partition of $\bar{X}_A/X(H)$ corresponding to the partition $\mathcal{P}$ of T^r and P' be the projection of $\bar{X}_A$ onto $\bar{X}_A/X(H)$, then $P'^{-1}(\mathcal{P}')$ is an weak Bernoulli partition on $\bar{X}_A$ for $\bar{\sigma}$. Because we have $\bigcap_{j \in Z} \bar{\sigma}^j X(H) = X_n$, $\bigvee_{\mathcal{P}} \bigvee_{j \in Z} \bar{\sigma}^j P'^{-1}(\mathcal{P}')$ is the partition of $\bar{X}_A$ into cosets of X_n. By Ornstein 's theorem $(\bar{X}_A/X_n, \bar{\sigma})$ has the Bernoulli partitions.

Therefore, for $n \geqslant 1$ we have showed that $(\bar{X}_A/X_n, \bar{\sigma})$ has the Bernoulli properties. Since we have

$$X_1 \supset X_2 \supset \cdots \supset \bigcap_{n=1}^{\infty} X_n = \{e\},$$

Ornstein 's theorem implies that σ on $\bar{X}_A$ is Bernoullian. (X_A, σ) has the Bernoulli properties.

We can conclude that (X, σ) has the Bernoulli properties.

Using Theorem 1, I can prove the result for the case of non-abelian. Today I do not discusse it here, but the proof of it is found in [13].

References

1. R.L. Adler and B. Weiss, Similarity of automorphisms of the torus, Mem. Amer. Math. Soc. 98 (1970).
2. N. Aoki and H. Totoki, Ergodic automorphisms of T^{∞} are Bernoulli transformations, Pub. RIMS, Kyoto Univ. 10 (1975), 535 - 544.
3. N.A. Friedman and D.S. Ornstein, On isomorphism of weak Bernoulli transformations, Advances in Math. 5 (1971), 365 - 394.
4. Y. Katznelson, Ergodic automorphisms of T^n are Bernoulli shifts, Israel J. Math. 10 (1971), 186 - 195.
5. D.A. Lind, Ergodic automorphisms of the infinite torus are Bernoulli, Israel J. Math. 2 (1974), 162 - 168.
6. D.S. Ornstein, Bernoulli shifts with the same entropy are isomorphic, Advances in Math. 4 (1970), 337 - 352.
7. ———— , Two Bernoulli shifts with infinite entropy are isomorphic, ibid. 5 (1971), 339 - 348.
8. ———— , Factors of Bernoulli shifts are Bernoulli shifts, ibid. 5 (1971), 349 - 364.
9. L. Pontrjagin, Topological Groups, Princeton Univ. Press, Princeton, 1946.
10. V.A. Rohlin, Metric Properties of endomorphisms of compact commutative groups, Amer. Math. Soc. Transl. 64 (1967), 244 - 272.
11. Ya. G. Sinai, Markov partitions and C-diffeomorphisms, Func. Anal. and its Appl. 2 (1968), 61 - 82.
12. S.A. Yuzvinskii, Metric properties of endomorphisms of compact groups, Amer. Math. Transl. 66 (1968), 63 - 98.
13. N. Aoki, Ergodic automorphisms of compact metric groups are Bernoulli shifts, to appears.