RENDICONTI
del
SEMINARIO MATEMATICO
della
UNIVERSITÀ DI PADOVA

J. P. SINGHAL

Operational formulae for certain classical polynomials

Rendiconti del Seminario Matematico della Università di Padova, tome 38 (1967), p. 33-40

<http://www.numdam.org/item?id=RSMUP_1967__38__33_0>

© Rendiconti del Seminario Matematico della Università di Padova, 1967, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (http://rendiconti.math.unipd.it/) implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/conditions). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM
Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
http://www.numdam.org/

OPERATIONAL FORMULAE FOR CERTAIN CLASSICAL POLYNOMIALS

J. P. SINGHAL *)

1. Recently Srivastava [7, p. 43] has defined a set of polynomials $A_n^{(\alpha)}(x)$ related to the Laguerre polynomials by means of the relations

$$\sum_{r=0}^{n} A_r^{(\alpha)}(x)\, L_{n-r}^{(\alpha+r)}(x) = 0, \quad n \geq 1, \tag{1.1}$$

$$A_0^{(\alpha)}(x) = 1.$$

He also gave the generating function, hypergeometric representation and the Rodrigues' formula for these polynomials [7, pp. 44-45] in the forms :

$$(1+t)^{-1-\alpha} e^{xt} = \sum_{r=0}^{\infty} t^r A_r^{(\alpha)}(x), \tag{1.2}$$

$$A_n^{(\alpha)}(x) = \frac{1}{(n)!} \sum_{r=0}^{\infty} \frac{(-n)_r (1+\alpha)_r}{(r)!} x^{n-r}, \tag{1.3}$$

and

$$A_n^{(\alpha)}(x) = \frac{x^{n+\alpha+1}}{(n)!} D^n \{x^{-\alpha-1} e^x\}, \quad \left(D = \frac{d}{dx}\right), \tag{1.4}$$

respectively.

---

*) Indirizzo dell'A. : Deptt. of Maths. — University of Jodhpur — Jodhpur — India.

In this paper we give some operational formulae for these as well as Laguerre polynomials and employ them to derive many interesting results.

The first operational formula to be proved is

$$\prod_{j=1}^{n} (xD + x - \alpha - j) = (n)! \sum_{r=0}^{n} \frac{x^r}{(r)!} A_{n-r}^{(\alpha)}(x) D^r . \tag{1.5}$$

Note that the formula (1.5) corresponds to the one given by Carlitz [3, p. 219] in the case of Laguerre polynomials [8, p. 428].

To prove (1.5) we observe that if

$$\Omega_n = \prod_{j=1}^{n} (xD + x - \alpha - j) , \qquad \Omega_0 = 1,$$

it can be proved very easily by the method of induction that

$$\Omega_n(y) = x^{n+\alpha+1} e^{-x} D^n \{e^x x^{-\alpha-1} y\},$$

where y is some differentiable function of x.

Next since

$$D^n \{e^x x^{-\alpha-1} \cdot y\} = \sum_{r=0}^{n} \binom{n}{r} D^{n-r} \{e^x x^{-\alpha-1}\} D^r y$$

$$= \sum_{r=0}^{n} x^{-n-\alpha-1} e^x (n)! \frac{x^r}{(r)!} A_{n-r}^{(\alpha)}(x) D^r y,$$

(1.5) follows immediately.

In (1.5) if we take $y = 1$, we obtain

$$\prod_{j=1}^{n} (xD + x - \alpha - j) \cdot 1 = (n)! A_n^{(\alpha)}(x) . \tag{1.6}$$

As an application of (1.5) and (1.6), let us consider

$$(m+n)! A_{m+n}^{(\alpha)}(x) = \prod_{j=1}^{m} (xD + x - \alpha - n - j) \prod_{j=1}^{n} (xD + x - \alpha - j) \cdot 1$$

$$= (n)! \prod_{j=1}^{m} (xD + x - \alpha - n - j) A_n^{(\alpha)}(x)$$

$$= (m)!\,(n)! \sum_{r=0}^{n} \frac{x^r}{(r)!} A_{m-r}^{(\alpha+n)}(x) D^r A_n^{(\alpha)}(x).$$

But since

$$D^r A_n^{(\alpha)}(x) = A_{n-r}^{(\alpha)}(x),$$

we readily get

$$\binom{m+n}{n} A_{m+n}^{(\alpha)}(x) = \sum_{r=0}^{\min(m,n)} \frac{x^r}{(r)!} A_{m-r}^{(\alpha+n)}(x) A_{n-r}^{(\alpha)}(x). \tag{1.7}$$

Further, from (1.7) we have

$$\sum_{m=0}^{\infty} \binom{m+n}{n} t^m A_{m+n}^{(\alpha)}(x) = \sum_{r=0}^{n} \frac{(xt)^r}{(r)!} A_{n-r}^{(\alpha)}(x) \sum_{m=0}^{\infty} t^m A_m^{(\alpha+n)}(x)$$

$$= \sum_{r=0}^{n} \frac{(xt)^r}{(r)!} A_{n-r}^{(\alpha)}(x) \cdot e^{xt} (1+t)^{-(\alpha+n+1)},$$

and making use of the relation [7, p. 45]

$$A_n^{(\alpha)}(x+y) = \sum_{r=0}^{n} \frac{y^r}{(r)!} A_{n-r}^{(\alpha)}(x),$$

we get the known formula [6, p. 7]

$$\sum_{m=0}^{\infty} \binom{m+n}{n} t^m A_{m+n}^{(\alpha)}(x) = e^{xt}(1+t)^{-(\alpha+n+1)} A_n^{(\alpha)}\{x(1+t)\}. \tag{1.8}$$

Another operational formula for the polynomials $A_n^{(\alpha)}(x)$ is

$$(1+D)^{-1-\alpha} x^n = (n)!\, A_n^{(\alpha)}(x), \qquad \left(D = \frac{d}{dx}\right). \tag{1.9}$$

To prove it we note that

$$(1+D)^{-1-\alpha} x^n = \sum_{r=0}^{n} (-1)^r \frac{(1+\alpha)_r}{(r)!} D^r x^n = \sum_{r=0}^{n} \frac{(1+\alpha)_r (-n)_r}{(r)!} x^{n-r},$$

which evidently yields the formula (1.9).

From (1.9) we have

$$(n)!\, A_n^{(\alpha+\beta)}(x) = (1+D)^{-1-\alpha-\beta}\, x^n,$$

$$A_n^{(\alpha+\beta)}(x) = (1+D)^{-\beta}\, A_n^{(\alpha)}(x); \tag{1.10}$$

the last formula gives us

$$A_n^{(\alpha+\beta)}(x) = \sum_{r=0}^{n} (-1)^r \frac{(\beta)^r}{(r)!}\, A_{n-r}^{(\alpha)}(x). \tag{1.11}$$

Further let us operate on the identity

$$e^{xt} = \sum_{n=0}^{\infty} \frac{t^n}{(n)!}\, x^n,$$

by $(1+D)^{-1-\alpha}$; the familiar shift rule then gives us

$$(1+D)^{-1-\alpha}\, e^{xt} = e^{xt}\,(1+t)^{-1-\alpha}\left\{1+\frac{D}{1+t}\right\}^{-1-\alpha}\cdot 1 = (1+t)^{-1-\alpha}\, e^{xt}.$$

On the other hand the second member yields

$$\sum_{n=0}^{\infty} t^n\, A_n^{(\alpha)}(x)$$

with the help of (1.9).

We thus arrive at the familiar generating function (1.2).

Now replace by tD_y, $\left(D_y = \frac{d}{dy}\right)$ in (1.2) and operate on both sides by $(1+D_y)^{-1-\beta}$. The left-hand side gives us

$$(1+D_y)^{-1-\beta}\, e^{xty}\,(1+ty) = e^{xty}\,\{1+xt+D_y\}^{-1-\beta}\,(1+yt)^{-1-\alpha}$$

$$= e^{xyt}\,(1+xt)^{-1-\beta} \sum_{r=0}^{\infty} (-1)^r \frac{(1+\beta)_r}{(r)!}\cdot\frac{D_y^r}{(1+xt)^r}\,(1+yt)^{-1-\alpha}$$

$$\simeq e^{xyl}\,(1+xt)^{-1-\beta}\,(1+yt)^{-1-\alpha}\,{}_2F_0\left[1+\alpha, 1+\beta; -; \frac{t}{(1+xt)(1+yt)}\right]$$

and the right-hand side yields

$$\sum_{n=0}^{\infty} (n)!\, t^n\, A_n^{(\alpha)}(x)\, A_n^{(\beta)}(y)\,.$$

Combining these two sides we finally get

$$(1.12)\qquad \sum_{n=0}^{\infty} (n)!\, t^n\, A_n^{(\alpha)}(x)\, A_n^{(\beta)}(y)$$

$$\simeq e^{xyt}(1+xt)^{-1-\beta}(1+yt)^{-1-\alpha}\,{}_2F_0\left[1+\alpha, 1+\beta; -; \frac{t}{(1+xt)(1+yt)}\right].$$

From the relation (1.12) we have

$$(1+xt)^{-1-\beta}(1+yt)^{-1-\alpha}\,{}_2F_0\left[1+\alpha, 1+\beta; -; \frac{t}{(1+xt)(1+yt)}\right]$$

$$= \sum_{r,s,k=0}^{\infty} \frac{(1+\alpha)_r (1+\beta)_r (1+\alpha+r)_s (1+\beta+r)_k}{(r)!\ (s)!\ (k)!} (-1)^{s+k} x^k y^s \cdot t^{r+s+k}$$

$$= \sum_{n=0}^{\infty} (-t)^n \sum_{k=0}^{n} \sum_{r=0}^{k} (-1)^r \frac{(1+\alpha)_{n+r-k}\ (1+\beta)_k}{(r)!\ (k-r)!\ (n-k)!} x^{k-r} y^{n-k}$$

$$= \sum_{n=0}^{\infty} (-t)^n \sum_{k=0}^{n} \frac{(1+\beta)_k (1+\alpha)_{n-k}}{(n-k)!} y^{n-k} \sum_{r=0}^{k} (-1)^r \frac{(1+\alpha+n-k)_r}{(r)!\ (k-r)!} x^{k-r}$$

$$= \sum_{n=0}^{\infty} (-t)^n \sum_{k=0}^{n} \frac{(1+\beta)_k (1+\alpha)_{n-k}}{(n-k)!} y^{n-k} A_k^{(\alpha+n-k)}(x).$$

Thus (1.12) is equivalent to

$$(1.13)\qquad \sum_{r=0}^{n} \frac{(r)!}{(n-r)!} (-1)^r (xy)^{n-r} A_n^{(\alpha)}(x)\, A_n^{(\beta)}(y)$$

$$= \sum_{k=0}^{n} \frac{(1+\beta)_k (1+\alpha)_{n-k}}{(n-k)!} y^{n-k} A_k^{(\alpha+n-k)}(x).$$

2. Making use of the relation [7, p. 45]

$$L_n^{-(\alpha+n+1)}(-x) = A_n^{(\alpha)}(x)$$

and our formula (1.8), we get

$$(1-D)^{\alpha+n}(-x)^n = (n)!\, L_n^{(\alpha)}(x), \tag{2.1}$$

which yields the following interesting result

$$L_n^{(\alpha+\beta)}(x) = (1-D)^{\beta} L_n^{(\alpha)}(x). \tag{2.2}$$

Now consider

$$L_n^{(\alpha+\beta+1)}(x) = (1-D)^{\beta+1} L_n^{(\alpha)}(x) = \left[1 + \frac{D}{1-D}\right]^{-\beta-1} L_n^{(\alpha)}(x)$$

$$= \sum_{r=0}^{n} (-1)^r \frac{(\beta+1)_r}{(r)!} (1-D)^{-r} D^r L_n^{(\alpha)}(x),$$

and it follows that

$$L_n^{(\alpha+\beta+1)}(x) = \sum_{r=0}^{n} \frac{(\beta+1)_r}{(r)!} L_{n-r}^{(\alpha)}(x). \tag{2.3}$$

Formula (2.3) was proved earlier in a different way by Al-Salam [1, p. 131], and our proof differs markedly with that of Rainville [5, p. 209].

Next let us consider the expression,

$$(-1)^n e^x D^n [e^{-x} L_n^{(\alpha)}(x)]$$

which by the usual shift rule gives us

$$(-1)^n e^x D^n [e^{-x} L_n^{(\alpha)}(x)] = (1-D)^n L_n^{(\alpha)}(x).$$

On making use of the relation (2.2) we obtain

$$(-1)^n e^x D^n [e^{-x} L_n^{(\alpha)}(x)] = L_n^{(\alpha+n)}(x),$$

which may be put in the form

$$R_n(1+\alpha, x) = (-1)^n e^x D^n [e^{-x} L_n^{(\alpha)}(x)], \tag{2.4}$$

where $R_n(a, x)$ is the pseudo Laguerre set defined by Shively [5, p. 298] as

$$R_n(a, x) = \frac{(a)_{2n}}{(n)!\,(a)_n} {}_1F_1(-n; a+n; x).$$

The formula (2.4) has been proved recently by Khandekar in a different way (see [4], p. 2).

Further, from (2.1) we have

$$\frac{(-x)^n}{(n)!} = (1-D)^{-\alpha-n} L_n^{(\alpha)}(x) = \sum_{r=0}^{n} \frac{(\alpha+n)_r}{(r)!} D^r L_n^{(\alpha)}(x)$$

which gives

$$\frac{(-x)^n}{(n)!} = \sum_{r=0}^{n} (-1)^r \frac{(\alpha+n)_r}{(r)!} L_{n-r}^{(\alpha+r)}(x). \tag{2.5}$$

Next consider the identity

$$e^{-xt} = \sum_{n=0}^{\infty} (-x)^n \frac{t^n}{(n)!},$$

operate on both sides by $(1-D)^\alpha$, make use of (2.1) and proceed as in the cases of (1.12) and (1.13). We then get the generating function

$$(1+t)^\alpha e^{-xt} = \sum_{n=0}^{\infty} t^n L_n^{(\alpha-n)}(x), \tag{2.6}$$

due to Erdélyi, and the known formula [2, p. 151]

$$\sum_{n=0}^{\infty} (n)!\, t^n L_n^{(\alpha-n)}(x) L_n^{(\beta-n)}(y) \tag{2.7}$$

$$= \begin{cases} e^{xyt}(1-yt)^{\alpha-\beta} t^\beta (\beta)!\, L_\beta^{(\alpha-\beta)}\left(-\dfrac{(1-xt)(1-yt)}{t}\right) \\ e^{xyt}(1-xt)^{\beta-\alpha} t^\alpha (\alpha)!\, L_\alpha^{(\beta-\alpha)}\left(-\dfrac{(1-xt)(1-yt)}{t}\right). \end{cases}$$

ACKNOWLEDGEMENT

I am grateful to Dr. H. M. Srivastava of Jodhpur University for his kind guidance and invaluable help during the preparation of this paper.

REFERENCES

1. W. A. Al-Salam, *Operational representations for the Laguerre and other polynomials,* Duke Math. J., **31** (1964), 127-142.

2. H. Buchholz, *Die konfluente hypergeometrische funktionen,* Berlin (1953).

3. L. Carlitz, *A note on the Laguerre polynomials,* Michigan Math. J., **7** (1960), 219-223.

4. P. R. Khandekar, *Rodrigues' formula for some generalized hypergeometric polynomials,* Math. Stu. of the Indian Math. Soc. (1) **34** (1966), 1-3.

5. E. D. Rainville, *Special Functions,* Macmillan, New York (1960).

6. R. P. Singh, *Some polynomials of Srivastava related to the Laguerre polynomials,* Math. Japonicae, **9** (1964), 5-9.

7. K. N. Srivastava, *Some polynomials related to the Laguerre polynomials,* J. Indian Math. Soc. (2), **28** (1964), 43-50.

8. H. M. Srivastava, *On Bessel, Jacobi and Laguerre polynomials,* Rend. Semin Mat. Univ. Padova, **35** (1965), 424-432.

Manoscritto pervenuto in redazione il 1° settembre 1966.