

RENDICONTI
del
SEMINARIO MATEMATICO
della
UNIVERSITÀ DI PADOVA

BIAGIO RICCERI

**Carathéodory's selections for multifunctions
with non-separable range**

Rendiconti del Seminario Matematico della Università di Padova,
tome 67 (1982), p. 185-190

http://www.numdam.org/item?id=RSMUP_1982__67__185_0

© Rendiconti del Seminario Matematico della Università di Padova, 1982, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

Carathéodory's Selections for Multifunctions with Non-Separable Range.

BLAGIO RICCERI (*)

Introduction.

Let (S, \mathcal{E}) be a measurable space, X a paracompact topological space, Y a Banach space and F a multifunction from $S \times X$ into Y . Our problem is to find a Carathéodory selection for F , that is a single-valued function f from $S \times X$ into Y such that: i) $f(t, x) \in F(t, x)$ for each $(t, x) \in S \times X$; the function $t \rightarrow f(t, x)$ is \mathcal{E} -measurable for each $x \in X$; ii) the function $x \rightarrow f(t, x)$ is continuous for each $t \in S$.

There are already some papers devoted to this problem (see [1], [2], [3], [4]). Nevertheless, in all of these papers the separability of X is assumed and then, from it, the separability of $F(S \times X)$ follows (see Proposition 2.2).

The aim of the present paper is to establish two theorems in which no hypothesis of separability of X neither of $F(S \times X)$ is needed. Moreover, some particular values of the multifunction F are allowed to be non-convex.

In Section 1 we put notations and basic definitions, while Section 2 contains the main results.

(*) Indirizzo dell'A.: Seminario Matematico, Città Universitaria - Viale A. Doria 6 - 95125 Catania.

Lavoro eseguito nell'ambito del G.N.A.F.A. del C.N.R.

1. Notations and basic definitions.

Given two sets $A', A'' \neq \emptyset$, we denote by $\mathcal{F}(A', A'')$ the set of all functions from A' into A'' . Given a function F from A' into the family of all non-empty subsets of A'' , we say that F is a multifunction from A' into A'' and, for each $A \subseteq A'$ and each $\Omega \subseteq A''$, we put: $F(A) = \bigcup_{t \in A} F(t)$ and $F^-(\Omega) = \{t \in A' : F(t) \cap \Omega \neq \emptyset\}$. If A', A'' are topologized, we say that a multifunction F from A' into A'' is lower semicontinuous whenever the set $F^-(\Omega)$ is open in A' for each open set $\Omega \subseteq A''$. Moreover, if \mathcal{E} is a σ -algebra of subsets of A' , we say that the multifunction F is \mathcal{E} -measurable whenever the set $F^-(\Omega)$ belongs to \mathcal{E} for each open set $\Omega \subseteq A''$. Now let (Σ, d) be a metric space. For each $x \in \Sigma$, $\Omega, \Omega', \Omega'' \subseteq \Sigma$ and $r \in \mathbb{R}^+$, we put: $B_d(\Omega, r) = \{y \in \Sigma : \exists z \in \Omega : d(y, z) < r\}$; $d(x, \Omega'') = \inf \{d(x, y) : y \in \Omega''\}$; $d^*(\Omega', \Omega'') = \sup \{d(z, \Omega'') : z \in \Omega'\}$. If $(\Sigma, \|\cdot\|)$ is a normed space, we will consider Σ with the metric d induced by the norm $\|\cdot\|$. Furthermore, given a set $A \neq \emptyset$, we put $\mathcal{F}_b(A, \Sigma) = \{f \in \mathcal{F}(A, \Sigma) : \sup_{t \in A} \|f(t)\| < +\infty\}$ and we consider $\mathcal{F}_b(A, \Sigma)$ as a normed space with the usual norm: $\|f\|_b = \sup_{t \in A} \|f(t)\|$, $f \in \mathcal{F}_b(A, \Sigma)$. Finally, we denote by d_b the metric induced by the norm $\|\cdot\|_b$.

2. The main results.

We begin this section by proving an useful proposition.

PROPOSITION 2.1. *Let (S, \mathcal{E}) be a measurable space, (Σ, d) a metric space and F, G two \mathcal{E} -measurable multifunctions from S into Σ such that $F(S)$ and $G(S)$ are separable. If $r \in \mathbb{R}^+$ is such that $I_r(t) = F(t) \cap B_d(G(t), r) \neq \emptyset$ for each $t \in S$, then the multifunction I_r is \mathcal{E} -measurable.*

PROOF. Put $P(t) = F(t) \times G(t)$ for each $t \in S$. Thus, for each $\Omega', \Omega'' \subseteq \Sigma$, we have

$$(1) \quad P^-(\Omega' \times \Omega'') = F^-(\Omega') \cap G^-(\Omega'')$$

Moreover, for each $\Omega \subseteq \Sigma$, we have

$$(2) \quad I_r^-(\Omega) = P^-(\{(x, y) \in F(S) \times G(S) : d(x, y) < r\} \cap (\Omega \times \Sigma)).$$

Now our claim follows from (2) and (1) since any open set in $F(S) \times G(S)$ is the union of a countable family of sets of type $\Omega' \times \Omega''$, with Ω', Ω'' open sets, respectively, in $F(S)$ and $G(S)$. \blacktriangle

We now prove the following selection theorem:

THEOREM 2.1. *Let $(S, \mathfrak{E}), X, Y, F$ be as in the Introduction and let $\{S_n\}$ be a sequence in \mathfrak{E} such that $S = \bigcup_{n=1}^{\infty} S_n$; Z a subset of X , with $\dim_X(Z) \leq 0$ ⁽¹⁾; M an \mathfrak{E} -measurable function from S into \mathbb{R}^+ ; $\{\lambda_n\}$ a sequence in $\mathcal{F}(X \times X, \mathbb{R}^+)$, with $\lim_{y \rightarrow x} \lambda_n(x, y) = 0$ for each $x \in X$ and each $n \in \mathbb{N}$. Moreover, let the following conditions be satisfied:*

- 1) *the set $F(t, x)$ is closed for each $(t, x) \in S \times X$ and convex for each $(t, x) \in S \times (X - Z)$;*
- 2) *the set $F(S_n, x)$ is separable and bounded for each $x \in X, n \in \mathbb{N}$;*
- 3) *the multifunction $t \rightarrow F(t, x)$ is \mathfrak{E} -measurable for each $x \in X$;*
- 4) *we have $d^*(F(t, x), F(t, y)) \leq M(t)\lambda_n(x, y)$ for each $t \in S_n, x, y \in X, n \in \mathbb{N}$.*

Under such hypotheses, there exists a Carathéodory selection for F .

PROOF. For each $m, n \in \mathbb{N}$ put: $S_{m,n} = \{t \in S_n : M(t) \leq m\}$. Rearrange the non-empty members of the family $\{S_{m,n}\}_{m,n \in \mathbb{N}}$ as a single sequence $\{S_k^*\}$. Obviously, $S_k^* \in \mathfrak{E}$ for each $k \in \mathbb{N}$ and $S = \bigcup_{k=1}^{\infty} S_k^*$. Fix $k \in \mathbb{N}$ and denote by \mathfrak{E}_k the family of all subsets of S_k^* belonging to \mathfrak{E} . Furthermore, put $\mathcal{A}_k = \{\varphi \in \mathcal{F}_b(S_k^*, Y) : \varphi \text{ is } \mathfrak{E}_k\text{-measurable and } \varphi(S_k^*) \text{ is separable}\}$ and observe that \mathcal{A}_k is a closed linear subspace of the Banach space $\mathcal{F}_b(S_k^*, Y)$. For each $x \in X$ put $H_k(x) = \{\varphi \in \mathcal{A}_k : \varphi(t) \in F(t, x), \forall t \in S_k^*\}$. From 1), 2), 3) and from the Kuratowski and Ryll-Nardzewski selection theorem, it follows that $H_k(x) \neq \emptyset$, and so we can consider the multifunction $x \rightarrow H_k(x)$ from X into \mathcal{A}_k . We claim that this multifunction is lower semicontinuous. Therefore, let Ω be an open subset of \mathcal{A}_k and let $\bar{x} \in H_k^-(\Omega)$. We must prove that \bar{x} is an interior point of $H_k^-(\Omega)$. For this purpose, choose $\varphi \in H_k(\bar{x}) \cap \Omega$ and $r \in \mathbb{R}^+$ such that $B_{d_b}(\varphi, r) \subseteq \Omega$. Now, let $m^*, n^* \in \mathbb{N}$ be such that $S_k^* = S_{m^*, n^*}$. Since $\lim_{x \rightarrow \bar{x}} \lambda_{n^*}(\bar{x}, x) = 0$, it follows that there exists a

⁽¹⁾ $\dim_X(Z) \leq 0$ means that $\dim(T) \leq 0$ for each $T \subseteq Z$ which is closed in X , where $\dim(T)$ denotes the covering dimension of T .

neighbourhood V of \bar{x} such that $\lambda_{n^*}(\bar{x}, x) < r/2m^*$ for each $x \in V$, $x \neq \bar{x}$. Therefore, by virtue of 4), for each $t \in S_k^*$ and each $x \in V$, we have

$$(3) \quad d^*(F(t, \bar{x}), F(t, x)) \leq m^* \lambda_{n^*}(\bar{x}, x) < m^* \frac{r}{2m^*} = \frac{r}{2}$$

On the other hand, since $\varphi \in H_k(\bar{x})$, we have $\varphi(t) \in F(t, \bar{x})$ for each $t \in S_k^*$, and so, by (3), we have $F(t, x) \cap B_d(\varphi(t), r/2) \neq \emptyset$ for each $t \in S_k^*$ and each $x \in V$. Now fix $x \in V$. For each $t \in S_k^*$, put $I_{r/2}(t) = F(t, x) \cap B_d(\varphi(t), r/2)$ and $\bar{I}_{r/2}(t) = \bar{I}_{r/2}(\bar{t})$. From 1), 2), 3), from the fact that $\varphi \in \mathcal{A}_k$ and from Proposition 2.1, it follows that the multifunction $I_{r/2}$ is δ_k -measurable and so the multifunction $\bar{I}_{r/2}$, since we have $I_{r/2}(W) = \bar{I}_{r/2}(W)$ for each open set $W \subseteq Y$. Thus, by virtue of the Kuratowski and Ryll-Nardzewski selection theorem again, there exists $\psi \in \mathcal{A}_k$ such that $\psi(t) \in \bar{I}_{r/2}(t)$ for each $t \in S_k^*$. Given $\varepsilon \in]0, r/2[$, obviously, for each $t \in S_k^*$, we have $\bar{I}_{r/2}(t) \subseteq F(t, x) \cap B_d(\varphi(t), r/2 + \varepsilon)$. Therefore, $\psi \in H_k(x)$ and $d_b(\varphi, \psi) \leq r/2 + \varepsilon < r$ and so $\psi \in H_k(x) \cap \Omega$. Hence, the neighbourhood V of \bar{x} is contained in $H_k^-(\Omega)$ and so our claim is proved. Furthermore, from 1) it follows that the set $H_k(x)$ is closed for each $x \in X$ and convex for each $x \in X - Z$. Therefore, by virtue of Theorem 1.1 of [5], there exists a continuous selection h_k for H_k . Now fix $\bar{t} \in S_k^*$ and put $h_{k,\bar{t}}(x) = h_k(x)(\bar{t})$ for each $x \in X$. Moreover, given $\bar{z} \in Y$, for each $x \in X$, define the function $g_{k,x}: S_k^* \rightarrow Y$ as follows

$$g_{k,x}(t) = \begin{cases} \bar{z} & \text{if } t = \bar{t} \\ h_k(x)(t) & \text{if } t \neq \bar{t}, t \in S_k^* \end{cases}$$

Obviously, $g_{k,x} \in \mathcal{F}_b(S_k^*, Y)$. For each $\varrho \in \mathbb{R}^+$, we have

$$(4) \quad h_{k,\bar{t}}^{-1}(B_d(\bar{z}, \varrho)) = \bigcup_{x \in X} h_k^{-1}(B_{d_b}(g_{k,x}, \varrho) \cap \mathcal{A}_k).$$

From (4) it follows that the function $x \rightarrow h_{k,\bar{t}}(x)$ is continuous, since the function $x \rightarrow h_k(x)$ is continuous. Now put $f_k(t, x) = h_k(x)(t)$ for each $(t, x) \in S_k^* \times X$. Thus, the function f_k is a Carathéodory selection for $F|_{S_k^* \times X}$. This holds for any $k \in N$. Finally, put:

$$f(t, x) = \begin{cases} f_1(t, x) & \text{if } (t, x) \in S_1^* \times X \\ f_k(t, x) & \text{if } (t, x) \in \left(S_k^* - \bigcup_{j=1}^{k-1} S_j^* \right) \times X, k \geq 2. \end{cases}$$

Obviously, the function f is the claimed Carathéodory selection for F . ▲

In a very similar way it is possible to prove the following further result:

THEOREM 2.2. *Let (S, \mathcal{E}) be a measurable space and $\{S_n\}$ a sequence in \mathcal{E} such that $S = \bigcup_{n=1}^{\infty} S_n$ and S_n is finite for each $n \in \mathbb{N}$; let X, Y, Z be as in Theorem 2.1 and let F be a multifunction from $S \times X$ into Y which satisfies conditions 1), 3) of Theorem 2.1 and the following ones:*

- 2) *the set $F(S, x)$ is separable for each $x \in X$;*
- 4) *the multifunction $x \rightarrow F(t, x)$ is lower semicontinuous for each $t \in S$.*

Under such hypotheses, there exists a Carathéodory selection for F .

Finally, we want to justify our assertion, made in the Introduction, that the range of the multifunction F in Theorem 1 of [1] is separable (in fact, in [2], [3], [4] the whole space Y is already assumed to be separable).

Indeed, we have the following

PROPOSITION 2.2. *Let S be a non-empty set and X, Y two topological spaces, with X separable. Furthermore, let A be a countable dense subset of X and F a multifunction from $S \times X$ into Y such that:*

- 1) *the set $F(S, x)$ is separable for each $x \in A$;*
- 2) *the multifunction $x \rightarrow F(t, x)$ is lower semicontinuous for each $t \in S$.*

Under such hypotheses, the set $F(S \times X)$ is separable.

PROOF. For each $x \in A$ choose a countable dense subset D_x of $F(S, x)$. We claim that the countable set $\bigcup_{x \in A} D_x$ is dense in $F(S \times X)$.

Indeed, let $\bar{y} \in F(S \times X)$ and let W be an open neighbourhood of \bar{y} . Let $(\bar{t}, \bar{x}) \in S \times X$ be such that $\bar{y} \in F(\bar{t}, \bar{x})$. From 2) it follows that there exists a neighbourhood V of \bar{x} such that $F(\bar{t}, x) \cap W \neq \emptyset$ for each $x \in V$. Since A is dense in X , there exists $x^* \in A \cap V$. But then, since $F(\bar{t}, x^*) \cap W \neq \emptyset$ and D_{x^*} is dense in $F(S, x^*)$, it follows that $D_{x^*} \cap W \neq \emptyset$, and so our claim is proved. ▲

REFERENCES

- [1] A. CELLINA, *A selection theorem*, Rend. Sem. Mat. Univ. Padova, **55** (1976), pp. 143-149.
- [2] C. CASTAING, *Sur l'existence des sections séparément mesurables et séparément continues d'une multi-application*, Séminaire d'Analyse Convexe, Montpellier, Exposé n. 14 (1975).
- [3] C. CASTAING, *A propos de l'existence des sections séparément mesurables et séparément continues d'une multiapplication séparément mesurable et séparément semi-continue inférieurement*, Séminaire d'Analyse Convexe, Montpellier, Exposé n. 6 (1976).
- [4] A. FRYSZKOWSKI, *Carathéodory type selectors of set-valued maps of two variables*, Bull. Acad. Polon. Sci., Sér. Sci. Math. Astronom. Phys., **25** (1977), pp. 41-46.
- [5] E. MICHAEL - C. PIXLEY, *A unified theorem on continuous selections*, Pacific J. Math., **87** (1980), pp. 187-188.

Manoscritto pervenuto in redazione il 15 luglio 1981.