RENDICONTI
del
SEMINARIO MATEMATICO
della
UNIVERSITÀ DI PADOVA

C. MENINI

Errata-Corrige : "Linearly compact rings and strongly quasi-injective modules"

Rendiconti del Seminario Matematico della Università di Padova, tome 69 (1983), p. 305-306

<http://www.numdam.org/item?id=RSMUP_1983__69__305_0>

© Rendiconti del Seminario Matematico della Università di Padova, 1983, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (http://rendiconti.math.unipd.it/) implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/conditions). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM
Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
http://www.numdam.org/

Rend. Sem. Mat. Univ. Padova, 69 (1983)

ERRATA - CORRIGE

« Linearly Compact Rings and Strongly Quasi-Injective Modules ».

C. MENINI (*)

Introduction.

I) Statement (d) of THE MAIN THEOREM of $[M]$ must be changed in the following form:

« (d) *Let* ${}_RU$ *be the minimal cogenerator of* $\mathcal{C}_{\mathcal{F}}$, $T = \mathrm{End}({}_RU)$. *Then* ${}_RU_T$ *is faithfully balanced and the module* U_T *is s.q.i.* »

Consequently the last assertion of the same theorem has to be erased.

The proof of $(a) \Rightarrow (d)$ of this theorem is now the following:

« Let us remark, first of all, that in view of Lemma 5, R is linearly compact in its U-topology. Thus, since ${}_RU$ is a selfcogenerator, by Corollary 7.4 [2], $R = \mathrm{End}(U_T)$ and hence ${}_RU_T$ is faithfully balanced. Moreover, since we already proved that $(a) \Rightarrow (b)$, U_T is q.i.. Thus it is enough to prove that R separates points and submodules of U_T. Let $L \leqslant U_T$ and let $x \in U_T$. Assume that $\mathrm{Ann}_R(x) \geqslant \mathrm{Ann}_R(L)$. Since R is linearly compact in the U-topology, Rx is linearly compact discrete. Thus, since $Rx \leqslant {}_RU$, Rx is finitely embedded. Hence, by Lemma 8, there is a finite subset $\{y_1, \ldots, y_n\} \subseteq L$ such that $\mathrm{Ann}_R(x) \geqslant \bigcap_{j=1}^{n} \mathrm{Ann}_R(y_j)$. Let $S_1, \ldots, S_m \in \mathcal{S}_{\mathcal{F}}$ such that the R-submodule of ${}_RU$ spanned by x and y_j's, $j = 1, \ldots, n$, is contained in ${}_RM = \bigoplus_{i=1}^{m} t_{\mathcal{F}}(E(S_i)) \leqslant {}_RU$. Since

(*) Indirizzo dell'A.: Istituto di Matematica dell'Università di Ferrara, Via Machiavelli 35, 44100 Ferrara (Italy).

${}_RM$ is quasi-injective, by the proof of Lemma 7, there exists a morphism $f: {}_RM^n \to {}_RM$ such that $x = ((y_1, \dots, y_n))f$. Since ${}_RM$ is a direct summand of ${}_RU$, f extends to a morphism $g: {}_RU^n \to {}_RU$. Thus there exist $t_1, \dots, t_n \in T$ such that $x = \sum_{i=1}^{n} t_i y_i$ and hence $x \in L$. »

II) The last assertions of Theorem 10, concerning the explicit form of ${}_RK$ and K_A are false, while equivalence of statements (a), (b) and (c) is true and is also true that if (a), (b), (c) hold then A is linearly compact in its K-topology.

The proof of $(c) \Rightarrow (b)$ rules as follows.

« Let $x \in K$. Rx is linearly compact discrete and hence $\mathrm{Soc}(Rx)$ is a direct sum of a finite number of left simple R-modules $S_1, \dots, S_n$. By hypothesis, $\mathrm{Soc}({}_RK)$ is essential in ${}_RK$. Hence $\mathrm{Soc}(Rx)$ is essential in Rx. It follows that

$$Rx \leqslant \bigoplus_{1=i}^{n} t_{\mathcal{F}}(E(S_i)) \,. \tag{1}$$

Let us prove that R separates points and submodules of K_A. Let $L \leqslant K_A$ and let $x \in K$. Assume that $\mathrm{Ann}_R(x) \geqslant \mathrm{Ann}_R(L)$. Note that, by (1), $R/\mathrm{Ann}_R(x) \cong Rx$ is finitely embedded. Hence, by Lemma 8, there is a finite subset $F \subseteq L$ such that $\mathrm{Ann}_R(x) \geqslant \bigcap_{l \in F} \mathrm{Ann}_R(l)$. Thus, by Lemma 7, x belongs to the submodule of K_A spanned by F and hence $x \in L$.

III) Finally, the first part of the proof of Theorem 14 is modified as follows.

« Proof. Let $\mathcal{F}$ be the filter of open left ideals of τ and let U_R be the minimal injective cogenerator of $\mathcal{C}_{\mathcal{F}}$: ${}_RU = t_{\mathcal{F}}\Big(E\Big(\bigoplus_{S \in \mathcal{S}_{\mathcal{F}}} S\Big)\Big)$. Set $A = \mathrm{End}({}_RU)$. By theorem 10 and Lemma 6, ${}_RU_A$ is faithfully balanced and both the modules ${}_RU$ and U_A are s.q.i. »

At this point the remaining part of the proof works.

Manoscritto pervenuto in redazione il 7 dicembre 1982.

REFERENCES

[M] C. Menini, *Linearly compact rings and strongly quasi-injective modules.*