RENDICONTI
del
SEMINARIO MATEMATICO
della
UNIVERSITÀ DI PADOVA

PIERO MONTECCHIARI

Multiplicity results for a class of semilinear elliptic equations on $\mathbb{R}^m$

Rendiconti del Seminario Matematico della Università di Padova, tome 95 (1996), p. 217-252

<http://www.numdam.org/item?id=RSMUP_1996__95__217_0>

© Rendiconti del Seminario Matematico della Università di Padova, 1996, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (http://rendiconti.math.unipd.it/) implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/conditions). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM
Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
http://www.numdam.org/

REND. SEM. MAT. UNIV. PADOVA, Vol. 95 (1996)

Multiplicity Results for a Class of Semilinear Elliptic Equations on $\mathbb{R}^m$.

PIERO MONTECCHIARI (*)

ABSTRACT - We prove the existence of an uncountable set of bounded solutions for a class of semilinear elliptic equations on $\mathbb{R}^m$.

1. - Introduction.

In this work we are concerned with the study of existence and multiplicity of solutions to the problem

$$(P) \qquad -\Delta u + u = f(x, u), \qquad u \in H^1(\mathbb{R}^m)$$

where $m \geq 1$ and f satisfies the assumptions

(f1) $f \in C(\mathbb{R}^m \times \mathbb{R}, \mathbb{R})$.

(f2) $f(x, 0) = f_z(x, 0)$ for any $x \in \mathbb{R}^m$.

(f3) $\exists b_1, b_2 > 0$ such that $|f(x, z)| \leq b_1 + b_2 |z|^s$, $\forall (x, z) \in \mathbb{R}^m \times \mathbb{R}$, where $s \in (1, 2^* - 1)$ with $2^* = 2m/(m-2)$ if $m > 2$ and s is not restricted if $m = 1, 2$.

The hypotheses (f1)-(f3) are the ones studied in [10] assuming also that $f(x, z)$ is periodic in x and superquadratic in z. Here we consider the following more general case.

We say that a set $A \subset \mathbb{R}^m$ is large at infinity if $\forall R > 0$ $\exists x \in A$ such that $B_R(x) = \{ y \in \mathbb{R}^m / |y - x| < R \} \subset A$. Clearly any cone in $\mathbb{R}^m$ is large at infinity. Another example is a cone minus the union of the annuli centered in zero and with radii $(2n)^2$, $(2n+1)^2$.

(*) Indirizzo dell'A.: S.I.S.S.A., Via Beirut 2/4, 34013 Trieste.

We assume that there exists a function $f_\infty : \mathbb{R}^m \times \mathbb{R} \to \mathbb{R}$ verifying (f1)-(f3), and a set $A \subset \mathbb{R}^m$ large at infinity such that

(f4) $\exists \beta > 2$ and $\alpha \in [0, \mu/2 - 1)$ such that $\beta F_\infty(x, z) = \beta \int_0^z f_\infty(x, t)\,dt \leqslant f_\infty(x, z)z + \alpha|z|^2$, $\forall (x, z) \in \mathbb{R}^m \times \mathbb{R}$, and $F_\infty(x_0, z_0) > \alpha/(\beta - 2)z_0^2$ for an $(x_0, z_0) \in \mathbb{R}^m \times \mathbb{R}$.

(f5) $f_\infty(x + p, z) = f_\infty(x, z)$ for any $p \in \mathbb{Z}^m$, $(x, z) \in \mathbb{R}^m \times \mathbb{R}$.

(f6) $\forall \varepsilon > 0\ \exists R > 0$ such that $\sup_{x \in A \setminus B_R(0)} |f(x, z) - f_\infty(x, z)| \leqslant \varepsilon(|z| + |z|^s)$ $\forall z \in \mathbb{R}$.

Putting $F(x, z) = \int_0^z f(x, t)\,dt$ we define on $X = H^1(\mathbb{R}^m)$ the functionals $\varphi(u) = (1/2)\|u\|^2 - \int_{\mathbb{R}^m} F(x, u)\,dx$, $\varphi_\infty(u) = (1/2)\|u\|^2 - \int_{\mathbb{R}^m} F_\infty(x, u)\,dx$, where $\|u\|^2 = \int_{\mathbb{R}^m} |\nabla u|^2 + |u|^2\,dx$, and we look for solutions of (P) as critical points of φ.

As we will see the assumptions (f1)-(f5) are sufficient to guarantee the existence of at least one non zero critical point of the «periodic» functional φ_∞. By (f5), if u is a critical point of φ_∞, then also $p * u = u(\cdot - p)$ is a critical point of φ_∞ for any $p \in \mathbb{Z}^m$. If we translate this u in a region where f and f_∞ are very close one to the other, one could expect that nearby this translate of u there is a critical point of φ. In general this is not true as the following example shows.

Let $f(x, z) = a(x_1)|z|^2 z$ with $a \in C^1(\mathbb{R}, \mathbb{R})$, $a(t) \geqslant a_0 > 0$, $\dot{a}(t) > 0$ $\forall t \in \mathbb{R}$, and assume that u is a solution of (P). By standard bootstrap argument we get that $u \in H^2(\mathbb{R}^m)$, therefore $\varphi'(u)\,\partial_1 u = 0$. But, if $e_1 = (1, 0, \ldots, 0)$, we have

$$\varphi'(u)\,\partial_1 u = \frac{d}{ds}\,\varphi(u(\cdot + se_1))\big|_{s=0} = \int \dot{a}(x_1)\,\frac{|u|^4}{4}\,dx = 0$$

which implies $u = 0$ (see [12]).

To avoid this situations we make a discreteness assumption on the set of the critical points of the functional at infinity.

We note first of all that φ_∞ satisfies the geometrical hypotheses of the mountain pass theorem. Letting $\Gamma = \{\gamma \in C([0, 1], X);\ \gamma(0) = 0, \varphi_\infty(\gamma(1)) < 0\}$, we put $c = \inf_{\gamma \in \Gamma} \max_{t \in [0, 1]} \varphi_\infty(\gamma(t))$.

Setting $K_\infty = \{u \in X\setminus\{0\} / \varphi'_\infty(u) = 0\}$, we assume

(*) $\exists c^* > c$ such that $K_\infty^{c^*} = K_\infty \cap \{\varphi_\infty < c^*\}$ is denumerable.

We note that the hypothesis (*) excludes the asymptotically autonomous cases, because if f_∞ does not depends on x and $u \in K_\infty^{c^*}$ then $p * u \in K_\infty^{c^*}$ for any $p \in \mathbb{R}^m$.

In this setting we are able to prove the following

THEOREM 1.1. *If* (f1)-(f6) *and* (*) *hold then* (P) *admits infinitely many distinct solutions.*

Precisely there exists $u \in X$, *solution to the equation* $-\Delta u + u = f_\infty(x, u)$ *for which we have that* $\forall r > 0$ *there exist* $M = M(r) \in \mathbb{N}$ *and* $R = R(r) > 0$ *such that for any finite sequence* $\{p_1, \ldots, p_k\} \subset \mathbb{Z}^m$ *that verifies*

i) $|p_1| \geq R$ and $|p_i| \geq |p_{i-1}| + 2M \quad i = 2, \ldots, k$,

ii) $B_M(p_i) \subset A\setminus B_R(0) \quad i = 1, \ldots, k$,

there exists a solution v *to* (P) *such that if we put* $|p_{k+1}| = +\infty$ *then*

$$\|v - p_1 * u\|_{B_{(1/2)(|p_1| + |p_2|)}(0)} \leq r,$$

$$\|v - p_i * u\|_{B_{(1/2)(|p_i| + |p_{i+1}|)}(0)\setminus B_{(1/2)(|p_i| + |p_{i-1}|)}(0)} \leq r \quad i = 2, \ldots, k,$$

where if $A \subset \mathbb{R}^m$ *is measurable then* $\|u\|_A^2 = \int_A |\nabla u|^2 + |u|^2 dx$.

In particular, for $k = 1$ we get that if $p \in \mathbb{Z}^m$ verifies $B_M(p) \subset A\setminus B_R(0)$ then there is a solution v to (P) which is near $u(\cdot - p)$. Moreover for $k > 1$ if we choose any set of k disjoint annuli centered in zero, each of which intersects the set $A\setminus B_R(0)$ in a ball of radius M centered in a point of $\mathbb{Z}^m$, then there is a solution to (P) which is near a translate of u in each of this balls. We call this type of solution k-bump solution.

The first proof via variational methods of the existence of 2-bump solutions was given by E. Séré in [22] solving the homoclinic existence problem for first order periodic and convex Hamiltonian systems. This paper inspired the work of V. Coti Zelati and P. H. Rabinowitz [9] on second order periodic Hamiltonian systems where was proved the existence of infinitely many k-bump homoclinic solutions for any $k \in \mathbb{N}$. In [10] they adapted these techniques to find the existence of infinitely many k-bump solutions for any $k \in \mathbb{N}$ for the problem (P) in the case in which $f(x, z)$ is periodic in x and superquadratic in z. S. Alama and Y.

Y. Lee in [3] studied the problem (P) assuming f asymptotic as $|x| \to \infty$ to a function f_∞ of the type considered in [10]. In that paper they were able to prove that the problem (P) admits infinitely many k-bump solutions. All this results are based on assuming that there exists $c^* > c$ such that $K_\infty^{c^*}/\mathbb{Z}^m$ is finite (clearly, in the periodic case, the functional φ_∞ is φ itself).

The existence of k-bump homoclinic solutions with minimum distance between the bumps independent from k was first proved by E. Séré in [23] for first order convex and periodic Hamiltonian systems. As a consequence, considering the C^1_{loc}-closure of the set of the multibump homoclinic orbits, using the Ascoli Arzelà Theorem, he finds solutions with possibly infinitely many bumps.

Also in Theorem 1.1 the minimum distance between the bumps of any k-bump solution depends only on r (being given by $M(r)$). Using the Ascoli Arzelà theorem it is therefore possible to prove as in [23] the existence of a class of bounded solutions of the equation $-\Delta u + u = f(x, u)$. Precisely we have:

THEOREM 1.2. *Under the same assumptions of Theorem* 1.1, *it holds that for any* $r > 0$ *there exist* $M = M(r) \in \mathbb{N}$ *and* $R = R(r) > 0$ *such that for any sequence* $\{p_j\}_{j \in \mathbb{N}} \subset \mathbb{Z}^m$ *that verifies*

$$\text{i)} \quad |p_1| \geq R \text{ and } |p_i| \geq |p_{i-1}| + 2M \qquad i \geq 2,$$

$$\text{ii)} \quad B_M(p_i) \subset A \setminus B_R(0) \qquad i \in \mathbb{N},$$

and for every sequence $\sigma = (\sigma_j)_{j \in \mathbb{N}} \in \{0, 1\}^{\mathbb{N}}$ *there exists* $v_\sigma \in H^1_{\text{loc}}(\mathbb{R}^m)$ *satisfying* $-\Delta v_\sigma + v_\sigma = f(x, v_\sigma)$ *such that*

$$\|v_\sigma - \sigma_1(p_1 * u)\|_{B_{(1/2)(|p_1| + |p_2|)}(0)} \leq r,$$

$$\|v_\sigma - \sigma_i(p_i * u)\|_{B_{(1/2)(|p_i| + |p_{i+1}|)}(0) \setminus B_{(1/2)(|p_i| + |p_{i-1}|)}(0)} \leq r \qquad i \geq 2.$$

The tools used in the proof of Theorem 1.1 are related to the ones developed in [18] and then improved with P. Caldiroli in [7] and with S. Abenda and P. Caldiroli in [1] studying the homoclinic existence problem for second order Hamiltonian systems. These arguments, inspired by [23] and [9], permits us to strengthens the results contained in [3] in a more general setting. In fact the superquadratic assumption (f4) is verified also by functions f_∞ which change sign. Moreover the assumption (*), is satisfied if the functional φ_∞ is for example a Morse functional. In the one dimensional case ($m = 1$), it is possible to verify this condition via the Melnikov theory when f_∞ is a periodic perturbation of particular autonomous problems (see [1]).

Another differences with the work of S. Alama and Y. Y. Lee [3] is the fact that f is not assumed to be asymptotic to f_∞ as $|x| \to \infty$ but only on a set large at infinity. This permits us to consider the problem (P) when f is assumed to be asymptotic in different sets large at infinity to different functions. Precisely we consider the hypothesis

(f7) $\exists A_1, \ldots, A_l \subset \mathbb{R}^m$, large at infinity, $f_1, \ldots, f_l$ satisfying (f1)-(f5) for which $\forall \varepsilon > 0$ $\exists R > 0$ such that $\sup_{x \in A_\iota \setminus B_R(0)} |f(x, z) - f_\iota(x, z)| \leq \varepsilon(|z| + |z|^s)$ $\forall z \in \mathbb{R}$, $\forall \iota \in \{1, \ldots, l\}$.

If for any $\iota \in \{1, \ldots, l\}$, we define $\varphi_\iota(u) = (1/2)\|u\|^2 - \int_{\mathbb{R}^m} F_\iota(x, u(x))\,dx$, $\mathcal{K}_\iota = \{u \in X \setminus \{0\};\ \varphi_\iota'(u) = 0\}$, c_ι the mountain pass level of φ_ι and we assume

($*_\iota$) $\exists c_\iota^* > c_\iota$ such that $\mathcal{K}_\iota^{c_\iota^*} = \mathcal{K}_\iota \cap \{\varphi_\iota < c_\iota^*\}$ is denumerable.

then, by Theorem 1.2, we have l different sets of multibump solutions, each constructed with a suitable critical point of the functional φ_ι.

In fact, we prove that there are also multibump solutions of (P) of mixed type, as said in the following theorem.

THEOREM 1.3. *Assume that* (f1)-(f5), (f7) *and* ($*_\iota$) *hold. There exists* $u_1, \ldots, u_l \in X$, *satisfying* $-\Delta u_\iota + u_\iota = f_\iota(x, u_\iota)$ *for which we have that for any* $r > 0$ *there exist* $M = M(r) \in \mathbb{N}$ *and* $R = R(r) > 0$ *such that for any sequences* $\{p_i\}_{i \in \mathbb{N}} \subset \mathbb{Z}^m$, $\{j_i\}_{i \in \mathbb{N}} \subset \{1, \ldots, l\}^{\mathbb{N}}$, *that verify*

i) $|p_1| \geq R$ and $|p_i| \geq |p_{i-1}| + 2M \quad i \geq 2$,

ii) $B_M(p_i) \subset A_{j_i} \setminus B_R(0) \quad i \in \mathbb{N}$,

and for every sequence $\sigma = (\sigma_i)_{i \in \mathbb{N}} \in \{0, 1\}^{\mathbb{N}}$ *there exists* $v_\sigma \in H^1_{\mathrm{loc}}(\mathbb{R}^m)$ *satisfying* $-\Delta v_\sigma + v_\sigma = f(x, v_\sigma)$ *such that*

$$\|v_\sigma - \sigma_1(p_1 * u_{j_1})\|_{B_{(1/2(|p_1| + |p_2|)}(0)} \leq r,$$

$$\|v_\sigma - \sigma_i(p_i * u_{j_i})\|_{B_{(1/2)(|p_i| + |p_{i+1}|)}(0) \setminus B_{(1/2)(|p_i| + |p_{i-1}|)}(0)} \leq r \quad i \geq 2.$$

If $\sigma_i \neq 0$ *only for a finite number of indices then* v_σ *is actually a solution to* (P).

As last remark we point out that an analogous result was proved by S. Angenent in [2] in a different setting ($z - f(x, z)$ is assumed to be periodic in x and bounded together with its derivatives), using essen-

tially fixed point arguments. He proved his result under the assumption that the solution u was such that the operator $-\Delta + I - f_z(x, u(x))$ had a bounded inverse. He was able to verify this hypothesis for periodic perturbation of particular autonomous problem which admits a unique (up to translations) radial solution, using a bifurcation theorem due to A. Weinstein [24]. It is known that the problem (P) when $f(x, z) = z^p$ admits a unique positive solution (see [13]) and it should be interesting to check if the hypothesis (*) holds for periodic perturbations of this f.

The work is divided into four parts. In the first section we give some preliminaries. The second is devoted to the study of some properties of the periodic problem which we use in section three to construct a particular pseudogradient field of φ. Finally in section four we prove the existence theorem.

2. – A local compactness property.

In this section we study some properties of the functional φ which are independent on the assumptions on the asymptotic behaviour of f. All the results contained here are true under the hypotheses (f1)-(f3). In the proofs that follow we shall always consider the case $m \geq 3$, the proofs for $m = 1$ or 2 being not more difficult.

We have to note first of all that (f1)-(f3) imply

$$(2.1) \qquad \forall \varepsilon > 0 \ \exists A_\varepsilon > 0 / |f(x, z)h| \leq \varepsilon |z| |h| + A_\varepsilon |z|^s |h| \quad \text{for all } (x, z) \in \mathbb{R}^m \times \mathbb{R}$$

and obviously an analogous estimate holds also for $F(x, z)$. This permits us to say that φ is well defined on X because of the Sobolev Immersion Theorem. Actually the following holds:

PROPOSITION 2.2. $\varphi \in C^1(X, \mathbb{R})$.

PROOF. We prove first that φ is Gateaux differentiable. Given $h \in X$, by (2.1), we get that

$$\frac{1}{t} |F(x, u + th) - F(x, u)| = \frac{1}{t} \left| \int_0^t f(x, u + sh) h \, ds \right| \leq$$

$$\leq |h(x)|(|u(x)| + |h(x)|) + c_1 A_1 |h(x)|(|u(x)|^s + |h(x)|^s).$$

Being this last function in $L^1(\mathbb{R}^m)$ we can use the dominated conver-

gence theorem to get that

$$\lim_{t\to 0}\frac{1}{t}\,|(u+th)-\varphi(u)| = \langle u, h\rangle - \int_{\mathbb{R}^m} f(x,u)\,h\,dx = \varphi'_G(u)\,h\,.$$

We prove now that φ'_G is continuous. Let $u_n \to u$ and $\{u_{n_k}\} \subset \{u_n\}$. By the Sobolev Immersion theorem there exists $\{u_{n_{k_j}}\} \subset \{u_{n_k}\}$ and a function $v \in L^2(\mathbb{R}^m) \cap L^{s+1}(\mathbb{R}^m)$ such that $u_{n_{k_j}}(x) \to u(x)$ a.e. in $\mathbb{R}^m$ and $|u_{n_{k_j}}(x)| \leqslant v(x)$ a.e. in $\mathbb{R}^m$. Using again the dominated convergence theorem we get $\varphi'_G(u_{n_{k_j}}) \to \varphi'_G(u)$ in X^*. Since this can be done for any subsequence of $\{u_n\}$ the proposition is proved. ∎

LEMMA 2.3. *$\varphi(u) = (1/2)\|u\|^2 + o(\|u\|^2)$ and $\varphi'(u)\,u = \|u\|^2 + o(\|u\|^2)$ as $u \to 0$.*

PROOF. If $\varepsilon > 0$ then, by (2.1), $\left|\int_{\mathbb{R}^m} f(x,u)\,u\,dx\right| \leqslant (\varepsilon + c_2 A_\varepsilon \cdot \|u\|^{s+1-2})\|u\|^2$ from which $\int_{\mathbb{R}^m} f(x,u)\,u\,dx = o(\|u\|^2)$. Analogously $\int_{\mathbb{R}^m} F(x,u)\,dx = o(\|u\|^2)$. ∎

As a consequence we get a first compactness property of φ:

(2.4) *$\exists \varrho > 0$ such that if $\|u_n\| \leqslant 2\varrho$ and $\varphi'(u_n) \to 0$ then $u_n \to 0$.*

The hypotheses (f1)-(f3) are not sufficient to guarantee that the Palais Smale sequences are bounded in X. In the following we study the behavior of the bounded Palais-Smale sequences of φ. If $M \subset \mathbb{R}^m$ is measurable then we put $\|u\|^2_M = \int_M |\nabla u|^2 + |u|^2\,dx$.

LEMMA 2.5. *If $u_n \to u$ weakly in X is such that $\varphi(u_n) \to b$, $\varphi'(u_n) \to 0$ then $\varphi'(u) = 0$, $\varphi(u_n - u) \to b - \varphi(u)$ and $\varphi'(u_n - u) \to 0$.*

PROOF. We prove first that $\varphi'(u) = 0$. Let $h \in X$ with $\|h\| = 1$. Fixing $\varepsilon > 0$ there exists $R > 0$ such that $\|h\|_{|x|>R} \leqslant \varepsilon$. Let $\{u_{n_k}\} \subset \{u_n\}$ and $v \in L^2(B_R(0)) \cap L^{s+1}(B_R(0))$ be such that $u_{n_k}(x) \to u(x)$ a.e. on $B_R(0)$, $|u_{n_k}(x)| \leqslant v(x)$ a.e. on $B_R(0)$. By (2.1) and the dominated convergence theorem we get

$$|\varphi'(u)\,h| = |\varphi'(u_{n_k})\,h + \langle u - u_{n_k}, h\rangle - \int_{\mathbb{R}^m} (f(x,u) - f(x,u_{n_k}))\,h\,dx| \leqslant$$

$$\leqslant o(1) + \left| \int_{|x| > R} (f(x, u) - f(x, u_{n_k}))\, h\, dx \right| \leqslant$$

$$\leqslant o(1) + c_3 \int_{|x| > R} (|u| + |u_{n_k}|)|h| + (|u|^s + |u_{n_k}|^s)|\,|h|\, dx \leqslant$$

$$\leqslant o(1) + c_4(\|f\|_{|x| > R} + \|h\|^{s+1}_{|x| > R}) \leqslant o(1) + c_4(\varepsilon + \varepsilon^{s+1}).$$

Since ε is arbitrary the claim follows.

Let's now prove that $\varphi'(u_n - u) \to 0$. Since

$$\varphi'(u_n - u)\, h = \varphi'(u_n)\, h - \int_{\mathbf{R}^m} (f(x, u_n - u) - f(x, u_n) + f(x, u))\, h\, dx,$$

it is sufficient to show that

$$\sup_{\|h\| = 1} \left| \int_{\mathbf{R}^m} (f(x, u_n - u) - f(x, u_n) + f(x, u))\, h\, dx \right| \to 0.$$

Given $\varepsilon > 0$ fix $R > 0$ such that $\|u\|_{|x| > R} < \varepsilon$. Then

$$\sup_{\|h\| = 1} \left| \int_{\mathbf{R}^m} (f(x, u_n - u) - f(x, u_n) + f(x, u))\, h\, dx \right| \leqslant$$

$$\leqslant \sup_{\|h\| = 1} \left| \int_{|x| \leqslant R} (\ldots)\, h\, dx \right| + \sup_{\|h\| = 1} \left| \int_{|x| > R} (\ldots)\, h\, dx \right|.$$

Consider the first addendum. Given $\{u_{n_k}\} \subset \{u_n\}$ there exists $\{u_{n_{k_j}}\} \subset \{u_{n_k}\}$ and a function $v \in L^2(B_R(0)) \cap L^{s+1}(B_R(0))$ such that $u_{n_{k_j}}(x) \to u(x)$ a.e. on $B_R(0)$ and $|u_{n_{k_j}}(x)| \leqslant v(x)$ a.e. on $B_R(0)$. We get

$$\left| \int_{B_R(0)} (f(x, u_{n_{k_j}} - u) - f(x, u_{n_{k_j}}) + f(x, u))\, h\, dx \right| \leqslant$$

$$\leqslant c_5 \left(\int_{B_R(0)} |f(x, u_{n_{k_j}}) - f(x, u_{n_{k_j}}) + f(x, u)|^{(s+1)/s}\, dx \right)^{s/(s+1)} \|h\|.$$

Since

$$|f(x, u_{n_{k_j}} - u) - f(x, u_{n_{k_j}}) + f(x, u)|^{(s+1)/s} \leqslant$$

$$\leqslant c_6(|v|^{(s+1)/s} + |v|^{s+1} + |u|^{(s+1)/s} + |u|^{s+1}) \in L^1(B_R(0))$$

we can use the dominated convergence theorem to get that

$$\int_{B_R(0)} |f(x, u_{n_{k_j}} - u) - f(x, u_{n_{k_j}}) + f(x, u)|^{(s+1)/s} dx = o(1).$$

Considering that this can be done for any subsequence of $\{u_n\}$ we actually get that

$$\sup_{\|h\|=1} \int_{B_R(0)} |f(x, u_n - u) - f(x, u_n) + f(x, u)|\,|h|\,dx = o(1).$$

For the second addendum we note first, by the choice of R,

$$\int_{|x|>R} f(x, u)\,h\,dx \leqslant c_6 \int_{|x|>R} |u|\,|h| + |u|^s|h|\,dx \leqslant c_7(\varepsilon + \varepsilon^s)\|h\|.$$

Since $f_x(x, 0) = 0$ we also infer that

$$\int_{|x|>R} |f(x, u_n - u) - f(x, u_n)|\,|h|\,dx \leqslant$$

$$\leqslant c_8 \int_{|x|>R} (1 + |u_n - u|^{s-1} + |u_n|^{s-1})|u|\,|h|\,dx \leqslant$$

$$\leqslant c_9 \left(\|u\|_{|x|>R} + \left(\int_{|x|>R} (|u_n - u|^{s-1}|u|)^{(s+1)/s} dx \right)^{s/(s+1)} + \right.$$

$$\left. + \left(\int_{|x|>R} (|u_n|^{s-1}|u|)^{(s+1)/s} dx \right)^{s/(s+1)} \right) \|h\| \leqslant$$

$$\leqslant c_9 \left(\|u\|_{|x|>R} + \left(\int_{|x|>R} |u_n - u|^{s+1} dx \right)^{(s-1)/(s+1)} \|u\|_{|x|>R} + \right.$$

$$\left. + \left(\int_{|x|>R} |u_n|^{s+1} dx \right)^{(s-1)/(s+1)} \|u\|_{|x|>R} \right) \|h\| \leqslant c_{10}\varepsilon\|h\|.$$

The proof that $\varphi(u_n - u) \to b - \varphi(u)$ is analogous. ■

LEMMA 2.6. *Let $\{u_n\}$ be a bounded sequence in X and suppose that there exists $\varrho > 0$ such that $\sup_{y \in \mathbb{R}^m} \int_{B_\varrho(y)} |u_n|^2 dx \to 0$ as $n \to \infty$. Then $u_n \to 0$ in $L^q(\mathbb{R}^n)$ for any $q \in (2, 2^*)$ as $n \to \infty$.*

PROOF. We give for completeness the proof given in [10] (see Lemma 2.18).

Given $q \in (2, 2^*)$ let $\theta \in (0, 1)$ be such that $q = 2 + \theta(2^* - 2)$. By the Holder inequality

$$\int_{B_\varrho(y)} |u|^q dx = \int_{B_\varrho(y)} |u|^{2(1-\theta)} |u|^{2^*\theta} dx \leqslant$$

$$\leqslant \left(\int_{B_\varrho(y)} |u|^2 dx \right)^{1-\theta} \left(\int_{B_\varrho(y)} |u|^{2^*} dx \right)^{\theta},$$

$\forall u \in X$. So

$$\|u\|_{L^q(B_\varrho(y))} \leqslant \|u\|^{(2/q)(1-\theta)}_{L^2(B_\varrho(y))} \|u\|^{(2^*/q)\theta}_{L^{2^*}(B_\varrho(y))} = \|u\|^{1-\bar{\alpha}}_{L^2(B_\varrho(y))} \|u\|^{\bar{\alpha}}_{L^{2^*}(B_\varrho(y))}$$

where

$$\alpha = \frac{2^*}{q}\theta = \frac{2^*}{q}\frac{(q-2)(m-2)}{4} = \frac{q-2}{q}m.$$

By the Sobolev Immersion theorem there exists $A_1 = A_1(q, m, \varrho)$ such that

$$\|u\|_{L^q(B_\varrho(y))} \leqslant A_1 \|u\|^{1-\bar{\alpha}}_{L^2(B_\varrho(y))} \|u\|^{\bar{\alpha}}_{B_\varrho(y)} \qquad \forall u \in X, \ \forall y \in \mathbb{R}^m.$$

Assume now that $\alpha q \geqslant 2$ that is $q \geqslant 4/m + 2$. We have

$$\int_{B_\varrho(y)} |u|^q dx \leqslant A_1^q \|u\|^{q(1-\alpha)}_{L^2((B_\varrho(y))} \|u\|^{\alpha q - 2} \int_{B_\varrho(y)} |\nabla u|^2 + |u|^2 dx.$$

Choosing a family of balls $\{B_\varrho(y_i)\}_{i \in \mathbb{N}}$ such that each point of $\mathbb{R}^m$ is contained in at least one and at most k of such balls, summing over this

family, we obtain

$$\|u\|_{L^q(\mathbb{R}^m)} \leqslant$$

$$\leqslant \sum_i \|u\|^q_{L^q(B_\varrho(y_i))} \leqslant A_1^q \|u\|^{\alpha q - 2} \sup_{y \in \mathbb{R}^m} \left(\int_{B_\varrho(y)} |u|^2 dx \right)^{q(1-\alpha)} \sum_i \|u\|^2_{B_\varrho(y_i)} \leqslant$$

$$\leqslant k A_1^q \|u\|^{\alpha q} \sup_{y \in \mathbb{R}^m} \left(\int_{B_\varrho(y)} |u|^2 dx \right)^{q(1-\alpha)} \qquad \forall u \in X.$$

Setting in the above formula $u = u_n$ we get $u_n \to 0$ in $L^q(\mathbb{R}^m)$ for all $q \in [4/m + 2, 2^*)$. The proof now go on with another interpolation inequality.

If $q \in (2, \bar{q})$ $(\bar{q} = 4/m + 2)$, we have $q = 2\theta + \bar{q}(1 - \theta)$ for some $\theta \in (0, 1)$. By the Holder inequality we have $\|u_n\|^q_{L^q(\mathbb{R}^m)} \leqslant \leqslant \|u_n\|^{2\theta}_{L^2(\mathbb{R}^m)} \|u_n\|^{\bar{q}(1-\theta)}_{L^{\bar{q}}(\mathbb{R}^m)}$, for any natural n. The lemma follows from the fact that $u_n \to 0$ in $L^{\bar{q}}(\mathbb{R}^m)$. ∎

LEMMA 2.7. *Let $u_n \to 0$ weakly in X such that $\varphi'(u_n) \to 0$. Then, for any $R > 0$ we have $\|u_n\|_{|x| < R} \to 0$ and the sequence $\{u_n\}$ verifies either*

$$a)\ u_n \to 0 \quad \text{or} \quad b)\ \exists \varrho, \eta > 0,\ \{y_n\} \subset \mathbb{R}^m / \limsup_{n \to \infty} \|u_n\|^2_{L^2(B_\varrho(y_n))} \geqslant \eta .$$

PROOF. Let $R > 0$ and let $g_R \in C^\infty(\mathbb{R}^m, \mathbb{R})$ be such that $g_R(x) \geqslant 0$ for any $x \in \mathbb{R}^m$, $g_R(x) = 1$ if $x \in B_R(0)$, $\operatorname{supp} g_R \subset B_{2R}(0)$. Clearly $\|u_n\|^2_{B_R(0)} = \langle u_n, g_R u_n \rangle - \langle u_n, g_R u_n \rangle_{|x| \geqslant R}$. We prove first that $\langle u_n, g_R u_n \rangle \to 0$.

Since $\langle u_n, g_R u_n \rangle = \varphi'(u_n) g_R u_n + \int_{\mathbb{R}^m} f(x, u_n) g_R u_n dx$ and since $\|g_R u_n\| \leqslant c_{10}$, it's enough to show that $\int_{\mathbb{R}^m} f(x, u_n) g_R u_n dx \to 0$. But this is a consequence of the Lebesgue dominated convergence theorem since $\operatorname{supp} g_R \subset B_{2R}(0)$.

Therefore we have

$$\|u_n\|^2_{B_R(0)} = o(1) - \int_{|x| \geqslant R} \nabla g_R \nabla u_n u_n dx - \int_{|x| \geqslant R} g_R (|\nabla u_n|^2 + u_n^2) dx .$$

Since $\int_{|x| \geqslant R} g_R (|\nabla u_n|^2 + u_n^2) dx \geqslant 0$, to prove that $\|u_n\|_{B_R(0)} \to 0$ it is suf-

ficient to show that $\int_{|x| \geq R} \nabla g_R \nabla u_n u_n dx \to 0$. But this is true because

$$\left| \int_{|x| \geq R} \nabla g_R \nabla u_n u_n dx \right| \leq$$

$$\leq \left(\int_{R \leq |x| \leq 2R} |\nabla g_R \nabla u_n|^2 dx \right)^{1/2} \left(\int_{R \leq |x| \leq 2R} |u_n|^2 dx \right)^{1/2}$$

and because $u_n \to 0$ in $L^2(B_{2R}(0) \setminus B_R(0))$.

The first part of the lemma is so proved. It's easy to prove the alternative.

Assume that (b) does not hold. In that case $\lim_{n \to \infty} \sup_{y \in \mathbb{R}^m} \|u\|_{B_\varrho(y)} = 0$ for any $\varrho > 0$. Since $\{u_n\}$ is bounded, by Lemma 2.6 we get that $u_n \to 0$ in $L^{s+1}(\mathbb{R}^m)$. From this, since $\int f(x, u_n) u_n \, dx \leq \varepsilon \|u_n\|^2_{L^2(\mathbb{R}^m)} +$ $+ A_\varepsilon \|u_n\|^{s+1}_{L^{s+1}(\mathbb{R}^m)}$, we get $\int_{\mathbb{R}^m} f(x, u_n) u_n dx \to 0$. Therefore $\|u_n\|^2 =$ $= \varphi'(u_n) u_n + \int_{\mathbb{R}^m} f(x, u_n) u_n dx \to 0$ as we claimed. ■

Therefore if u_n is a Palais-Smale sequence which converges weakly to a certain point u, then u_n converges to u in $H^1_{\text{loc}}(\mathbb{R}^m)$. Moreover if u_n does not converge to u in X, then fixed any $R > 0$ we have $\limsup_{n \to \infty} \|u_n\|_{|x| > R} \geq r > 0$. This mass r cannot be smaller than a certain positive fixed value as the next lemma says.

LEMMA 2.8. *Let $u_n \to u$ weakly in X, $\varphi'(u_n) \to 0$. If there exists $R > 0$ such that $\limsup_{n \to \infty} \|u_n\|_{|x| > R} \leq \varrho$ then $u_n \to u$ in X.*

PROOF. Fix $T > 0$ such that $\|u\|_{|x| \geq T} \leq \varrho/2$. Putting $M =$ $= \max\{R, T\}$ we have by Lemmas 2.5, 2.7, that $\|u_n - u\|_{|x| \leq M} \to 0$. Therefore

$$\|u_n - u\|^2 = o(1) + \|u_n - u\|^2_{|x| > M} =$$

$$= o(1) + \frac{\varrho^2}{4} + \varrho \|u_n\|_{|x| > M} + \|u_n\|^2_{|x| > M}$$

from which we get $\limsup \|u_n - u\| < 2\varrho$. Since $\varphi'(u_n - u) \to 0$ we derive from (2.4) that $u_n \to u$. ■

This is a first local compactness property of the functional which will be useful in the following. From it we derive easily

LEMMA 2.9. *If* $\operatorname{diam}\{u_n\} < \varrho$ *and* $\varphi'(u_n) \to 0$ *then* $\{u_n\}$ *has an accumulation point.*

PROOF. Let $\operatorname{diam}\{u_n\} = \varrho_0$ and $T > 0$ such that $\|u_1\|_{|x|>T} \leq \varrho - \varrho_0$. In that case $\|u_n\|_{|x|>T} \leq \|u_n - u_1\|_{|x|>T} + \varrho - \varrho_0 \leq \varrho$. Since $\{u_n\}$ is bounded it has a subsequence $\{u_{n_k}\}$ which converges weakly in X to a certain point u. Then, by Lemma 2.8, $u_{n_k} \to u$. ■

3. – The periodic case.

Here we will study some properties of the functional φ_∞. Obviously all the results given in the previous section remain valid for φ_∞. First of all we see how the further hypothesis (f4) implies that the functional φ_∞ satisfies the geometrical hypotheses of the mountain pass theorem.

By Lemma 2.3 we just know that there exists $r > 0$ such that $\varphi_\infty(u) \geq (1/4) r^2$ for any $u \in \partial\mathcal{B}_r(0)$. Then we note that the assumption (f4) gives information about the behavior of F_∞ at infinity with respect to z. In fact, one can infer that given $z_1 \neq 0$ then if $z/z_1 \geq 1$ we have

$$F_\infty(x, z) \geq \left[F_\infty(x, z_1) - \frac{\alpha}{\beta - 2} z_1^2\right] \left|\frac{z}{z_1}\right|^\beta + \frac{\alpha}{\beta - 2} z^2 \tag{3.1}$$

$$\forall x \in \mathbb{R}^m, \quad \frac{z}{z_1} \geq 1.$$

LEMMA 3.2. *There exists* $u_1 \in E$ *such that* $\varphi_\infty(u_1) < 0$.

PROOF. Let $(x_0, z_0) \in \mathbb{R}^m \times \mathbb{R}$ be given by (f4), then $\delta_0 = F_\infty(x_0, z_0) - (\alpha/(\beta - 2)) z_0^2 > 0$. By continuity there exists $\varepsilon > 0$ such that $F_\infty(x, z_0) - (\alpha/(\beta - 2)) z_0^2 \geq (1/2)\delta_0$ for any $x \in B_\varepsilon(z_0)$. Chosen $\varrho \in C^\infty(\mathbb{R}^m, \mathbb{R}^+)$ with $\operatorname{supp}\varrho \subset B_\varepsilon(x_0)$, we define $u_0(t) = z_0 \varrho(t)$. Then $\varphi_\infty(\lambda u_0) = (\lambda^2/2)\|u_0\|^2 - \int_{A_\lambda} F_\infty(x, \lambda u_0)\,dx - \int_{B_\lambda} F_\infty(x, \lambda u_0)\,dx$ where $A_\lambda = \{x: \lambda\varrho(x) < 1\}$ and $B_\lambda = \mathbb{R}^m \setminus A_\lambda$. Then $\int_{A_\lambda} |F_\infty(t, \lambda u_0)| \leq$

$\leqslant |B_\varepsilon(x_0)| \max\{|F_\infty(x, z)| : x \in \mathbb{R}^m,\ |z| \leqslant |z_0|\}$, whereas, by (3.1),

$$\int_{B_\lambda} F_\infty(x, \lambda u_0)\,dx \geqslant \lambda^\beta \int_{B_\lambda} \left[F_\infty(x, z_0) - \frac{\alpha}{\beta - 2} z_0^2\right] |\varrho|^\beta\,dx +$$

$$+\lambda^2 \frac{\alpha}{\beta - 2} \int_{B_\lambda} u_0^2\,dx \geqslant \frac{1}{2}\delta_0 \|\varrho\|^\beta_{L^\beta(B_\lambda)} \lambda^\beta .$$

Therefore $\varphi_\infty(\lambda u_0) \to -\infty$ as $\lambda \to +\infty$ and the thesis follows. ∎

This shows that the functional at infinity verifies the geometrical hypotheses of the mountain pass theorem. Then, if we define $\Gamma = \{\gamma \in \in C([0, 1], E)\colon \gamma(0) = 0,\ \varphi_\infty(\gamma(1)) < 0\}$ and $c = \inf_{\gamma \in \Gamma} \max_{s \in [0, 1]} \varphi_\infty(\gamma(s))$, we infer that c is a positive, asymptoticall critical value for φ_∞.

Now we use again (f4) to show that the Palais-Smale sequences of φ_∞ are in fact bounded sequences in X.

LEMMA 3.3. *If* $\{u_n\} \subset X$ *is such that* $\varphi'_\infty(u_n) \to 0$ *and* $\limsup \varphi_\infty(u_n) < +\infty$, *then* $\{u_n\}$ *is bounded in* E *and* $\liminf \varphi(u_n) \geqslant \geqslant 0$. *In particular any Palais-Smale sequence for* φ_∞ *is bounded in* E.

PROOF. From (f4), we easily get that

$$\left(\frac{1}{2} - \frac{1}{\beta} - \frac{\alpha}{\beta}\right)\|u\|^2 - \frac{1}{\beta}\|\varphi'_\infty(u)\|\|u\| \leqslant \varphi_\infty(u) \qquad \forall u \in X. \tag{3.4}$$

Now, given a sequence $\{u_n\} \subset X$ such that $\varphi'_\infty(u_n) \to 0$ and $\limsup \varphi_\infty(u_n) < +\infty$, from (3.4) we obtain $\|u_n\| \leqslant C$ for all $n \in \mathbb{N}$, C being a positive constant. Consequently we have that $\varphi_\infty(u_n) \geqslant \geqslant -C\|\varphi'_\infty(u_n)\|$ and this implies that $\liminf \varphi_\infty(u_n) \geqslant 0$. ∎

Using the periodicity we can now prove that the problem at infinity always admits a non zero solution which is obtained as weak limit of a suitable translated of the Palais-Smale sequence given by the mountain pass theorem.

THEOREM 3.5. *The problem*:

$$-\Delta u + u = f_\infty(x, u), \qquad u \in H^1(\mathbb{R}^m), \tag{P_∞}$$

admits a non zero solution.

PROOF. Let $\{u_n\}$ be the Palais-Smale sequence given by the mountain pass theorem. By Lemma 3.3 we can assume that $u_n \to u$ weakly in

X. If $u \neq 0$ then by Lemma 2.5 the theorem is proved. Assume $u_n \to 0$ weakly in X. Since $\varphi_\infty(u_n) \to c > 0$ it cannot be $u_n \to 0$ so the alternative (b) of Lemma 2.7 holds and (up to a subsequence) $\exists \alpha_1\, \alpha_2 > 0$, $\{y_n\} \subset \subset \mathbb{R}^m$ for which $\lim_{n\to\infty} \|u_n\|^2_{L^2(B_{\alpha_1}(y_n))} \geq \alpha_2$. Define $v_n = u_n(\cdot - [y_n])$ where $[y_n] = ([y_{n,1}], \ldots, [y_{n,m}])$. Then $\varphi_\infty(v_n) \to c$, $\varphi'_\infty(v_n) \to 0$ and $\|v_n\| \leq C$. Let $v_n \to v$ (up to a subsequence) weakly in X. By Lemma 2.7 we have $\|v_n - v\|_{L^2(B_{\alpha_1+1}(0))} \to 0$ therefore $\|v\|^2_{L^2(B_{\alpha_1+1}(0))} \geq \alpha_2 > 0$. So v is a non zero critical point of φ_∞. ■

Therefore $K_\infty = \{v \in X\backslash\{0\}/\varphi'_\infty(v) = 0\} \neq \emptyset$ and using 2.4 we have also that

$$\inf_{v\in K_\infty} \|v\| = \lambda > 0\,. \tag{3.6}$$

As we have point out in the introduction, the fact that the set of critical points of the functional at infinity is not empty does not guarantee that the problem (P) has non trivial solutions. We will prove that if the set of the critical points of the functional at infinity is numerable then this forces the functional φ itself to have infinitely many critical points.

To study better the Palais-Smale sequences, following [23], [7], we introduce two sets of real numbers. Letting

$$\mathcal{S}^b_{PS} = \{(u_n) \subset E\colon \lim \varphi'_\infty(u_n) = 0,\ \limsup \varphi_\infty(u_n) \leq b\}$$

we define

$$\Phi^b = \{l \in \mathbb{R}\colon \exists (u_n) \in \mathcal{S}^b_{PS} \text{ s.t. } \varphi_\infty(u_n) \to l\}$$

the set of the asymptotic critical values lower than b and

$$D^b = \{r \in \mathbb{R}\colon \exists (u_n),\ (\overline{u}_n) \in \mathcal{S}^b_{PS} \text{ s.t. } \|u_n - \overline{u}_n\| \to r\}$$

the set of the asymptotic distances between two Palais-Smale sequences under b.

The sets Φ^b and D^b are actually closed subsets of $\mathbb{R}$ (see [7]) and we have:

(3.7) given $b > 0$, for any $r \in \mathbb{R}^+\backslash D^b$ there exists $d_r > 0$ such that $[r - 3d_r, r + 3d_r] \subset \mathbb{R}^+\backslash D^b$ and there exists $\mu_r > 0$ such that $|\varphi'_\infty(u)| \geq \mu_r$ for any $u \in \mathcal{A}_{r-3d_r, r+3d_r}(K^b_\infty) \cap \{\varphi_\infty \leq b\}$,

(3.8) given $b > 0$, for any $l \in (0, b)\backslash\Phi^b$ there exists $\delta > 0$ such that $[l - \delta, l + \delta] \subset (0, b)\backslash\Phi^b$ and there exists $\nu > 0$ such that $|\varphi'_\infty(u)| \geq \nu$ for any $u \in \{l - \delta \leq \varphi_\infty \leq l + \delta\}$,

where, if $S \subset X$ and $0 \leq r_1 \leq r_2$, $\mathcal{A}_{r_1, r_2}(S) = \bigcup_{x\in S} B_{r_2}(x)\backslash B_{r_1}(x)$.

Using Lemmas 2.5, 2.7, 3.3 and (3.6) together with the periodicity assumption it is possible to characterize the Palais-Smale sequences of φ_∞. As in [10] we prove the following.

LEMMA 3.9. *Let $\{u_n\} \subset X$ be such that $\varphi_\infty(u_n) \to b$ and $\varphi'_\infty(u_n) \to 0$. Then there are $v_0 \in K_\infty \cup \{0\}, v_1, \ldots, v_k \in K_\infty$, a subsequence of $\{u_n\}$, denoted again $\{u_n\}$, and corresponding sequences $\{y_n^1\}, \ldots, \{y_n^k\} \in \mathbb{Z}^m$ such that, as $n \to \infty$:*

$$\|u_n - [v_0 + v_1(\cdot - y_n^1) + \ldots + v_k(\cdot - y_n^k)]\| \to 0,$$

$$\varphi_\infty(v_0) + \ldots + \varphi_\infty(v_k) = b,$$

$$|y_n^j| \to +\infty \qquad (j = 1, \ldots, k),$$

$$|y_n^i - y_n^j| \to +\infty \qquad (i \neq j).$$

PROOF. By Lemma 3.3 $\{u_n\}$ is a bounded sequence in X and we can assume that $\exists \lim_{n\to\infty} \|u_n\| \in \mathbb{R}$ and that u_n converges weakly to some v_0 in X. If $u_n \to v_0$ the lemma follows. Otherwise the case (b) of the alternative in Lemma 2.7 holds for the sequence $u_n - v_0$: $\exists \varrho, \eta > 0$, $\{y_n\} \subset \mathbb{Z}^m$, such that, up to a subsequence, $\lim_{n\to\infty} \|u_n - v_0\|^2_{L^2(B_\varrho(y_n))} \geq \eta$. Putting $u_n^1 = (u_n - v_0)(\cdot + y_n)$ we observe that there exists $v_1 \in K_\infty$ such that $u_n^1 \to v_1$ weakly in X, $\lim_{n\to\infty} \|u_n - v_0 - v_1(\cdot - y_n^1)\|^2 = \lim_{n\to\infty} \|u_n\|^2 - \|v_0\|^2 - \|v_1\|^2$. If $\lim_{n\to\infty} \|u_n\|^2 = \|v_0\|^2 + \|v_1\|^2$ the lemma follows because in that case $0 = \lim_{n\to\infty} \varphi_\infty(u_n - v_0 - v_1(\cdot - y_n^1)) = \lim_{n\to\infty} \varphi_\infty(u_n - v_0) - \varphi_\infty(v_1) = b - \varphi_\infty(v_0) - \varphi_\infty(v_1)$. If $\lim_{n\in\mathbb{N}} \|u_n\|^2 > \|v_0\|^2 + \|v_1\|^2$ we have that the sequence $\{u_n - v_0 - v_1(\cdot - y_n^1)\}$ verifies the case (b) of the alternative in Lemma 2.7 and we can continue as above. After a number of steps not greater than $\lim_{n\to\infty} \|u_n\| / \inf \sqrt{K_\infty}\|u\|$, the lemma follows. ■

This characterization reflects on the structure of the sets Φ^b and D^b. In fact, as in [7] (see Lemma 3.10), one can prove now that

$$\Phi^b = \{\textstyle\sum \varphi_\infty(v_i)\colon\ v_i \in K_\infty\} \cap [0, b],$$

$$D^b = \left\{ \left(\sum_{j=1}^{k} \|v_j - \bar{v}_j\|^2 \right)^{1/2} : \right.$$

$$\left. k \in \mathbb{N},\ v_i, \bar{v}_i \in K_\infty \cup \{0\},\ \sum_1^k \varphi_\infty(v_i) \leq b, \sum_1^k \varphi_\infty(\bar{v}_i) \leq b \right\}.$$

Using the hypothesis (*) given in the introduction this permits us to bound from below the norm of the gradient φ'_∞ in large regions of X. In fact if we assume

(*) *there exists* $c^* > c$ *such that* $K_\infty^{c^*}$ *is countable*

then both the sets D^{c^*} and Φ^{c^*} are countable too. Being D^{c^*} and Φ^{c^*} also closed we then have that

(3.10) D^{c^*} does not contain any neighborhood of 0 in $\mathbb{R}^+$,

(3.11) $]0, c^*[\setminus \Phi^{c^*}$ is open and dense in $[0, c^*]$.

By (3.7) and (3.10) we have that around $K_\infty^{c^*}$ there is a sequence of annuli of radii smaller and smaller on which there are not Palais-Smale sequences at a level less then or equal to c^*. Analogously, by (3.8) and (3.11), fixed any $\varepsilon \in (0, c^* - c)$ there exist two closed intervals $[a_1, b_1] \subset (c - \varepsilon, c)$ and $[a_2, b_2] \subset (c, c + \varepsilon)$ such that the sets $\{a_1 \leqslant \leqslant \varphi_\infty \leqslant b_1\}$ and $\{a_2 \leqslant \varphi_\infty \leqslant b_2 b\}$ do not contain Palais-Smale sequences.

Using (3.7), (3.10) and the local compactness property given by Lemma 2.9 it is possible to show as in [7] that the functional φ_∞ admits a local mountain pass type critical point (see [19]).

DEFINITION 3.12. Let f be a functional of class C^1 on a Banach space X and let Ω be a nonempty open subset of X.

We say that two points $x_0, x_1 \in \Omega$ are connectible in $\Omega \cap \{f < c\}$ if there is a path $p \in C([0, 1], X)$ joining x_0 and x_1, with range $p \subset \Omega$ and such that $\max_p f < c$.

A critical point $\bar{x} \in X$ for f is called of *local mountain pass-type* for f on Ω if $\bar{x} \in \Omega$ and for any neighborhood $\mathcal{N}$ of $\bar{x}$ subset of Ω the set $\{f < f(\bar{x})\} \cap \mathcal{N}$ contains two points not connectible in $\Omega \cap \{f < f(\bar{x})\}$.

We refer to Section 4 of [7] for the proof of the following

LEMMA 3.13. *If* φ_∞ *verifies* (*) *then it admits a non zero critical point of mountain pass type. In particular there exist* $\bar{c} \in [c, c^*)$ *and* $\bar{r} \in (0, \varrho)$ *such that for any sequence* $(r_n) \subset \mathbb{R}_+ \setminus D^*$, $r_n \to 0$ *there is a sequence* $(v_n) \subset K_\infty(\bar{c})$, $v_n \to \bar{v} \in K_\infty(\bar{c})$ *having this property: for any* $n \in \mathbb{N}$

and for any $h > 0$ *there is a path* $\gamma_n \in C([0, 1], X)$ *satisfying*:

(i) $\gamma_n(0),\ \gamma_n(1) \in \partial B_{r_n}(v_n)$;

(ii) $\gamma_n(0)$ *and* $\gamma_n(1)$ *are not connectible in* $B_{\bar{r}}(\bar{v}) \cap \{\varphi_\infty < \bar{c}\}$;

(iii) *range* $\gamma_n \subseteq \overline{B}_{r_n}(v_n) \cap \{\varphi_\infty \leq \bar{c} + h\}$;

(iv) *range* $\gamma_n \cap A_{r_n - (1/2)d_{r_n}, r_n}(v_n) \subseteq \{\varphi_\infty \leq \bar{c} - h_n\}$;

(v) $\operatorname{supp} \gamma_n(s) \subset [-R_n, R_n]$ *for any* $s \in [0, 1]$,

where $R_n > 0$ *is independent of* s, $h_n = (1/8)\, d_{r_n} \mu_{r_n}$ *and* d_{r_n} *and* μ_{r_n} *are defined by* (3.7).

REMARK 3.14. Clearly the property given in this sections are true for all the functionals φ_ι ($\iota = 1, \ldots, l$) if the assumptions $(*_\iota)$ are verified. In the following we write c_ι as the mountain pass level of φ_ι, $\bar{c}_\iota$ as the local mountain pass level near $\bar{v}_\iota$ etc.

4. – The construction of a pseudogradient vector field.

In this section we will study some consequences of the assumption (f7) with which we ask that there exist $A_1, \ldots, A_l \subset \mathbb{R}^m$, large at infinity on where f is asymptotic to l different periodic and superquadratic functions $f_1, \ldots, f_l$ as $|x| \to \infty$.

First of all we show that by (f7) if a function u is translated in a region where f and f_ι are close one to the other then $\varphi'(u)$ is near $\varphi'_\iota(u)$ (here $\varphi_\iota(u) = (1/2)\|u\|^2 - \int_{\mathbb{R}^m} F_\iota(x, u)\, dx$ is the functional associated to the function f_ι ($\iota = 1, \ldots, l$)).

LEMMA 4.1. *For any* $\delta > 0$ *and* $C > 0$ *there exists* $R > 0$ *such that if* $\|u\| \leq C$ *and* $\operatorname{supp} u \subset A_\iota \backslash B_R(0)$ *then* $\|\varphi'(u) - \varphi'_\iota(u)\| \leq \delta$.

PROOF. For any $\varepsilon > 0$ we choose $R > 0$ such that, if $\operatorname{supp} u \subset A_\iota \backslash B_R(0)$, we have $|f(x, u(x)) - f_\iota(x, u(x))| \leq \varepsilon(|u(x)| + |u(x)|^s)$ for almost every $x \in \mathbb{R}^m$. Then $|\varphi'(u)h - \varphi'_\iota(u)h| \leq \varepsilon \int_{A_\iota} |u|\,|h| + |u|^s |h|\, dx \leq \varepsilon c_{10} \|u\| \|h\|$ and the lemma follows. ∎

Given $k, N \in N$ we say that $p = (p_1, \ldots, p_k) \in P(k, N)$ if $p_j \in \mathbb{Z}^k$ for any j and $|p_j| \geq |p_{j-1}| + 4N^2 + 6N$ for $j \geq 2$.

If $p \in P(k, N)$ we define the annuli

$$\mathcal{U}_1 = \left\{ |x| \leqslant \frac{1}{2}(|p_1| + |p_2|) \right\},$$

$$\mathcal{U}_j = \left\{ \frac{1}{2}(|p_j| + |p_{j-1}|) \leqslant |x| \leqslant \frac{1}{2}(|p_j| + |p_{j+1}|) \right\} \quad (j = 2, \ldots, k)$$

and

$$M_i = \{ |p_i| + 2N(N+1) < |x| < |p_{i+1}| - 2N(N+1) \} \quad (i = 1, \ldots, k),$$

where $|p_{k+1}| = +\infty$. Since $p \in P(k, N)$ the thickness of the annulus M_i is alway greater than or equal to $2N$.

Given $r > 0$, $p \in P(k, N)$, $V = (V_1, \ldots, V_l) \in X^l$, $J = (j_1, \ldots, j_k) \in \in \{1, \ldots, l\}^k$ we define the set

$$\mathcal{B}_r(V; p; J) = \left\{ u \in X / \max_{i=1,\ldots,k} \|u - V_{j_i}(\cdot - p_i)\|_{\mathcal{U}_i} < r \right\}.$$

It is easy to see that if N is sufficiently large (depending on V and r) then $\mathcal{B}_r(V; p; J)$ is a nonempty open subset of X. Moreover the elements of $\mathcal{B}_r(V; p; J)$ are multibump functions of the mixed type. In fact if $u \in \mathcal{B}_r(V; p; J)$ then u is near the function $V_{j_i}(\cdot - p_i)$ on the annulus $\mathcal{U}_i$.

Defining

$$\varphi_{\iota, i}(u) = (1/2)\|u\|^2_{\mathcal{U}_i} - \int_{\mathcal{U}_i} F_\iota(x, u(x))\, dx,$$

$$u \in X, \quad (i = 1, \ldots, k), \qquad (\iota = 1, \ldots, l),$$

we investigate some properties of the functionals φ and $\varphi_{\iota, i}$ on the set $\mathcal{B}_r(V; p; J)$.

We note that for any given $V \in E^l$, $r > 0$, if $\tilde{N} = \tilde{N}(V, r) \in \mathbb{N}$ is such that $\|V_\iota\|_{|x| > \tilde{N}} \leqslant r$ for any $\iota \in \{1, \ldots, l\}$ then given $k \in \mathbb{N}$, $N > \tilde{N}$, $p \in \in P(k, N)$ and $J \in \{1, \ldots, l\}^k$ then $\forall u \in \mathcal{B}_r(V; p; J)$ and $\forall i \in \{1, \ldots, k\}$ there exists $j \in \{N+2, \ldots, 2N+1\}$ such that

$$\|u\|^2_{jN \leqslant ||x| - |p_i|| \leqslant (j+1)N} \leqslant \frac{4r^2}{N}. \tag{4.2}$$

Therefore if $u \in \mathcal{B}_r(V; p; J)$ then for any $i \in \{1, \ldots, k\}$, the annulus $\mathcal{U}_i$ contains two annular regions of thickness N, symmetric with respect to p_i, over which the norm of u is small as we want if N is sufficiently

large. Moreover, by construction, M_i never intersects any one of these annuli.

We call $j_{u,i}$ the smallest index in $\{N+2, \ldots, 2N+1\}$ which verifies (4.2).

For any $\varepsilon \in (0, r)$ there exists $N_\varepsilon \in \mathbb{N}$, $N_\varepsilon \geq \max\{\tilde{N}(V, r), 2\}$ such that

$$\max_{\iota=1,\ldots,l} \left\{ \|V_\iota\|^2_{|x|>N_\varepsilon}, \frac{4r^2}{N_\varepsilon} \right\} < \frac{\varepsilon}{2}.$$

So if $k \in \mathbb{N}$, $N > N_\varepsilon$ and $p \in P(k, N)$, then $\forall u \in \mathcal{B}_r(V; p; J)$ and $\forall i \in \in \{1, \ldots, k\}$ we get that

$$\|u\|^2_{j_{u,i}N \leq ||x| - |p_i|| \leq (j_{u,i}+1)N} < \frac{\varepsilon}{2}. \tag{4.3}$$

Then, fixed $u \in \mathcal{B}_r(V; p; J)$, we define the following subsets of $\mathbb{R}^m$:

$$E_{u,i} = \{|p_i| + (j_{u,i}+1)N \leq |x| \leq |p_{i+1}| - (j_{u,i+1}+1)N\} \qquad (i = 1, \ldots, k),$$

$$E_u = \bigcup_{i=1}^{k} E_{u,i},$$

$$\tilde{E}_{u,i} = \{x \in \mathbb{R}^m / \mathrm{dist}(x, E_{u,i}) \leq N\} \qquad (i = 1, \ldots, k),$$

$$\tilde{E}_u = \bigcup_{i=1}^{k} \tilde{E}_{u,i},$$

$$\mathcal{F}_{u,i} = \mathcal{U}_i \cap (\tilde{E}_u \setminus E_u) \qquad (i = 1, \ldots, k).$$

With this notation (4.3) can be rewritten in the form

$$\|u\|^2_{\mathcal{F}_{u,i}} \leq \frac{\varepsilon}{2} \qquad \forall u \in \mathcal{B}_r(V; p; J), \ \forall i \in \{1, \ldots, k\}. \tag{4.4}$$

We plainly recognize also that

$$\|u\|^2_{\tilde{E}_{u,i} \setminus E_{u,i}} \leq \varepsilon \qquad \forall u \in \mathcal{B}_r(V; p; J), \ \forall i \in \{1, \ldots, k\}. \tag{45}$$

By construction $M_i \subset E_{u,i}$, therefore the thickness of $E_{u,i}$ is greater than or equal to N $\forall i \in \{1, \ldots, k\}$, $\forall u \in \mathcal{B}_r(V; p; J)$. This is true also for the connected parts of the sets $\mathcal{F}_{u,i}$ and $\tilde{E}_{u,i} \setminus E_{u,i}$.

For $i \in \{1, \ldots, k\}$, we define the cut-off functions:

$$\beta_{u,i}(x) = \begin{cases} 1 & x \in E_{u,i}, \\ 0 & x \notin \tilde{E}_{u,i}, \end{cases}$$

with $\beta_{u,i}$ continuous on $\mathbb{R}^m$ and linear if restricted on the connected parts of $\tilde{E}_u \setminus E_u$ intersected with any straight line passing through the origin. We put also $\beta_{u,0} \equiv 0$.

Then, for $i \in \{1, \ldots, k\}$, we set:

$$\bar{\beta}_{u,i}(x) = \begin{cases} 0 & x \notin \mathcal{U}_i, \\ 1 - \beta_{u,i-1} - \beta_{u,i} & x \in \mathcal{U}_i. \end{cases}$$

If β is any one of the above cut-off functions then $|\nabla\beta(x)| \leqslant 1/N$, for a.e. $x \in \mathbb{R}^m$, therefore since $N \geqslant 2$, it is easy to see that, if A is measurable $\subset \mathbb{R}^m$ then $\|\beta u\|_A^2 \leqslant 2\|u\|_A^2$, $\forall u \in X$. Moreover if $u \in \mathcal{B}_r(V; p; J)$ and $i \in \{1, \ldots, k\}$, then by (4.5), we get

$$(4.6) \quad \langle u, \beta_{u,i} u\rangle = \|u\|_{E_{u,i}}^2 + \int_{\tilde{E}_{u,i} \setminus E_{u,i}} [\nabla\beta_{u,i} \nabla u u + \beta_{u,i}(|\nabla u|^2 + |u|^2)]\, dx \geqslant$$

$$\geqslant \|u\|_{E_{u,i}}^2 - \frac{1}{4}\|u\|_{\tilde{E}_{u,i} \setminus E_{u,i}}^2 \geqslant \|u\|_{E_{u,i}}^2 - \frac{1}{4}\varepsilon.$$

Now we define, for $i \in \{1, \ldots, k\}$, the functions

$$\chi_i(u) = \begin{cases} 1 & \|u\|_{E_{u,i}}^2 \geqslant \varepsilon, \\ \dfrac{1}{k} & \text{otherwise}, \end{cases}$$

and we set finally

$$W_u = \sum_{i=1}^{k} \chi_i(u) \beta_{u,i} u.$$

If we define the finite cone $C = \{y \in \mathbb{R}^m;\ |y| < 1/2,\ 1/4 < y_1 < 1/2\}$ then the embedding constant relative to the immersion $H^1(\Omega) \to L^{s+1}(\Omega)$ can be chosen to be independent of Ω if Ω is an open set of $\mathbb{R}^m$ which verifies the cone property with respect to C.

This implies, by (f2) (f3), that we can fix $r_0 \in (0, \min\{\bar{r}, \sqrt{2} - 1\})$

such that if $u, w \in X$ then

$$(4.7) \qquad \|u\|_A \leq r_0 \Rightarrow \int_A F(x, u)\,dx \leq \frac{1}{8}\|u\|_A^2$$

$$\text{and} \quad \int_A f(x, u)\,w\,dx \leq \frac{1}{8}\|u\|_A \|w\|_A$$

for any open set $A \subset \mathbb{R}^m$ which satisfies the cone property with respect to C. We can assume that r_0 is such that (4.7) holds also if we consider f_ι $(\iota = 1, \ldots, l)$ instead of f.

Using (4.6), (4.7), we can prove now that:

LEMMA 4.8. *Let* $r \in (0, (1/4)r_0)$ *and* $0 < \varepsilon < r^2$. *Then* $\forall u \in \in \mathcal{B}_r(V; p; J)$ *we have*

$$\varphi'(u)\,W_u \geq \frac{1}{2}\sum_{j=1}^{k} \chi_j(u)(\|u\|_{E_{u,j}}^2 - \varepsilon),$$

$$\varphi'_{\iota,i}(u)\,W_u \geq \frac{1}{2}\sum_{j=1}^{k} \chi_j(u)(\|u\|_{\mathcal{U}_i \cap E_{u,j}}^2 - \varepsilon).$$

PROOF. We have that $N > 2$, and the thickness of the annuli $E_{u,j}$ and of the ones whose union is $\bar{E}_{u,j} \setminus E_{u,j}$ is greater than or equal to N. Therefore these sets satisfy the cone property with respect to C. Moreover $\|u\|_{E_{u,j}} \leq 4r \leq r_0$ (in fact $\|u\|_{E_{u,j} \cap \mathcal{U}_i} \leq 2r$ $\forall i \in \{1, \ldots, k\}$) and $\|u\|_{\bar{E}_{u,j} \setminus E_{u,j}} \leq \varepsilon^{1/2} < r_0$. Therefore, by (4.6), and (4.7), we get

$$\varphi'(u)\,W_u \geq \sum_{j=1}^{k} \chi_j(u)\left(\|u\|_{E_{u,j}}^2 - \frac{\varepsilon}{4} - \right.$$

$$\left. - \int_{E_{u,j}} f(x, u)\,u\,dx - \int_{\bar{E}_{u,j} \setminus E_{u,j}} f(x, \beta_{u,i}u)\,\beta_{u,i}\,u\,dx\right) \geq$$

$$\geq \sum_{j=1}^{k} \chi_j(u)\left(\frac{7}{8}\|u\|_{E_{u,i}}^2 - \frac{1}{4}\varepsilon - \frac{1}{4}\varepsilon\right) \geq \frac{1}{2}\sum_{j=1}^{k} \chi_j(u)(\|u\|_{E_{u,j}}^2 - \varepsilon).$$

The computation is perfectly analogous for $\varphi_{\iota,i}$. ∎

By Lemma 4.8 we always have that

$$\varphi'(u)W_u \geq \frac{1}{2}\sum_{j=1}^{k}\chi_j(u)(\|u\|^2_{E_{u,j}} - \varepsilon) \geq$$

$$\geq \frac{1}{2}\sum_{\{j/\|u\|^2_{E_{u,j}} < \varepsilon\}}\chi_j(u)(\|u\|^2_{E_{u,j}} - \varepsilon) \geq -\frac{\varepsilon}{2}$$

and analogously

$$\varphi'_{\iota,i}(u)W_u \geq -\frac{\varepsilon}{2} \qquad \forall i \in \{1, \ldots, k\}$$

for all $u \in \mathcal{B}_r(V; p; J)$.

Moreover if $\|u\|_{\mathcal{U}_i \cap E_{u,j}}$ is greater then $2\varepsilon^{1/2}$, for a certain couple of index (i, j), then W_u indicates an increasing direction both for φ and $\varphi_{\iota,i}$. We note also that W_u has support in a region where each $V_{j_i}(\cdot - p_i)$ is small and it holds that $\langle W_u, u\rangle_{M_i} \geq (1/k)\|u\|^2_{M_i}$ for any $u \in \in B_r(V; p; J)$ and for any $i \in \{1, \ldots, k\}$.

Given $J = (j_1, \ldots, j_k) \in \{1, \ldots, l\}^k$, $R > 0$ we say that $p \in \in P_R(k, N, J)$ if $p \in P(k, N)$ and $B_{N(N+1)}(p_i) \subset A_{j_i} \setminus B_R(0)$ $(i = 1, \ldots, k)$.

Let b_ι be any nonzero critical level of φ_ι, and $r \in \in (0, (1/8)r_0) \setminus \bigcup_{\iota=1}^{l} D_\iota^{c_\iota^*}$, r_1, r_2, r_3 be such that $r - 3d_r < r_1 < r_2 < < r_3 < r + 3d_r < (1/4)r_0$ where $d_r = \min\{d_r^\iota;\ \iota = 1, \ldots, l\}$ $(\iota = 1, \ldots, l)$.

Let also $b_{-,\iota}$, $b_{+,\iota}$ and δ be such that $[b_{-,\iota} - \delta, b_{-,\iota} + 2\delta] \subset \subset]0, b_\iota[\setminus \Phi^{c_\iota^*}$ and $[b_{+,\iota} - \delta, b_{+,\iota} + 2\delta] \subset]b_\iota, c_\iota^*[\setminus \Phi^{c_\iota^*}$.

PROPOSITION 4.9. *There exists $\mu = \mu(r) > 0$ and $\varepsilon_1 = \varepsilon_1(r, b_{+,\iota}, b_{-,\iota}, \delta) > 0, R > 0$ such that: $\forall v_\iota \in \mathcal{K}_\iota(b_\iota)$ $(\iota = 1, \ldots, l)$, $\forall \varepsilon \in]0, \varepsilon_1[$ there exists $N \in \mathbb{N}$, such that, for any $k \in \mathbb{N}$, $J = (j_1, \ldots, j_k) \in \{1, \ldots, l\}^k$ and $p \in P_R(k, N, J)$, there exists a locally lipschitz continuous function $W: X \to X$ which verifies*

(W_0) $\|W(u)\|_{\mathcal{U}_j} \leq 2$ $\forall u \in X$, $j = 1, \ldots, k$, $\varphi'(u)W(u) \geq 0$ $\forall u \in X$, $W(u) = 0$ $\forall u \in X \setminus \mathcal{B}_{r_3}(V; p; J)$ $(V = (v_1, \ldots, v_l))$,

(W_1) $\varphi'_{j_i,i}(u)W(u) \geq \mu$ *if* $r_1 \leq \|u - v_{j_i}(\cdot - p_i)\|_{\mathcal{U}_i} \leq r_2$, $u \in \in \mathcal{B}_{r_2}(V; p; J) \cap \bigcap_{i=1}^{k} \varphi^{b_{+,j_i}+\delta}_{j_i,i}$,

(W_2) $\varphi'_{j_i,i}(u)W(u) \geq 0$ $\forall u \in (\varphi^{b_{+,j_i}+\delta}_{j_i,i} \setminus \varphi^{b_{+,j_i}}_{j_i,i}) \cup (\varphi^{b_{-,j_i}+\delta}_{j_i,i} \setminus \varphi^{b_{-,j_i}}_{j_i,i})$,

(W_3) $\langle u, W(u)\rangle_{M_j} \geq 0$ $\forall j \in \{1, \ldots, k\}$ *if* $\max\|u\|^2_{M_i} \geq 4\varepsilon$.

Moreover if $\mathcal{K} \cap \mathcal{B}_{r_2}(V; p; J) = \emptyset$ *then there exists* $\mu_k > 0$ *such that*

(W_4) $\quad \varphi'(u) W(u) \geq \mu_k \ \forall u \in \mathcal{B}_{r_2}(V; p; J).$

PROOF. Let $\tilde{r}_1 = r_1 - (1/2)(r_1 - r + 3d_r)$, $\tilde{r}_3 = r_3 + (1/2)(r + 3d_r - r_3)$ and let μ_r be given by 3.8.

Let also

$$\nu = \inf\{\|\varphi_\iota'(u)\| / u \in (\varphi_\iota^{b_{+,\iota}+2\delta} \setminus \varphi_\iota^{b_{+,\iota}-\delta}) \cup (\varphi_\iota^{b_{-,\iota}+2\delta} \setminus \varphi_\iota^{b_{-,\iota}-\delta}); \quad \iota = 1, \ldots, l\};$$

by Remark 3.9 we have that $\nu > 0$.

Let $C = 2 \sup\{\|u\|; u \in \mathcal{K}_\iota(b_\iota), \ \iota = 1, \ldots, l\} + r_0$; by Lemma 4.1 there exists $R > 0$ such that if $\|u\| \leq C$ and $\operatorname{supp} u \subset A_\iota \setminus B_R(0)$ then $\|\varphi_\iota'(u) - \varphi'(u)\| \leq (1/4) \min\{\nu, \mu_r\}$.

Let

$$\varepsilon_1^{1/2} = \min\left\{\frac{(r_1 - r + 3d_r)}{12}, \frac{(r + 3d_r - r_3)}{12}, \frac{\mu_r}{16}, \frac{\nu}{16}, \frac{\delta^{1/2}}{6}\right\}.$$

Let's fix $v_\iota \in \mathcal{K}_\iota(b_\iota)$ $(\iota = 1, \ldots, l)$, $\varepsilon \in (0, \varepsilon_1)$, $k \in \mathbb{N}$, $N > N_\varepsilon$, $J = (j_1, \ldots, j_k) \subset \{1, \ldots, l\}^k$ and $(p_1, \ldots, p_k) \in P_R(k, N, J)$.

We construct the vector field W_u on $\mathcal{B}_{r_3}(v; p; J)$, using Lemma 4.8 with $r = r_3$. We will now define another vector field analyzing the different cases.

Case 1) $\quad u \in (\mathcal{B}_{r_3}(V; p; J) \setminus \mathcal{B}_{r_1}(V; p; J)) \cap \bigcap_{i=1}^{k} \varphi_{j_i, i}^{b_{+,j_i}+3\delta/2}.$

We set $\mathfrak{I}_1(u) = \{i \in \{1, \ldots, k\} / \|u - v_{j_i}(\cdot - p_i)\|_{\mathcal{U}_i} \geq r_1\}$. Obviously $\mathfrak{I}_1(u) \neq \emptyset$.

Let $i \in \mathfrak{I}_1(u)$ and $\xi_1 = (1/2) \min\{r_1 - \tilde{r}_1, \tilde{r}_3 - r_3\}$.

We consider the two possible subcases:

$$\|u\|_{\mathcal{U}_i \cap E_u} \geq \xi_1 \quad \text{or} \quad \|u\|_{\mathcal{U}_i \cap E_u} < \xi_1.$$

In the first one, using Lemma 4.8 and the fact that $\varepsilon_1^{1/2} \leq \xi_1/3$, we get (putting $E_0 = \emptyset$)

$$(4.10) \quad \varphi'(u) W_u \geq \frac{1}{2}(\|u\|^2_{E_{u,i-1}} + \|u\|^2_{E_{u,i}} - 2\varepsilon) - \sum_{\{l / \|u\|^2_{E_{u,l}} < \varepsilon\}} \chi_l(u) \frac{\varepsilon}{2} \geq$$

$$\geq \frac{1}{2}(\|u\|^2_{\mathcal{U}_i \cap E_u} - 2\varepsilon) - \sum_{\{l / \|u\|^2_{E_{u,l}} < \varepsilon\}} \chi_l(u) \frac{\varepsilon}{2} \geq \frac{\xi_1^2}{2} - 2\varepsilon \geq \frac{\xi_1^2}{4},$$

and analogously

$$\varphi'_{j_i,i}(u)\,W_u \geqslant \frac{\xi_1^2}{2} - 2\varepsilon \underset{k}{\geqslant} \frac{\xi_1^2}{4}. \tag{4.11}$$

For all $u \in (\mathcal{B}_{r_3}(V; p; J) \setminus \mathcal{B}_{r_1}(V; p; J)) \cap \bigcap_{i=1} \varphi_{j_i,i}^{b_+,j_i+\delta}$ if $\|u\|_{\mathcal{U}_i \cap E_u} \geqslant \xi_1$ and $i \in \mathfrak{I}_1(u)$ we put $\mathcal{W}_{u,i} = 0$.

In the second subcase we firstly note that, arguing as in (4.3), there exists $j_u \in \{1, \ldots, N\}$ such that $\|u\|^2_{j_u N \leqslant |x-p_i| \leqslant (j_u+1)N} < \varepsilon/2$. We put $\tilde{B}_u = B_{(j_u+1)N}(p_i)$ and $B_u = B_{j_u N}(p_i)$ noting that $\operatorname{dist}(\tilde{B}_u, \tilde{E}_u) \geqslant N$ and therefore that the set $\mathcal{U}_i \setminus (\tilde{B}_u \cup \tilde{E}_u)$ has the cone property with respect to C.

We define also the cut-off function $\eta_u \in C(\mathrm{R}^m, \mathrm{R})$ such that $\eta_u(x) = 1$ if $x \in B_u$, $\eta_u(x) = 0$ if $x \notin \tilde{B}_u$, and in such a way η_u is linear if restricted on the connected parts of $\tilde{B}_u \setminus B_u$ intersected with any line passing through p_i.

Consider the following alternative:

i) $\|u\|_{\mathcal{U}_i \setminus (\tilde{B}_u \cup \tilde{E}_u)} \geqslant \xi_1/2$, or

ii) $\|u\|_{\mathcal{U}_i \setminus (\tilde{B}_u \cup \tilde{E}_u)} < \xi_1/2$.

If i) holds we put $\mathcal{W}_{u,i} = (1 - \eta_u)\,\bar{\beta}_{u,i} u$ and by (4.7) we get

$$\varphi'(u)\,\mathcal{W}_{u,i} \geqslant \|u\|^2_{\mathcal{U}_i \setminus (\tilde{B}_u \cup \tilde{E}_u)} - 3\varepsilon - \frac{1}{8}\|u\|^2_{\mathcal{U}_i \setminus (\tilde{B}_u \cup \tilde{E}_u)} \geqslant \frac{\xi_1^2}{8} - 3\varepsilon \geqslant \frac{\xi_1^2}{16}.$$

Analogously we get

$$\varphi'_{j_i,i}(u)\,\mathcal{W}_{u,i} \geqslant \frac{\xi_1^2}{16}.$$

We finally observe that

$$\min\{\varphi'_{j_i,i}(u)(\mathcal{W}_{u,i} + W_u),\ \varphi'(u)(\mathcal{W}_{u,i} + W_u)\} \geqslant \frac{\xi_1^2}{16} - \frac{\varepsilon}{2} \geqslant \frac{\xi_1^2}{32}. \tag{4.12}$$

If ii) holds we claim that $\eta_u u \in \mathcal{C}_{r-3d_r,\,r+3d_r}(v_{j_i}(\cdot - p_i)) \cap \varphi_{j_i}^{c_{j_i}^*}$.

In fact we firstly note that since $(5/4)\,\xi_1^2 - 3\varepsilon^{1/2}\xi_1 - 7\varepsilon \leqslant r_1^2 - (r - 3d_r)^2$ we have

$$\|\eta_u u - v_{j_i}(\cdot - p_i)\|^2 \geqslant \|u - v_{j_i}(\cdot - p_i)\|^2_{B_u} =$$

$$= \|\ldots\|^2_{\mathcal{U}_i} - \|\ldots\|^2_{E_u \cap \mathcal{U}_i} - \|\ldots\|^2_{\mathcal{F}_{u,i}} + - \|\ldots\|^2_{\mathcal{U}_i \setminus (\tilde{E}_u \cup \tilde{B}_u)} - \|\ldots\|^2_{\tilde{B}_u \setminus B_u} \geqslant$$

$$\geqslant r_1^2 - (\xi_1^2 + 2\varepsilon^{1/2}\xi_1 + \varepsilon) - 3\varepsilon - \left(\frac{\xi_1^2}{4} + \varepsilon^{1/2}\xi_1 + \varepsilon\right) - 3\varepsilon =$$

$$= r_1^2 - \frac{5}{4}\xi_1^2 - 3\varepsilon^{1/2}\xi_1 - 7\varepsilon \geqslant (r - 3d_r)^2.$$

On the other hand since $(3/2)\xi_1 + r_3 + 3\varepsilon^{1/2} \leqslant r + 3d_r$ we get

$$\|\eta_u u - v_{j_i}(\cdot - p_i)^2 \leqslant \|\eta_u u - v_{j_i}(\cdot - p_i)\|^2_{\mathcal{U}_i} + \varepsilon \leqslant$$

$$\leqslant (\|(1-\eta_u)u\|_{\mathcal{U}_i} + \|u - v_{j_i}(\cdot - p_i)\|_{\mathcal{U}_i})^2 + \varepsilon \leqslant$$

$$\leqslant (\|u\|_{\mathcal{U}_i \cap E_u} + \|u\|_{\mathcal{F}_{u,i}} + \|u\|_{\mathcal{U}_i \setminus (\bar{E}_u \cup \bar{B}_u)} + 2^{1/2}\|u\|_{\bar{B}_u \setminus B_u} + r_3)^2 + \varepsilon \leqslant$$

$$\leqslant \left(\frac{3}{2}\xi_1 + r_3 + 3\varepsilon^{1/2}\right)^2 \leqslant (r + 3d_r)^2 .$$

To end the proof of the claim we note that, since $\|u\|_{\mathcal{U}_i \setminus B_u} \leqslant (1/2) r_0$, by (4.7) we have that $(1/2)\|u\|^2_{\mathcal{U}_i \setminus B_u} - \int_{\mathcal{U}_i \setminus B_u} F_{j_i}(x, u)\, dx \geqslant 0$. Therefore $\varphi_{j_i}(\eta_u u) = \varphi_{j_i, i}(\eta_u u) \leqslant v_{j_i, i}(u) + \|u\|^2_{\bar{B}_u \setminus B_u} \leqslant \varphi_{j_i, i}(u) + \varepsilon \leqslant b_{+, j_i} + 2\delta < c^*_{j_i}$ as we claimed.

Therefore there exists $Z_{u,i} \in X$, $\|Z_{u,i}\| \leqslant 1$ such that

$$\varphi'_{j_i, i}(\eta_u u) Z_{u,i} = \varphi'_{j_i}(\eta_u u) Z_{u,i} \geqslant \frac{\mu_r}{2} .$$

Since $p \in P_R(k, N, J)$ we have $\operatorname{supp} \eta_u u \subset A_{j_i} \setminus B_R(0)$. Moreover $\|\eta_u u\| \leqslant 2^{1/2}\|u\|_{\mathcal{U}_i} \leqslant 2^{1/2}(\|v_{j_i}\| + r_0) \leqslant C$, which implies by the choice of R that

$$\varphi'(\eta_u u) Z_{u,i} = \varphi'_{j_i}(\eta_u u) Z_{u,i} + (\varphi'(\eta_u u) - \varphi'_{j_i}(\eta_u u)) Z_{u,i} \geqslant \frac{\mu_r}{4} .$$

As last step we note that

$$|\varphi'_{j_i, i}(\eta_u u) Z_{u,i} - \varphi'_{j_i, i}(u)\eta_u Z_{u,i}| =$$

$$= \Big| \langle \eta_u u, Z_{u,i}\rangle_{\bar{B}_u \setminus B_u} - \langle u, \eta_u Z_{u,i}\rangle_{\bar{B}_u \setminus B_u} -$$

$$- \int_{\bar{B}_u \setminus B_u} (f_{j_i}(x, \eta_u u) - f_{j_i}(x, u)\eta_u) Z_{u,i}\, dx \Big| \leqslant$$

$$\leqslant \frac{2}{N}\|u\|_{\bar{B}_u \setminus B_u} + \frac{1}{4}\|u\|_{\bar{B}_u \setminus B_u} \leqslant \varepsilon^{1/2} \leqslant \frac{\mu_r}{8}$$

and the same argument gives also

$$|\varphi'(\eta_u u) Z_{u,i} - \varphi'(u)\eta_u Z_{u,i}| \leqslant \frac{\mu_r}{8}.$$

From the two above inequality it follows that

$$\min\{\varphi'(u)\eta_u Z_{u,i}, \varphi'_{j_i,i}(u)\eta_u Z_{u,i}\} \geqslant \frac{\mu_r}{8}.$$

In this case we put $\mathcal{W}_{u,i} = (1/2)\eta_u Z_{u,i}$ observing that

$$\min\{\varphi'(u)(\mathcal{W}_{u,i} + W_u), \varphi'_{j_i,i}(u)(\mathcal{W}_{u,i} + W_u)\} \geqslant \frac{\mu_r}{16} - \frac{\varepsilon}{2} \geqslant \frac{\mu_r}{32}. \tag{4.13}$$

We now set $2\mu = \min\{(\mu_r/32, \xi_1^2/32\}$ and

$$\mathcal{V}_{u,1} = \begin{cases} W_u + \sum\limits_{i\in\mathfrak{I}_1(u)} \mathcal{W}_{u,i} & u \in (\mathcal{B}_{r_3}(V; p; J)\backslash\mathcal{B}_{r_1}(V; p; J)) \cap \bigcap\limits_{i=1}^{k} \varphi^{b_{+,i}+3\delta/2}_{j_i,i}, \\ 0 & \text{otherwise}, \end{cases}$$

obtaining by (4.10)-(4.13) that $\forall u \in (\mathcal{B}_{r_3}(V; p; J)\backslash\mathcal{B}_{r_1}(V; p; J)) \cap \cap \bigcap\limits_{i=1}^{k} \varphi^{b_{+,j_i}+3\delta/2}_{j_i,i}$

$$\begin{cases} \varphi'(u)\mathcal{V}_{u,1} \geqslant 2\mu, \\ \varphi'_{j_i,i}(u)\mathcal{V}_{u,1} \geqslant 2\mu \quad \forall i \in \mathfrak{I}_1(u), \\ \langle u, \mathcal{V}_{u,1}\rangle_{M_l} = \langle u, W_u\rangle_{M_l} \geqslant \frac{1}{k}\|u\|^2_{M_l} \quad l = 0, \ldots, k. \end{cases} \tag{4.14}$$

We also note that $\|\mathcal{V}_{u,1}\|_{\mathcal{U}_i} \leqslant \|\mathcal{W}_{u,i}\|_{\mathcal{U}_i} + \|W_u\|_{\mathcal{U}_i} \leqslant 1/\sqrt{2}(1 + r_0) < 1$ for any $i \in \{1, \ldots, k\}$.

Case 2) $u \in \mathcal{B}_{r_3}(V; p; J) \cap \left(\bigcup\limits_{i=1}^{k} (\varphi_{j_i,i})^{b_{+,j_i}+\delta}_{b_{+,j_i}}\right)$.

We put $\mathfrak{I}_2^+(u) = \{i \in \{1, \ldots, k\}/u \in (\varphi_{j_i,i})^{b_{+,j_i}+\delta}_{b_{+,j_i}}\}$ and fix $i \in \in \mathfrak{I}_2^+(u)$.

Fixing also $\xi_2^2 = \delta/4$ it can be either

$$\|u\|_{\mathcal{U}_i \cap E_u} \geqslant \xi_2 \quad \textit{or} \quad \|u\|_{\mathcal{U}_i \cap E_u} < \xi_2.$$

In the first subcase, considering that $\varepsilon_1 \leqslant \xi_2^2/9$, we get as above that

$$\varphi'(u)W_u \geqslant \frac{1}{2}\xi_2^2 - 2\varepsilon \geqslant \frac{1}{4}\xi_2^2 \quad \text{and} \quad \varphi'_{j_i,i}(u)W_u \geqslant \frac{1}{2}\xi_2^2 - 2\varepsilon \geqslant \frac{1}{4}\xi_2^2.$$

For all

$$u \in \mathcal{B}_{r_3}(V; p; J) \cap \left(\bigcup_{i=1}^{k} (\varphi_{j_i,i})_{b_+^{j_i}}^{b_+^{j_i}+\delta} \right)$$

and $i \in \mathfrak{I}_2(u)$, if $\|u\|_{\mathcal{U}_i \cap E_u} \geqslant \xi_2$ we put $\tilde{\mathcal{W}}_{u,i} = 0$.

In the second subcase we proceed as in the case 1 considering the alternative

i) $\|u\|_{\mathcal{U}_i \setminus (\bar{B}_u \cup \bar{E}_u)} \geqslant \xi_2/2$, or

ii) $\|u\|_{\mathcal{U}_i \setminus (\bar{B}_u \cup \bar{E}_u)} < \xi_2/2$.

If i) holds then putting $\mathcal{W}_{u,i} = (1 - \eta_u)\,\tilde{\beta}_{u,i}u$ arguing as in case 1 we get

$$\min\{\varphi'(u)(\mathcal{W}_{u,i} + W_u),\ \varphi'_{j_i}(u)(\mathcal{W}_{u,i} + W_u)\} \geqslant \frac{\xi_2^2}{32}. \tag{4.15}$$

If ii) holds then we claim that $\eta_u u \in (\varphi_{j_i})_{b_+^{j_i}-\delta}^{b_+^{j_i}+2\delta}$.

In fact

$$\|u\|_{\mathcal{U}_i}^2 - \|\eta_u u\|_{\mathcal{U}_i}^2 \leqslant \|u\|_{\mathcal{U}_i \cap E_u}^2 + \|u\|_{\mathcal{T}_{u,i}}^2 + \|u\|_{\mathcal{U}_i \setminus (\bar{E}_u \cup \bar{B}_u)}^2 +$$

$$+ \|u\|_{\bar{B}_u \setminus B_u}^2 + \|\eta_u u\|_{\bar{B}_u \setminus B_u}^2 \leqslant \frac{5}{4}\xi_2^2 + 2\varepsilon$$

and

$$\int_{\mathcal{U}_i} F_{j_i}(x, u) - F_{j_i}(x, \eta_u u)\,dx =$$

$$= \int_{\mathcal{U}_i \setminus \bar{B}_u} F_{j_i}(x, u)\,dx + \int_{\bar{B}_u \setminus B_u} F_{j_i}(x, u) - F_{j_i}(x, \eta_u u)\,dx \leqslant$$

$$\leqslant \frac{1}{8}\|u\|_{\mathcal{U}_i \setminus \bar{B}_u}^2 + \frac{1}{2}\|u\|_{\bar{B}_u \setminus B_u}^2 \leqslant \frac{1}{2}(\xi_2^2 + \varepsilon).$$

We finally derive that

$$|\varphi_{j_i,i}(u) - \varphi_{j_i,i}(\eta_u u)| \leqslant \xi_2^2 + \varepsilon < \delta$$

which implies $\eta_u u \in (\varphi_{j_i,i})^{b_{+,j_i}+2\delta}_{b_{+,j_i}-\delta}$ as we claimed. So there exists $Z_{u,i} \in X$, $\|Z_{u,i}\| \leqslant 1$ such that $\varphi'_{j_i}(\eta_u u) Z_{u,i} = \varphi'_{j_i,i}(\eta_u u) Z_{u,i} \geqslant \nu/2$.

As in case 1), since $\operatorname{supp} \eta_u u \subset A_{j_i} \setminus B_R(0)$, we get $\varphi'(\eta_u u) Z_{u,i} \geqslant \geqslant \nu/4$.

Moreover, again as in the case 1), since $\varepsilon^{1/2} < \nu/16$

$$\min\{\varphi'(u)\eta_u Z_{u,i}, \varphi'_{j_i,i}(u)\eta_u Z_{u,i}\} \geqslant \frac{\nu}{8}$$

and we put in this case $\tilde{\mathcal{W}}_{u,i} = (1/2)\eta_u Z_{u,i}$ noting that

$$\min\{\varphi'(u)(W_u + \tilde{\mathcal{W}}_{u,i}), \varphi'_{j_i,i}(u)(W_u + \tilde{\mathcal{W}}_{u,i})\} \geqslant \tag{4.16}$$

$$\geqslant \frac{\nu}{16} - \frac{\varepsilon}{2} \geqslant \frac{\nu}{32}.$$

Let now $\nu^+ = \min\{\nu/32, \xi_2^2/32\}$ and

$$\mathcal{V}_{u,2} = \begin{cases} W_u + \sum\limits_{i \in \mathfrak{I}_2^+(u)} \tilde{\mathcal{W}}_{u,i} & u \in \bigcup\limits_{i=1}^{k} (\varphi_{j_i,i})^{b_{+,j_i}+\delta}_{b_{+,j_i}} \cap \mathcal{B}_{r_3}(V; p; J), \\ 0 & \text{otherwise}, \end{cases}$$

obtaining, as in the case 1), that

$$\forall u \in \mathcal{B}_{r_3}(V; p; J) \cap \left(\bigcup_{i=1}^{k} (\varphi_{j_i,i})^{b_{+,j_i}+\delta}_{b_{+,j_i}} \right),$$

$$\begin{cases} \varphi'(u)\, \mathcal{V}_{u,2} \geqslant \nu^+, \\ \varphi'_{j_i,i}(u)\, \mathcal{V}_{u,2} \geqslant \nu^+ \qquad \forall i \in \mathfrak{I}_2^+(u), \\ \langle u, \mathcal{V}_{u,2} \rangle_{M_l} = \langle u, W_u \rangle_{M_l} \geqslant \dfrac{1}{k} \|u\|^2_{M_l} \qquad l \in \{0, \ldots, k\}. \end{cases} \tag{4.17}$$

As in the case 1) it is easy to prove that $\max\limits_i \|\mathcal{V}_{u,2}\|_{\mathcal{U}_i} \leqslant 1$.

Case 3) $u \in \mathcal{B}_{r_3}(V; p; J) \cap \left(\bigcup\limits_{i=1}^{k} (\varphi_{j_i,i})^{b_{-,j_i}+\delta}_{b_{-,j_i}} \right)$.

As in case 2) we put $\xi_3 = \delta/4$, $\nu^- = \min\{\nu/32, \xi_3^2/32\}$, and $\mathfrak{I}_2^-(u) = \{i \in \{1, \ldots, k\} / u \in (\varphi_{j_i,i})^{b_{-,j_i}+\delta}_{b_{-,j_i}}\}$, getting that $\forall u \in \mathcal{B}_{r_3}(V; p; J) \cap$

$\cap \left(\bigcup_{i=1}^{k} (\varphi_{j_i, i})_{b_-, j_i}^{b_-, j_i + \delta} \right)$ there exists $\nabla_{u,3} \in X$ such that $\max_i \|\nabla_{u,3}\|_{\mathcal{U}_i} \leq 1$ and

$$(4.18) \quad \begin{cases} \varphi'(u)\,\nabla_{u,3} \geq \nu^-, \\ \varphi'_{j_i,i}(u)\,\nabla_{u,3} \geq \nu^- \qquad \forall i \in \mathfrak{I}_2^-(u) \\ \langle u, \nabla_{u,3}\rangle_{M_l} = \langle u, W_u\rangle_{M_l} \geq \dfrac{1}{k}\|u\|_{M_l}^2 \qquad l \in \{1, \ldots, k\}. \end{cases}$$

We put $\nabla_{u,3} = 0$ if $u \notin \mathcal{B}_{r_3}(V; p; J) \cap \left(\bigcup_{i=1}^{k} (\varphi_{j_i, i})_{b_-, j_i}^{b_-, j_i + \delta} \right)$.

Case 4) $u \in \mathcal{B}_{r_1}(V; p; J)$.

In this case we distinguish between the two subcases:

$$\max_{1 \leq l \leq k} \|u\|_{M_l}^2 \geq 4\varepsilon \quad \text{or} \quad \max_{0 \leq l \leq k} \|u\|_{M_l}^2 < 4\varepsilon.$$

In the first case, if we have $\|u\|_{M_{\bar{l}}}^2 = \max_{0 \leq l \leq k} \|u\|_{M_l}^2 \geq 4\varepsilon$, we get using Lemma 4.8 that

$$\varphi'(u)\,W_u \geq \frac{1}{2}(\|u\|_{E_{u,\bar{l}}}^2 - \varepsilon) - \frac{1}{2}\varepsilon \geq \frac{1}{2}(\|u\|_{M_{\bar{l}}}^2 - \varepsilon) - \frac{1}{2}\varepsilon \geq \varepsilon$$

and we set $\nabla_{u,4} = W_u$.

In the second case, by the local compactness property of φ (Lemma 2.8), we obtain that if $K \cap \mathcal{B}_{r_1}(V; p; J) = \emptyset$ then there exists $V_u \in X$, $\|V_u\| \leq 1$ and there exists $\mu'_k > 0$, independent of u, such that $\varphi'(u)\,V_u \geq \mu'_k/2$. We set $\nabla_{u,4} = V_u$.

Let also $\nabla_{u,4} = 0$ if $u \notin \mathcal{B}_{r_1}(V; p; J)$.

We can conclude that if put $2\mu_k = \min\{\varepsilon, \mu'_k/2\}$ we have $\forall u \in \mathcal{B}_{r_1}(V; p; J)$ that $\varphi'(u)\,\nabla_{u,4} \geq 2\mu_k$ and if $\max_{j=1,\ldots,k} \|u\|_{M_j}^2 \geq 4\varepsilon$ then

$$(4.19) \qquad \langle u, \nabla_{u,4}\rangle_{M_l} = \langle u, W_u\rangle_{M_l} \geq \frac{1}{k}\|u\|_{M_l}^2 \qquad l \in \{0, \ldots, k\}.$$

For $u \in X$ we put $\nabla_u = \sum_{i=1}^{4} \nabla_{u,i}$ noting that $\max_i \|\nabla_u\|_{\mathcal{U}_i} \leq 2$. Then the proposition follows with a classical pseudogradient construction, by using a suitable partition of unity and suitable cutoff functions.

5. – Multiplicity result.

In this section we will state and prove the main Theorem.

THEOREM 5.1. *Assume that* (f1)-(f5), (f7) *and* $(*_\iota)$ $(\iota = 1, \dots, l)$ *hold. Let* v_ι *be the critical point of* φ_ι $(\iota = 1, \dots, l)$ *given by* 3.14. *Then for any* $r > 0$ *there is* $N \in \mathbb{N}$, $R > 0$, *such that for every* $k \in \mathbb{N}$ $J \in \{1, \dots, k\}^l$ *and* $p \in P_R(k, N, J)$ *we have* $\mathcal{K} \cap \mathcal{B}_r(V; p; J) \neq \emptyset$.

PROOF. Suppose the contrary, then there exists $\tilde{r} \in (0, r_0/8)$ such that for any $\tilde{N} \in \mathbb{N}$, $\tilde{R} > 0$ there are $k \in \mathbb{N}$, $J \in \{j_1, \dots, j_k\}^l$ and $p \in P_{\tilde{R}}(k, \tilde{N}, J)$ for which $\mathcal{K} \cap \mathcal{B}_{\tilde{r}}(V; p; J) = \emptyset$ $(V = (v_1, \dots, v_l))$. Let $(v_n^\iota) \subset \mathcal{K}_\iota(\bar{c}_\iota)$ and $(r_n) \subset \mathbb{R}^+$, be the sequences given by 3.14. Since $v_n^\iota \to v_\iota$ and $r_n \to 0$ we can choose $n \in \mathbb{N}$ such that $\|v_n^\iota - v_\iota\| < \tilde{r}/2$, $r_n < \tilde{r}/2 - 3d_{r_n}$ and $\mathcal{B}_{2r_n}(v_n^\iota) \subset \mathcal{B}_{\tilde{r}}(\bar{v}_\iota)$. In particular we have that $\mathcal{B}_{r_n}(V_n; p; J) \subset \mathcal{B}_{\tilde{r}}(V; p; J)$ $(V_n = (v_n^1, \dots, v_n^l))$.

Fixing this n, fix also any r_-, r, r_+ such that $r_n - 3d_{r_n} < r_- < r < r_+ < r_n + 3d_{r_n}$ and fix $c_{-,\iota}$, $c_{+,\iota}$, δ such that $]c_{-,\iota} - \delta, c_{-,\iota} + 2\delta[\subset \subset \bar{c}_\iota - 1/4 \min\{h_n, \mu(r - r_-)\}, \bar{c}[\setminus \Phi^{c_\iota^*}$ and $]c_{+,\iota} - \delta, c_{+,\iota} + 2\delta[\subset \subset]\bar{c}_\iota, \min\{c_\iota^*, \bar{c}_\iota + (1/4)\mu(r - r_-)\}[\setminus \Phi^{c_\iota^*}$.

By 3.14 we can choose $\gamma_\iota \in C([0, 1], X)$ such that

(i) $\gamma_\iota(0)$, $\gamma_\iota(1) \in \partial\mathcal{B}_{r_n}(v_n^\iota) \cap \varphi_\iota^{\bar{c}_\iota - (1/2)h_n}$;

(ii) $\gamma_\iota(0)$ and $\gamma_\iota(1)$ are not connectible in $\{\varphi_\iota < \bar{c}_\iota\} \cap \mathcal{B}_{\tilde{r}}(v_\iota)$;

(iii) range $\gamma_\iota \subset \bar{\mathcal{B}}_{r_n}(v_n^\iota) \cap \varphi_\iota^{c_{+,\iota}}$;

(iv) range $\gamma_\iota \cap \mathcal{C}_{r_n - (1/2)d_{r_n}, r_n}(v_n^\iota) \subset \varphi_\iota^{\bar{c}_\iota - (1/2)h_n}$;

(v) $\operatorname{supp} \gamma_\iota(s) \subseteq [-T, T]$ for any $s \in [0, 1]$, being $T > 0$ independent on s.

Fix $0 < \varepsilon < \min\{\varepsilon_1, (1/9)(\bar{c} - c_{-,\iota}), (1/2)d_{r_n}^2\}$. We can also assume, enlarging T if necessary, that $\|v_n^\iota\|_{|x| \geq T}^2 \leq \varepsilon/4$ and, we fix an integer $N_1 \geq \max\{T, N, 4\}$ where N is given by Proposition 4.9 for these value of r, $b_{+,\iota} = c_{+,\iota}$, $b_{-,\iota} = c_{-,\iota}$, $\varepsilon/4$ instead of ε and V_n instead of V.

If $R > 0$ is given by the Proposition 4.9, since $\mathcal{K} \cap \mathcal{B}_{r_n}(V; p; J) = \emptyset$ for a $p \in P_R(k, N, J)$, there exists a locally Lipschitz continuous function $W: X \to X$ which satisfies the properties (W_0)-(W_4). Let us consider the flow associated to the following Cauchy problem

$$\frac{d\eta}{ds}(s, u) = -W(\eta(s, u)), \qquad \eta(0, u) = u.$$

Plainly, by $(\mathcal{W}_0)$ for any $u \in X$ this Cauchy problem admits a unique solution $\eta(\cdot, u)$ defined on $\mathbb{R}^+$ and the function η is continuous on $\mathbb{R}^+ \times E$. Moreover, the function $s \mapsto \varphi(\eta(s, u))$ is nonincreasing.

We define the function $G\colon Q=[0,1]^k\to X$ by setting $G(\theta)=\sum_{i=1}^{k}\gamma_{j_i}(\theta_i)(\cdot-p_i)$ for $\theta=(\theta_1,\ldots,\theta_k)\in Q$. We put $F_i^0=\{\theta\in Q\colon \theta_i=0\}$ and $F_i^1=\{\theta\in Q\colon \theta_i=1\}$ and we note that $G(\theta)|_{\mathcal{U}_i}=\gamma_{j_i}(0)$ if $\theta\in F_i^0$, $G(\theta)|_{\mathcal{U}_i}=\gamma_{j_i}(1)$ if $\theta\in F_i^1$.

Moreover $G(\theta)|_{\mathcal{U}_i}=\gamma_{j_i}(\theta_i)(\cdot-p_i)$ and $\operatorname{supp}\gamma_{j_i}(\theta_i)(\cdot-p_i)\subseteq[-R+p_i,R+p_i]\subset\mathcal{U}_i\setminus(M_i\cup M_{i-1})$. Therefore $\varphi_{j_i,i}(G(\theta))=\varphi_{j_i}(\gamma_{j_i}(\theta_i))$ for any $i\in\{1,\ldots,k\}$ and for any $\theta=(\theta_1,\ldots,\theta_k)\in Q$.

To prove the theorem, we make the following claim.

CLAIM. *There exists $\tau>0$ such that the continuous function $\bar G\colon Q\to X$ given by $\bar G(\theta)=\eta(\tau,G(\theta))$ satisfies*:

(vi) $\bar G=G$ *on* ∂Q;

(vii) $\max_i\|\bar G(\theta)\|^2_{M_i}\le\varepsilon$ *for any* $\theta\in Q$;

(viii) *there is a path ξ inside Q joining two opposite faces $F_{\bar i}^0$ and $F_{\bar i}^1$ such that, along ξ, the function $\varphi_{j_{\bar i}}\circ\bar G$ takes values under $c_{-,j_{\bar i}}+\varepsilon$; namely*: $\exists\,\bar i\in\{1,\ldots,k\}$ *and* $\xi=(\xi_1,\ldots,\xi_k)\in C([0,1],Q)$ *such that* $\xi_{\bar i}(0)=0$, $\xi_{\bar i}(1)=1$ *and* $\bar G(\xi(s))\in\varphi_{j_{\bar i}}^{c_{-,j_{\bar i}}+\varepsilon}$ *for any* $s\in[0,1]$.

Assume the claim holds and introduce a cut-off function $\chi\in C(\mathbb{R}^m,\mathbb{R})$, such that $\chi(x)=0$ if $x\notin\mathcal{U}_{\bar i}$, $\chi(t)=1$ if $t\in\mathcal{U}_{\bar i}\setminus(M_{\bar i}\cup M_{\bar i-1})$ and linear on the connected parts of the intersection of any line passing through the origin with the set $\mathcal{U}_{\bar i}\cap(M_{\bar i}\cup M_{\bar i-1})$.

Define $g\in C([0,1],X)$ by setting $g(s)=\chi\bar G(\xi(s))$ for $s\in[0,1]$. We observe that, because of (vi) and (viii) we have $g(0)=\chi\bar G(\xi(0))=\chi G(\xi(0))=\gamma_{j_{\bar i}}(0)(\cdot-p_{\bar i})$ and similarly $g(1)=\gamma_{j_{\bar i}}(1)(\cdot-p_{\bar i})$. We have also that the path g is contained in the ball $B_{\bar r}(v_{j_{\bar i}}(\cdot-p_{\bar i}))$. Indeed, first of all $\|v_n^{j_{\bar i}}-v_{j_{\bar i}}\|\le r_0/16<\bar r/2$. Secondly, since $\operatorname{supp}g_{j_{\bar i}}(s)\subset\mathcal{U}_{\bar i}$, $\varepsilon<(1/2)d^2_{r_n}$ and since $B_{r_+}(V_n,p,J)$ is invariant under η, we get

$$\|g_{j_{\bar i}}(s)-v_n^{j_{\bar i}}(\cdot-p_{\bar i})\|^2\le\|g_{j_{\bar i}}(s)-v_n^{j_{\bar i}}(\cdot-p_{\bar i})\|^2_{\mathcal{U}_{\bar i}}+\varepsilon\le$$

$$\le\max_{\theta\in Q}\big(\|\chi(\bar G(\theta)-v^{j_{\bar i n}}(\cdot-p_{\bar i}))\|_{\mathcal{U}_{\bar i}}+2\varepsilon\big)^2\le 2(r_++\varepsilon)^2+\varepsilon\le 4r_n^2\le\frac{\bar r^2}{4}.$$

Translating by $-p_{\bar i}$ the path g, we get a curve joining $\gamma_{j_{\bar i}}(0)$ with $\gamma_{j_{\bar i}}(1)$ in $\mathcal{B}_{\bar r}(v_{j_{\bar i}})$. Showing that on g the functional $\varphi_{j_{\bar i}}$ remains under the level $\bar c_{j_{\bar i}}$ we will get a contradiction with the property (ii) of $\gamma_{j_{\bar i}}$.

To prove this, we notice that

$$(5.2)\qquad \varphi_{j_{\bar i}}(g(s)) = \varphi_{j_{\bar i},\bar i}(g(s)) \leqslant \varphi_{j_{\bar i},\bar i}(\bar G(\xi(s))) + \frac{1}{2}\|g(s)\|^2_{\mathcal{U}_{\bar i}\cap(M_{\bar i}\cup M_{\bar i-1})} + $$

$$+ \int_{\mathcal{U}_{\bar i}\cap(M_{\bar i}\cup M_{\bar i-1})} F_{j_{\bar i}}(x, \bar G(\xi(s))) - \int_{\mathcal{U}_{\bar i}\cap(M_{\bar i}\cup M_{\bar i-1})} F_{j_{\bar i}}(x, g(s)).$$

By (viii), $\varphi_{j_{\bar i},\bar i}(\bar G(\xi(s))) \leqslant c_{-,j_{\bar i}} + \varepsilon$. Moreover, by (vii), $(1/2)\|g(s)\|^2_{\mathcal{U}_{\bar i}\cap(M_{\bar i}\cup M_{\bar i-1})} \leqslant \|\bar G(\xi(s))\|^2_{M_{\bar i}} + \|\bar G(\xi(s))|^2_{M_{\bar i-1}} \leqslant 2\varepsilon$. Finally,

$$\int_{\mathcal{U}_{\bar i}\cap(M_{\bar i}\cup M_{\bar i-1})} |V(t, \bar G(\xi(s)))| \leqslant \|\bar G(\xi(s))\|^2_{\mathcal{U}_{\bar i}\cap(M_{\bar i}\cup M_{\bar i-1})} \leqslant 2\varepsilon$$

and

$$\int_{\mathcal{U}_{\bar i}\cap(M_{\bar i}\cup M_{\bar i-1})} |F_{j_{\bar i},\bar i}(x, g(s))| \leqslant 2\|\bar G(\xi(s))\|^2_{\mathcal{U}_{\bar i}\cap(M_{\bar i}\cup M_{\bar i-1})} \leqslant 4\varepsilon.$$

Putting together all these estimates in (5.2), and considering that $\varepsilon < (1/9)(\bar c - c_-)$, we finally get that $\varphi_{\bar i}(g(s)) \leqslant c_{-,j_{\bar i}} + 9\varepsilon < \bar c_{j_{\bar i}}$, which contradicts (ii).

To check the claim we firstly note that the properties (vi) and (vii) are true for any $\tau > 0$, and follows easily from (W_2) and (W_3).

We divide the proof of (viii) in some lemmas.

LEMMA 5.3. *There is $\tau > 0$ such that for any $u \in \mathcal{B}_{r_-}(V_n; p; J) \cap \bigcap_{i=1}^{k} \varphi_{j_i,i}^{c_+,j_i}$ there exists $\bar i \in \{1, \ldots, k\}$ for which $\eta(\tau, u) \in \varphi_{j_{\bar i},\bar i}^{c_-,j_{\bar i}}$.*

PROOF. Set $\sigma = 2 \operatorname{diam} \varphi(\mathcal{B}_{r_+}(V_n; p; J))$. Since $\varphi(\mathcal{B}_{r_+}(V_n; p; J))$ is a bounded set, $\sigma < +\infty$. Put $\tau = \sigma/\mu_k$ and let $u \in \mathcal{B}_{r_-}(V_n; p; J) \cap \bigcap_{i=1}^{k} \varphi_{j_i,i}^{c_+,j_i}$. By (W_2) the curve $s \mapsto \eta(s, u)$ remains in $\bigcap_{i=1}^{k} \varphi_{j_i,i}^{c_+,j_i}$. Moreover it must goes out of $\mathcal{B}_r(V_n; p; J)$ at some $\bar s \in]0, \tau[$, otherwise, by (W_4),

$$\varphi(u) - \varphi(\eta(\tau, u)) = \int_0^\tau \varphi'(\eta(s, u))\, \mathcal{W}(\eta(s, u))\, ds \geqslant \mu_k \tau = \sigma$$

in contrast with the definition of σ. Then, there are $\bar i \in \{1, \ldots, k\}$ and an interval $[s_1, s_2] \subset]0, \tau[$ such that $\|\eta(s_1, u) - v_n^{j_{\bar i}}(\cdot - p_{\bar i})\|_{\mathcal{U}_{\bar i}} = r_-$, $\|\eta(s_2, u) - v_n^{j_{\bar i}}(\cdot - p_{\bar i})\|_{\mathcal{U}_{\bar i}} = r$ and $r_- < \|\eta(s, u) - v_n^{j_{\bar i}}(\cdot - p_{\bar i})\|_{\mathcal{U}_j} < r$ for any $s \in]s_1, s_2[$. Then by (W_1) and since by (W_2) $\eta(s, u) \in \varphi_{j_{\bar i},\bar i}^{c_+,\bar i}$ for any

$s \geqslant 0$, we get

$$\varphi_{j_{\bar{i}},\bar{i}}(\eta(s_2, u)) \leqslant \varphi_{j_{\bar{i}},\bar{i}}(\eta(s_1, u)) -$$

$$- \int_{s_1}^{s_2} \varphi'_{j_{\bar{i}},\bar{i}}(\eta(s, u))\, \mathcal{W}(\eta(s, u))\, ds \leqslant c_{+,j_{\bar{i}}} - \mu(s_2 - s_1).$$

But since $\|\mathcal{W}(\eta(s, u))\|_{\mathcal{U}_{\bar{i}}} \leqslant 2$ for any $s \geqslant 0$ we get also

$$r - r_- \leqslant \|\eta(s_2, u) - \eta(s_1, u)\|_{\mathcal{U}_{\bar{i}}} \leqslant \int_{s_1}^{s_2} \|\mathcal{W}(\eta(s, u))\|_{\mathcal{U}_{\bar{i}}} \leqslant 2(s_2 - s_1)$$

from which $\varphi_{j_{\bar{i}},\bar{i}}(\eta(s_2, u)) \leqslant c_{+,j_{\bar{i}}} - (1/2)\mu(r - r_-) < c_{-,j_{\bar{i}}}$. By (W_2) we get that $\eta(s, u) \in \varphi_{j_{\bar{i}},\bar{i}}^{c_{-,j_{\bar{i}}}}$ for any $s \geqslant s_2$ and in particular that $\eta(\tau, u) \in \varphi_{j_{\bar{i}},\bar{i}}^{c_{-,j_{\bar{i}}}}$. ∎

LEMMA 5.4. *For any $\theta \in Q$ there is $\bar{i} \in \{1, \ldots, k\}$ such that* $\varphi_{j_{\bar{i}},\bar{i}}(\bar{G}(\theta)) \leqslant c_{-,j_{\bar{i}}}$.

PROOF. Assume first that $G(\theta) \in \mathcal{B}_{r_-}(V_n; p; J)$. Then, since by construction $G(\theta) \in \bigcap_{i=1}^{k} \varphi_{j_i,i}^{c_{+,j_i}}$, we obtain the result by Lemma 5.3.

In the other case there exists $\bar{i} \in \{1, \ldots, k\}$ such that

$$r_n - \frac{1}{2} d_{r_n} \leqslant r_- \leqslant \|G(\theta) - v_n^{j_{\bar{i}}}(\cdot - p_{\bar{i}})\|_{\mathcal{U}_{\bar{i}}} =$$

$$= \|\gamma_{j_{\bar{i}}}(\theta_{\bar{i}})(\cdot - p_{\bar{i}}) - v_n^{j_{\bar{i}}}(\cdot - p_{\bar{i}})\|_{\mathcal{U}_{\bar{i}}} \leqslant \|\gamma_{j_{\bar{i}}}(\theta_{\bar{i}}) - v_n^{j_{\bar{i}}}\|$$

so using the properties (iii) and (iv) of $\gamma_{j_{\bar{i}}}$ we get

$$\varphi_{j_{\bar{i}},\bar{i}}(G(\theta)) = \varphi_{j_{\bar{i}},\bar{i}}(\gamma(\theta_{\bar{i}})(\cdot - p_{\bar{i}})) = \varphi_{j_{\bar{i}}}(\gamma(\theta_{\bar{i}})) \leqslant \bar{c}_{j_{\bar{i}}} - \frac{1}{2} h_n \leqslant c_{-,j_{\bar{i}}}.$$

By Lemma 5.3 we then have that $\eta(s, G(\theta)) \in \varphi_{j_{\bar{i}},\bar{i}}^{c_{-,j_{\bar{i}}}}$ for any $s \geqslant 0$ and the lemma follows. ∎

Now we can conclude the proof of property (viii). We proceed by contradiction assuming the contrary. That is, for any $i \in \{1, \ldots, k\}$ the set $D_i = (\varphi_{j_i,i} \circ \bar{G})^{-1}([c_{-,j_i} + \varepsilon, +\infty[)$ separates F_i^0 from F_i^1 in Q. Let C_i be the component of $Q \setminus D_i$ containing F_i^1 and let $\sigma_i : Q \to \mathrm{R}$ be the fun-

ction given by

$$\sigma_i(\theta) = \begin{cases} \operatorname{dist}(\theta, D_i) & \text{if } \theta \in Q \backslash C_i, \\ -\operatorname{dist}(\theta, D_i) & \text{if } \theta \in C_i. \end{cases}$$

Then σ_i is a continuous function on Q such that $\sigma_i|_{F_i^0} \geq 0$, $\sigma_i|_{F_i^1} \leq 0$ and $\sigma_i(\theta) = 0$ if and only if $\theta \in D_i$. Using a theorem by Carlo Miranda (see [17]) we get that there exists $\theta \in Q$ such that $\sigma_i(\theta) = 0$ for all $i \in$ $\in \{1, \ldots, k\}$ which means that $\bigcap_{i=1}^{k} D_i \neq \emptyset$. But this is in contrast with Lemma 5.4. ■

Aknowledgment. I wish to thank Vittorio Coti Zelati and Paolo Caldiroli for the useful comments and suggestions.

REFERENCES

[1] S. ABENDA - P. CALDIROLI - P. MONTECCHIARI, *Multibump solutions for Duffing-like systems*, Preprint S.I.S.S.A. (1994).

[2] R. A. ADAMS, *Sobolev Spaces*, Academic Press, New York (1975).

[3] S. ALAMA - Y.Y. LI, *On «Multibump» Bound States for Certain Semilinear Elliptic Equations*, Research Report No. 92-NA-012, Carnegie Mellon University (1992).

[4] S. ALAMA - G. TARANTELLO, *On semilinear elliptic equations with indefinite nonlinearities*, Calc. Var., 1 (1993), pp. 439-475.

[5] A. AMBROSETTI, *Critical points and nonlinear variational problems*, Bul. Soc. Math. France, 120 (1992).

[6] S. ANGENENT, *The shadowing lemma for elliptic PDE*, in *Dynamics of Infinite Dimensional Systems*, S. N. Chow and J. K. Hale eds., F37(1987).

[7] P. CALDIROLI - P. MONTECCHIARI, *Homoclinic orbits for second order Hamiltonian systems with potential changing sign*, Comm. Appl. Nonlinear Anal., 1 (1994), pp. 97-129.

[8] V. COTI ZELATI - I. EKELAND - E. SÉRÉ, *A Variational approach to homoclinic orbits in Hamiltonian systems*, Math. Ann., 288 (1990), pp. 133-160.

[9] V. COTI ZELATI - P. H. RABINOWITZ, *Homoclinic orbits for second order Hamiltonian systems possessing superquadratic potentials*, J. Amer. Math. Soc., 4 (1991), pp. 693-727.

[10] V. COTI ZELATI - P. H. RABINOWITZ, *Homoclinic type solutions for a semilinear elliptic PDE on* $\mathbb{R}^n$, Comm. Pure Appl. Math., 45 (1992), pp. 1217-1269.

[11] W. Y. DING - W. M. NI, *On the existence of positive entire solutions of a semilinear elliptic equation*, Arch. Rat. Mech. Anal., 91 (1986), pp. 283-308.

[12] M. J. ESTEBAN - P. L. LIONS, *Existence and nonexistence results for semilinear elliptic problems in unbounded domains*, Proc. Roy. Soc. Edinburgh, **93** (1982), pp. 1-14.

[13] M. K. KWONG, *Uniqueness positive solutions of* $\Delta u - u + u^p = 0$ *in* R^n, Arch. Rat. Mech. Anal., **105** (1985), pp. 243-266.

[14] L. LASSOUED, *Periodic solution of a second order superquadratic system with change of sign of potential*, J. Diff. Eq., **93**(1991), pp. 1-18.

[15] P.L. LIONS, *The concentration-compactness principle in the calculus of variations. The locally compact case*, part 1, Ann. Inst. Henri Poincaré, **1** (1984), pp. 109-145.

[16] P. L. LIONS, *The concentration-compactness principle in the calculus of variations. The locally compact case*, part 2, Ann. Inst. Henri Poincaré, **1** (1984), pp. 223-283.

[17] C. MIRANDA, *Un'osservazione su un teorema di Brouwer*, Boll. Unione Mat. Ital., **3** (1940), pp. 5-7.

[18] P. MONTECCHIARI, *Existence and multiplicity of homoclinic solutions for a class of asymptotically periodic second order Hamiltonian systems*, Ann. Mat. Pura Appl. (to appear). See also: *Multiplicity of homoclinic solutions for a class of asymptotically periodic second order Hamiltonian systems*, Rend. Mat. Acc. Lincei, s. 9, 4 (1993), pp. 265-271.

[19] P. PUCCI - J. SERRIN, *The structure of the critical set in the mountain pass theorem*, Tran. Am. Math. Soc., **299** (1987), pp. 115-132.

[20] P. H. RABINOWITZ, *Homoclinic orbits for a class of Hamiltonian systems*, Proc. Roy. Soc. Edinburgh, **114**-A (1990), pp. 33-38.

[21] P.H. RABINOWITZ, *A note on a semilinear elliptic equation on* R^m, in *A tribute in honour of Giovanni Prodi*, A. Ambrosetti and A. Marino eds., Quaderni Scuola Normale Superiore, Pisa (1991).

[22] E. SÉRÉ, *Existence of infinitely many homoclinic orbits in Hamiltonian systems*, Math. Z., **209** (1992), pp. 27-42.

[23] E. SÉRÉ, *Looking for the Bernoulli shift*, Ann. Inst. H. Poincaré. Anal. Non Linéaire, **10** (1993), pp. 561-590.

[24] A. WEINSTEIN, *Bifurcations and Hamilton's principle*, Math. Z., **159** (1978), pp. 235-248.

Manoscritto pervenuto in redazione il 22 febbraio 1995.