

Test sets for factorization properties of modules

JAN ŠAROCH (*) – JAN TRLIFAJ (**)

To László Fuchs in honor of his 95th birthday

ABSTRACT – Baer’s Criterion of injectivity implies that injectivity of a module is a factorization property with respect to a single monomorphism. Using the notion of a cotorsion pair, we study generalizations and dualizations of factorization properties in dependence on the algebraic structure of the underlying ring R and on additional set-theoretic hypotheses. For R commutative noetherian of Krull dimension $0 < d < \infty$, we show that the assertion ‘projectivity is a factorization property with respect to a single epimorphism’ is independent of ZFC + GCH. We also show that if R is any ring and there exists a strongly compact cardinal $\kappa > |R|$, then the category of all projective modules is κ -accessible.

MATHEMATICS SUBJECT CLASSIFICATION (2010). Primary: 16D40, 03E35; Secondary: 16E30, 16B70, 03E55, 16D10.

KEYWORDS. Projective module, factorization class, cotorsion pair, Dedekind domain, λ -purity, Weak Diamond Principle, strongly compact cardinal.

Introduction

Various classes of modules studied in homological algebra are defined by factorization properties with respect to proper classes of monomorphisms or epimorphisms. A natural question arises of whether these defining classes of morphisms

(*) *Indirizzo dell’A.*: Department of Algebra, Faculty of Mathematics and Physics, Charles University, Sokolovská 83, 186 75 Prague 8, Czech Republic
E-mail: saroch@karlin.mff.cuni.cz

(**) *Indirizzo dell’A.*: Department of Algebra, Faculty of Mathematics and Physics, Charles University, Sokolovská 83, 186 75 Prague 8, Czech Republic
E-mail: trlifaj@karlin.mff.cuni.cz

can be replaced by sets, and hence by single morphisms. In many cases, this is possible. We start with the easier setting of injective modules, and more in general, right-hand classes of complete cotorsion pairs. Then we will turn to the more difficult setting of projective modules, and left-hand classes of cotorsion pairs, where solutions often require extra algebraic and/or set-theoretic assumptions. Our notation will follow [7] and [14].

1. Injective modules and their generalizations

Injective modules and their various generalizations are defined by factorization properties with respect to proper classes of monomorphisms. Denote by Ψ the class of all monomorphisms in $\text{Mod-}R$. Then a module M is *injective*, if and only if $\text{Hom}_R(\psi, M)$ is surjective for each $\psi \in \Psi$, i.e., if and only if for each $\psi \in \Psi$ with $\psi \in \text{Hom}_R(N, N')$, each homomorphism from N to M factorizes through ψ . The classic Baer Criterion for injectivity [4] says that in the definition above, one can replace the proper class Ψ by a subset, and hence by a single element. Namely, if S denotes the set of all right ideals of R , then M is injective, if and only if $\text{Hom}_R(\varphi, M)$ is surjective, where $\varphi: \bigoplus_{I \in S} I \hookrightarrow \bigoplus_{I \in S} R$. By dimension shifting, it follows that for each $n < \omega$, there is a single monomorphism φ_n such that factorization through φ_n defines the property of having injective dimension $\leq n$.

Given a monomorphism ψ in $\text{Mod-}R$, we call a class of modules \mathcal{C} a *factorization class* of ψ provided that $\mathcal{C} = \{M \in \text{Mod-}R \mid \text{Hom}_R(\psi, M) \text{ is surjective}\}$. So, for example, the class \mathcal{I}_n of all modules of injective dimension $\leq n$ is the factorization class of φ_n . Using Enochs' proof of the Flat Cover Conjecture (i.e., the fact that the class \mathcal{F}_0 of all flat modules is deconstructible, see definition below), we infer that there is a monomorphism ν such that the class \mathcal{EC} of all Enochs cotorsion modules is the factorization class of ν .

In order to see that these results are particular instances of a more general phenomenon, we need to recall several definitions:

Let \mathcal{S} be a class of modules. A module M is \mathcal{S} -*filtered* provided that M is the union of an increasing well-ordered continuous chain of its submodules $(M_\alpha, \nu_{\beta\alpha} \mid \alpha < \beta < \sigma)$ such that $\nu_{\beta\alpha}: M_\alpha \rightarrow M_\beta$ is the inclusion for all $\alpha < \beta < \sigma$, $M_0 = 0$, and for each $\alpha < \sigma$, $\text{Coker}(\nu_{\alpha+1\alpha})$ is isomorphic to an element of \mathcal{S} . The class of all \mathcal{S} -filtered modules is denoted by $\text{Filt}(\mathcal{S})$. The class of all direct summands of \mathcal{S} -filtered modules is denoted by $\text{sFilt}(\mathcal{S})$. Note that $\text{Filt}(\text{Filt}(\mathcal{S})) = \text{Filt}(\mathcal{S})$ and $\text{Filt}(\text{sFilt}(\mathcal{S})) = \text{sFilt}(\mathcal{S})$. In particular, both $\text{Filt}(\mathcal{S})$ and $\text{sFilt}(\mathcal{S})$ are closed under arbitrary direct sums and extensions.

For example, if $\mathcal{S} = \{R\}$, then $\text{Filt}(\mathcal{S})$ is the class of all free modules, and $\text{sFilt}(\mathcal{S})$ the class of all projective modules.

A class of modules \mathcal{C} is *deconstructible* provided that there is a subset $\mathcal{S} \subseteq \mathcal{C}$ such that $\mathcal{C} = \text{Filt}(\mathcal{S})$.

For a class of modules \mathcal{C} , we define

$$\mathcal{C}^\perp = \{M \in \text{Mod-}R \mid \text{Ext}_R^1(C, M) = 0 \text{ for all } C \in \mathcal{C}\}$$

and ${}^\perp\mathcal{C} = \{M \in \text{Mod-}R \mid \text{Ext}_R^1(M, C) = 0 \text{ for all } C \in \mathcal{C}\}$. A pair of classes of modules $\mathcal{C} = (\mathcal{A}, \mathcal{B})$ is a *cotorsion pair* in case $\mathcal{A} = {}^\perp\mathcal{B}$ and $\mathcal{B} = \mathcal{A}^\perp$. \mathcal{C} is said to be generated (cogenerated) by a class \mathcal{C} , if $\mathcal{B} = \mathcal{C}^\perp$ ($\mathcal{A} = {}^\perp\mathcal{C}$). \mathcal{C} is *hereditary*, if $\text{Ext}_R^i(A, B) = 0$ for all $A \in \mathcal{A}$, $B \in \mathcal{B}$ and all $i \geq 2$.

LEMMA 1.1. *Let $\mathcal{C} = (\mathcal{A}, \mathcal{B})$ be a cotorsion pair in $\text{Mod-}R$. Then the following conditions are equivalent:*

1. \mathcal{C} is generated by a set;
2. there is a subset $\mathcal{S} \subseteq \mathcal{A}$ such that $\mathcal{A} = \text{sFilt}(\mathcal{S})$;
3. \mathcal{A} is deconstructible;
4. there is a monomorphism $f: Q \hookrightarrow P$ with P projective such that \mathcal{B} is the factorization class of f .

PROOF. (1) \Rightarrow (3) by the Kaplansky Theorem for cotorsion pairs [14, 7.13]. Since \mathcal{A} is closed under direct summands, (3) \Rightarrow (2). That (2) \Rightarrow (1) follows by the Eklof Lemma [14, 6.2].

Assume (1). Then there is a set $\mathcal{S} \subseteq \mathcal{A}$ such that $\mathcal{B} = \mathcal{S}^\perp$. For each $A \in \mathcal{S}$, there is a presentation $0 \rightarrow Q_A \xrightarrow{\nu_A} P_A \rightarrow A \rightarrow 0$ with P_A projective. We define a monomorphism f by $f = \bigoplus_{A \in \mathcal{S}} \nu_A$. Then $f \in \text{Hom}_R(Q, P)$ where $Q = \bigoplus_{A \in \mathcal{S}} Q_A$ and $P = \bigoplus_{A \in \mathcal{S}} P_A$ is projective. Moreover, for each $N \in \text{Mod-}R$, each homomorphism from Q to N factorizes through f , if and only if $\text{Ext}_R^1(A, N) = 0$ for each $A \in \mathcal{S}$, if and only if $N \in \mathcal{B}$. So (4) holds.

Assume (4). Then for each $N \in \text{Mod-}R$, one has $N \in \mathcal{B}$ if and only if $\text{Ext}_R^1(P/Q, N) = 0$. That is, \mathcal{C} is generated by the one element set $\{P/Q\}$, proving (1). \square

For integral domains, other variants of cotorsion modules are studied. However, Lemma 1.1 applies to them, too. Recall that a module M over an integral domain R is called *Matlis cotorsion* if $\text{Ext}_R^1(Q, M) = 0$ where Q denotes the quotient field of R .

COROLLARY 1.2. *Let R be an integral domain. Then there exist monomorphisms $g_0 \in \text{Hom}_R(Q_0, P_0)$ and $g_1 \in \text{Hom}_R(Q_1, P_1)$ with P_0 and P_1 projective, such that the class of all Matlis cotorsion (Warfield cotorsion) modules is the factorization class of g_0 (g_1).*

PROOF. The first claim follows immediately by Lemma 1.1. For the second claim, one uses the characterization of Warfield cotorsion modules as the Matlis cotorsion modules of injective dimension ≤ 1 , [14, 7.47]. \square

A different, and more involved, argument shows that there is a single pure monomorphism μ such that the class of all pure-injective modules is the factorization class of μ :

LEMMA 1.3. *Let $\kappa = |R| + \aleph_0$. Then a module L is pure-injective, if and only if for each pure embedding $v \in \text{Hom}_R(N, M)$ with $|M| \leq \kappa$ and each $f \in \text{Hom}_R(N, L)$ there exists $g \in \text{Hom}_R(M, L)$ with $gv = f$.*

PROOF. The only-if part is trivial, let us prove the if part. By [7, Theorem V.1.2], it is enough to show that each system \mathcal{S} consisting of R -linear equations (in any number, finite or infinite, of variables) with parameters in L has a solution in L provided that $|\mathcal{S}| \leq \kappa$ and each finite $\mathcal{F} \subseteq \mathcal{S}$ has a solution in L .

Let \mathcal{S} be such a system. Fix a submodule $N \leq L$ of cardinality $\leq \kappa$ such that N contains all the parameters from \mathcal{S} and each finite $\mathcal{F} \subseteq \mathcal{S}$ has a solution in N . Let us denote by f the inclusion $N \subseteq L$. Following the proof of [7, Theorem V.1.2 (2) \Rightarrow (3)], we put $M = (N \oplus F)/K$ where F is the free module whose basis is the set of variables in the system \mathcal{S} , and K is the submodule generated by all the elements $(-b, \sum_{j=1}^m y_j r_j)$ where $\sum_{j=1}^m y_j r_j = b$ belongs to \mathcal{S} . Let $\iota: N \hookrightarrow M$ be the canonical embedding. Using finite solvability of \mathcal{S} in N , it is straightforward to check that ι is actually a pure embedding. Since $|M| \leq \kappa$, we can use our assumption to find $g \in \text{Hom}_R(M, L)$ such that $g\iota = f$. By construction, \mathcal{S} has a solution in M . The g -image of this solution is a desired solution of \mathcal{S} in L . \square

The notions of purity and pure-injectivity naturally generalize as follows. Given a regular infinite cardinal λ , a monomorphism $f: B \rightarrow A$ is λ -pure if the homomorphism $\text{Hom}_R(C, A) \rightarrow \text{Hom}_R(C, \text{Coker}(f))$ induced by the canonical projection $A \rightarrow \text{Coker}(f)$ is surjective for each $< \lambda$ -presented module C . A module is called λ -pure-injective if it is injective with respect to all λ -pure monomorphisms. The case $\lambda = \aleph_0$ amounts to the classical notions of purity/pure-injectivity.

Going back to Lemma 1.3, if we consider λ -purity and λ -pure-injective modules for a regular uncountable cardinal λ , the situation is rather unclear. The difficulty here stems from the fact that $\text{Mod-}R$ often does not possess *enough λ -pure-injective modules*, i.e. there exist R -modules which cannot be λ -purely embedded into λ -pure-injective ones.

The following proposition shows in particular that, over a non-right perfect ring, the class of \aleph_1 -pure-injective modules does not precisely recognize \aleph_1 -pure monomorphisms, i.e. there is a pure and not \aleph_1 -pure monomorphism v such that $\text{Hom}_R(v, N)$ is surjective for each \aleph_1 -pure-injective module N .

PROPOSITION 1.4. *Let R be a ring which is not right perfect. Then $\text{Mod-}R$ does not have enough \aleph_1 -pure-injective objects.*

PROOF. Let us denote by \mathcal{C} the class of all \aleph_1 -pure-injective modules. Since R is not right perfect, there exists a countably presented flat module M which is not projective. This module is the direct limit of a countable directed system $F_0 \rightarrow F_1 \rightarrow F_2 \rightarrow \dots$ consisting of finite rank free modules. In particular, we have a pure, albeit not \aleph_1 -pure, short exact sequence $0 \rightarrow R^{(\omega)} \xrightarrow{\varepsilon} R^{(\omega)} \rightarrow M \rightarrow 0$.

Each flat Mittag-Leffler module (being \aleph_1 -projective, cf. [14, 3.19]) is an \aleph_1 -pure-epimorphic image of a projective module, whence it belongs to ${}^\perp\mathcal{C}$. By [5, Theorem 6], $M \in {}^\perp\mathcal{C}$ as well.¹ It immediately follows that the module $R^{(\omega)}$ cannot be \aleph_1 -purely embedded into an \aleph_1 -pure-injective module since this would imply that ε is \aleph_1 -pure and thus splits. \square

Under the additional set-theoretic assumption that $0^\#$ does not exist (a weak form of $V = L$, cf. [17, §18] or [7, VI.3.16]), we can even show that:

PROPOSITION 1.5 ($0^\#$ does not exist). *Let R be a ring which is not right perfect. There does not exist a regular uncountable cardinal λ such that $\text{Mod-}R$ has enough λ -pure-injective modules.²*

¹ **ADDED IN PROOF.** This argument shows that, over any ring, all countably presented flat modules belong to ${}^\perp\mathcal{C}$. Since each flat module F is the direct limit of an \aleph_1 -directed system consisting of countably presented flat modules, it is an \aleph_1 -pure-epimorphic image of a direct sum of countably presented flat modules. This direct sum belongs to ${}^\perp\mathcal{C}$, and \mathcal{C} consists of \aleph_1 -pure-injective modules. We conclude that $F \in {}^\perp\mathcal{C}$ as well, whence each \aleph_1 -pure-injective module is cotorsion.

² **ADDED IN PROOF.** In a forthcoming paper of the first author and Manuel Cortés-Izurdiaga, we show that the conclusion of Proposition 1.5 actually characterizes the right non-pure-semisimple rings (under the same additional set-theoretical assumption).

PROOF. For the sake of contradiction, assume that there exists such a λ . Let $f: R^{(\omega)} \rightarrow B$ be a λ -pure embedding into a λ -pure-injective module B . Let κ be a strong limit singular cardinal with uncountable cofinality such that $\kappa \geq |B| + \lambda$ and fix a countably presented non-projective flat module N together with a pure short exact sequence $0 \rightarrow R^{(\omega)} \xrightarrow{\varepsilon} R^{(\omega)} \rightarrow N \rightarrow 0$.

It follows from our assumption on $0^\#$ and [7, VI.3.16 and VI.3.6] that $E(\kappa^+)$ holds which allows us to use [7, Theorem VII.1.4] to obtain a κ^+ -free module M which is not projective. In particular, M is a λ -pure-epimorphic image (even κ^+ -pure) of a free module, whence $M \in {}^\perp B$. Moreover, by the proof of [7, Theorem VII.1.4], there are a stationary subset E of κ^+ consisting of ordinals with countable cofinality and a filtration $\mathcal{M} = (M_\alpha \mid \alpha < \kappa^+)$ of M consisting of κ -generated free modules such that $M_{\nu+1}/M_\nu \cong N$ for each $\nu \in E$.

We have $2^\kappa = \kappa^+$ by [17, Corollary 18.33]. Using [23], we get $\diamondsuit_{\kappa^+}(E)$, in particular $\Phi_{\kappa^+}(E)$. By Lemma 2.9, there exists a $\nu \in E$ such that $N \cong M_{\nu+1}/M_\nu \in {}^\perp B$. In particular, f can be factorized through ε yielding that ε is λ -pure, and thus split, in contradiction with the choice of N . \square

REMARK 1.6. If $\lambda = \aleph_n$ for $0 < n < \omega$, then the proof of Proposition 1.5 can be carried out in ZFC using the tools from [7, Section VII.3].

EXAMPLE 1.7. If R is a countable ring and \mathcal{C} the class of all \aleph_1 -pure-injective modules, then ${}^\perp \mathcal{C}$ is just the class of all flat modules (cf. [5, Theorem 6]). As a consequence, \aleph_1 -pure-injective modules over countable rings are cotorsion.

Suppose that R is a countable von Neumann regular ring which is not semisimple. Then all the classes of cotorsion, pure-injective, \aleph_1 -pure-injective and injective modules coincide. Subsequently, there exists a single monomorphism μ such that a module M is \aleph_1 -pure-injective if and only if $\text{Hom}_R(\mu, M)$ is surjective. On the other hand, there are strictly more pure monomorphisms than \aleph_1 -pure monomorphisms. Also, since all cyclic modules are countable, it immediately follows from the Baer test lemma that \aleph_1 -pure submodules of (\aleph_1 -pure-)injective modules are just the direct summands. Thus not only $\text{Mod-}R$ does not have enough \aleph_1 -pure-injective modules, even more so: no non-injective module \aleph_1 -purely embeds into an \aleph_1 -pure-injective module.

Consider now the class \mathcal{T} of all \aleph_1 -pure embeddings into projective modules. By [14, Remark 10.7], $\{\text{Coker}(\nu) \mid \nu \in \mathcal{T}\}$ is precisely the class of all flat Mittag-Leffler modules, hence (again by [5, Theorem 6]) \mathcal{T} is the test class for (\aleph_1 -pure-)injectivity. However, we cannot pick a subset $\mathcal{S} \subset \mathcal{T}$ with the same property, since for each set \mathcal{F} of flat Mittag-Leffler modules \mathcal{F}^\perp contains a non-injective module.

2. Projective modules and their generalizations

We turn to the dual setting of projective modules and their generalizations. These classes of modules are defined by factorization properties with respect to proper classes of epimorphisms. Here, the problem of replacing a proper class of morphisms by a set is more complex: we will see that its solution may depend on the underlying ring, and on the extension of ZFC that we work in.

Given an epimorphism φ in $\text{Mod-}R$, we call a class of modules \mathcal{D} a *cofactorization class* of φ provided that $\mathcal{D} = \{N \in \text{Mod-}R \mid \text{Hom}_R(N, \varphi) \text{ is surjective}\}$.

Our first aim is to dualize Lemma 1.1. To this purpose, we need more definitions. Let \mathcal{S} be a class of modules. A module M is \mathcal{S} -cofiltered provided that M is isomorphic to an inverse limit of a well-ordered continuous inverse system $(M_\alpha, \pi_{\alpha\beta} \mid \alpha < \beta < \sigma)$ such that $\pi_{\alpha\beta}: M_\beta \rightarrow M_\alpha$ is an epimorphism for all $\alpha < \beta < \sigma$, $M_0 = 0$, and for each $\alpha < \sigma$, $\text{Ker}(\pi_{\alpha\alpha+1})$ is isomorphic to an element of \mathcal{S} . The class of all \mathcal{S} -cofiltered modules is denoted by $\text{Cofilt}(\mathcal{S})$. The class of all direct summands of \mathcal{S} -cofiltered modules is denoted by $\text{sCofilt}(\mathcal{S})$.

Note that $\text{Cofilt}(\text{Cofilt}(\mathcal{S})) = \text{Cofilt}(\mathcal{S})$ and also $\text{Cofilt}(\text{sCofilt}(\mathcal{S})) = \text{sCofilt}(\mathcal{S})$. In particular, both $\text{Cofilt}(\mathcal{S})$ and $\text{sCofilt}(\mathcal{S})$ are closed under arbitrary direct products and extensions.

A class of modules \mathcal{C} is *codeconstructible* provided that there is a subset $\mathcal{S} \subseteq \mathcal{C}$ such that $\mathcal{C} = \text{Cofilt}(\mathcal{S})$.

Here is a partial dual of Lemma 1.1:

LEMMA 2.1. *Let $\mathfrak{C} = (\mathcal{A}, \mathcal{B})$ be a cotorsion pair in $\text{Mod-}R$. Consider the following conditions:*

1. \mathfrak{C} is cogenerated by a set;
2. there is a subset $\mathcal{S} \subseteq \mathcal{B}$ such that $\mathcal{B} = \text{sCofilt}(\mathcal{S})$;
3. \mathcal{B} is codeconstructible;
4. there is an epimorphism $g \in \text{Hom}_R(I, J)$ with I injective, such that \mathcal{A} is the cofactorization class of g .

Then (1) \iff (4). Moreover, (3) \implies (2), and (2) \implies (1).

PROOF. The proof of the equivalence of (1) and (4) is dual to the proof of the corresponding equivalence in Lemma 1.1. Moreover, (3) \implies (2) because \mathcal{B} is closed under direct summands. That (2) \implies (1) follows by the Lukas Lemma [14, 6.37]. \square

The following example shows that (2) need not imply (3) even for the trivial cotorsion pair $\mathfrak{I} = (\text{Mod-}R, \mathfrak{I}_0)$, i.e., there is no dual to Kaplansky's theorem for cotorsion pairs for \mathfrak{I} .

EXAMPLE 2.2. (i) For any ring R , \mathfrak{I} is cogenerated by any set consisting of injective modules. Consider such a set \mathcal{S} . Then $\text{Cofilt}(\mathcal{S})$ is the class of all modules isomorphic to (arbitrary) direct products of modules from \mathcal{S} . In particular, if $\mathfrak{I}_0 = \{W\}$ where W is an injective cogenerator for $\text{Mod-}R$, then $\mathfrak{I}_0 = \text{sCofilt}(\mathfrak{I}_0)$. So condition (2) is satisfied.

(ii) Now, assume that R is a commutative noetherian ring. For $M \in \text{Mod-}R$, $\text{Ass } M$ will denote the set of all associated prime ideals of M , that is, all $p \in \text{Spec}(R)$ such that R/p embeds into M . Recall that for each module M , $\text{Ass } M = \text{Ass } E(M)$, where $E(M)$ denotes the injective envelope of M . Also, \mathfrak{I}_0 coincides with the class of all modules I of the form $I \cong \bigoplus_{p \in P} E(R/p)^{(\alpha_p)}$ where $P \subseteq \text{Spec}(R)$ and α_p ($p \in P$) are non-zero ordinals; for such I , we have $\text{Ass } I = P$, so P is uniquely determined by I , and so are the ordinals α_p ($p \in P$). In particular, the indecomposable injective modules are the modules isomorphic to $E(R/p)$ for some $p \in \text{Spec}(R)$, and $\text{Ass } E(R/p) = \{p\}$.

Assume moreover that R possesses a maximal ideal m satisfying the conditions $\bigcap_{n < \omega} m^n = 0$ and $m \notin \text{Ass } R$ (this occurs when R is a noetherian domain which is not a field, or R is a local Gorenstein ring of Krull dimension ≥ 1). Suppose that $\mathfrak{I}_0 = \text{Cofilt}(\mathcal{S})$ for a set $\mathcal{S} \subseteq \mathfrak{I}_0$. Then $\mathfrak{I}_0 = \text{Prod } \mathcal{S}$, the class of all modules isomorphic to direct products of copies of modules in \mathcal{S} . In particular, \mathcal{S} must contain copies of all indecomposable injective modules. Since $\bigcap_{n < \omega} m^n = 0$, if \mathcal{D} is any infinite sequence of non-zero elements of \mathcal{S} such that for each $D \in \mathcal{D}$, $m \in \text{Ass } D$, then $I = \prod_{D \in \mathcal{D}} D$ contains a copy of R . Then $E(R)$ is isomorphic to a direct summand in I . As $m \notin \text{Ass } R = \text{Ass } E(R)$, I has an indecomposable direct summand isomorphic to $E(R/p)$ for some prime ideal $p \neq m$. So for each infinite cardinal κ , if $E(R/m)^{(\kappa)} \in \text{Prod } \mathcal{S}$, then $E(R/m)^{(\kappa)}$ is isomorphic to a module in \mathcal{S} , in contradiction with \mathcal{S} being a set. Thus $\text{Cofilt}(\mathcal{S}) \subsetneq \mathfrak{I}_0$ for any set \mathcal{S} of injective modules, i.e., condition (3) fails.

Our next example generalizes part (i) of Example 2.2 to 1-cotilting cotorsion pairs. Hence, it also covers the setting of Dedekind domains and cotorsion pairs $(\mathcal{A}, \mathcal{B})$ such that $\mathcal{A} \supseteq \mathcal{F}_0$ (see Theorem 2.8 below):

EXAMPLE 2.3. Let R be a ring and $\mathfrak{C} = (\mathcal{A}, \mathcal{B})$ be a cotilting cotorsion pair cogenerated by a 1-cotilting module C , i.e., a module C such that $\text{Cogen } C = {}^\perp C$. Then \mathfrak{C} satisfies the condition (2) of Lemma 2.1.

Indeed, using [14, 15.20] and a dual version of [14, 6.13], we see that \mathcal{B} coincides with the class of all modules that are direct summands of the modules M that fit in an exact sequence $0 \rightarrow C^\lambda \rightarrow M \rightarrow I \rightarrow 0$ for an injective module I and a cardinal λ . Let W denote an injective cogenerator for $\text{Mod-}R$. Then $I \oplus J \cong W^\kappa$ for a cardinal κ , and we have the exact sequence $0 \rightarrow C^\lambda \rightarrow M \oplus J \rightarrow W^\kappa \rightarrow 0$. Let $\mathcal{S} = \{C, W\}$ ($\subseteq \mathcal{B}$). Then the module $M \oplus J$ is \mathcal{S} -cofiltered, whence $\mathcal{B} = \text{sCofilt}(\mathcal{S})$.³

Lemma 2.1 is sufficient to settle our problem for flat and torsion-free modules:

COROLLARY 2.4. *Let R be a ring and $n < \omega$. Then there exist epimorphisms $g_n \in \text{Hom}_R(I_n, J_n)$ and $h \in \text{Hom}_R(I, J)$ with I_n and I injective, such that the class \mathcal{F}_n of all modules of flat dimension $\leq n$ (the class of all torsionfree modules) is a cofactorization class of g_n (h).*

PROOF. The first claim follows by Lemma 2.1, since a module M has flat dimension $\leq n$, if and only if $\text{Tor}_{n+1}^R(M, C) = 0$ for each cyclic left R -module C , if and only if $\text{Tor}_1^R(M, \Omega^n(C)) = 0$ for each cyclic left R -module C , where $\Omega^n(C)$ denotes the n th syzygy module of C in a projective resolution of C . It follows that the cotorsion pair $(\mathcal{F}_n, \mathcal{F}_n^\perp)$ is cogenerated by the set of dual modules of modules $\Omega^n(C)$ where C is a cyclic left R -module. For the second claim, one uses the characterization of torsion-free modules as the modules M such that $\text{Tor}_1^R(M, R/Rr) = 0$ for each non-zero divisor $r \in R$. \square

In particular, we have a solution for modules of bounded projective dimension in the case when projectivity and flatness of modules coincide, that is, for right perfect rings:

COROLLARY 2.5. *Let R be a right perfect ring and $n < \omega$. Then there exists an epimorphism $g_n \in \text{Hom}_R(I_n, J_n)$ with I_n injective, such that the class \mathcal{P}_n of all modules of projective dimension $\leq n$ is a cofactorization class of g_n .*

REMARK 2.6. Note that, over each right perfect ring R , there are simpler test epimorphisms than g_0 from Corollary 2.4 available:

1. By [20] (or [18]), the Dual Baer Criterion (DBC) for projectivity holds in $\text{Mod-}R$, that is, one can use the epimorphism $g: R^T \rightarrow \prod_{I \in T} R/I$, where T is the set of all right ideals of R : \mathcal{P}_0 is the cofactorization class of g .

³ ADDED IN PROOF. Positselski has recently extended this result to arbitrary n -cotilting cotorsion pairs $\mathcal{C} = (\mathcal{A}, \mathcal{B})$, $n \geq 0$, by constructing for each such pair \mathcal{C} a set \mathcal{S} consisting of $n + 1$ modules, such that $\mathcal{B} = \text{sCofilt}(\mathcal{S})$. See Theorem 2.19 in his preprint “An explicit construction of complete cotorsion pairs in the relative context”, [19].

2. Let π be the finite direct product of the projections $\pi_S: E(S) \rightarrow E(S)/S$ where S runs over a representative set of all simple modules. Then \mathcal{P}_0 is the cofactorization class of π by [14, 8.8].

For non-right perfect rings, the following consistency result gives a barrier to testing for projectivity using *sets* of epimorphisms:

LEMMA 2.7. *Assume Shelah's Uniformization Principle (SUP). Let R be a non-right perfect ring.*

Then for each set $\mathcal{S} \subset \text{Mod-}R$, there exists a module M with $\text{proj. dim } M = 1$ and $\text{Ext}_R^1(M, S) = 0$ for all $S \in \mathcal{S}$. So the cotorsion pair $(\mathcal{P}_0, \text{Mod-}R)$ is not cogenerated by any set of modules.

Moreover, there is no epimorphism π such that \mathcal{P}_0 is the cofactorization class of π .

PROOF. For the first claim, see e.g. [25, 2.4]. The second claim follows from the first one for $\mathcal{S} = \{\text{Ker}(\pi)\}$. \square

Note that if ZFC is consistent, then so is ZFC + SUP, see [8].

We now pause to look at other cotorsion pairs, but in the particular setting of Dedekind domains. Once again, (SUP) comes as a handy tool here. The following theorem generalizes [9, 1.3]:

THEOREM 2.8. *Let R be a Dedekind domain with the quotient field Q , and $\mathfrak{C} = (\mathcal{A}, \mathcal{B})$ be a cotorsion pair.*

a. *Assume $Q \in \mathcal{A}$ (or, equivalently, $\mathcal{F}_0 \subseteq \mathcal{A}$). Then \mathfrak{C} is a cotilting cotorsion pair, i.e., \mathfrak{C} is cogenerated by a pure-injective module C such that $\text{Cogen } C = {}^\perp C (= \mathcal{A})$. In particular, \mathfrak{C} is generated by a set, and there is a unique subset $P \subseteq \text{mSpec}(R)$ such that*

$$\mathcal{A} = \{M \in \text{Mod-}R \mid \text{for all } p \in P: \text{Hom}_R(R/p, M) = 0\}.$$

b. *Assume (SUP). Moreover, assume that $Q \notin \mathcal{A}$, Q is a countably generated module, and \mathfrak{C} is generated by a set.*

Then there is no epimorphism π such that \mathcal{A} is the cofactorization class of π .

PROOF. (a) This is proved in [14, 16.29] (see also [14, 16.21]).

(b) In [9, 1.3], it is proved (in a different notation) that under the same assumptions, \mathfrak{C} is not cogenerated by a set, i.e., there is no epimorphism $\pi: I \rightarrow J$ with I injective such that \mathcal{A} is the cofactorization class of π (see Lemma 2.1). In order to prove our stronger claim, we will follow the pattern of the proof of [9, 1.3], indicating only the necessary changes needed to strengthen the claim.

Assume there is an epimorphism π in $\text{Mod-}R$ such that \mathcal{A} is the cofactorization class of π . Let $K = \text{Ker}(\pi)$.

Since Q is countably generated, there is a short exact sequence $0 \rightarrow R^{(\omega)} \xrightarrow{f} R^{(\omega)} \rightarrow Q \rightarrow 0$ with $f(1_i) = 1_i - \rho_i 1_{i+1}$ where $\{\rho_i \mid i < \omega\}$ is a set of non-invertible non-zero elements of R . As in [9], we use this presentation and (SUP) to define for a cardinal $\mu = \tau^+$, where τ is an uncountable singular cardinal of cofinality ω , a non-projective module $H = F/G$, where F and G are free modules of rank μ , such that $\text{Ext}_R^1(H, N) = 0$ for each module N of cardinality $< \tau$.

As in [9], we distinguish the following two cases:

1. \mathfrak{C} is generated by a cotorsion module B ;
2. there is no cotorsion module generating \mathfrak{C} (but, by our assumption, there is nevertheless a non-cotorsion module B such that $\mathcal{B} = B^\perp$).

In the case (2), we take a singular cardinal τ of cofinality ω bigger than the cardinalities of R , B and K and $\mu = \tau^+$. Taking the appropriate $\{B\}$ -filtered module $A \in B^\perp = \mathcal{B}$, we infer as in the proof of case (2) in [9, 1.3] that for the module H as above, $\text{Ext}_R^1(H, K) = 0$, so $H \in \mathcal{A}$, but $\text{Ext}_R^1(H, A) \neq 0$, in contradiction with $A \in \mathcal{B}$.

In the case (1), we will first prove the following claim: $\mathcal{B} = B^\perp = (E(T(B)) \oplus F(B))^\perp$ where $T(B)$ is the torsion part of B , $E(T(B))$ is its injective hull, and $F(B)$ is the torsion-free part of B . Notice that $F(B)$ is a cotorsion module since B is such.

Consider the push-out of $T(B) \subseteq B$ and $T(B) \subseteq E(T(B))$ in Figure 1. Since the second row splits, it suffices to prove that $B^\perp = X^\perp$. As R is hereditary, the second column gives $X^\perp \subseteq B^\perp$, and the first row yields $B^\perp \subseteq T(B)^\perp$. Let $P = \text{Ass } T(B)$. Then,

$$\begin{aligned} E(T(B))^\perp &= T(B)^\perp \\ &= \left(\bigoplus_{p \in P} R/p \right)^\perp \\ &= \{M \in \text{Mod-}R \mid \text{for all } p \in P: M.p = M\}, \end{aligned}$$

$$\begin{array}{ccccccc}
& & 0 & & 0 & & \\
& & \downarrow & & \downarrow & & \\
0 & \longrightarrow & T(B) & \xrightarrow{\subseteq} & B & \longrightarrow & F(B) \longrightarrow 0 \\
& & \subseteq \downarrow & & \downarrow & & \parallel \\
0 & \longrightarrow & E(T(B)) & \longrightarrow & X & \longrightarrow & F(B) \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \\
& & E(T(B))/T(B) & \equiv & E(T(B))/T(B) & & \\
& & \downarrow & & \downarrow & & \\
& & 0 & & 0 & &
\end{array}$$

Figure 1

since $E(T(B))$ is $\{R/p \mid p \in P\}$ -filtered. Moreover, there is a subset $P' \subseteq P$ such that $E(T(B))/T(B)$ is $\{R/p \mid p \in P'\}$ -filtered. Hence

$$T(B)^\perp \subseteq (E(T(B))/T(B))^\perp.$$

The second column gives $B^\perp \cap (E(T(B))/T(B))^\perp \subseteq X^\perp$. However, $B^\perp \subseteq T(B)^\perp \subseteq (E(T(B))/T(B))^\perp$, whence $B^\perp \subseteq X^\perp$, and our claim is proved.

Furthermore, $F(B)$ is a torsion-free cotorsion module, which is reduced (otherwise Q is a direct summand in $F(B)$, and since $F(B) \in \mathcal{A}$, also $Q \in \mathcal{A}$, a contradiction). So $F(B)$ is isomorphic to a product of completions of localizations of free R_q -modules where q ranges over some subset $S \subseteq \text{mSpec}(R)$, see [10, 5.3.28]. If $q \in P \cap S$, then $E(R/q) \in \mathcal{A}$, and the q -adic module J_q belongs to \mathcal{A} (because J_q is a direct summand in $F(B)$, and $F(B) \in \mathcal{A}$). Since \mathcal{A} is closed under extensions, also $E(J_q) \in \mathcal{A}$, whence $Q \in \mathcal{A}$, a contradiction. Thus, $P \cap S = \emptyset$. Since the completion of the localizations of a free R_q -module is p -divisible for all $p \neq q$, we infer that $F(B).p = F(B)$ for all $p \in P$.

The rest of the proof is the same as for the case (1) in [9, 1.3], i.e., we take a singular cardinal τ of cofinality ω bigger than the cardinalities of R , B and K and let $\mu = \tau^+$, and take the appropriate $\{F(B)\}$ -filtered module $A \in B^\perp = \mathcal{B}$ so that for the module H as above, $\text{Ext}_R^1(H, K) = 0$, whence $H \in \mathcal{A}$, but $\text{Ext}_R^1(H, A) \neq 0$, in contradiction with $A \in \mathcal{B}$. \square

Next we show that in the extensions of ZFC with the Weak Diamond Principle Φ , for many non-right perfect rings, the class of all projective modules is the cofactorization class of a suitable (single) epimorphism. The point is that one can

combine the methods developed (in ZFC) in infinite dimensional tilting theory [14, §8.2] with the following lemma employing Φ which generalizes [12, Lemma A.7] and [7, Theorem XII.1.10].

First, however, we have to recall the statement of the principle Φ . Given a set X , we say that a system $(X_\alpha \mid \alpha < \kappa)$ is a κ -filtration of X if $X = \bigcup_{\alpha < \kappa} X_\alpha$, $|X_\alpha| < \kappa$, $X_\alpha \subseteq X_{\alpha+1}$ for each $\alpha < \kappa$, and $X_\alpha = \bigcup_{\beta < \alpha} X_\beta$ whenever $\alpha < \kappa$ is a limit ordinal.

Given a regular uncountable cardinal κ and a stationary subset $S \subseteq \kappa$, we denote by $\Phi_\kappa(S)$ the following statement. let X be a set with $|X| \leq \kappa$ and $(X_\alpha \mid \alpha \leq \kappa)$ be a κ -filtration of X ; moreover, for each $\alpha \in S$ let a mapping $P_\alpha: \mathcal{P}(X_\alpha) \rightarrow \{0, 1\}$ be given. Then there exists $\varphi: S \rightarrow \{0, 1\}$ such that, for each $Y \subseteq X$, the set $\{\alpha \in S \mid P_\alpha(Y \cap X_\alpha) = \varphi(\alpha)\}$ is stationary in κ .

The Weak Diamond Principle Φ asserts that “ $\Phi_\kappa(S)$ holds for each regular uncountable cardinal κ and stationary subset $S \subseteq \kappa$ ”. It is consistent with ZFC since it follows, for instance, from the axiom of constructibility $V = L$. It is however much weaker than $V = L$, and also weaker than the famous Jensen’s diamond principle \diamond .

LEMMA 2.9. *Let A, B be R -modules with $|B| \leq \kappa$ where κ is a regular uncountable cardinal. Assume that A is the direct limit of a continuous well-ordered directed system $\mathcal{A} = (A_\alpha, h_{\beta\alpha}: A_\alpha \rightarrow A_\beta \mid \alpha < \beta < \kappa)$, where A_α is $< \kappa$ -generated and $\text{Ext}_R^1(A_\alpha, B) = 0$ for each $\alpha < \kappa$. If the set $S \subseteq \{\alpha < \kappa \mid \text{Ext}_R^1(\text{Coker}(h_{\alpha+1\alpha}), B) \neq 0\}$ is stationary in κ and $\Phi_\kappa(S)$ holds, then one has $\text{Ext}_R^1(A, B) \neq 0$.*

PROOF. First, we extend the system \mathcal{A} into a continuous well-ordered system of short exact sequences $\mathcal{E}_\alpha: 0 \rightarrow K_\alpha \xrightarrow{\subseteq} F_\alpha \xrightarrow{\pi_\alpha} A_\alpha \rightarrow 0$ where F_α is a free module of rank $< \kappa$ and the three components of connecting maps $\varepsilon_{\alpha+1\alpha}: \mathcal{E}_\alpha \rightarrow \mathcal{E}_{\alpha+1}$ are inclusion, split inclusion and $h_{\alpha+1\alpha}$, respectively. Using the assumptions that κ is infinite regular and A_α is $< \kappa$ -generated for $\alpha < \kappa$, this is easy.

As the direct limit, we get a short exact sequence $0 \rightarrow K \xrightarrow{\subseteq} F \rightarrow A \rightarrow 0$ where F is free of rank $\leq \kappa$. Thus we obtain also a κ -filtration $(V_\alpha \mid \alpha < \kappa)$ of a set V of free generators of F where V_α is a set of free generators of F_α for each $\alpha < \kappa$.

Since $\text{Ext}_R^1(A_\alpha, B) = 0$ for $\alpha < \kappa$, we can fix for each homomorphism $f: K_\alpha \rightarrow B$ one of its extensions $f^e \in \text{Hom}_R(F_\alpha, B)$. Furthermore, we fix, for each $\alpha \in S$, a $k_\alpha \in \text{Hom}_R(A_\alpha, B)$ which cannot be factorized through $h_{\alpha+1\alpha}$.

Consider any κ -filtration $(B_\alpha \mid \alpha < \kappa)$ of the set B (the B_α need not be submodules of B , just subsets) and put $X = V \times B$ and $X_\alpha = V_\alpha \times B_\alpha$ for

each $\alpha < \kappa$. Let $\alpha \in S$ be arbitrary. We define the mapping $P_\alpha: \mathcal{P}_\alpha \rightarrow \{0, 1\}$ as follows: if $Z \subseteq X_\alpha$ is not a mapping from V_α to B_α , we put $P_\alpha(Z) = 0$; otherwise, we fix a unique extension $z \in \text{Hom}_R(F_\alpha, B)$ of Z and put $y = (z \upharpoonright K_\alpha)^e$. Then $y - z$ is zero on K_α and thus it defines a unique homomorphism from A_α to B . We put $P_\alpha(Z) = 1$ if and only if this homomorphism can be factorized through $h_{\alpha+1\alpha}$.

Using $\Phi_\kappa(S)$, we get $\varphi: S \rightarrow \{0, 1\}$ for our choice of the mappings P_α . To show that $\text{Ext}_R^1(A, B) \neq 0$, we recursively construct a homomorphism $f: K \rightarrow B$ which cannot be extended to an element of $\text{Hom}_R(F, B)$. We start with $f_0: K_0 \rightarrow B$ the zero map. If $\alpha \leq \kappa$ is a limit ordinal, we put $f_\alpha = \bigcup_{\beta < \alpha} f_\beta$. Let us assume that f_α is already constructed and $\alpha < \kappa$. We define $f_{\alpha+1}: K_{\alpha+1} \rightarrow B$ as follows:

Let us put $f'_\alpha = f_\alpha^e$ if $\alpha \notin S$ or $\varphi(\alpha) = 0$; otherwise, we put $f'_\alpha = f_\alpha^e + k_\alpha \pi_\alpha$. We extend f'_α arbitrarily to a homomorphism $f_\alpha^+: F_{\alpha+1} \rightarrow B$ and define $f_{\alpha+1}$ as $f_\alpha^+ \upharpoonright K_{\alpha+1}$.

Finally, we put $f = f_\kappa: K \rightarrow B$. For the sake of contradiction, we assume that there exists $g \in \text{Hom}_R(F, B)$ such that $g \upharpoonright K = f$. It is easy to see that the set $C = \{\alpha < \kappa \mid g(V_\alpha) \subseteq B_\alpha\}$ is closed and unbounded. Using the property of φ for $Y = g \upharpoonright V$, we obtain a $\delta \in C \cap S$ such that $P_\delta(g \upharpoonright V_\delta) = \varphi(\delta)$. Obviously, $f_\delta^+ - g \upharpoonright F_{\delta+1}$ is zero on $K_{\delta+1}$. Thus, it defines a unique $h \in \text{Hom}_R(A_{\delta+1}, B)$. Then $k = hh_{\delta+1,\delta}$ where $k: A_\delta \rightarrow B$ is such that $k\pi_\delta = f'_\delta - g \upharpoonright F_\delta$.

If $\varphi(\delta) = 0$, then $k\pi_\delta = f_\delta^e - g \upharpoonright F_\delta$ in the contradiction with $P_\delta(g \upharpoonright V_\delta) = 0$ which has meant that k cannot be factorized through $h_{\delta+1\delta}$.

If on the other hand $\varphi(\delta) = 1$, then $k\pi_\delta = k_\delta\pi_\delta + f_\delta^e - g \upharpoonright F_\delta$. Since $P_\delta(g \upharpoonright V_\delta) = 1$, we know that $k - k_\delta$ can be factorized through $h_{\delta+1\delta}$ which immediately implies that k_δ has this property as well, in contradiction with its choice. \square

A typical application of Lemma 2.9 is the following: given a filtration $\mathcal{A} = (A_\alpha \mid \alpha < \kappa)$ of A consisting of $< \kappa$ -generated modules such that $A, A_\alpha \in {}^\perp B$ for each $\alpha < \kappa$ and $|B| \leq \kappa$, there exists a subfiltration of \mathcal{A} with consecutive factors in ${}^\perp B$ provided that Φ holds true.

In the sequel, given a $\mathcal{C} \subseteq \text{Mod-}R$ and a cardinal μ , we denote by $\mathcal{C}^{\leq \mu}$ (or $\mathcal{C}^{< \mu}$, respectively) the subclass of \mathcal{C} consisting of all the modules which are μ -presented ($< \mu$ -presented, respectively).

Also, we will say that a module M has \mathcal{C} -resolution dimension $\leq n$ provided that there exists an exact sequence $\mathcal{E}: 0 \rightarrow C_n \rightarrow C_{n-1} \rightarrow \cdots \rightarrow C_1 \rightarrow C_0 \rightarrow M \rightarrow 0$ with $C_i \in \mathcal{C}$ for all $i \leq n$. The sequence \mathcal{E} is called a \mathcal{C} -resolution of M of length n .

Finally, recall that given a module M and a cotorsion pair $\mathfrak{C} = (\mathcal{A}, \mathcal{B})$ where $\mathcal{A} = \text{Filt}(\mathcal{A}^{\leq \mu})$ for a cardinal μ , there exists a *special \mathcal{B} -preenvelope of M* , i.e. a monomorphism $f: M \rightarrow B$ where $B \in \mathcal{B}$ and $\text{Coker}(f) \in \mathcal{A}$. This follows from [14, Theorem 6.11] and the fact that \mathfrak{C} is generated by a representative set of μ -presented modules from \mathcal{A} .

THEOREM 2.10. *Assume that Φ holds true. Let $\mathfrak{C} = (\mathcal{A}, \mathcal{B})$ be a hereditary cotorsion pair such that $\mathcal{A} \cap \mathcal{B} = \text{Add } K$ for some $K \in \text{Mod-}R$. Assume that $\mu \geq |K| + |R|$ is an infinite cardinal such that $\mathcal{A} = \text{Filt}(\mathcal{A}^{\leq \mu})$.*

Let M be a module of \mathcal{A} -resolution dimension $\leq n$ (e.g., assume $M \in \mathcal{P}_n$). Then $M \in \mathcal{A}$ if and only if $\text{Ext}_R^i(M, K^{(\mu)}) = 0$ for all $0 < i \leq n$.

PROOF. The only-if part follows immediately since $K^{(\mu)} \in \mathcal{B}$ and \mathfrak{C} is hereditary. Let us concentrate on the if part. We shall prove it by induction on n . The result is trivial for $n = 0$, so let $n = 1$.

We thus have a short exact sequence $\mathcal{E}: 0 \rightarrow A_1 \rightarrow A_0 \xrightarrow{\pi} M \rightarrow 0$ with $A_0, A_1 \in \mathcal{A}$. Considering a special \mathcal{B} -preenvelope of A_1 and subsequently forming the obvious push-out, we can w.l.o.g. assume that $A_1 \in \text{Add } K$. By further adding a suitable direct summand from $\text{Add } K$ to A_0 and A_1 , we can moreover assume that $A_1 = K^{(\kappa)}$ for some cardinal κ . We argue by induction on κ that $M \in \mathcal{A}$.

If $\kappa \leq \mu$, then the short exact sequence \mathcal{E} splits by our assumption, whence $M \in \mathcal{A}$. Assume that $\kappa = |A_1| > \mu$. Then A_1 is trivially $\{K\}$ -filtered and A_0 is $\mathcal{A}^{\leq \mu}$ -filtered by one of our assumptions. If κ is regular, we use [24, Theorem 3.4] to obtain a filtration $(\mathcal{E}_\alpha: 0 \rightarrow A_{\alpha,1} \rightarrow A_{\alpha,0} \rightarrow M_\alpha \rightarrow 0 \mid \alpha < \kappa)$ of \mathcal{E} where $A_{\alpha,1}$ is a canonical direct summand of A_1 and $A_{\alpha,0} \in \mathcal{A}$, whilst $|A_{\alpha,0}| < \kappa$, for each $\alpha < \kappa$. Consequently, since $\text{Hom}_R(-, K^{(\mu)})$ turns \mathcal{E} into a short exact sequence by our assumption, we see that $\text{Ext}_R^1(M_\alpha, K^{(\mu)}) = 0$ for each $\alpha < \kappa$. Applying Lemma 2.9, we easily obtain a subfiltration of $(M_\alpha \mid \alpha < \kappa)$ with consecutive factors in ${}^\perp K^{(\mu)}$. Using the property of $M_{\alpha+1}/M_\alpha$ guaranteed by [24, Theorem 3.4], we see that the consecutive factors belong to \mathcal{A} by inductive assumption, whence $M \in \mathcal{A}$ by the Eklof lemma.

If $\kappa > \mu$ is singular, we easily construct, for each regular $\lambda < \kappa$ such that $\mu < \lambda$, a system \mathcal{S}_λ consisting of subobjects of the short exact sequence \mathcal{E} whose first term is a canonical direct summand of A_1 of cardinality $< \lambda$, the second term is in $\mathcal{A}^{< \lambda}$, and such that $\bigcup \mathcal{S}_\lambda = \mathcal{E}$ and \mathcal{S}_λ is closed under unions of chains of cardinality $< \lambda$. The third terms of short exact sequences in \mathcal{S}_λ then belong to ${}^\perp K^{(\mu)}$, and thus to $\mathcal{A} = \text{Filt}(\mathcal{A}^{\leq \mu})$ by the inductive hypothesis. We use the Singular Compactness Theorem, [14, Theorem 7.29], to deduce that $M \in \mathcal{A}$. This settles the case $n = 1$.

Finally, assume that $n > 1$. Then the first syzygy, $\Omega(M)$, in an \mathcal{A} -resolution of M of length n has \mathcal{A} -resolution dimension $\leq n - 1$. By the inductive hypothesis, $\Omega(M) \in \mathcal{A}$ which immediately implies that $M \in \mathcal{A}$, too. \square

REMARK 2.11. Using dimension shifting, the conclusion of Theorem 2.10 can be generalized to the statement “if $0 \leq d < n$, then M has \mathcal{A} -resolution dimension $\leq d$, if and only if $\text{Ext}_R^i(M, K^{(\mu)}) = 0$ for all $d < i \leq n$.” The original statement amounts to $d = 0$.

Also, it follows from Theorem 2.10 that, if M is a module of $\text{Add } K$ -resolution dimension $\leq n$, then $M \in \text{Add } K$, if and only if $\text{Ext}_R^i(M, K^{(\mu)}) = 0$ for all $0 < i \leq n$.

All tilting cotorsion pairs, including, of course, the trivial cotorsion pair $(\mathcal{P}_0, \text{Mod-}R)$, satisfy the assumptions of Theorem 2.10. More generally, all hereditary cotorsion pairs with the right-hand class closed under direct limits (cf. [3, Lemma 5.4]). Other examples include, for instance, the cotorsion pair $\mathfrak{PGF} = (\mathcal{PGF}, \mathcal{PGF}^\perp)$ from [22]. In particular, if R is a ring over which each module has finite \mathcal{PGF} -resolution dimension, then Φ implies the existence of a test module for projectivity in $\text{Mod-}R$: indeed, by Theorem 2.10, \mathfrak{PGF} is cogenerated by a single module. At the same time, the flat cotorsion pair $(\mathcal{F}_0, \mathcal{EC})$ is also cogenerated by a single module, and the flat modules in \mathcal{PGF} are precisely the projective ones.

As the following proposition shows, the Weak Diamond Principle can help us also in another special case which covers the setting of Theorem 2.8 and complements part (b) therein.

PROPOSITION 2.12. *Let $\mathfrak{C} = (\mathcal{A}, \mathcal{B})$ be a cotorsion pair with $\mathcal{B} \subseteq \mathcal{I}_1$. Assuming Φ , \mathfrak{C} is generated by a set if and only if \mathfrak{C} is cogenerated by a set.*

PROOF. The if part follows e.g. by [14, Theorem 11.2] where Φ and our Lemma 2.9 play the role of Ψ and [14, Lemma 11.1]. Let us prove the only-if part.

Denote by κ the least infinite cardinal such that $\kappa \geq |R|$ and $\mathcal{A} = \text{Filt}(\mathcal{A}^{\leq \kappa})$. For each infinite cardinal μ , fix a module $W_\mu \in \mathcal{B}$ such that a $< \mu$ -presented module M belongs to \mathcal{A} if and only if $M \in {}^\perp W_\mu$. The module W_μ can be defined e.g. as $\prod_{M \in \mathcal{S}} W_M$ where \mathcal{S} is the representative set of all $< \mu$ -presented modules not belonging to \mathcal{A} and $W_M \in \mathcal{B}$ is such that $\text{Ext}_R^1(M, W_M) \neq 0$.

We construct a continuous increasing sequence $(\lambda_\alpha \mid \alpha \leq \kappa^+)$ of cardinals: we start by putting $\lambda_0 = \kappa^+$; if α is limit, we put $\lambda_\alpha = \sum_{\beta < \alpha} \lambda_\beta$. Finally, if λ_α is defined and $\alpha < \kappa^+$, we set $\lambda_{\alpha+1} = (\lambda_\alpha + |W_{\lambda_\alpha}|)^\kappa$. Put $\lambda = \lambda_{\kappa^+}$. By the definition, we see that $\lambda^\kappa = \lambda$.

Let W denote the submodule of $\prod_{\alpha < \kappa^+} W_{\lambda_\alpha}$ consisting of elements of bounded support (i.e., of support of cardinality $\leq \kappa$). Then $W \in \mathcal{B}$ since $\mathcal{A} = \text{Filt}(\mathcal{A}^{\leq \kappa})$. Also $|W| = \lambda^\kappa = \lambda$. We claim that $\mathcal{A} = {}^\perp W$. Clearly, $\mathcal{A} \subseteq {}^\perp W$. Let $M \in {}^\perp W$ be arbitrary. We shall prove that $M \in \mathcal{A}$ by induction on $\theta = |M|$.

If $\theta < \lambda$, then $M \in \mathcal{A}$ immediately follows by the definition of modules W_{λ_α} , $\alpha < \kappa^+$. Assume that $N \in {}^\perp W \Rightarrow N \in \mathcal{A}$ holds for all modules N of cardinality $< \theta$ for some $\theta \geq \lambda$.

First, consider the case of a regular cardinal $\theta = |M|$ where $M \in {}^\perp W$. Since $W \in \mathcal{J}_1$ by our assumption, ${}^\perp W$ is closed under submodules, and we can use Lemma 2.9 to infer that there is a ${}^\perp W$ -filtration of M consisting of modules of cardinality $< \theta$. By the inductive assumption, this is actually an \mathcal{A} -filtration, whence $M \in \mathcal{A}$ by the Eklof lemma.

If θ is singular, we use the inductive hypothesis and $W \in \mathcal{J}_1$ to deduce that, for each regular $\mu < \theta$ with $\kappa < \mu$, the system \mathcal{S}_μ of submodules in M of cardinality $< \mu$ consists of modules from $\mathcal{A} = \text{Filt}(\mathcal{A}^{\leq \kappa})$. Consequently, we can use Singular Compactness Theorem [14, Theorem 7.29], to infer that $M \in \mathcal{A}$. \square

Combining the results above, we obtain

COROLLARY 2.13. *Let R be a non-right perfect ring such that each flat module has finite projective dimension. Then the assertion “There exists an epimorphism π such that \mathcal{P}_0 is the cofactorization class of π ” is independent of ZFC.*

In particular, this is the case when

- *R is a commutative noetherian ring of non-zero finite Krull dimension,*
- *R is an n -Iwanaga–Gorenstein ring for $n \geq 1$,*
- *R is an almost perfect ring.*

PROOF. Consistency of the failure of the assertion under SUP follows by Lemma 2.7.

Assuming Φ , we can apply Theorem 2.10 for $K = R$ and $\mathcal{C} = (\mathcal{P}_0, \text{Mod-}R)$ to obtain the epimorphism $\pi: \bigoplus_{0 < i \leq n} E(\Omega^{-i}(R^{(R)})) \rightarrow E(\Omega^{-i}(R^{(R)})) / \Omega^{-i}(R^{(R)})$, where $\Omega^{-i}(R^{(R)})$ denotes the i th cosyzygy of $R^{(R)}$, which tests for projectivity of modules of finite projective dimension. From our assumption that all flat modules have finite projective dimension it follows that \mathcal{P}_0 is the cofactorization class of the epimorphism $\pi \oplus g_0$, where g_0 is the epimorphism from Corollary 2.4.

Finally, flat modules over a commutative noetherian ring of Krull dimension $d < \omega$ have projective dimension $\leq d$ by a classic result of Gruson and Jensen (cf. [27, 4.2.8]). If R is n -Iwanaga–Gorenstein, then $\text{proj. dim } M \leq n$ for each

module of finite flat dimension by [10, 9.1.10]. If R is almost perfect, then all flat modules have projective dimension ≤ 1 by [13, 7.1] (see also [11]). \square

REMARK 2.14. By Remark 2.6(1), the Dual Baer Criterion (DBC) holds for each right perfect ring, that is, $g: R^T \rightarrow \prod_{I \in T} R/I$, where T is the set of all right ideals of R , is a test epimorphism for projectivity.

By Lemma 2.7, it is consistent with ZFC that the DBC fails for each non-right perfect ring. In fact, in contrast with Corollary 2.13(2), DBC fails (in ZFC) for all commutative noetherian rings that are not perfect (i.e., have Krull dimension ≥ 1) by [15]. However, there do exist commutative hereditary rings for which DBC is independent of ZFC, [26].

3. Projective modules and large cardinals

The question of whether it is consistent with ZFC that there exists a test epimorphism for projectivity over any ring remains open. Apart from using the Weak Diamond Principle, another possible approach here is to utilize large cardinals; in particular the strongly compact ones. An uncountable cardinal κ is called a *strongly compact cardinal* provided that each κ -complete filter (on any set I) can be extended to a κ -complete ultrafilter; recall that a filter \mathcal{F} is κ -complete if $\bigcap \mathcal{S} \in \mathcal{F}$ for each $\mathcal{S} \subseteq \mathcal{F}$ of cardinality $< \kappa$. (Each filter is \aleph_0 -complete by definition.)

In [21], it is shown that, assuming the existence of a strongly compact $\kappa > |R|$, there exists a free R -module F such that all projective modules belong to $\text{Prod}(F)$. It is straightforward to check then that a module M of finite projective dimension such that $\text{Ext}_R^i(M, F) = 0$ for all $i > 0$ is necessarily projective. As a result, in the proof above, one can alternatively assume that there exists a proper class of strongly compact cardinals, instead of Φ . We should note, however, that consistency results not relying on large cardinals are usually considered more valuable.

Nonetheless, using strongly compact cardinals, we can get to a situation which resembles the case of right perfect rings where a test module for projectivity always exists. We need just one more preparatory lemma inspired by [1, Proposition 2.31]. In its proof, we use the characterization of λ -pure monomorphisms via solvability of systems of R -linear equations, cf. [12, Exercise XIII.7.2]. Also recall that, given a module B and a filter \mathcal{F} on I , a reduced power B^I/\mathcal{F} is defined as $B^I/Z_{\mathcal{F}}$ where

$$Z_{\mathcal{F}} = \{f \in B^I \mid \{i \in I \mid f(i) = 0\} \in \mathcal{F}\}.$$

If \mathcal{F} is actually an ultrafilter, then B^I/\mathcal{F} is called *an ultrapower* of B .

LEMMA 3.1. *Let λ be an infinite regular cardinal, $f: A \rightarrow B$ a λ -pure monomorphism which is the colimit of a λ -directed system $(f'_i: A'_i \rightarrow B'_i \mid i \in I)$ of morphisms between $<\lambda$ -presented modules. Let \mathcal{F} be an arbitrary λ -complete filter on I containing all the sets $\uparrow i = \{j \in I \mid i \leq j\}$ where i runs through I . Then there exists a λ -pure embedding of $\text{Coker}(f)$ into the reduced power B^I/\mathcal{F} .*

PROOF. As in the proof of [1, Proposition 2.30(ii)], we use push-outs and λ -purity of f to get a λ -directed system $\mathcal{S} = (f_i: A \rightarrow B_i, b_{ji}: B_i \rightarrow B_j \mid i \leq j \in I)$ in the comma-category $A \downarrow \text{Mod-}R$ consisting of split monomorphisms and such that $\lim \mathcal{S} = f$. Let us denote by $b_i: B_i \rightarrow B$, $i \in I$, the colimit maps (so $b_i f_i = f$) and by $g_i: B_i \rightarrow A$ the retractions, i.e. $g_i f_i = 1_A$.

Let $p: B \rightarrow B^I/\mathcal{F}$ be the diagonal embedding of B into the reduced power. We also define the morphism $q: B \rightarrow B^I/\mathcal{F}$ by first picking, for each $b \in B$, an index $i \in I$ and $c \in B_i$ such that $b_i(c) = b$, and then setting $q(b) = [(u_j)_{j \in I}]_{\mathcal{F}}$ where $u_j = b_j f_j g_j b_{ji}(c)$ if $i \leq j$ and $u_j = 0$ otherwise. By the assumption on \mathcal{F} , this is a correctly defined homomorphism.

It is easy to check that $p f = q f$, whence we get a unique morphism $\nu: \text{Coker}(f) \rightarrow B^I/\mathcal{F}$ such that $p - q = \nu \pi$ where $\pi: B \rightarrow \text{Coker}(f)$ is the canonical projection. It remains to verify that ν is a λ -pure monomorphism.

Let Ω be a system of R -linear equations with parameters from $\text{Coker}(f)$ such that $\tau := |\Omega| < \lambda$. Since \mathcal{S} is λ -directed, there exists $i \in I$ such that each parameter in Ω is of the form $\pi b_i(c)$ for some $c \in B_i$. Let us enumerate in $(c_\alpha \mid \alpha < \tau)$ all these elements c . Assume that the system Ω with parameters $(\nu \pi b_i(c_\alpha) \mid \alpha < \tau)$ has a solution in B^I/\mathcal{F} . By λ -completeness of \mathcal{F} and the other assumption we have put on it, there exists $j \in I$, $i \leq j$ such that Ω with parameters $(b_i(c_\alpha) - b_j f_j g_j b_{ji}(c_\alpha) \mid \alpha < \tau)$ has a solution in B . Taking the π -image of this solution (note that $\pi b_j f_j = \pi f = 0$), we obtain a desired solution of Ω in $\text{Coker}(f)$, showing that ν is a λ -pure monomorphism. \square

REMARK 3.2. (1) Given a λ -pure monomorphism f , there is always a λ -directed system $(f'_i \mid i \in I)$ consisting of morphisms between $<\lambda$ -presented modules such that its directed colimit is f . Also note that the filter on I generated by the sets $\uparrow i$ is λ -complete, whence there always exists a filter \mathcal{F} satisfying the assumption of Lemma 3.1.

(2) Since Lemma 3.1 holds also for $\lambda = \aleph_0$, it gives a direct proof of the well-known fact that each definable class of modules (i.e. closed under directed colimits, pure submodules and products) is closed under pure-epimorphic images. Note that the reduced power B^I/\mathcal{F} is the directed colimit of powers of B and canonical epimorphisms between them.

Now we are ready to prove the promised result. Recall that, given a regular infinite cardinal κ , a category \mathcal{K} is called κ -accessible, if it has κ -directed colimits and there is a set \mathcal{S} of $< \kappa$ -presented objects such that every object is a κ -directed colimit of objects from \mathcal{S} . For instance, $\text{Mod-}R$ is an \aleph_0 -accessible category for any ring R since each module is a direct limit of finitely presented modules.

THEOREM 3.3. *Let κ be a strongly compact cardinal and R a ring such that $|R| < \kappa$. Then each κ -pure epimorphism $g: B \rightarrow C$ with B projective splits. In particular, the category \mathcal{P}_0 of all projective R -modules and R -homomorphisms is κ -accessible.*

PROOF. We can assume without loss of generality that B is actually a free module. Using the lemma above for $\kappa = \lambda$, we can find a κ -pure embedding $h: C \rightarrow B^I/\mathcal{U}$ with \mathcal{U} a κ -complete ultrafilter; remember that each κ -complete filter extends to a κ -complete ultrafilter. By [7, Theorem II.3.8], the ultrapower B^I/\mathcal{U} is a free module as well. We repeat the process with $\text{Coker}(h)$, and so on, and eventually obtain an unbounded pure-exact complex consisting of projective (even free) modules which is well known to be contractible, cf. [6, Theorem 2.5]. In particular, C is projective and g splits.

Finally, since each κ -directed colimit of projective modules is a κ -pure-epimorphic image of a direct sum of projective modules, we see that the class \mathcal{P}_0 is closed under taking κ -directed colimits. To see that \mathcal{P}_0 is κ -accessible, observe that each projective module is a directed union of a κ -directed system of $< \kappa$ -generated projective modules. \square

Another consequence of Lemma 3.1 is the following closure property of the class \mathcal{D} of all flat Mittag-Leffler ($= \aleph_1$ -projective) right modules.

PROPOSITION 3.4. *Let R be a left coherent ring such that any (finite or infinite) intersection of finitely generated submodules of ${}_R R^{(\omega)}$ is finitely generated. Then \mathcal{D} is closed under \aleph_1 -pure-epimorphic images.*

PROOF. Recall that \mathcal{D} is always closed under pure submodules and directed colimits of \aleph_1 -continuous directed systems, cf. [16, Lemma 4.1, Proposition 2.2]. Furthermore, by [16, Theorem 4.7], \mathcal{D} is closed under products precisely over rings satisfying our assumption. Let $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ be an \aleph_1 -pure short exact sequence with $B \in \mathcal{D}$. It follows from Lemma 3.1 that C is a pure submodule of a reduced power B^I/\mathcal{F} where \mathcal{F} is an \aleph_1 -complete filter. Consequently, B^I/\mathcal{F} is the directed colimit of \aleph_1 -continuous directed system consisting of direct powers of B and epimorphisms. As such $B^I/\mathcal{F} \in \mathcal{D}$, and hence also $C \in \mathcal{D}$. \square

Below, we show that the additional assumption on R is necessary.

EXAMPLE 3.5. In this example, by “countable,” we mean “of cardinality $< \aleph_1$.” Consider the boolean ring R consisting of all the subsets of ω_1 which are either countable, or have countable complement, with the usual operations of symmetric difference and intersection. The ideal I of all the countable subsets of ω_1 is easily seen to be maximal, and it is not a direct summand in R_R . It follows that the simple module R/I is not (flat) Mittag-Leffler. On the other hand, the inclusion $I \hookrightarrow R_R$ is the \aleph_1 -directed (but not \aleph_1 -continuous!) colimit of split inclusions $eR \hookrightarrow R_R$ where e runs through countable subsets of ω_1 , thus $I \hookrightarrow R_R$ is \aleph_1 -pure and R/I is an \aleph_1 -pure-epimorphic image of a free module which is not (flat) Mittag-Leffler.

Acknowledgment. Research supported by GAČR 17-23112S.

REFERENCES

- [1] J. ADÁMEK – J. ROSICKÝ, *Locally presentable and accessible categories*, London Mathematical Society Lecture Note Series, 189, Cambridge University Press, Cambridge, 1994.
- [2] F. W. ANDERSON – K. R. FULLER, Rings and categories of modules, Second edition, Graduate Texts in Mathematics, 13, Springer-Verlag, New York, 1992.
- [3] L. ANGELERI HÜGEL – J. ŠAROCH – J. TRLIFAJ, *Approximations and Mittag-Leffler conditions—the applications*, Israel J. Math. 226 (2018), no. 2, pp. 757–780.
- [4] R. BAER, *Abelian groups that are direct summands of every containing abelian group*, Bull. Amer. Math. Soc. 46 (1940), pp. 800–806.
- [5] S. BAZZONI – J. ŠTOVÍČEK, *Flat Mittag-Leffler modules over countable rings*, Proc. Amer. Math. Soc. 140 (2012), no. 5, pp. 1527–1533.
- [6] D. J. BENSON – K. R. GOODEARL, *Periodic flat modules, and flat modules for finite groups*, Pacific J. Math. 196 (2000), no. 1, pp. 45–67.
- [7] P. C. EKLOF – A. H. MEKLER, *Almost free modules*, Set-theoretic methods, revised edition, North-Holland Mathematical Library, 65, North-Holland Publishing Co., Amsterdam, 2002.
- [8] P. C. EKLOF – S. SHELAH, *On Whitehead modules*, J. Algebra 142 (1991), no. 2, pp. 492–510.
- [9] P. C. EKLOF – S. SHELAH – J. TRLIFAJ, *On the cogeneration of cotorsion pairs*, J. Algebra 277 (2004), no. 2, pp. 572–578.
- [10] E. E. ENOCHS – O. M. G. JENDA, *Relative homological algebra*, Volume 1, Second revised and extended edition, De Gruyter Expositions in Mathematics, 30, Walter de Gruyter GmbH & Co. KG, Berlin, 2011.

- [11] A. FACCHINI – Z. NAZEMIAN, *Equivalence of some homological conditions for ring epimorphisms*, J. Pure Appl. Algebra 223 (2019), no. 4, pp. 1440–1455.
- [12] L. FUCHS – L. SALCE, *Modules over non-Noetherian domains*, Mathematical Surveys and Monographs, 84. American Mathematical Society, Providence, R.I., 2001.
- [13] L. FUCHS – L. SALCE, *Almost perfect commutative rings*, J. Pure Appl. Algebra 222 (2018), no. 12, pp. 4223–4238.
- [14] R. GÖBEL – J. TRLIFAJ, *Approximations and endomorphism algebras of modules*, Volume 1. Approximations. Second revised and extended edition. De Gruyter Expositions in Mathematics, 41. Walter de Gruyter GmbH & Co. KG, Berlin, 2012.
- [15] R. M. HAMSHER, *Commutative, noetherian rings over which every module has a maximal submodule*, Proc. Amer. Math. Soc. 17 (1966), pp. 1471–1472.
- [16] D. HERBERA – J. TRLIFAJ, *Almost free modules and Mittag-Leffler conditions*, Adv. Math. 229 (2012), no. 6, pp. 3436–3467.
- [17] TH. JECH, *Set theory*, The third millennium edition, revised and expanded, Springer Monographs in Mathematics, Springer-Verlag, Berlin etc., 2003.
- [18] R. D. KETKAR – N. VANAJA, *R-projective modules over a semiperfect ring*, Canad. Math. Bull. 24 (1981), no. 3, pp. 365–367.
- [19] L. POSITSELSKI, *An explicit construction of complete cotorsion pairs in the relative context*, preprint, 2020. [arXiv:2006.01778v4](https://arxiv.org/abs/2006.01778v4) [math.RA]
- [20] F. SANDOMIERSKI, *Relative injectivity and projectivity*, Ph.D. thesis, Penn State University, State College, PA, 1964.
- [21] J. ŠAROCH, *Σ -algebraically compact modules and $\mathcal{L}_{\omega_1\omega}$ -compact cardinals*, MLQ Math. Log. Q. 61 (2015), no. 3, pp. 196–201.
- [22] J. ŠAROCH – J. ŠTOVÍČEK, *Singular compactness and definability for Σ -cotorsion and Gorenstein modules*, Selecta Math. (N.S.) 26 (2020), no. 2, paper no. 23, 40 pp.
- [23] S. SHELAH, *Diamonds*, Proc. Amer. Math. Soc. 138 (2010), no. 6, pp. 2151–2161.
- [24] A. SLÁVIK – J. TRLIFAJ, *Approximations and locally free modules*, Bull. Lond. Math. Soc. 46 (2014), no. 1, pp. 76–90.
- [25] J. TRLIFAJ, *Whitehead test modules*, Trans. Amer. Math. Soc. 348 (1996), no. 4, pp. 1521–1554.
- [26] J. TRLIFAJ, *The dual Baer Criterion for non-perfect rings*, Forum Math. 32 (2020), no. 3, pp. 663–672.
- [27] J. XU, *Flat covers of modules*, Lecture Notes in Mathematics, 1634. Springer-Verlag, Berlin etc., 1996.