Séminaire Delange-Pisot-Poitou. Théorie des nombres

Wladyslaw Narkiewicz

Numbers with good factorization properties

Séminaire Delange-Pisot-Poitou. Théorie des nombres, tome 13, n° 2 (1971-1972), exp. n° 13, p. 1-3

<http://www.numdam.org/item?id=SDPP_1971-1972__13_2_A1_0>

© Séminaire Delange-Pisot-Poitou. Théorie des nombres
(Secrétariat mathématique, Paris), 1971-1972, tous droits réservés.

L'accès aux archives de la collection « Séminaire Delange-Pisot-Poitou. Théorie des nombres » implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/conditions). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

Numdam
Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
http://www.numdam.org/

Séminaire DELANGE-PISOT-POITOU
(Théorie des nombres)
13e année, 1971/72, n° 13, 3 p.

7 février 1972

NUMBERS WITH GOOD FACTORIZATION PROPERTIES

by Władysław NARKIEWICZ

1. - F. ROGELS (1943, [2]) has shown that in $Q(\sqrt{-5})$, the simplest quadratic field with non-trivial class-group, almost no algebraic integer has unique factorization, that is to say, if $F(x)$ denotes the number of non-associated integers α with $|N(\alpha)| \leqslant x$, which have unique factorization, then $F(x)/x$ tends to zero.

He proved also, that the same holds for the number $H(x)$ of natural numbers $n \leqslant x$ with unique factorization in $Q(\sqrt{-5})$.

In fact, analogous results are true for all fields with non-trivial class-groups, as shown in [4],[6]. For $F(x)$, one gets evaluations of the form

$$F(x) \ll x/\log^{\alpha} x, \quad \alpha > 0,$$

whereas similar evaluations for $H(x)$ are obtained only for K/Q normal.

So we get the first question.

QUESTION I. - Is the evaluation $H(x) \ll x/\log^{\alpha} x$ $(\alpha = \alpha(K) > 0)$, true for all fields K with non-trivial class-group ?

2. - In 1960, L. CARLITZ [1] observed, that in an algebraic number field K with the class-number $h(K) \geqslant 3$, one can find integers α which have factorizations of different lengths, i. e. $\alpha = \pi_1 \dots \pi_r = \rho_1 \dots \rho_s$, with π_j, ρ_i irreductible, and $r \neq s$.

If $G(x)$ is the number of non-associated integers α with $|N(\alpha)| \leqslant x$, whose all factorizations are of the same length, then again (see [4]) one has

$$G(x) \ll x/\log^{\beta} x \quad (\beta = \beta(K) > 0),$$

and the analogue for natural numbers holds, provided K/Q is normal [5]. For non-normal K/Q, only $o(x)$ is proved at this moment [6].

So we have the second question.

QUESTION II. - Obtain the evaluation $O(x/\log^{\beta} x)$ $(\beta > 0)$ for natural numbers having all factorizations of the same length, in a given field K with $h(K) \geqslant 3$.

3. - We shall now indicate the main points of the proof of the following theorem.

THEOREM. - If $h = h(K) \geqslant 2$, then

$$F(x) = (C + o(1)) \frac{x(\log\log x)^M}{(\log x)^{1-(1/h)}},$$

where M is the maximal number of non-principal prime ideals, which can occur in a factorization of a number counted by $F(x)$ with the exponent one.

Let $X_1, \dots, X_{h-1}$ be the non-principal ideal classes in K. If I is any ideal, without principal prime ideal factors, write it in the form

$$I = \prod_i (\mathfrak{p}_{i1}^{\alpha_{i1}} \dots \mathfrak{p}_{ik_i}^{\alpha_{ik_i}}).\mathfrak{Q}, \quad 1 \leq i \leq h-1,$$

with $\mathfrak{p}_{ij} \in X_i$, $\alpha_{ij} > 0$.

We say, that the system

$$\tau = \tau(I) = \langle \{\alpha_{11}, \dots, \alpha_{1k_1}\}, \dots, \{\alpha_{h-1,1}, \dots, \alpha_{h-1,k_{h-1}}\} \rangle$$

is the type of I. If I has no prime divisors from a class X_i say, then we write $\emptyset$ in the place of $\{\alpha_{i1}, \dots\}$.

For a given type τ, let $d(\tau) = \pi\{\alpha_{ij} = 1\}$ be its depth.

The proof of the theorem is based on the following result.

PROPOSITION. - Let $\mathfrak{A}$ be any set of principal ideals subject to the following conditions :

(i) $I \in \mathfrak{A}$, $\tau(I) = \tau(J) \Longrightarrow J \in \mathfrak{A}$;

(ii) If all prime ideal factors of I are principal, then $I \in \mathfrak{A}$;

(iii) $\exists B$, $I \in \mathfrak{A} \Longrightarrow d(\tau(I)) \in B$, whenever $\tau(I)$ is defined.

Then

$$\pi\{I = N(I) \leq x ; \ I \in \mathfrak{A}\} = (\underset{\sim}{C} + o(1)) \frac{x(\log\log x)^M}{(\log x)^{1-(1/h)}}$$

with $\underset{\sim}{C} = \underset{\sim}{C}(\mathfrak{A}) > 0$ and $M = \max\{d(\tau(I)) ; \ I \in \mathfrak{A}\} \leq B$.

This implies immediately the theorem, as if α has a unique factorization, then α has at most $2h-1$ different prime ideal factors from a given class. Indeed, if it has $\geq 2h$, say $\mathfrak{p}_1, \dots, \mathfrak{p}_{2h}, \dots$ then

$$(\mathfrak{p}_1 \dots \mathfrak{p}_h)(\mathfrak{p}_{1+h} \dots \mathfrak{p}_{2h}) = (\mathfrak{p}_{1+h}\, \mathfrak{p}_2 \dots)(\mathfrak{p}_1\, \mathfrak{p}_{2+h} \dots).$$

The proof of the proposition is based on the tauberian theorem of DELANGE.

One starts with the following lemma.

LEMMA. - If $X \in \mathfrak{X}(K)$, and $F_1(t), \dots, F_n(t)$ are real and

$$0 < F_i(t) \ll t^{-2}, \quad d \geq 1,$$

for $\mathrm{Re}\, s > 1$ write

$$S(s) = \sum_{p_1,\dots,p_\alpha} 1/(N_{p_1}^s \dots N_{p_\alpha}^s) \sum_{q_1,\dots,q_n} F_1(Nq_1^s) \dots F_n(Nq_n^s) ,$$

with $p_1 , \dots , p_\alpha \in X$, and distinct ; $q_1 , \dots , q_n \in X$, distinct, and $q_i \neq p_j$. then $S(s) = P(\log(1/s - 1))$, $P \in \Omega[X]$, Ω ring of funtions regular in $\operatorname{Re} s \geqslant 1$, $\deg P = d$, and the leading coefficient is positive at $s = 1$.

Proof. - Induction in d . This allows to show, that il τ is given, and

$$S_\tau = \{I : I = I_1 I_2 , \tau(I_1) = \tau , I_2 \text{ has all prime factors principal}\} ,$$

then

$$\sum_{I \in S_\tau} N(I)^{-s} = (b(\log \frac{1}{s-1}))/(s-1)^{1/h} ,$$

$b \in \Omega[X]$, $\deg b = d(\tau)$, loading coefficient of b positive at 1 , and in fact the same result holds if we sum up this equality over any set of τ with $d(\tau)$ fixed.

There are also another applications of our proposition. Using it in the case $K = Q$, one regains a theorem of L. MIRSKY [3] :

$$\pi\{n \leqslant x : d(n) = k\} = (C + o(1)) \frac{x^{1/(p-1)} (\log \log x)^m}{\log x} ,$$

whose $p = \min\{p : p|k\}$, $p^m \| k$.

As well as analogues of this for all multiplicative functions with $f(p^t) = a_k$.

BIBLIOGRAPHY

[1] CARLITZ (L.). - A characterization of algebraic number fields with class number two, Proc. Amer. math. Soc., t. 11, 1960, p. 391-392.

[2] FOGELS (E.). - Zur Arithmetik quadratischer Zahlenkörper, Univ. Riga, Wiss. Abh., t. 1, 1943, p. 23-47.

[3] MIRSKY (L.). - On the distribution of integers having a prescribed number of divisors, Simon Stevin, Gent, t. 26, 1949, p. 168-175.

[4] NARKIEWICZ (W.). - On algebraic number fields with non-unique factorization, Coll. Math., Warszawa, t. 12, 1964, p. 59-68.

[5] NARKIEWICZ (W.). - On algebraic number fields with non-unique factorization, Coll. Math., Warszawa, t. 15, 1966, p. 49-58.

[6] NARKIEWICZ (W.). - Acta Arith, Warszawa, t. 20, 1972 (to appear).

(Texte reçu le 28 février 1972)

Wladyslaw NARKIEWICZ
Prof. Echange Univ. Bordeaux-I
Mathématiques
351 cours de la Libération
33405 TALENCE CEDEX